

12 Plaice in eastern English Channel

ple.27.7d – *Pleuronectes platessa* in Division 7.d

This stock is in category 1. This year, the assessment of plaice in Division 7.d was made following methodological information described in the Stock Annex revised during ICES WKBNSCS (2025) and WGNSSK 2025. During WGNSSK 2025, a new set of reference points were calculated for this stock. This calculation was carried out following the procedures set in ICES (2021a).

12.1 General

12.1.1 Stock definition

A summary of available information can be found in the Stock Annex.

12.1.2 Ecosystem aspects

No new information on ecosystem aspects was presented at the working group in 2025. All available information on ecological aspects can be found in the Stock Annex.

12.1.3 Fisheries

Plaice is mainly caught in two offshore fisheries, i.e. the beam trawl fishery for sole and the mixed demersal fishery using otter trawls. There is also a directed fishery during parts of the year by inshore trawlers and netters on the English and French coasts. All available information on the fisheries can be found in the Stock Annex.

12.1.4 ICES Advice for previous years

2023 advice: ICES advises that when the MSY approach is applied, catches in 2024 should be no more than 2367 tonnes.

The use of a combined TAC for plaice in divisions 7.d and 7.e prevents effective control of single-stock exploitation rates and could lead to the overexploitation of either stock. ICES advises that management should be implemented at the stock area level.

ICES notes the existence of a precautionary management plan, developed and adopted by one of the relevant management authorities for this stock.

2024 advice: ICES advises that when the MSY approach is applied, catches in 2025 should be no more than 2600 tonnes.

The use of a combined TAC for plaice in divisions 7.d and 7.e prevents effective control of stock exploitation rates and could lead to the overexploitation of either stock. ICES advises that management should be implemented at the stock area level.

ICES notes the existence of a precautionary management plan, developed and adopted by one of the relevant management authorities for this stock.

12.1.5 Management

There are no explicit management objectives for this stock.

The TACs have been set for the combined ICES divisions 7.d and 7.e until 2023. From 2024, separate TACs have been defined for plaice in 7.d and 7.e in line with area-based advice provided by ICES, which takes into account migration between stocks (EU, 2024a).

The minimum landing size for plaice is 27 cm, which is not in accordance with the minimum mesh size of 80 mm, permitted for catching plaice by beam and otter trawling. Fixed nets are required to use 90 mm mesh as an absolute minimum.

Demersal fisheries in the area are mixed fisheries, with many stocks exploited together in various combinations in the various fisheries. In these cases, management advice must consider both the state of individual stocks and their simultaneous exploitation in demersal fisheries. Stocks in the poorest condition, particularly those which suffer from reduced reproductive capacity, become the overriding concern for the management of mixed fisheries, where these stocks are exploited either as a targeted species or as bycatch.

12.2 Data available

12.2.1 Catch

Official landings data shown in Figure 12.2.1.1, Figure 12.2.1.2 as well as in Table 12.2.1.1 and Table 12.2.1.2 together with the total landings estimated by the Working Group. The 2024 official landings of 753 tonnes are the lowest since the beginning of the time series in 1976. France and Belgium are the highest contributors to the total 7.d landings in 2024, contributing for 26% and 54% respectively and UK contributing for 17%. The Belgian TBB and the French OTB recorded the highest landings.

Routine discard monitoring began following the introduction of the EU data collection regulations. Based on the sampling intensity (WKPLE; ICES, 2015), a discards time-series starting in 2006 has been included in the assessment. In 2024, discards of plaice in 7.d as reported to ICES decreased from 2956 tonnes to 2265 tonnes. Following the WKFLAT (ICES, 2010) and WKPLE (ICES, 2015) conclusions, 65% of the first quarter catches were removed. These 65% were estimated during WKFLAT (ICES, 2010), based on published tagging results and some previous studies (e.g. Burt *et al.*, 2006; Hunter *et al.*, 2004; Kell *et al.*, 2004) showing that 50% of the fish caught during the first quarter are fish coming from area 4 to spawn. The same study also shown that 15% of the fish caught during the first quarter were fishes from Division 7.e. Following the WKPLE (ICES, 2015) conclusions, only mature individuals are removed, both from landings and discards. Table 12.2.1.1 shows the Quarter 1 landings and discards and the corresponding removals. Removing this part of the catches allows for assessing the stock resident biomass. All the following figures will take into account this Quarter 1 removal. Of the plaice caught in Division 7.d, 12.9% is assumed to originate in the North Sea and 3.9% from the western English Channel, which are added to the predicted values for the Division 7.d plaice stock and applies to total catch, projected landings, and projected discards.

12.2.2 InterCatch

UK, France, the Netherlands and Belgium have been providing landings data under the ICES InterCatch format since 2011, and InterCatch was used to produce the input data. Age distributions were provided by France, Belgium and England, accounting for 85% of the landings

(Figure 12.2.2.1). Belgium has not always been able to provide landings data per quarter: for 2004, 2005, 2006, 2011, catch data were provided per semester or year. Since 2013, they were provided per year for the TBB fleet with at least quarter 1 landings data on a separate excel spreadsheet. For 2024, Belgium landings data were transmitted per quarter. Allocations to calculate age structures for the remaining landings were done per quarter, using the groups below:

Unsampled fleet*	Sampled fleet**
All nets (GNS, GTR)	All nets
All OTB, OTT, TBB and Seines	All OTB, OTT, TBB and Seines
Others (MIS, OTM, DRB, FPO and LLS)	All métiers

* Unsampled fleet are those fleets for which no age structure is known.

** Sampled fleet are those fleets for which the age structure is known.

Discards data have also been provided under the ICES InterCatch format by France, Belgium, and the UK since WKPLE (ICES, 2015). In 2024, 69 % of landings had associated discards data imported to InterCatch. The discard volumes of the remaining strata have been raised annually using the grouping below (all quarters were pooled). This grouping was updated during 2025 benchmark WKBNSCS (2025).

Unsampled fleet*	Sampled fleet**
TBB	TBB
OTB, OTT	OTB, OTT
GNS, GTR	GNS, GTR
Others (MIS, OTM, DRB, FPO and LLS, SDN, SSC)	All métiers

* Unsampled fleet are those fleets for which no discards data have been provided.

** Sampled fleet are those fleets for which the discards volumes are known.

As a result, the raised discards account for 31% of the total discards. Allocations to calculate age structures of discards were done per quarter, using the groups below:

Unsampled fleet*	Sampled fleet**
All nets	All nets
All OTB, OTT, TBB and Seines	All OTB, OTT and TBB
Others (MIS, OTM, DRB, FPO and LLS)	All métiers

* Unsampled fleet are those fleets for which no age structure is known.

** Sampled fleet are those fleets for which the age structure is known.

Age distributions were provided by France, Belgium and England, accounting for 66% of the total discards (imported + raised).

Due to a lack of samples, a different approach was used to calculate age structures for the remaining discards for the following quarters:

- Discards Q2_OTT-OTB-TBB-Seines raised by samples from Q1, Q2, and Q3

- Discards Q3_OTT-OTB-TBB-Seines raised by samples from Q2, Q3 and Q4

12.2.3 Age compositions

Age compositions of the landings and of the discards are presented in Table 12.2.3.1 and Figure 12.2.3.1, and Table 12.2.3.2 and Figure 12.2.3.2 respectively.

Figures 12.2.3.3 and 12.2.3.4 present the discards at age ratios (i.e., discards numbers/landings numbers) per age and through time over the sampled period 2014–2024. Since 2014, the ratio is increasing for ages 4, 5, 6 and 7+. The ratio for ages 1 to 3 remains stable between 2019 and 2024.

12.2.4 Weight-at-age

Weights-at-age in the landings, discards and stock are presented in tables 12.2.4.1, 12.2.4.2 and 12.2.4.3, respectively and in Figure 12.2.4.1. Stock weights are assumed to be the Q2 landings weights. These weights-at-age do not show specific trends, apart from a general decrease in landing weights in 2013–2018 for ages 5, 6 and 7.

12.2.5 Maturity and natural mortality

The maturity ogive used in the assessment is given in the table below. The maturity ogive was estimated using quantitative histological analysis for plaice in 7d (Sauger *et al.*, 2023). More details are available in the Stock Annex.

Age	1	2	3	4	5	6	7
Proportion of mature	0.2609	0.8929	0.9298	1	1	1	1

A scaled Lorenzen mortality (Lorenzen, 1996) have been estimated during 2025 WKBNSCS benchmark (ICES, 2025), as detailed in the Stock Annex.

Age	1	2	3	4	5	6	7
Natural mortality	0.3618	0.3185	0.2999	0.2816	0.2653	0.2504	0.2389

12.2.6 Surveys

The survey series used in the assessment are the French Ground Fish Survey (FR-GFS) [G3425], 1990–present and the UK beam trawl survey (UK-BTS) [B2453], 1989–present (Figure 12.2.6.1 and Tables 12.2.6.1 and 12.2.6.2). They are fully described in the Stock Annex. Both time-series were re-calculated in 2016 and the impact of those changes were assessed during 2016 WGNSSK meeting (ICES, 2016). A new delta-GAM index was presented during 2022 WGNSSK meeting.

In 2022, five stations within the 6 nm in UK waters were not sampled during the FR-GFS survey. This area is an important habitat for plaice. The effect of this issue has been investigated during 2022 WGNSSK (ICES, 2022) meeting by testing the impact of removing the missing stations in the calculation of the index and although one year of missing five sampling stations did not have a substantial impact on the current assessment outputs, it may become a problem if the issue persists in future. In addition, there has been no UK-BTS survey in the eastern English Channel in 2022.

The consistencies between ages are good for the UK-BTS survey (Figure 12.2.6.2). However, for the FR GFS survey, the internal consistency between ages is slightly less robust compared with UK-BTS survey.

12.3 Assessment

The model used is SAM model (Nielsen and Berg, 2014; Berg and Nielsen, 2016; for more details, please refer to the Stock Annex).

The SAM model inputs and configuration used in the WGNSSK 2025 assessment are shown in the table below. This configuration is based on the model approved at WKBNSCS (ICES, 2025) benchmark and revised during the WGNSSK 2025 meeting (more details about model revision are available in the section 2.6 Quality of the assessment).

Configuration setting	
Model	SAM
Assessment age range	1 – 7+
Is maximum considered a plus group	1 0 0
Coupling of the fishing mortality states processes for each age	0 1 2 3 4 5 6
Correlation of fishing mortality across ages	2 AR(1)
Coupling of the survey catchability parameters	-1 -1 -1 -1 -1 -1 -1 0 1 2 3 4 4 -1 5 6 6 7 7 8 -1
Coupling of process variance parameters for log(F)-process	0 1 1 1 1 1 1
Coupling of process variance parameters for N	0 1 1 1 1 1 1
Coupling of the variance parameters for observations	0 1 1 1 1 1 1 2 2 2 2 2 -1 3 3 3 3 3 -1
Covariance structure for each fleet	"ID" "AR" "AR"
Coupling of correlation parameters	[2,] 0 1 1 2 2 -1 [3,] 3 4 4 5 5 -1
Stock recruitment code	Random walk (0)
Fbar Range	3 – 6
Landings data:	1980–2024
Discards data	1980–2024
Fleets	UKBTS Age range : 1-6, Year : 1989-owards

Configuration setting	
	FRGFS
	Age range : 1-6,
	Year : 1990-owards

12.3.1 Results

The catch time series estimated by the model is shown in Figure 12.3.1.1. Observed catches were well captured by the model and lie within the confidence bounds. Figure 12.3.1.2 shows that the model provides a good fit to the catch-at-age data. However, the fit to the UK-BTS and FR-GFS survey indices reveals some deviations for younger age groups particularly for age 1 (Figure 12.3.1.2).

Figure 12.3.1.3 presents the Standardized one-observation-ahead residuals for catch data as well as the UK-BTS and FR-GFS surveys. Overall, the residuals do not show any clear pattern, with the exception of age 2 in the UK-BTS survey, where the model consistently overestimates in recent years. This pattern could be explained by opposite signals between surveys indices UK-BTS and FR-GFS for age 2 (Figure 12.2.6.1). The process residuals do not indicate any obvious pattern (Figure 12.3.1.4). The leave-one-out runs are mostly within the pointwise 95% confidence intervals (Figure 12.3.1.5). The analysis revealed conflicting trends in F and SSB from 2005. The removal of FR-GFS index resulted in an increase of SSB and a decrease of F.

The final outputs are given in Table 12.3.1.1 (fishing mortalities) and Table 12.3.1.2 (stock numbers). A summary of the assessment results is given in Table 12.3.1.3 and trends in fishing mortality, recruitment, spawning-stock biomass and total catches are shown in Figure 12.3.1.6. Retrospective patterns for the final run are shown in Figure 12.3.1.7. Results from the retrospective analysis suggest a robust model configuration, as Mohn’s rho values for SSB and F and recruitment remain within limits as defined at WKFORBIAS (ICES, 2020) ($\rho_{SSB} = 0.1168$, $\rho_{F_{bar(3-6)}} = -0.1298$, $\rho_{recruitment(age1)} = 0.1809$).

The 1986 year class dominated the history of this stock until the late 2000s (Figure 12.3.1.8 and 12.3.1.3). A second peak occurred with the 1997-year class, although estimated to be at 75% of the 1986-year class. Between 2010 and 2019, a series of high recruitments occurred, reaching a maximum in 2014, which caused the biomass to increase until 2014 then stabilize and decrease from 2020 (Figure 12.3.1.6). The recruitment seems decreasing since 2019.

12.4 Biological reference points

Reference points were estimated during WKBNSCS benchmark (2025) and then updated and presented during WGNSSK (2025) following the benchmark guidelines due to a minor revision of the model (For more details, see Section 2.6 Quality of the assessment and the Annex 8 Working Documents 3 and 4).

Plaice in Division 7.d. Reference points, values, and their technical basis.

Framework	Reference point	Value	Technical basis	Source
MSY approach	MSY $B_{trigger}$	34942	B_{pa} in tonnes.	This report

Framework	Reference point	Value	Technical basis	Source
	F_{MSY}	0.203	EqSim analysis based on recruitment period 1980–2023.	This report
Precautionary approach	B_{lim}	25105	Break-point of hockey stick stock–recruit relationship, based on recruitment period 1980–2023; in tonnes.	This report
	B_{pa}	34942	$B_{lim} \times \exp(1.645 \times \sigma) \approx 1.4 \times B_{lim}$, $\sigma = 0.20$; in tonnes.	This report
	F_{pa}	0.246	The F that provides a 95% probability for SSB to be above B_{lim} ($F_{P,0.05}$ with advice rule).	This report
EU Management plan*	MAP MSY $B_{trigger}$	34942	MSY $B_{trigger}$; in tonnes.	This report
	MAP B_{lim}	25105	B_{lim} ; in tonnes.	This report
	MAP F_{MSY}	0.203	F_{MSY} .	This report
	MAP range F_{lower}	0.147-0.203	Consistent with ranges resulting in no more than 5% reduction in long-term yield compared with MSY.	This report
	MAP range F_{upper}	0.203-0.246	Consistent with ranges resulting in no more than 5% reduction in long-term yield compared with MSY.	This report

* EU multiannual plan (MAP) for the Western Waters (EU, 2019).

12.5 Short-term forecasts

Weight-at-age in the stock and in the catch were taken to be the average estimated weights over the last 3 years. The exploitation pattern, as well as the discards/landings numbers ratio, were taken to be the mean value of the last three years. The short-term forecast was conducted using the stockassessment R package.

The change in advice (-56%) is due to a downward revision in SSB following the benchmark updates (model, input data and reference points).

12.5.1 Recruitment estimates

The recruitment of the intermediate year and onwards is sampled with replacement over the period 1980 – 2024 (Figure 12.5.1.1). The analysis of the ratio between recruitment and SSB didn't show a regime shift to justify the use of a different period to calculate the geometric mean.

12.5.2 Calculation of the 7.d resident stock

This year, F for the intermediate year is set as F 2024 rescaled to the exploitation pattern for 2022–2024. Plaice in 7.d are under landing obligation since the 1st January 2019. To assess if the TAC in 2025 will be fully taken, we compared ICES catches of resident plaice in 7.d in 2025 to the 2025 TAC corresponding to resident plaice in 7.d (3113 t; Figure 12.5.2.1). Then the Q1 removal ratio was applied over the period 2003–2024 to account for migration of mature individuals of plaice from the 7.e and 4.c during Q1 (Figure 12.5.2.2). ICES catches were compared to 2025 TAC corresponding to resident plaice in 7.d by taking into account survivability exemption applied to OTB, OTT, OTM, GTR, GTN, TBB, SSC and SDN landings (EU, 2024b). ICES catches under landings obligation are significantly lower than the TAC (Figure 12.5.2.1), leading to opt for the status quo assumption for fishing mortality (F_{sq}).

12.5.3 Management options tested

12.5.3.1 Calculation of STF

Potential TACs for 2026 were calculated using the MSY approach. In 2026, since $SSB < MSY B_{trigger}$ a reduced F value was used for the forecast instead of F_{MSY} corresponding to $F_{MSY} \times SSB(2026)/MSY B_{trigger}$. Alternative options (e.g., $F_{MSY lower} \times SSB(2026)/MSY B_{trigger}$) were also tested. Results are presented in Tables 12.5.3.1.1 and 12.5.3.1.2 for the resident stock.

Following the MSY approach catches from the Division 7.d plaice stock in 2024 should be no more than 1151 tonnes. Assuming the same proportion of the Division 7.e and Subarea 4 plaice stocks is taken in Division 7.d as during 2003–2024, this will correspond to catches of plaice in Division 7.d in 2026 of no more than 1384 tonnes.

12.6 Quality of the assessment

The stock was benchmarked in 2025. This resulted in a new assessment model and updated catches, maturity and natural mortality (ICES, 2025). The SAM model, initially validated during the WKBNSCS benchmark was revised during WGNSSK the same year due to the emergence of a strong retrospective pattern following the inclusion of 2024 data (Mohn's rho $SSB = 0.2351$). This retrospective pattern mainly results from inconsistencies between survey and catch data for younger ages 1 and 2. The revision was discussed during the meeting and involved coupling all ages of the variance parameters for observations of the two fleets (UK-BTS and FR-GFS), more details about the model revision is available in the Annex 8 Working document 3. Figure 12.6.1 representing the standard deviation of the log observations as estimated in SAM, indicates that by coupling all ages the model gives less weight to survey data which reduced the conflict between catch and survey data.

The table below compares the configuration of the original model validated during the WKBNSCS benchmark with the revised version adopted at WGNSSK 2025. The changes introduced in the revision are highlighted in red.

Configuration setting	SAM Model (WKBNSCS 2025)			SAM Model (WGNSSK 2025)		
Data	Assessment year 2024			Assessment year 2024		
Assessment age range	1 – 7+			1 – 7+		
Is maximum considered a plus group	1 0 0			1 0 0		
Coupling of the fishing mortality states processes for each age	0	1	2 3 4 5 6	0	1	2 3 4 5 6
Correlation of fishing mortality across ages	2 AR(1)			2 AR(1)		
Coupling of the survey catchability parameters	-1	-1	-1 -1 -1 -1	-1	-1	-1 -1 -1 -1
	0	1	2 3 4 4 -1	0	1	2 3 4 4 -1
	5	6	6 7 7 8 -1	5	6	6 7 7 8 -1
Coupling of process variance parameters for log(F)-process	0	1	1 1 1 1 1	0	1	1 1 1 1 1
Coupling of process variance parameters for N	0	1	1 1 1 1 1	0	1	1 1 1 1 1
Coupling of the variance parameters for observations	0	1	1 1 1 1 1	0	1	1 1 1 1 1
	2	3	3 3 3 3 -1	2	2	2 2 2 2 -1
	4	5	5 5 5 5 -1	3	3	3 3 3 3 -1
Covariance structure for each fleet	"ID" "AR" "AR"			"ID" "AR" "AR"		
Coupling of correlation parameters	[2,] 0 1 1 2 2 -1			[2,] 0 1 1 2 2 -1		
	[3,] 3 4 4 5 5 -1			[3,] 3 4 4 5 5 -1		
Stock recruitment code	Random walk (0)			Random walk (0)		
Fbar Range	3 – 6			3 – 6		
Mohn's Rho	R(age1)	SSB	Fbar(3-6)	R(age1)	SSB	Fbar(3-6)
	-0.2835	0.2351	0.2255	0.1809	0.1167	-0.1298

The revision of the model configuration required an update of the reference points (Annex 8 Working Documents 3 and 4). Therefore, the revised model and updated reference points resulted in a different stock perception.

A fishery on the spawners takes place during the first quarter of the year, yielding an age distribution different from the rest of the year. It is unknown whether there is major interannual variability of the immigration from the North Sea to these spawning grounds, which could distort any catch-based analysis. Any migration events taking place in the first quarter cannot be represented in the surveys in the second semester.

Landings-at-age information are highly dependent on the accuracy of the spatial declaration of the fishing activity as an important component of the fisheries operates on the borderline to ICES Subdivision 4.c.

There is uncertainty about catches of the Division 7.d plaice stock because of migrations between Division 7.d and the North Sea and the western English Channel during the spawning period. The current assessment results depend on the proportion of first quarter removals estimated from a historical tagging survey (ICES, 2010).

12.7 Status of the stock

ICES assesses that fishing pressure on the stock is above F_{MSY} , and spawning-stock size is below MSY $B_{trigger}$, B_{pa} and B_{lim} (Figure 12.3.1.3).

		Fishing pressure			Stock size		
		2022	2023	2024	2023	2024	2025
Maximum sustainable yield	F_{MSY}	✖	✖	✖ Above	$MSY B_{trigger}$	✖	✖ Below trigger
Precautionary approach	F_{pa}	✖	✖	✖ Undefined	$B_{pa} B_{lim}$	⚠	✖ Reduced reproductive capacity
Management plan	F_{MGT}	✖	✖	✖ Above	B_{MGT}	✖	✖ Increased risk

12.8 Management considerations

The stock identity of plaice in the English Channel is unclear and may raise some issues.

The plaice stock in 7.d is mostly harvested in a mixed fishery with sole in 7.d.

Due to the minimum mesh size (80 mm) in the mixed beam and otter trawl fisheries, a large number of undersized plaice are discarded. The 80 mm mesh size is not matched to the minimum landing size of plaice (27 cm). Measures taken specifically to control sole fisheries will impact the plaice fisheries.

12.9 Issue for future benchmarks

12.9.1 Data

Migration data are required to update the Q1 migration proportion. Migration between plaice in Subarea 4 and Subdivision 20 and plaice in Division 7d is known to occur. However, current estimates are based on historical studies and may not reflect current dynamics. New studies may be needed to better quantify recent plaice migration. Moreover, migration is currently estimated based on the dynamics of the receiving stock in Division 7.d, rather than from the donor stocks in Subarea 4 and 7e, which may lead to a mismatch between current migration patterns and those assumed in the assessment.

12.9.2 Assessment

Residuals pattern in the UK-BTS index for age 2 in the recent years could be reduced by testing alternative model settings with updated data.

12.9.3 Short-term forecast

If FR-YFS indices are available, the use of RCT3 to estimate recruitment could be investigated. New information for age 0 could be introduced from YFS.

12.10 References

Burt, G., D. Goldsmith, and M. Armstrong. 2006. A summary of demersal fish tagging data maintained and published by Cefas. SCIENCE SERIES TECHNICAL REPORT-CENTRE FOR ENVIRONMENT FISHERIES AND AQUACULTURE SCIENCE 135.

EU. 2019. REGULATION (EU) 2019/472 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL establishing a multiannual plan for stocks fished in the Western Waters and adjacent waters, and for fisheries exploiting those stocks, amending Regulations (EU) 2016/1139 and (EU) 2018/973, and repealing Council Regulations (EC) No 811/2004, (EC) No 2166/2005, (EC) No 388/2006, (EC) No 509/2007 and

- (EC) No 1300/2008. 17pp. <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32019R0472>
- EU. 2024a. Council Regulation (EU) 2024/257 of 10 January 2024 fixing for 2024, 2025 and 2026 the fishing opportunities for certain fish stocks, applicable in Union waters and, for Union fishing vessels, in certain non-Union waters, and amending Regulation (EU) 2023/194. https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=OJ:L_202400257
- EU. 2024b. Commission Delegated Regulation (EU) 2023/2623 of 22 August 2023 Suppl. Regulation (EU) 2019/472 of the European Parliament and of the Council by specifying details of the landing obligation for certain fisheries in Western Waters for the period 2024–2027.
- Hunter, E., Metcalfe, J.D., O'Brien, C.M., Arnold, G.P., Reynolds, J.D., 2004. Vertical activity patterns of free-swimming adult plaice in the southern North Sea. *Marine Ecology Progress Series* 279, 261–273. <https://doi.org/10.3354/meps279261>
- ICES. 2010. Report of the Benchmark Workshop on Flatfish (WKFLAT), 25 February–4 March 2010, Copenhagen, Denmark. ICES CM 2010/ACOM:37. 270 pp.
- ICES. 2015. Report of the Benchmark Workshop on Plaice (WKPLE), 23–27 February 2015, ICES Headquarters, Copenhagen, Denmark. ICES CM 2015\ACOM:33. 200 pp.
- ICES 2016. Report of the Working Group on the Assessment of Demersal Stocks in the North Sea and Skagerrak (WGNSSK), 26 April–5 May 2016, Hamburg, Germany. ICES CM 2016/ ACOM:14. 19 pp.
- ICES. 2021a. “Technical Guidelines - ICES Fisheries Management Reference Points for Category 1 and 2 Stocks (2021).” <https://doi.org/10.17895/ICES.ADVISE.7891>.
- ICES. 2021b. Working Group on the Assessment of Demersal Stocks in the North Sea and Skagerrak (WGNSSK). ICES Scientific Reports. 3:66. 988 pp. <https://doi.org/10.17895/ices.pub.8211>
- ICES. 2022. Working Group on the Assessment of Demersal Stocks in the North Sea and Skagerrak (WGNSSK). ICES Scientific Reports. 4:43. 1367 pp. <http://doi.org/10.17895/ices.pub.19786285>
- ICES. 2025. Benchmark Workshop on Selected North Sea and Celtic Sea Stocks (WKBNSCS). ICES Scientific Reports. 07:42. 349 pp. <https://doi.org/10.17895/ices.pub.28715180>
- Kell, L.T., Scott, R., Hunter, E., 2004. Implications for current management advice for North Sea plaice: Part I. Migration between the North Sea and English Channel. *Journal of Sea Research, Proceedings of the Fifth International Symposium on Flatfish Ecology, Part II* 51, 287–299. <https://doi.org/10.1016/j.seares.2003.10.004>
- Lorenzen, K. 1996. The relationship between body weight and natural mortality in juvenile and adult fish: a comparison of natural ecosystems and aquaculture. *Journal of Fish Biology*, 49: 627–642.
- Nielsen, A. and Berg, C. W. 2014. Estimation of time-varying selectivity in stock assessments using state-space models. *Fisheries Research*, 158: 96–101. <https://doi.org/10.1016/j.fishres.2014.01.014>
- Sauger, C., Quinquis, J., Berthelin, C., Lepoittevin, M., Elie, N., Dubroca, L., and Kellner, K. 2023. A Quantitative Histologic Analysis of Oogenesis in the Flatfish Species *Pleuronectes platessa* as a Tool for Fisheries Management. *Animals*, 13: 2506.
- Travers-Trolet Morgane, Girardin Raphael, Coppin Franck. 2016. Calcul des indices d’abondance issus de CGFS

12.11 Tables and figures

Table 12.2.1.1. Plaice in Division 7.d. History of commercial landings; the official landings and below minimum size (BMS) landings are presented for each country participating in the fishery, along with the ICES estimates. All weights are in tonnes.

Year	Belgium	France	UK (E+W)	Others	Official landings in 7.d	BMS landings	Unallocated in 7.d	ICES estimated landings of plaice in 7.d	Quarter 1 removals^	ICES estimated landings for 7.d plaice^	ICES estimated landings for 7.e plaice	Agreed TAC for 7.d–e*
1976	147	1439	376		1962		1			1963	640	
1977	149	1714	302		2165		81			2246	702	
1978	161	1810	349		2320		156			2476	784	
1979	217	2094	278		2589		28			2617	977	
1980	435	2905	304		3644		-1421		427	2223	1178	
1981	815	3431	489		4735		-726		760	4009	1676	
1982	738	3504	541	22	4805		-765		825	4040	1878	
1983	1013	3119	548		4680		-587		950	4093	1714	
1984	947	2844	640		4431		-182		912	4249	1758	
1985	1148	3943	866		5957		-957		1022	5000	1677	
1986	1158	3288	828		5274		399		1161	5673	2078	
1987	1807	4768	1292		7867		-861	8366	1360	7006	2272	8300

Year	Belgium	France	UK (E+W)	Others	Official landings in 7.d	BMS landings	Unallocated in 7.d	ICES estimated landings of plaice in 7.d	Quarter 1 removals^	ICES estimated landings for 7.d plaice^	ICES estimated landings for 7.e plaice	Agreed TAC for 7.d-e*
1988	2165	5688	1250		9103		-318	10420	1635	8785	2835	9960
1989	2019	3713	1383		7115		-22	8758	1665	7093	2742	11700
1990	2149	4739	1479		8367		-1018	9047	1698	7349	2985	10700
1991	2265	4082	1566		7913		-1551	7813	1451	6362	2183	10700
1992	1560	3099	1572	1	6232		-1013	6337	1118	5219	1882	9600
1993	877	2792	1102		4771		-292	5331	852	4479	1614	8500
1994	1418	3199	1007	9	5633		-586	6121	1074	5047	1404	9100
1995	1157	2598	814		4569		-373	5130	934	4196	1247	8000
1996	1112	2630	856		4598		-168	5393	963	4430	1266	7530
1997	1161	3077	1078		5316		-136	6307	1127	5180	1583	7090
1998	854	3276	700		4830		1	5762	931	4831	1346	5700
1999	1306	3388	743		5437		-169	6326	1058	5268	1543	7400
2000	1298	3183	754		5235		-714	6014	1494	4521	1625	6500
2001	1346	2962	660		4968		-588	5266	886	4380	1310	6000
2002	1204	3450	841	1	5496		-378	6466	931	5118	1568	6700
2003	998	2893	756	3	4650		-1198	4033	476	3452	1411	5970

Year	Belgium	France	UK (E+W)	Others	Official landings in 7.d	BMS landings	Unallocated in 7.d	ICES estimated landings of plaice in 7.d	Quarter 1 removals^	ICES estimated landings for 7.d plaice^	ICES estimated landings for 7.e plaice	Agreed TAC for 7.d-e*
2004	954	2766	582	10	4312		-211	4624	544	4101	1332	6060
2005	832	2432	421	21	3706		-263	3867	506	3443	1300	5150
2006	1024	1935	550	16	3525		-613	3606	421	2912	1474	5151
2007	1355	2017	463	10	3845		-626	3928	620	3219	1167	5050
2008	1386	1740	471	12	3609		-69	3830	586	3540	1043	5050
2009	1002	1892	612	16	3522		-675	3442	436	2847	1061	4646
2010	1123	2190	517	62	3892		-236	4313	501	3656	1243	4274
2011	1067	1994	472	60	3593		-682	3641	358	2911	1503	4665
2012	1045	1962	542	63	3611		-564	3710	544	3047	1519	5062
2013	1295	2159	641	87	4182		-778	4157	523	3404	1525	6400
2014	1389	2229	633	76	4326		-1133	4357	645	3193	1609	5322
2015	1600	1702	392	54	3749		-766	3793	770	2983	1433	6223
2016	2247	1557	795	60	4658		-1038	4691	1020	3620	2025	12446
2017	2185	1487	823	86	4581		-903	4616	924	3678	2131	10022
2018	1879	2171	839	98	4986		-957	5104	1024	4029	1892	10360
2019	1277	1719	630	88	3714	< 1	-827	3819	885	2887	1736	10354

Year	Belgium	France	UK (E+W)	Others	Official landings in 7.d	BMS landings	Unallocated in 7.d	ICES estimated landings of plaice in 7.d	Quarter 1 removals^	ICES estimated landings for 7.d plaice^	ICES estimated landings for 7.e plaice	Agreed TAC for 7.d-e*
2020	744	984	346	50	2121	< 1	-331	2254	424	1790	1382	9154
2021	712	737	285	52	1757	< 1	-251	1832	280	1506	1413	11920
2022	596	630	341	46	1606	0	-366	1626	368	1240	1203	9138
2023* *	403	245	193	36	877	0	-222	884	193	655	1194	6775
2024* *	407	193	129	24	753	< 1	1	754	179	575	724	3930

* TACs for divisions 7.d and 7.e.

** Official values are preliminary.

^ Takes into account the "Quarter 1 removal" of 65% of the first quarter Division 7.d landings of plaice that originate from Division 7.e and Subarea 4.

Table 12.2.1.2 Plaice in Division 7.d. Official landings for plaice in Division 7.d and ICES catch estimates of Division 7.d plaice and for plaice in Division 7.d. All weights are in tonnes.

Year	Official landings of plaice in 7.d*	ICES landings of plaice in 7.d*	ICES landings of 7.d plaice	ICES discards of 7.d plaice	ICES discards of plaice in 7.d^
1987	7867	8366	7006		
1988	9103	10420	8785		
1989	7115	8758	7093		
1990	8367	9047	7349		
1991	7913	7813	6362		
1992	6232	6337	5219		
1993	4771	5331	4479		
1994	5633	6121	5047		
1995	4569	5130	4196		
1996	4598	5393	4430		
1997	5316	6307	5180		
1998	4830	5762	4831		
1999	5437	6326	5268		
2000	5235	6014	4521		
2001	4968	5266	4380		
2002	5496	6466	5118		
2003	4650	4033	3452		
2004	4312	4624	4101		
2005	3706	3867	3443		
2006	3525	3606	2912	761	928
2007	3845	3928	3219	1161	1342
2008	3609	3830	3540	682	724
2009	3522	3442	2847	966	1182
2010	3892	4313	3656	1956	2310
2011	3593	3641	2911	1523	2086
2012	3611	3710	3047	1823	2321
2013	4182	4157	3404	2210	2744

Year	Official landings of plaice in 7.d*	ICES landings of plaice in 7.d*	ICES landings of 7.d plaice	ICES discards of 7.d plaice	ICES discards of plaice in 7.d^
2014	4326	4357	3193	2730	3652
2015	3749	3793	2983	3507	4352
2016	4658	4691	3620	4149	5019
2017	4581	4616	3678	4085	5235
2018	4986	5104	4029	3736	4816
2019	3714	3819	2887	5446	6538
2020	2121	2254	1790	1537	1865
2021	1757	1832	1506	4198	5032
2022	1606	1626	1240	4600	6018
2023	877**	884	655	2124	2956
2024	753**	754	575	1513	2265

* Plaice in Division 7.d, taking into account fish caught in the first quarter in Division 7.d that come to spawn from Division 7.e and Subarea 4.

** Preliminary.

^ Including below minimum size (BMS) landings.

Table 12.2.3.1. Plaice in Division 7.d. Landings in numbers (thousands) as used in the assessment, taking into account the first quarter removal.

Year	1	2	3	4	5	6	7+
1980	53	2598	1253	370	324	50	133
1981	16	2403	5866	1643	192	106	238
1982	265	1369	5964	2262	505	138	179
1983	92	2977	2761	4048	617	151	214
1984	350	1838	6310	1928	1242	356	312
1985	142	5614	5347	3346	274	409	300
1986	679	4799	6072	2510	965	375	247
1987	25	8350	6481	2379	833	287	512
1988	16	4923	16239	3357	741	362	561
1989	826	3574	6238	6477	1770	392	497
1990	1632	2581	7550	4099	2386	535	572
1991	1542	5758	4700	3099	1614	1123	429
1992	1665	6085	3841	1183	786	697	745
1993	740	7473	3295	863	359	313	581
1994	1242	3570	6015	2131	563	280	781
1995	2592	4264	2532	2006	611	152	591
1996	1119	4762	3113	1060	951	326	585
1997	550	4168	6184	2382	724	506	722
1998	464	4323	7467	2335	360	94	289
1999	741	1737	10493	4583	696	121	223
2000	1383	6177	3432	3992	752	150	142
2001	2682	4070	3589	1385	1253	203	145
2002	924	4685	3812	1117	612	1227	530
2003	56	5279	1669	988	288	273	593
2004	477	2108	4801	873	557	259	841
2005	127	2692	1980	2107	445	290	528
2006	336	3058	2359	1025	669	102	217
2007	502	2636	2252	1129	471	338	275

Year	1	2	3	4	5	6	7+
2008	417	4236	2394	1070	615	214	284
2009	428	3208	2245	718	393	222	245
2010	615	3820	3183	1305	404	283	363
2011	102	3713	2546	877	286	57	120
2012	26	1647	3520	1612	507	211	179
2013	1	560	3327	3166	1013	284	220
2014	91	838	1950	2887	1455	327	135
2015	2	400	1183	1357	1868	1029	725
2016	7	139	1812	2107	1510	1883	1147
2017	2	98	1424	2941	1696	1003	1934
2018	19	225	1253	2994	2620	1353	2220
2019	25	227	1581	1197	1734	1412	1427
2020	23	454	464	925	816	863	1259
2021	8	189	1249	859	692	659	1015
2022	8	73	562	1677	745	463	429
2023	8	81	233	515	784	301	278
2024	1	96	370	273	340	356	280

Table 12.2.3.2. Plaice in Division 7.d. Discards in numbers (thousands) as used in the assessment, taking into account the first quarter removal.

Year	1	2	3	4	5	6	7+
1980	116	3144	830	80	23	3	1
1981	35	2908	3886	354	13	7	2
1982	582	1657	3951	488	35	9	3
1983	202	3603	1829	873	43	9	3
1984	769	2225	4180	416	86	22	6
1985	312	6794	3542	722	19	26	5
1986	1491	5808	4023	541	67	24	6
1987	55	10106	4294	513	58	18	11
1988	35	5959	10758	724	51	23	10

Year	1	2	3	4	5	6	7+
1989	1814	4326	4133	1397	123	25	8
1990	3583	3124	5002	884	166	34	8
1991	3386	6968	3114	668	112	70	7
1992	3656	7364	2545	255	55	44	17
1993	1625	9044	2183	186	25	20	10
1994	2727	4320	3985	460	39	18	10
1995	5691	5161	1677	433	42	10	9
1996	2457	5764	2062	228	66	20	6
1997	1208	5044	4097	514	50	32	12
1998	1019	5232	4947	504	25	6	4
1999	1627	2102	6951	988	48	8	3
2000	3037	7475	2274	861	52	9	2
2001	5889	4926	2378	299	87	13	2
2002	2028	5670	2525	241	43	77	10
2003	123	6389	1105	213	20	17	15
2004	1048	2552	3181	188	39	16	15
2005	279	3259	1312	454	31	18	5
2006	668	2589	1590	130	29	7	0
2007	1114	4501	1528	241	46	52	4
2008	782	2156	494	187	31	13	1
2009	1005	2523	1975	261	41	6	9
2010	1562	8409	2782	260	21	2	4
2011	1732	6789	1197	174	51	48	0
2012	1773	5870	2642	662	83	44	10
2013	3923	4913	4792	858	76	48	3
2014	6313	5955	2834	1612	352	73	3
2015	2256	8537	5999	2240	1058	396	17
2016	2845	10142	7850	2079	842	318	59
2017	4620	9163	6243	3414	602	348	45

Year	1	2	3	4	5	6	7+
2018	1537	6998	8432	7505	2500	868	462
2019	9895	10731	9748	2760	1146	303	254
2020	1326	3221	3962	2583	1468	582	293
2021	13116	8124	6654	4958	2521	1028	398
2022	10680	8494	7123	8940	2106	776	186
2023	3330	7259	3480	2423	2339	1092	240
2024	2131	3605	4559	1658	812	962	541

Table 12.2.4.1. Plaice in Division 7.d. Weights in the landings (kg).

Year	1	2	3	4	5	6	7+
1980	0.3144	0.3174	0.5077	0.6379	0.8007	1.1589	1.4387
1981	0.2305	0.2884	0.3598	0.4476	0.6868	0.8392	1.0318
1982	0.2374	0.2626	0.3421	0.4177	0.6202	0.7704	1.1933
1983	0.2537	0.2823	0.3328	0.4005	0.5169	0.7839	1.1775
1984	0.2111	0.2673	0.3044	0.3642	0.4603	0.6243	0.8525
1985	0.2412	0.264	0.2859	0.4056	0.4768	0.5414	0.8201
1986	0.2306	0.3123	0.3378	0.4143	0.5572	0.496	0.8226
1987	0.2501	0.281	0.3587	0.4753	0.5749	0.7802	0.9668
1988	0.2793	0.2564	0.3071	0.4133	0.5357	0.6285	0.9256
1989	0.1993	0.2658	0.3183	0.3669	0.469	0.6426	1.0734
1990	0.2086	0.2657	0.3384	0.3924	0.5014	0.6332	1.0912
1991	0.2235	0.2751	0.3089	0.3874	0.4509	0.5523	1.0089
1992	0.181	0.2755	0.3501	0.4267	0.5062	0.5818	0.7909
1993	0.2168	0.2681	0.3312	0.4258	0.4997	0.5825	0.8525
1994	0.2481	0.2757	0.2941	0.3635	0.4759	0.5882	0.9957
1995	0.2149	0.2672	0.3086	0.3845	0.4782	0.6784	0.9317
1996	0.2281	0.3097	0.2994	0.4088	0.4904	0.6638	1.1149
1997	0.2006	0.2541	0.3004	0.3347	0.4456	0.5817	1.0241
1998	0.1675	0.257	0.2812	0.4013	0.5288	0.8026	1.1748
1999	0.2037	0.2533	0.2429	0.3163	0.4766	0.7764	1.1331

Year	1	2	3	4	5	6	7+
2000	0.2165	0.2563	0.273	0.296	0.3923	0.6025	0.9526
2001	0.2328	0.2729	0.3281	0.4007	0.4841	0.6952	1.1326
2002	0.2663	0.3028	0.3477	0.4695	0.4995	0.6069	1.0438
2003	0.2469	0.2958	0.3784	0.4503	0.6027	0.5804	0.7894
2004	0.2413	0.2835	0.3704	0.4547	0.5606	0.6961	0.8558
2005	0.2939	0.3004	0.3432	0.4439	0.5251	0.6392	1.067
2006	0.2578	0.3059	0.3581	0.4124	0.5171	0.6281	0.9797
2007	0.2578	0.3276	0.3949	0.4781	0.5764	0.6998	1.0497
2008	0.2427	0.2936	0.3669	0.4599	0.6039	0.7028	1.0667
2009	0.2804	0.3086	0.3604	0.4539	0.5206	0.6714	1.0139
2010	0.2683	0.3091	0.3446	0.4052	0.4951	0.5785	0.885
2011	0.2833	0.3052	0.3766	0.4687	0.6541	0.8158	1.2027
2012	0.2828	0.307	0.3605	0.4297	0.5039	0.658	0.9921
2013	0.205	0.3226	0.3199	0.3801	0.5115	0.6636	1.1309
2014	0.1719	0.2796	0.3257	0.4029	0.5302	0.7174	1.0313
2015	0.153	0.2973	0.3274	0.36	0.4227	0.5677	0.8477
2016	0.1628	0.3253	0.3158	0.3477	0.4042	0.4636	0.6843
2017	0.2584	0.2307	0.2848	0.3309	0.3861	0.4215	0.6202
2018	0.2366	0.2458	0.2808	0.3008	0.346	0.3961	0.5736
2019	0.2079	0.2549	0.2898	0.3088	0.3581	0.3795	0.5879
2020	0.2227	0.2835	0.2704	0.2922	0.3353	0.3893	0.5171
2021	0.1603	0.248	0.2697	0.2761	0.3017	0.316	0.4604
2022	0.1117	0.2539	0.2623	0.2764	0.3111	0.3483	0.5046
2023	0.1939	0.2252	0.2587	0.2739	0.2856	0.305	0.4277
2024	0.2022	0.2223	0.2693	0.3043	0.3336	0.3616	0.4619

Table 12.2.4.2. Plaice in Division 7.d. Weights in the discards (Kg).

Year	1	2	3	4	5	6	7+
1980	0.1555	0.1626	0.26	0.2992	0.3188	0.4796	0.4736
1981	0.114	0.1477	0.1843	0.2099	0.2735	0.3473	0.3215

Year	1	2	3	4	5	6	7+
1982	0.1174	0.1345	0.1752	0.1959	0.247	0.3189	0.4209
1983	0.1255	0.1446	0.1705	0.1879	0.2058	0.3244	0.3759
1984	0.1044	0.1369	0.1559	0.1709	0.1833	0.2584	0.2646
1985	0.1193	0.1352	0.1464	0.1902	0.1899	0.2241	0.2553
1986	0.1141	0.16	0.173	0.1944	0.2219	0.2053	0.2641
1987	0.1237	0.1439	0.1837	0.2229	0.2289	0.3229	0.3123
1988	0.1382	0.1313	0.1573	0.1939	0.2133	0.2601	0.2948
1989	0.0986	0.1361	0.163	0.1721	0.1868	0.2659	0.3275
1990	0.1032	0.1361	0.1733	0.1841	0.1996	0.2621	0.3348
1991	0.1105	0.1409	0.1582	0.1817	0.1796	0.2286	0.3138
1992	0.0895	0.1411	0.1793	0.2001	0.2016	0.2408	0.2796
1993	0.1072	0.1373	0.1696	0.1997	0.199	0.2411	0.29
1994	0.1227	0.1412	0.1506	0.1705	0.1895	0.2434	0.3005
1995	0.1063	0.1369	0.1581	0.1804	0.1904	0.2808	0.256
1996	0.1128	0.1586	0.1533	0.1918	0.1953	0.2747	0.3261
1997	0.0992	0.1301	0.1539	0.157	0.1774	0.2408	0.3208
1998	0.0828	0.1316	0.144	0.1882	0.2105	0.3322	0.3564
1999	0.1007	0.1297	0.1244	0.1484	0.1898	0.3213	0.3184
2000	0.1071	0.1313	0.1398	0.1389	0.1562	0.2494	0.2921
2001	0.1151	0.1398	0.1681	0.1879	0.1927	0.2877	0.377
2002	0.1317	0.1551	0.1781	0.2202	0.1989	0.2512	0.3153
2003	0.1221	0.1515	0.1938	0.2112	0.24	0.2402	0.2771
2004	0.1193	0.1452	0.1897	0.2133	0.2232	0.2881	0.2897
2005	0.1453	0.1539	0.1758	0.2082	0.2091	0.2645	0.2969
2006	0.1242	0.1446	0.1694	0.2013	0.2181	0.2635	0.1194
2007	0.1336	0.1574	0.1638	0.1611	0.1353	0.1468	0.1182
2008	0.1527	0.1761	0.2332	0.2776	0.3036	0.3767	1.1644
2009	0.1323	0.1668	0.1771	0.191	0.1833	0.1515	0.223
2010	0.0997	0.1446	0.1882	0.201	0.2375	0.3925	0.334

Year	1	2	3	4	5	6	7+
2011	0.1175	0.1551	0.1767	0.1946	0.1849	0.2002	0.1224
2012	0.128	0.1547	0.1899	0.231	0.234	0.2274	0.3225
2013	0.0908	0.1642	0.1764	0.1775	0.2553	0.1992	0.255
2014	0.0907	0.1713	0.1909	0.2216	0.2194	0.2945	0.2749
2015	0.1207	0.153	0.1832	0.1939	0.2294	0.3555	0.4068
2016	0.1157	0.1754	0.1715	0.199	0.2075	0.237	0.325
2017	0.1092	0.1805	0.1755	0.1829	0.2027	0.2058	0.188
2018	0.1055	0.1088	0.1263	0.1432	0.1567	0.1919	0.1793
2019	0.1321	0.1471	0.1784	0.1797	0.1754	0.2067	0.232
2020	0.0896	0.1007	0.1056	0.1255	0.1401	0.1527	0.1723
2021	0.0938	0.1185	0.1276	0.1204	0.1299	0.145	0.1528
2022	0.0999	0.1342	0.1137	0.1258	0.1401	0.1619	0.1673
2023	0.1088	0.0908	0.1032	0.1157	0.1229	0.1285	0.1379
2024	0.0967	0.0991	0.1043	0.1087	0.1273	0.1274	0.1255

Table 12.2.4.3. Plaice in Division 7.d. Weights in the stock (kg).

Year	1	2	3	4	5	6	7+
1980	0.171	0.332	0.482	0.622	0.751	0.87	1.1955
1981	0.11	0.216	0.317	0.414	0.506	0.594	0.9231
1982	0.105	0.208	0.308	0.406	0.502	0.596	0.8672
1983	0.097	0.192	0.286	0.379	0.47	0.56	0.8517
1984	0.082	0.164	0.248	0.333	0.42	0.507	0.7361
1985	0.084	0.171	0.259	0.348	0.44	0.533	0.7757
1986	0.101	0.205	0.311	0.42	0.532	0.646	0.8482
1987	0.122	0.242	0.361	0.479	0.596	0.712	0.9276
1988	0.084	0.168	0.254	0.34	0.427	0.514	0.7128
1989	0.079	0.162	0.25	0.342	0.439	0.541	0.8523
1990	0.085	0.23	0.322	0.346	0.465	0.549	1.1145
1991	0.143	0.219	0.275	0.335	0.375	0.472	0.9545

Year	1	2	3	4	5	6	7+
1992	0.088	0.241	0.336	0.421	0.477	0.521	0.7232
1993	0.108	0.258	0.296	0.379	0.493	0.539	0.7253
1994	0.165	0.198	0.276	0.331	0.383	0.493	0.8638
1995	0.124	0.257	0.286	0.354	0.442	0.707	0.8508
1996	0.178	0.229	0.263	0.347	0.354	0.474	0.9313
1997	0.059	0.202	0.256	0.266	0.417	0.53	0.8983
1998	0.072	0.203	0.273	0.361	0.53	0.67	0.8701
1999	0.072	0.172	0.213	0.351	0.429	0.644	0.9022
2000	0.068	0.184	0.204	0.246	0.355	0.554	0.926
2001	0.093	0.206	0.274	0.338	0.404	0.624	1.1003
2002	0.2327	0.2865	0.3221	0.5658	0.5789	0.8948	1.1695
2003	0.2131	0.2767	0.3818	0.427	0.5657	0.4988	0.654
2004	0.2337	0.2958	0.405	0.4996	0.5833	0.7315	0.8266
2005	0.2283	0.283	0.3447	0.4413	0.564	0.6587	0.9395
2006	0.2234	0.2753	0.3468	0.4286	0.5185	0.6225	0.8406
2007	0.325	0.2781	0.3546	0.4791	0.5556	0.6247	0.7612
2008	0.1838	0.259	0.3034	0.3643	0.4791	0.4944	0.9297
2009	0.2668	0.2819	0.3279	0.3934	0.4562	0.6485	0.8364
2010	0.2394	0.2855	0.3431	0.3826	0.4972	0.4774	0.7799
2011	0.234	0.2943	0.3956	0.5237	0.705	0.7782	1.1143
2012	0.2131	0.2794	0.3415	0.3813	0.4253	0.5391	0.8307
2013	0.2131	0.2419	0.3126	0.391	0.5095	0.6419	1.0529
2014	0.172	0.2507	0.3176	0.406	0.5384	0.7306	1.006
2015	0.1781	0.2472	0.4025	0.4314	0.5369	0.6676	0.8444
2016	0.1808	0.275	0.3261	0.3888	0.4882	0.5349	0.7054
2017	0.2131	0.2628	0.2869	0.3221	0.411	0.4784	0.6374
2018	0.1743	0.1955	0.2442	0.2597	0.3104	0.3815	0.4803
2019	0.2131	0.2767	0.2549	0.2843	0.3087	0.3781	0.5127
2020	0.2131	0.2982	0.2935	0.3445	0.4095	0.4511	0.5861

Year	1	2	3	4	5	6	7+
2021	0.1578	0.2321	0.2581	0.2824	0.3369	0.334	0.4998
2022	0.1549	0.2294	0.2573	0.2762	0.2905	0.329	0.4005
2023	0.152	0.241	0.2551	0.2743	0.2921	0.3725	0.4451
2024	0.2131	0.1997	0.2421	0.2934	0.3303	0.3846	0.4605

Table 12.2.6.1. Plaice in Division 7.d. UK-BTS tuning fleet.

Year	1	2	3	4	5	6
1989	3.8	15.8	28.9	31.7	4	1.7
1990	9.2	9.4	11.1	11.7	12.6	1.5
1991	16.8	14.5	11.5	8.7	8.6	4.6
1992	22.4	21.3	6.6	6.6	7.2	5.4
1993	4.6	20.2	8	2.8	2.9	2.4
1994	9.3	8.5	10.1	5.9	2	0.6
1995	14.5	6.2	3.8	5.7	2.2	0.8
1996	22.1	17.3	1.7	1	2	1.3
1997	48.2	28.6	11	1.2	1.6	0.5
1998	30.6	37.9	12.1	5	0.6	0.6
1999	12.8	10.7	28.8	4.6	1.6	0.3
2000	19.5	30.2	18.8	20.5	5	1.3
2001	27.9	20.3	14.1	9.8	14.8	2.7
2002	37.9	25.9	12.5	5.5	2.6	5.3
2003	10.6	39.7	9.8	4.4	2.3	1.1
2004	52.9	22.5	20.7	4.8	1.1	0.3
2005	15.6	36.2	12.8	10	3.2	1.1
2006	30.1	28.8	16.8	5.9	4.3	1.3
2007	53.1	28.9	12.2	6.2	3.2	2.9
2008	39.6	40.6	10.5	4.3	3.8	1.8
2009	77.7	39.5	20.9	5.9	3.2	2.3
2010	64.2	64.7	17.7	9.1	3.1	1.7
2011	115.1	112.2	39.6	10.3	7	2.9

Year	1	2	3	4	5	6
2012	24.7	81.1	56	18.6	4.2	3.3
2013	32.3	61	88.2	45	10.2	3.4
2014	145.3	156.5	50.7	62.1	26.7	9
2015	38	178.7	63.2	30.2	33.4	15.7
2016	12.5	101.4	102.9	37.9	21.3	23.2
2017	50.1	102.1	83.2	56	16.6	8.4
2018	25.6	97	112.2	52.4	30.3	9.3
2019	117.5	81.7	55.3	37.3	18.2	11.7
2020	20.7	109.1	60.2	25.1	15.1	7.6
2021	17.9	72	93.1	71.6	11.8	7.9
2022	NA	NA	NA	NA	NA	NA
2023	10.5	66.6	41.8	28.7	25.6	15.6
2024	21.6	48.2	56.6	9.9	8.2	6.3

Table 12.2.6.2. Plaice in Division 7.d. FR-GFS tuning fleet.

Year	1	2	3	4	5	6
1990	402.507913	134.544569	186.637279	96.0263051	69.5488544	21.7670705
1991	380.810992	189.952783	78.3415568	75.5400497	26.2775966	11.734421
1992	1134.18141	403.225951	91.4909112	23.0090474	11.908858	17.2620626
1993	375.73873	506.507073	166.228505	53.4451961	21.8498204	18.5930351
1994	606.164377	270.149948	135.284923	49.9411463	25.8992156	40.8360588
1995	231.578842	87.5708008	47.5363112	31.5411667	13.5891628	47.9032626
1996	575.787757	236.005349	40.1508477	14.7147164	21.7434395	18.9716723
1997	911.634236	478.644567	215.526962	18.7864544	10.023066	38.6729736
1998	505.135094	853.781242	202.531603	35.5676144	0	10.9974993
1999	1009.43381	423.164612	656.914371	98.5763955	19.7806139	15.4827287
2000	292.845315	652.87708	205.053706	105.445921	25.265433	13.6076502
2001	451.683317	322.005778	137.59589	66.314114	33.8981272	15.5029803
2002	562.56506	421.889144	210.632489	75.3640568	41.8604558	24.4500345

Year	1	2	3	4	5	6
2003	513.01123	864.848487	166.228644	32.5278126	18.9573896	26.242738
2004	750.975483	607.421062	282.640517	81.0867902	7.6360552	21.6007803
2005	494.570485	741.199262	517.614829	134.689321	46.7391224	24.1145285
2006	544.667426	545.741052	246.291026	81.5475881	38.0921187	22.2850196
2007	552.757087	497.243704	276.15137	77.2547995	30.7067166	15.7352483
2008	438.204485	1208.30093	182.066181	62.1352622	27.7825702	15.6114596
2009	695.699061	523.223371	213.632252	51.6870784	26.4792183	19.1867842
2010	1915.24802	1250.38099	205.489985	41.4392933	27.1075468	15.1787884
2011	1152.38739	1911.85256	542.214419	60.7495993	17.4722392	16.7079269
2012	869.654097	1900.53445	1310.01744	266.039159	81.2824074	19.9516402
2013	309.554193	639.331164	903.256053	218.889767	63.3748293	17.0143809
2014	895.503117	765.664299	580.056612	785.390996	168.816467	22.4094154
2015	301.75832	802.59768	807.49035	252.926569	271.843782	85.7182112
2016	207.472148	544.603518	888.797683	329.369801	214.652341	106.760361
2017	369.734234	467.572202	538.307549	404.433965	115.06771	44.9524633
2018	179.580724	525.643416	308.454588	329.175298	138.397085	48.2734572
2019	1283.86626	859.47583	558.687595	205.692416	287.713582	174.027497
2020	1072.65426	2477.68082	528.714284	176.626508	120.803582	91.0972415
2021	1188.39268	1127.27979	1167.32784	164.335899	117.084138	49.5267116
2022	764.745879	408.419593	369.430716	281.377839	63.4631131	47.3830023
2023	468.45747	772.505204	230.633845	195.505826	165.878929	34.4260036
2024	429.712831	508.517836	414.810283	106.315037	66.5112899	30.2189656

Table 12.3.1.1. Plaice in Division 7.d. Fishing mortality (F) at age.

Year	1	2	3	4	5	6	7+
1980	0.00451203	0.17266284	0.43552471	0.4166408	0.25930958	0.16586317	0.12658651
1981	0.00579811	0.19995957	0.59548354	0.58363319	0.32701981	0.21864832	0.17185232
1982	0.00716743	0.20941464	0.61611645	0.65374634	0.37450473	0.26452472	0.19579653
1983	0.00768813	0.2138813	0.61191134	0.64256572	0.35158709	0.26102025	0.20496636
1984	0.00940894	0.23759128	0.68685618	0.71722127	0.40147655	0.31680824	0.24831614

Year	1	2	3	4	5	6	7+
1985	0.01180715	0.28450061	0.72181859	0.67108124	0.34543802	0.28610813	0.21724367
1986	0.01272402	0.30052734	0.78262421	0.71994785	0.42155852	0.33222418	0.23027839
1987	0.00909808	0.25043705	0.66133507	0.64942592	0.42716767	0.3413739	0.27073394
1988	0.01095114	0.25353943	0.62382431	0.62758969	0.46142443	0.38892509	0.32246188
1989	0.02126418	0.29808897	0.61548434	0.61512653	0.51757546	0.43629537	0.36521628
1990	0.04220188	0.39392331	0.70050792	0.61243196	0.50344509	0.45365839	0.39289377
1991	0.08340895	0.57047811	0.84769838	0.64791185	0.48788092	0.43265903	0.3709945
1992	0.08822022	0.56242041	0.79919554	0.54872966	0.4078679	0.36450917	0.33242911
1993	0.07757681	0.48393929	0.63872825	0.43578203	0.31677883	0.27661586	0.27149793
1994	0.10632437	0.56404428	0.80497278	0.58886835	0.42574743	0.34502835	0.35007883
1995	0.09317175	0.51698607	0.75215734	0.56949602	0.40696487	0.30836146	0.33993257
1996	0.062534	0.40830273	0.76927789	0.6694748	0.51357812	0.38936623	0.42870392
1997	0.05247184	0.37718227	0.87827528	1.00024835	0.74599284	0.5509552	0.57672187
1998	0.03742559	0.29004299	0.68218745	0.78300152	0.54741947	0.36793464	0.37763781
1999	0.04750541	0.33380976	0.67509881	0.76489088	0.53466906	0.36374325	0.35554322
2000	0.08098573	0.47403903	0.70985693	0.59284197	0.38741289	0.27075892	0.26958713
2001	0.0913788	0.53748892	0.78234415	0.53603578	0.32545226	0.24247824	0.24134924
2002	0.08673032	0.59987102	0.86398187	0.63821057	0.43157511	0.38108209	0.37813425
2003	0.03976883	0.40144087	0.59181896	0.46002518	0.3376142	0.32177062	0.32706593
2004	0.04070104	0.40316103	0.6438245	0.51992173	0.42700434	0.4160101	0.42360038
2005	0.03819969	0.38982875	0.60011395	0.51596431	0.42129309	0.39705247	0.39267234
2006	0.04017628	0.38308824	0.54836727	0.41547913	0.3106517	0.25622478	0.24481348
2007	0.05554964	0.45274004	0.64007017	0.47485173	0.35763446	0.2977166	0.2594323
2008	0.04737491	0.3873874	0.59863385	0.46072686	0.34504568	0.28144403	0.22504978
2009	0.04269432	0.34219412	0.54731322	0.42557141	0.31339196	0.25945043	0.19580352
2010	0.05209538	0.35601512	0.57702779	0.47349357	0.34912037	0.29723824	0.20778613
2011	0.02500948	0.19288103	0.28793354	0.25790364	0.19343283	0.16229139	0.10916723
2012	0.02346116	0.15792257	0.24205963	0.2509029	0.2111036	0.1842818	0.1179587
2013	0.02912258	0.15927121	0.23291539	0.24312332	0.20940872	0.18433359	0.11267793

Year	1	2	3	4	5	6	7+
2014	0.02955195	0.14841113	0.2114998	0.20854948	0.17169997	0.1445157	0.08650619
2015	0.03545827	0.16212309	0.24827385	0.26706443	0.23639479	0.21951654	0.13699853
2016	0.04470581	0.18000699	0.27123205	0.31049384	0.29134676	0.27942644	0.1671094
2017	0.05079436	0.18660414	0.28440881	0.34112473	0.33467256	0.3385923	0.19539537
2018	0.06568191	0.21632374	0.37507797	0.48514396	0.49490067	0.50533505	0.26890358
2019	0.073399	0.21378956	0.35437116	0.4213389	0.42981853	0.43776344	0.22145581
2020	0.03839826	0.13277715	0.22049042	0.31517515	0.36075572	0.39766014	0.18648103
2021	0.09909042	0.22486112	0.32343711	0.44496807	0.4893252	0.50831526	0.19602695
2022	0.14598746	0.29517812	0.41123472	0.51125884	0.47522704	0.45105443	0.14527802
2023	0.09570086	0.22380043	0.31847905	0.3930661	0.38206048	0.38189197	0.1194455
2024	0.06767623	0.17977961	0.25372574	0.31873335	0.31250139	0.34786218	0.11792836

Table 12.3.1.2. Plaice in Division 7.d. Stock numbers from the assessment (thousands).

Year	1	2	3	4	5	6	7+
1980	54740	32706	8506	1862	1364	411	1495
1981	32361	38411	19733	4130	937	797	1324
1982	53093	22015	23305	7847	1740	528	1371
1983	47086	37059	12880	9474	3030	913	1214
1984	60669	32284	21884	5167	3774	1626	1336
1985	70240	41814	18751	8161	1900	1925	1731
1986	128445	47948	22441	6819	3124	1060	2191
1987	79986	90227	25626	7481	2532	1553	1995
1988	54374	55528	51983	9733	2917	1272	2074
1989	35706	37812	31499	20720	3890	1410	1859
1990	40911	24250	21018	12715	8365	1752	1734
1991	54069	26945	12360	7780	5239	3812	1778
1992	65587	34715	10998	4004	3052	2476	2904
1993	35073	42176	14432	3673	1768	1557	2992
1994	34560	22515	18919	5731	1810	994	2750
1995	61351	21154	9423	6339	2382	904	2097

Year	1	2	3	4	5	6	7+
1996	65344	39464	9039	3359	2698	1223	1706
1997	90765	42166	19311	3129	1344	1222	1525
1998	38861	61288	21233	5909	877	486	1207
1999	43795	25124	35035	7985	2006	395	906
2000	39289	28847	13386	13332	2787	887	711
2001	38908	25354	12597	4942	5692	1437	949
2002	52837	24224	10974	4140	2188	3232	1491
2003	30298	34316	9450	3441	1614	1099	2551
2004	35091	20148	16601	3856	1651	874	2108
2005	29817	23151	9845	6408	1714	834	1545
2006	29786	20009	11252	4000	2871	846	1250
2007	40033	19583	9882	4767	1991	1609	1282
2008	38323	26314	8788	3833	2255	1064	1698
2009	64504	25212	12680	3545	1831	1220	1687
2010	97923	42892	12932	5324	1744	1029	1823
2011	108073	64947	21358	5312	2488	940	1732
2012	66954	74144	38605	11656	3103	1580	1841
2013	80005	44978	47281	22225	6697	1924	2294
2014	116848	54629	27583	28158	13065	4120	2816
2015	93358	79419	34131	16628	17218	8404	4895
2016	76146	63972	49425	19351	9673	10413	8651
2017	74533	52026	39376	27777	10546	5509	11980
2018	73914	49421	31975	22331	14766	5772	10890
2019	96981	49317	29111	16328	10284	6835	9239
2020	62163	62683	29392	14854	8168	5092	9205
2021	50429	41766	40365	17947	8101	4371	8526
2022	72074	31164	24203	22020	8756	3803	7359
2023	43959	43952	16452	11803	9907	4241	6830
2024	41608	27814	25930	8664	5974	5122	7088

Table 12.3.1.3 Plaice in Division 7.d. Assessment summary. Recruitment is in thousands; all weights are in tonnes. “Low” and “High” refer to 95% confidence intervals.

Year	Recruitment (age 1)			Spawning-stock biomass			Land-ings	Dis-cards*	Fishing pressure (ages 3–6)		
	Low	R	High	Low	SSB	High			Low	F	High
1980	41231	54740	72674	15581	20277	26388	2223	778	0.22	0.32	0.47
1981	25039	32361	41823	14334	18032	22685	4009	1231	0.31	0.43	0.61
1982	41370	53093	68137	14367	17781	22006	4040	1091	0.34	0.48	0.67
1983	36736	47086	60353	14359	17530	21401	4093	1035	0.34	0.47	0.64
1984	47092	60669	78160	13425	16186	19514	4248	1131	0.40	0.53	0.71
1985	54943	70240	89796	15506	18485	22037	5000	1622	0.39	0.51	0.65
1986	99381	128445	166009	21974	25720	30104	5673	1922	0.45	0.56	0.71
1987	63240	79986	101166	33092	38693	45243	7006	2387	0.41	0.52	0.66
1988	42602	54374	69398	24492	28484	33128	8785	2640	0.42	0.53	0.66
1989	28278	35706	45084	21216	24668	28681	7093	1714	0.43	0.55	0.69
1990	32612	40911	51323	19957	23363	27351	7349	1869	0.45	0.57	0.71
1991	43415	54069	67336	15800	18514	21694	6362	2008	0.49	0.60	0.75
1992	52477	65587	81973	16261	18944	22069	5219	1900	0.43	0.53	0.65
1993	28047	35073	43860	17140	19950	23219	4479	1836	0.34	0.42	0.52
1994	27029	34560	44190	13496	15779	18449	5047	1638	0.44	0.54	0.66
1995	48982	61351	76844	12950	15065	17526	4196	1668	0.42	0.51	0.62
1996	52907	65344	80705	15144	17603	20461	4430	1572	0.48	0.59	0.72
1997	69326	90765	118836	14590	17010	19831	5180	1508	0.64	0.79	0.98
1998	31297	38861	48254	18013	21202	24956	4831	1588	0.47	0.60	0.75
1999	34301	43795	55918	14015	16355	19086	5268	1460	0.46	0.59	0.75
2000	31227	39289	49433	11430	13394	15696	4521	1755	0.39	0.49	0.62
2001	30503	38908	49629	12455	14728	17416	4380	1844	0.38	0.47	0.59
2002	42398	52837	65845	17729	20936	24723	5118	1680	0.46	0.58	0.72
2003	24362	30298	37682	15355	18116	21372	3452	1255	0.33	0.43	0.55
2004	27924	35091	44097	15937	18984	22614	4101	1157	0.39	0.50	0.65
2005	23950	29817	37121	13679	16576	20087	3443	880	0.36	0.48	0.65
2006	23267	29786	38132	12319	15064	18421	2912	761	0.28	0.38	0.52

Year	Recruitment (age 1)			Spawning-stock biomass			Land-ings	Dis-cards*	Fishing pressure (ages 3–6)		
	Low	R	High	Low	SSB	High			Low	F	High
2007	31600	40033	50717	13824	16887	20628	3219	1161	0.32	0.44	0.61
2008	30119	38323	48761	11947	14984	18794	3540	682	0.30	0.42	0.59
2009	49846	64504	83470	15278	19134	23963	2847	966	0.27	0.39	0.55
2010	76237	97923	125776	20659	25993	32704	3656	1956	0.29	0.42	0.62
2011	83221	108073	140346	30345	38718	49403	2911	1523	0.149	0.23	0.34
2012	52480	66954	85419	33729	42623	53861	3047	1823	0.148	0.22	0.33
2013	63664	80005	100541	34228	43658	55686	3404	2210	0.148	0.22	0.32
2014	93342	116848	146274	39233	49927	63536	3193	2730	0.129	0.184	0.26
2015	73502	93358	118578	48658	60802	75977	2983	3507	0.175	0.24	0.34
2016	59496	76146	97455	46922	58205	72200	3620	4149	0.21	0.29	0.40
2017	59340	74533	93617	40395	50409	62905	3678	4085	0.24	0.33	0.45
2018	57041	73914	95779	29476	37064	46605	4029	3736	0.34	0.47	0.64
2019	76376	96981	123145	31397	39614	49982	2887	5446	0.29	0.41	0.58
2020	48609	62163	79495	34736	44321	56552	1790	1537	0.22	0.32	0.47
2021	37050	50429	68639	26405	33937	43619	1506	4198	0.31	0.44	0.63
2022	50304	72074	103263	21292	27910	36586	1240	4600	0.32	0.46	0.66
2023	28620	43959	67519	18497	25855	36138	655	2124	0.24	0.37	0.57
2024	22859	41608	75734	15288	22859	34178	575	1513	0.186	0.31	0.51
2025		55540**		14628	22798	35115					

* Since 2006, discard estimates are raised from observer programmes. Prior to 2006, discards are reconstructed. Discards include below minimum size (BMS) landings.

** Recruitment in 2025 is the geometric mean of resampled recruitment estimates from 1980 to 2024.

Table 12.5.3.1.1. Plaice in Division 7.d. Values in the forecast and for the interim year for Division 7.d plaice stock only.

Variable	Value	Notes
$F_{\text{ages } 3-6}$ (2025)	0.308	Average exploitation pattern (2022–2024), scaled to average $F_{\text{ages } 3-6}$ in 2024.
SSB (2026)	23993	Short-term forecast (STF); tonnes.
$R_{\text{age } 1}$ (2025–2027)	55540	Resampled recruitments from the years 1980–2024 (geometric mean shown); thousands
Total catch (2025)	2310	STF; tonnes.
Projected landings (2025)	601	STF; assuming average landings ratio by age, 2022–2024; tonnes.
Projected discards (2025)	1709	STF; assuming average discards ratio by age, 2022–2024; tonnes.

Table 12.5.3.1.2. Plaice in Division 7.d. Annual catch scenarios. All weights are in tonnes.

Basis	Division 7.d plaice stock						All plaice in Division 7.d [#]									
	Total catch (2026)	Projected landings (2026)	Projected discards* (2026)	F _{total} (ages 3–6) (2026)	F _{projected} landings (ages 3–6) (2026)	F _{projected} discards (ages 3–6) (2026)	SSB (2027)	% SSB change **	Probability of SSB (2027) < B _{lim} (%)	% advice change [§]	Total catch (2026)	Projected landings (2026)	Projected discards* (2026)	% change in projected catches [^]	% advice change ^{##}	% TAC change ^{##}
ICES advice basis																
MSY approach: F _{MSY} × SSB (2026)/MSY B _{trigger}	1151	284	867	0.139	0.029	0.110	27504	14.6	34	-56	1384	341	1042	-34	-56	-56
Other scenarios																
F = F _{MSY lower} × SSB (2026)/MSY B _{trigger}	846	210	636	0.101	0.021	0.080	28071	17.0	31	-67	1017	252	765	-51	-67	-67
F _{MSY lower}	1213	299	914	0.147	0.030	0.117	27375	14.1	35	-53	1458	359	1099	-30	-53	-53
F _{MSY}	1640	403	1237	0.203	0.042	0.161	26635	11.0	38	-37	1972	485	1487	-5.6	-37	-37
F = 0	0	0	0	0	0	0	29551	23	22	-100	0	0	0	-100	-100	-100
F _{pa}	1960	480	1480	0.246	0.050	0.196	26016	8.4	44	-25	2357	577	1779	12.9	-24	-24
SSB (2027) = B _{lim}	2501	611	1890	0.322	0.066	0.256	25105	4.6	50	-3.8	3007	735	2272	44	-3.4	-3.4
SSB (2027) = B _{pa} = MSY B _{trigger} ^^																
F = F ₂₀₂₅	2405	587	1818	0.308	0.063	0.245	25291	5.4	49	-7.5	2892	706	2186	38.5	-7.1	-7.1

* Including below minimum size (BMS) landings, assuming recent discard rate.

** SSB 2027 relative to SSB 2026.

\$ Total advised catch of plaice stock in 2026 relative to the advice value 2025 (2600 tonnes).

^ Projected catch of plaice caught in Division 7d in 2026 relative to the ICES estimates of catches in 2024 (2088 tonnes).

^^ B_{pa} and $MSY_{Btrigger}$ cannot be achieved in 2027, even with zero catches.

All plaice caught in Division 7.d, including plaice originating from the North Sea and the western English Channel. Of the plaice caught in Division 7.d, 12.86% are assumed to originate from the North Sea and 3.86% from the western English Channel (average ratio calculated over the years 2003–2024). Both of these predicted catches are added to the predicted values for the ple.27.7d stock and applied to projected catch, projected landings, and projected discards. Following ICES advice basis, catches in Division 7.d will comprise 1151 tonnes coming from the ple.27.7d stock, plus 178 tonnes coming from the ple.27.420 stock and 54 tonnes from the ple.27.7e stock.

#² Projected catch of plaice caught in division 7d in 2026 relative to the advice value and the TAC in 2025 (both 3113 tonnes).




Figure 12.2.1.1. Plaice in Division 7.d. Official landings in 7.d and 7.e compared to the TAC: since 2019, the advice was given on catch rather than landings.




Figure 12.2.1.2. Plaice in Division 7.d. Official landings by country.




Figure 12.2.2.1. Plaice in Division 7.d. Proportions of landings and discards per country with and without age distribution provided.




Figure 12.2.3.1. Plaice in Division 7.d. Age composition of the landings.




Figure 12.2.3.2. Plaice in Division 7.d. Age composition of the discards (discards before 2006 are reconstructed).




Figure 12.2.3.3. Plaice in Division 7.d. Discards-at-age ratio (discards numbers/landings numbers) per age.




Figure 12.2.3.4. Plaice in Division 7.d. Discards-at-age ratio (discards numbers/landings numbers) through time.




Figure 12.2.4.1. Plaice in Division 7.d. Stock, Landing and discard weights.




Figure 12.2.6.1. Plaice in Division 7.d. Survey Consistency: mean standardized indices by surveys for each age.




Figure 12.2.6.2. Plaice in Division 7.d. UK-BTS (top) and FR-GFS (bottom) indices consistencies.




Figure 12.3.1.1. Plaice in Division 7.d. Catch time-series: observed (cross) vs. modelled (line).




Figure 12.3.1.2. Plaice in Division 7.d. Catch (left), UK-BTS (middle) and FR-GFS (right) time-series: observed (dots) vs. modelled (line), and per age.




Figure 12.3.1.3. Plaice in Division 7.d. The standardized one-observation-ahead residuals for catch data, UK-BTS and FRGFS surveys from the final SAM assessment.



Figure 12.3.1.4. Plaice in Division 7.d. The process residuals from the final SAM assessment.




Figure 12.3.1.5. Plaiice in Division 7.d. Leave-one-out analysis results for spawning-stock biomass (SSB; tonnes), fishing mortality (F_{3-6}), and recruitment (age 1; thousands) from the final SAM assessment.




Figure 12.3.1.6. Plaice in Division 7.d. Summary of assessment results for spawning-stock biomass (SSB; tonnes), fishing mortality (F_{3-6} , ages 3–6), and recruitment (age 1; thousands) from the final SAM assessment.




Figure 12.3.1.7. Plaice in Division 7.d. Retrospective patterns for spawning-stock biomass (SSB; tonnes), fishing mortality (F_{3-6}), and recruitment (age 1; thousands) from the final SAM assessment.




Figure 12.3.1.8. Plaiice in Division 7.d. Estimated stock numbers-at-age.




Figure 12.5.1.1. Plaiice in Division 7.d. Number of individuals of age 1 as estimated by the assessment model (black), with the geometric mean over the period 1980–2024 (blue), the geometric mean over 2012–2023 (red).



Figure 12.5.2.1. Plaice in Division 7.d. ICES catches of resident plaice in 7.d (green line) and agreed TAC for 7.d and 7.e plaice (orange line). The orange dot corresponds to 7d 2025 TAC, the dark orange dot is the resident plaice in 7d of 2025 TAC, the green dot corresponds to the catch of the resident plaice 7d in the intermediate year 2025 (without considering the landing obligation) and the light green dot is the landings of resident plaice in 7d in 2024 taking into account the landing obligation in 2024 (ICES catches 2024 minus discards from exempted fleets).




Figure 12.5.2.2. Plaice in Division 7.d. The Q1 removal ratio over the period 2003–2024 to account for migration of mature individuals of plaice from the 7.e and 4.c during Q1.




Figure 12.6.1. Plaice in Division 7.d. The standard deviation of the log observations as estimated in SAM for the model approved during the benchmark (WKBNSCS 2025) (left) and the model revised during (WGNSSK 2025) (right).

Contents

12	Plaice in eastern English Channel	429
12.1	General.....	429
12.1.1	Stock definition	429
12.1.2	Ecosystem aspects	429
12.1.3	Fisheries	429
12.1.4	ICES Advice for previous years.....	429
12.1.5	Management.....	430
12.2	Data available	430
12.2.1	Catch	430
12.2.2	InterCatch	430
12.2.3	Age compositions.....	432
12.2.4	Weight-at-age	432
12.2.5	Maturity and natural mortality	432
12.2.6	Surveys	432
12.3	Assessment	433
12.3.1	Results.....	434
12.4	Biological reference points	434
12.5	Short-term forecasts	435
12.5.1	Recruitment estimates.....	436
12.5.2	Calculation of the 7.d resident stock	436
12.5.3	Management options tested	436
12.6	Quality of the assessment.....	436
12.7	Status of the stock	437
12.8	Management considerations	438
12.9	Issue for future benchmarks	438
12.9.1	Data.....	438
12.9.2	Assessment	438
12.9.3	Short-term forecast	438
12.10	References	438
12.11	Tables and figures	440