

# SchistoIBM

Matthew Malishev<sup>1\*</sup> and David J Civitello<sup>1</sup>

<sup>1</sup> *Department of Biology, Emory University, 1510 Clifton Road NE, Atlanta, GA, USA, 30322*

## Contents

Overview . . . . .	2
Environment . . . . .	3
Resources . . . . .	3
Miracidia . . . . .	3
Cercariae . . . . .	3
Individual . . . . .	4
Host . . . . .	4
Host length . . . . .	4
Host reserve . . . . .	4
Host development (maturity) . . . . .	4
Host reproduction . . . . .	4
Parasite . . . . .	4
Parasite biomass (growth) . . . . .	4
Parasite reproduction . . . . .	4
Damage density . . . . .	4
Cumulative hazard for host . . . . .	5
References . . . . .	6

Date: 2018-09-24

R version: 3.5.0

\*Corresponding author: [matthew.malishev@gmail.com](mailto:matthew.malishev@gmail.com)

DOI: <https://github.com/darwinanddavis/SchistoIBM>

Model from [Civitello et al 2018](#) Bioenergetic theory predicts infection dynamics of human schistosomes in intermediate host snails across ecological gradients

## Overview

Individual-based model of snail and schistosome population ecology driven by energetic consequences of host and parasites in response to resources and environmental temperature.

Population model of individual hosts (snails) and parasites (schistosomes) in a lake system. Host and parasite population sizes and production of different parasite life stages are driven by a population model. Parasites are released into the habitat according to reproduction rules of a full energy and mass budget model based on Dynamic Energy Budget (DEB) theory. Hosts feed on resource patches in the habitat and are exposed to schistosomes depending on the type of resource patch they feed on.

The host population is size and age structured, where parasite loading varies depending on the biomass of the host. Host biomass, including energy reserves, maintenance, and reproductive potential, is modelling through a DEB model.

The energetics model determines the intake and production of parasites. The simulation runs over a year and emulates a lake system in Kenya where schistosomes are found.

The energy reserves, somatic maintenance, and growth of hosts. The reproductive capacity of hosts is the parasite biomass produced.

## Environment

### Resources

$$\frac{dR}{dt} = r \left( 1 - \frac{R}{K} \right) R - \sum -i_M L^2 f_H$$

$\sum -i_M L^2 f_H$  = feeding by snails

### Miracidia

$$\frac{dM}{dt} = M(t) - \sum uptake - death$$

$M(t)$  = miracidia in environment

$\sum uptake$  = infection of hosts by miracidia. **Not a DEB parameter**

### Cercariae

$$\frac{dC}{dt} = \sum \frac{dRp}{dt} - death$$

$\sum \frac{dRp}{dt}$  = cumulative release of cercariae (parasite reproduction)

**Individual**

**Host**

**Host length**

$$\frac{dL}{dt} = \frac{gY_{VE}}{3\chi} \cdot \left( \frac{\kappa^* a_M e - (m_V + m_R E_M \delta) \chi L}{e + g} \right)$$

**Host reserve**

$$\frac{de}{dt} = (f_H - e) \frac{a_M}{\chi E_M L} - p \left( \frac{i_{PM} f_P}{E_M} \right)$$

**Host development (maturity)**

$$\frac{dD}{dt} = \begin{cases} \text{if } D < D_R, & (1 - \kappa^*)C - m_D D \\ \text{if } D \geq D_R, & 0 \end{cases}$$

**Host reproduction**

$$\frac{dR_H}{dt} = \begin{cases} \text{if } D < D_R, & 0 \\ \text{if } D \geq D_R, & (1 - \kappa^*)C - m_D D \end{cases}$$

**Parasite**

**Parasite biomass (growth)**

$$\frac{dP}{dt} = P(Y_{PE} i_{PM} f_P (1 - r_P) - m_P)$$

**Parasite reproduction**

$$\frac{dR_P}{dt} = \gamma_{RP} Y_{PE} i_{PM} f_P r_P$$

**Damage density**

$$\frac{d\delta}{dt} = \frac{\Theta}{\chi L^3} \cdot \frac{dR_P}{dt} + k_R (1 - e) - k_R \delta - \frac{3\delta}{L} \cdot \frac{dL}{dt}$$

$\frac{3\delta}{L} \cdot \frac{dL}{dt}$  = growth dilution. Growth of host reduces damage density  
 $\frac{\Theta}{\chi L^3}$  = damage intensity  
 $k_R$  = damage repair rate (via reserve depletion)

### Cumulative hazard for host

$$\frac{dH}{dt} = h_b + h_\delta \max(\delta - \delta_0, 0)$$

$\delta_0$  = linear damage density function

$h_b$  = background hazard rate

$h_\delta$  = hazard coefficient

## References

Civitello et al 2018 Bioenergetic theory predicts infection dynamics of human schistosomes in intermediate host snails across ecological gradients