

Matrix Forensics

*A brief guide to matrix math
and its efficient implementation*

RICHARD BARNES

Git Hash: 0ce356ce3f3bef1b495bfa9e508dfef86380157a
Compiled on: 2018/11/01 at 16:29:50

github.com/r-barnes/MatrixForensics

Contents

1	Introduction	4
2	Nomenclature	5
3	Basics	6
3.1	Fundamental Theorem of Linear Algebra	7
3.2	Matrix Properties	9
3.3	Matrix Multiplication	9
3.4	Time Complexities	10
4	Derivatives	11
4.1	Useful Rules for Derivatives	11
5	Matrix Rogue Gallery	12
5.1	Non-Singular vs. Singular Matrices	12
5.2	Diagonal Matrix	12
5.3	Dyads	13
5.4	Hermitian Matrix	13
5.5	Idempotent Matrix	14
5.6	Orthogonal Matrix	15
5.7	Permutation Matrix	16
5.8	Positive Definite	16
5.9	Positive Semi-Definite	16
5.10	Projection Matrix	17
5.11	Singular Matrix	17
5.12	Symmetric Matrix	17
5.13	Skew-Hermitian	18
5.14	Toeplitz Matrix, General Form	18
5.15	Toeplitz Matrix, Discrete Convolution	19
5.16	Triangular Matrix	19
5.17	Vandermonde Matrix	20
6	Matrix Decompositions	21
6.1	LLT/UTU: Cholesky Decomposition	21
6.2	LDLT	21
6.3	PCA: Principle Components Analysis	21
6.4	QR: Orthogonal-triangular	22
6.5	SVD: Singular Value Decomposition	22
6.6	Eigenvalue Decomposition for Diagonalizable Matrices	24
6.7	Eigenvalue (Spectral) Decomposition for Symmetric Matrices	24
6.8	Schur Complements	25
7	Transpose Properties	26
8	Determinant Properties	27
9	Trace Properties	28

<i>CONTENTS</i>	3
10 Inverse Properties	29
11 Pseudo-Inverse Properties	30
11.1 Moore-Penrose Pseudoinverse	30
12 Hadamard Identities	31
13 Eigenvalue Properties	32
14 Norms	33
14.1 Matrices	33
14.1.1 Frobenius norm	33
14.1.2 Operator Norms	33
14.1.3 Spectral Radius	34
14.2 Vectors	34
15 Bounds	35
15.1 Matrix Gain	35
15.2 Norms	35
15.3 Rayleigh quotients	35
16 Linear Equations	36
16.1 Least-Squares	36
16.1.1 Regularized least-squares with low-rank data	36
16.2 Minimum Norm Solutions	37
17 Updates	38
17.1 Removing a row from $\mathbf{A}^T \mathbf{A}$ ($\mathbf{A}^T \mathbf{A} \rightarrow \mathbf{A}_{\setminus i}^T \mathbf{A}_{\setminus i}$)	38
17.2 $\mathbf{1}_r^T \mathbf{A} \mathbf{1}_c$	38
17.3 $\mathbf{x}^T \mathbf{A} \mathbf{x}$	38
18 Optimization	40
18.1 Standard Forms	40
18.2 Transformations	41
18.2.1 Linear-Fractional to Linear	41
19 Algorithms	42
19.1 Gram-Schmidt	42

1 | Introduction

Goals: TODO

Contributing: Please contribute on Github at <https://github.com/r-barnes/MatrixForensics> either by opening an issue or making a pull request. If you are not comfortable with this, please send your contribution to rijard.barnes@gmail.com.

Contributors: Richard Barnes

Funding: TODO

2 | Nomenclature

\mathbf{A}	Matrix.
\mathbf{a}	(Column) vector.
a	Scalar.
\mathbf{A}_{ij}	Matrix indexed. Returns i th row and j th column.
$\mathbf{A} \circ \mathbf{B}$	Hadamard (element-wise) product of matrices \mathbf{A} and \mathbf{B} .
$\mathcal{N}(\mathbf{A})$	Nullspace of the matrix \mathbf{A} .
$\mathcal{R}(\mathbf{A})$	Range of the matrix \mathbf{A} .
$\det(\mathbf{A})$	Determinant of the matrix \mathbf{A} .
$\text{eig}(\mathbf{A})$	Eigenvalues of the matrix \mathbf{A} .
\mathbf{A}^H	Conjugate transpose of the matrix \mathbf{A} .
\mathbf{A}^T	Transpose of the matrix \mathbf{A} .
\mathbf{A}^+	Pseudoinverse of the matrix \mathbf{A} .
$\mathbf{x} \in \mathbb{R}^n$	The entries of the n -vector \mathbf{x} are all real numbers.
$\mathbf{A} \in \mathbb{R}^{m,n}$	The entries of the matrix \mathbf{A} with m rows and n columns are all real numbers.
$\mathbf{A} \in \mathbb{S}^n$	The matrix \mathbf{A} is symmetric and has n rows and n columns.
\mathbf{I}_n	Identity matrix with n rows and n columns.
$\{0\}$	The empty set

3 | Basics

3.1 Fundamental Theorem of Linear Algebra






Figure 3.4 The true action $Ax = A(x_{\text{row}} + x_{\text{null}})$ of any m by n matrix.



Matrix A converts n -tuples into m -tuples $\mathbb{R}^n \rightarrow \mathbb{R}^m$.
That is, linear transformation T_A is a map between rows and columns



Fundamental Subspaces


$C(A)$: Column space (image)
 $R(A)$: Row space (coimage)
 $N(A)$: Null space (kernel)
 $N(A^T)$: Left null space (cokernel)

Identities

$\dim(C) \equiv \text{rank}(A)$
 $\dim(N) \equiv \text{nullity}(A)$

Theorems

$\dim(C) + \dim(N) = n$
 $\dim(R) = \dim(C)$

Figure 1: Dimensions and orthogonality for any m by n matrix A of rank r .

3.2 Matrix Properties

$$\begin{array}{lll}
 \mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC} & \text{(left distributivity)} & (1) \\
 (\mathbf{B} + \mathbf{C})\mathbf{A} = \mathbf{BA} + \mathbf{CA} & \text{(right distributivity)} & (2) \\
 \mathbf{AB} \neq \mathbf{BA} & \text{(in general)} & (3) \\
 (\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC}) & \text{(associativity)} & (4)
 \end{array}$$

3.3 Matrix Multiplication

$$(\mathbf{AB})_{kl} = \sum_m \mathbf{A}_{km} \mathbf{B}_{ml} \quad \mathbf{A} \in \mathbb{R}^{k,m}, \mathbf{B} \in \mathbb{R}^{m,l} \quad (5)$$

3.4 Time Complexities

Operation	Input	Output	Algorithm	Time
Matmult	$A, B \in n \times n$	$n \times n$	Schoolbook	$O(n^3)$
			Strassen [1]	$O(n^{2.807})$
			Best	$O(n^\omega)$
Matmult	$A \in n \times m, B \in m \times p$	$n \times p$	Schoolbook	$O(nmp)$
Inversion	$A \in n \times n$	$n \times n$	Gauss–Jordan elimination	$O(n^3)$
			Strassen [1]	$O(n^{2.807})$
			Best	$O(n^\omega)$
SVD	$A \in m \times n$	$m \times m, m \times n, n \times n$ $m \times r, r \times r, n \times r$		$O(mn^2)$ ($m \geq n$)
Determinant	$A \in n \times n$	Scalar	Laplace expansion	$O(n!)$
			Division-free [2]	$O(n!)$
			LU decomposition	$O(n^3)$
			Integer preserving [3]	$O(n^3)$
Back substitution	A triangular	n solutions	Back substitution	$O(n^2)$

A comment on ω

The lower bound on matmult time complexity is $O(n^\omega)$, where ω is an unknown constant bounded by $2 \leq \omega \leq 2.373$. Algorithms achieving lower values of ω tend to be less efficient in practice for all but the largest matrices. Of the algorithm with times of less than $O(n^3)$, only the Strassen algorithm has seen serious attempts at optimized implementation. Most matmult implementations use highly optimized variants of the standard $O(n^3)$ algorithm. At this point, memory and bus speeds dominate the performance of implementations, so simple Big-O notation cannot be used to reliably compare matmult performances.

Name	Year	ω
Standard	-	3
Strassen [1]	1969	2.807
Pan [4]	1978	2.796
Bini et al. [5]	1979	2.78
Schönhage [6]	1981	2.548
Schönhage [6]	1981	2.522
Romani [7]	1982	2.517
Coppersmith and Winograd [8]	1982	2.496
Strassen [9]	1986	2.479
Coppersmith and Winograd [10]	1990	2.376
Williams [11]	2012	2.37294
Le Gall [12]	2014	2.3728639
Williams [11]	2012	2.3727

4 | Derivatives

4.1 Useful Rules for Derivatives

For general \mathbf{A} and \mathbf{X} (no special structure):

$$\partial \mathbf{A} = 0 \quad \text{where } \mathbf{A} \text{ is a constant} \quad (6)$$

$$\partial(c\mathbf{X}) = c\partial\mathbf{X} \quad (7)$$

$$\partial(\mathbf{X} + \mathbf{Y}) = \partial\mathbf{X} + \partial\mathbf{Y} \quad (8)$$

$$\partial(\text{tr}(\mathbf{X})) = \text{tr}(\partial\mathbf{X}) \quad (9)$$

$$\partial(\mathbf{XY}) = (\partial\mathbf{X})\mathbf{Y} + \mathbf{X}(\partial\mathbf{Y}) \quad (10)$$

$$\partial(\mathbf{X} \circ \mathbf{Y}) = (\partial\mathbf{X}) \circ \mathbf{Y} + \mathbf{X} \circ (\partial\mathbf{Y}) \quad (11)$$

$$\partial(\mathbf{X}^{-1}) = -\mathbf{X}^{-1}(\partial\mathbf{X})\mathbf{X}^{-1} \quad (12)$$

$$\partial(\det(\mathbf{X})) = \text{tr}(\text{adj}(\mathbf{X})\partial\mathbf{X}) \quad (13)$$

$$\partial(\det(\mathbf{X})) = \det(\mathbf{X}) \text{tr}(\mathbf{X}^{-1}\partial\mathbf{X}) \quad (14)$$

$$\partial(\ln(\det(\mathbf{X}))) = \text{tr}(\mathbf{X}^{-1}\partial\mathbf{X}) \quad (15)$$

$$\partial(\mathbf{X}^T) = (\partial\mathbf{X})^T \quad (16)$$

$$\partial(\mathbf{X}^H) = (\partial\mathbf{X})^H \quad (17)$$

5 | Matrix Rogue Gallery

5.1 Non-Singular vs. Singular Matrices

For $\mathbf{A} \in \mathbb{R}^{n,n}$ (initially drawn from [13, p. 574]):


Non-Singular

\mathbf{A} is invertible
 The columns are independent
 The rows are independent
 $\det(\mathbf{A}) \neq 0$
 $\mathbf{Ax} = 0$ has one solution: $\mathbf{x} = 0$
 $\mathbf{Ax} = \mathbf{b}$ has one solution: $\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$
 \mathbf{A} has n nonzero pivots
 \mathbf{A} has full rank $r = n$
 The reduced row echelon form is $\mathbf{R} = \mathbf{I}$
 The column space is all of \mathbb{R}^n
 The row space is all of \mathbb{R}^n
 All eigenvalues are nonzero
 $\mathbf{A}^T \mathbf{A}$ is symmetric positive definite
 \mathbf{A} has n positive singular values

Singular

\mathbf{A} is not invertible
 The columns are dependent
 The rows are dependent
 $\det(\mathbf{A}) = 0$
 $\mathbf{Ax} = 0$ has infinitely many solutions
 $\mathbf{Ax} = \mathbf{b}$ has either no or infinitely many solutions
 \mathbf{A} has $r < n$ pivots
 \mathbf{A} has rank $r < n$
 \mathbf{R} has at least one zero row
 The column space has dimension $r < n$
 The row space has dimension $r < n$
 Zero is an eigenvalue of \mathbf{A}
 $\mathbf{A}^T \mathbf{A}$ is only semidefinite
 \mathbf{A} has $r < n$ singular values

5.2 Diagonal Matrix



$$A = \text{diag}(a_1, \dots, a_n) = \begin{bmatrix} a_1 & & \\ & \ddots & \\ & & a_n \end{bmatrix} \quad (18)$$

Square matrix. Entries above diagonal are equal to entries below diagonal.

Number of “free entries”: $\frac{n(n+1)}{2}$.


Special Properties

$$\text{eig}(A) = a_1, \dots, a_n \quad (19)$$

$$\det(A) = \prod_i a_i \quad (20)$$

$$A^{-1} = \begin{bmatrix} \frac{1}{a_1} & & \\ & \ddots & \\ & & \frac{1}{a_n} \end{bmatrix} \quad (21)$$

5.3 Dyads



$\mathbf{A} \in \mathbb{R}^{m,n}$ is a dyad if it can be written as

$$\mathbf{A} = \mathbf{u}\mathbf{v}^T \quad \mathbf{u} \in \mathbb{R}^m, \mathbf{v} \in \mathbb{R}^n \quad (22)$$

Special Properties

- The columns of \mathbf{A} are copies of \mathbf{u} scaled by the values of \mathbf{v} .
- The rows of \mathbf{A} are copies of \mathbf{u}^T scaled by the values of \mathbf{v} .
- If \mathbf{A} is a dyad, it acts on a vector \mathbf{x} as $\mathbf{A}\mathbf{x} = (\mathbf{u}\mathbf{v}^T)\mathbf{x} = (\mathbf{v}^T\mathbf{u})\mathbf{x}$.
- $\mathbf{A}\mathbf{x} = c\mathbf{u}$ (\mathbf{A} scales \mathbf{x} and points it along \mathbf{u}).
- $\mathbf{A}_{ij} = \mathbf{u}_i\mathbf{v}_j$.
- If $\mathbf{u}, \mathbf{v} \neq 0$, then $\text{rank}(\mathbf{A}) = 1$.
- If $m = n$, \mathbf{A} has one eigenvalue $\lambda = \mathbf{v}^T\mathbf{u}$ and eigenvector \mathbf{u} .
- A dyad can always be written in a normalized form $c\tilde{\mathbf{u}}\tilde{\mathbf{v}}^T$.

5.4 Hermitian Matrix

$\mathbf{H} \in \mathbb{C}^{m,n}$ is Hermitian iff

$$\mathbf{H} = \mathbf{H}^H \quad (23)$$

where \mathbf{H}^H is the conjugate transpose of \mathbf{H} .

For $\mathbf{H} \in \mathbb{R}^{m,n}$, Hermitian and symmetric matrices are equivalent.

Special Properties

$$\mathbf{H}_{ii} \in \mathbb{R} \quad (24)$$

$$\mathbf{H}\mathbf{H}^H = \mathbf{H}^H\mathbf{H} \quad (25)$$

$$\mathbf{x}^H\mathbf{H}\mathbf{x} \in \mathbb{R} \quad \forall \mathbf{x} \in \mathbb{C} \quad (26)$$

$$\mathbf{H}_1 + \mathbf{H}_2 = \text{Hermitian} \quad (27)$$

$$\mathbf{H}^{-1} = \text{Hermitian} \quad (28)$$

$$\mathbf{A} + \mathbf{A}^H = \text{Hermitian} \quad (29)$$

$$\mathbf{A} - \mathbf{A}^H = \text{Skew-Hermitian} \quad (30)$$

$$\mathbf{A}\mathbf{B} = \text{Hermitian iff } \mathbf{A}\mathbf{B} = \mathbf{B}\mathbf{A} \quad (31)$$

$$\det(\mathbf{H}) \in \mathbb{R} \quad (32)$$

$$\text{eig}(\mathbf{H}) \in \mathbb{R} \quad (33)$$

5.5 Idempotent Matrix

A matrix \mathbf{A} is idempotent iff

$$\mathbf{A}\mathbf{A} = \mathbf{A} \quad (34)$$

Special Properties

$$\mathbf{A}^n = \mathbf{A} \quad \forall n \quad (35)$$

$$\mathbf{I} - \mathbf{A} \text{ is idempotent} \quad (36)$$

$$\mathbf{A}^H \text{ is idempotent} \quad (37)$$

$$\mathbf{I} - \mathbf{A}^H \text{ is idempotent} \quad (38)$$

$$\text{rank}(\mathbf{A}) = \text{tr}(\mathbf{A}) \quad (39)$$

$$\mathbf{A}(\mathbf{I} - \mathbf{A}) = 0 \quad (40)$$

$$\mathbf{A}^+ = \mathbf{A} \quad (41)$$

$$f(s\mathbf{I} + t\mathbf{A}) = (\mathbf{I} - \mathbf{A})f(s) + \mathbf{A}f(s + t) \quad (42)$$


$$\mathbf{A}\mathbf{B} = \mathbf{B}\mathbf{A} \implies \mathbf{A}\mathbf{B} \text{ is idempotent} \quad (43)$$

$$\text{eig}(\mathbf{A})_i \in \{0, 1\} \quad (44)$$

$$\mathbf{A} \text{ is always diagonalizable} \quad (45)$$

$\mathbf{A} - \mathbf{I}$ may not be idempotent.

5.6 Orthogonal Matrix



(Not much visible structure)

$$U = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix} \quad (46)$$

A matrix \mathbf{U} is orthogonal iff:

$$\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I} \quad (47)$$

Square matrix. The columns form an orthonormal basis of \mathbb{R}^n .

Special Properties

- The eigenvalues of \mathbf{U} are placed on the unit circle.
- The eigenvectors of \mathbf{U} are unitary (have length one).
- \mathbf{U}^{-1} is orthogonal.

$$\mathbf{U}^T = \mathbf{U}^{-1} \quad (48)$$

$$\mathbf{U}^{-T} = \mathbf{U} \quad (49)$$

$$\mathbf{U}^T \mathbf{U} = \mathbf{I} \quad (50)$$

$$\mathbf{U} \mathbf{U}^T = \mathbf{I} \quad (51)$$


$$\det(\mathbf{U}) = \pm 1 \quad (52)$$

Orthogonal matrices preserve the lengths and angles of the vectors they operator on. The converse is true: any matrix which preserves lengths and angles is orthogonal.

$$\|\mathbf{U}\mathbf{x}\|_2^2 = (\mathbf{U}\mathbf{x})^T (\mathbf{U}\mathbf{x}) = \mathbf{x}^T \mathbf{U}^T \mathbf{U} \mathbf{x} = \mathbf{x}^T \mathbf{x} = \|\mathbf{x}\|_2^2 \quad \forall \mathbf{x} \quad (53)$$

$$\|\mathbf{U}\mathbf{A}\mathbf{V}\|_F = \|\mathbf{A}\|_F \quad \forall \mathbf{A}, \mathbf{U}, \mathbf{V} \text{ with } U, V \text{ orthogonal} \quad (54)$$

5.7 Permutation Matrix



TODO

5.8 Positive Definite

$\mathbf{A} \in \mathbb{S}^n$ is positive definite (denoted $\mathbf{A} \succ 0$) if any of the following are true:

- $\mathbf{x}^T \mathbf{A} \mathbf{x} > 0, \forall \mathbf{x} \in \mathbb{R}^n$.
- $\text{eig}(\mathbf{A}) > 0$

Special Properties

- If \mathbf{A} is PD and invertible, \mathbf{A}^{-1} is also PD.
- If \mathbf{A} is PD and $c \in \mathbb{R}$ then $c\mathbf{A}$ is PD.
- The diagonal entries \mathbf{A}_{ii} are real and non-negative, so $\text{tr}(\mathbf{A}) \geq 0$.
- $\det(\mathbf{A}) > 0$
- For $\mathbf{A} \in \mathbb{R}^{m,n}$, $\mathbf{A}^T \mathbf{A} \succ 0 \iff \mathbf{A}$ is full-column rank ($\text{rank}(\mathbf{A}) = n$)
- For $\mathbf{A} \in \mathbb{R}^{m,n}$, $\mathbf{A} \mathbf{A}^T \succ 0 \iff \mathbf{A}$ is full-row rank ($\text{rank}(\mathbf{A}) = m$)
- $\mathbf{P} \succ 0$ defines a full-dimensional, bounded ellipsoid centered at the origin and defined by the set $\mathcal{E} = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{x}^T \mathbf{P}^{-1} \mathbf{x} \leq 1\}$. The eigenvalues λ_i and eigenvectors u_i of \mathbf{P} define the orientation and shape of the ellipsoid. u_i are the semi-axes while the lengths of the semi-axes are given by $\sqrt{\lambda_i}$. Using the Cholesky decomposition, $\mathbf{P}^{-1} = \mathbf{A}^T \mathbf{A}$, an equivalent definition of the ellipsoid is $\mathcal{E} = \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{A} \mathbf{x}\|_2 \leq 1\}$.

5.9 Positive Semi-Definite

\mathbf{A} is positive semi-definite (denoted $\mathbf{A} \succeq 0$) if any of the following are true:

- $\mathbf{x}^T \mathbf{A} \mathbf{x} \geq 0, \forall \mathbf{x} \in \mathbb{R}^n$.
- $\text{eig}(\mathbf{A}) \geq 0$

Special Properties

- For $\mathbf{A} \in \mathbb{R}^{m,n}$, $\mathbf{A}^T \mathbf{A} \succeq 0$
- For $\mathbf{A} \in \mathbb{R}^{m,n}$, $\mathbf{A} \mathbf{A}^T \succeq 0$
- The positive semi-definite matrices \mathbb{S}_+^n form a convex cone. For any two PSD matrices $\mathbf{A}, \mathbf{B} \in \mathbb{S}_+^n$ and some $\alpha \in [0, 1]$:

$$\mathbf{x}^T (\alpha \mathbf{A} + (1 - \alpha) \mathbf{B}) \mathbf{x} = \alpha \mathbf{x}^T \mathbf{A} \mathbf{x} + (1 - \alpha) \mathbf{x}^T \mathbf{B} \mathbf{x} \geq 0 \quad \forall \mathbf{x} \quad (55)$$

$$\alpha \mathbf{A} + (1 - \alpha) \mathbf{B} \in \mathbb{S}_+^n \quad (56)$$

- For $\mathbf{A} \in \mathbb{S}_+^n$ and $\alpha \geq 0$, $\alpha \mathbf{A} \succeq 0$, so \mathbb{S}_+^n is a cone.

5.10 Projection Matrix

A square matrix \mathbf{P} is a projection matrix that projects onto a vector space \mathcal{S} iff

$$\mathbf{P} \text{ is idempotent} \quad (57)$$

$$\mathbf{P} \mathbf{x} \in \mathcal{S} \quad \forall \mathbf{x} \quad (58)$$


$$\mathbf{P} \mathbf{z} = \mathbf{z} \quad \forall \mathbf{z} \in \mathcal{S} \quad (59)$$

5.11 Singular Matrix

A square matrix that is not invertible.

$\mathbf{A} \in \mathbb{R}^{n,n}$ is singular iff $\det \mathbf{A} = 0$ iff $\mathcal{N}(\mathbf{A}) \neq \{0\}$.

5.12 Symmetric Matrix



$\mathbf{A} \in \mathbb{S}^n$ is a symmetric matrix if $\mathbf{A} = \mathbf{A}^T$ (entries above diagonal are equal to entries below diagonal).

$$\begin{bmatrix} a & b & c & d & e & f \\ b & g & l & m & o & p \\ c & l & h & n & q & r \\ d & m & n & i & s & t \\ e & o & q & s & j & u \\ f & p & r & t & u & k \end{bmatrix} \quad (60)$$

Special Properties

$$\mathbf{A} = \mathbf{A}^T \quad (61)$$

$$\text{eig}(A) \in \mathbb{R}^n \quad (62)$$

Number of “free entries”: $\frac{n(n+1)}{2}$.

If \mathbf{A} is real, it can be decomposed into $\mathbf{A} = \mathbf{Q}^T \mathbf{D} \mathbf{Q}$ where \mathbf{Q} is a real orthogonal matrix (the columns of which are eigenvectors of \mathbf{A}) and \mathbf{D} is real and diagonal containing the eigenvalues of \mathbf{A} .


For a real, symmetric matrix with non-negative eigenvalues, the eigenvalues and singular values coincide.

5.13 Skew-Hermitian

A matrix $\mathbf{H} \in \mathbb{C}^{m,n}$ is Skew-Hermitian iff

$$\mathbf{H} = -\mathbf{H}^H \quad (63)$$


5.14 Toeplitz Matrix, General Form



Constant values on descending diagonals.

$$\begin{bmatrix} a_0 & a_{-1} & a_{-2} & \dots & \dots & a_{-(n-1)} \\ a_1 & a_0 & a_{-1} & \ddots & & \vdots \\ a_2 & a_1 & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & a_{-1} & a_{-2} \\ \vdots & & \ddots & a_1 & a_0 & a_{-1} \\ a_{n-1} & \dots & \dots & a_2 & a_1 & a_0 \end{bmatrix} \quad (64)$$


5.15 Toeplitz Matrix, Discrete Convolution



Constant values on main and subdiagonals.

$$\begin{bmatrix}
 h_m & 0 & 0 & \dots & 0 & 0 \\
 \vdots & h_m & 0 & \dots & 0 & 0 \\
 h_1 & \vdots & h_m & \dots & 0 & 0 \\
 0 & h_1 & \ddots & \ddots & 0 & 0 \\
 0 & 0 & h_1 & \ddots & h_m & 0 \\
 0 & 0 & 0 & \ddots & \vdots & h_m \\
 0 & 0 & 0 & \dots & h_1 & \vdots \\
 0 & 0 & 0 & \dots & 0 & h_1
 \end{bmatrix} \tag{65}$$

5.16 Triangular Matrix



$$\begin{bmatrix} a & b & c & d & e & f \\ & g & h & i & j & k \\ & & l & m & n & o \\ & & & p & q & r \\ & & & & s & t \\ & & & & & u \end{bmatrix} \quad \begin{bmatrix} a & & & & & \\ b & g & & & & \\ c & h & l & & & \\ d & i & m & p & & \\ e & j & n & q & s & \\ f & k & o & r & t & u \end{bmatrix} \quad (66)$$

Square matrices in which all elements either above or below the main diagonal are zero. An upper (left) and a lower (right) triangular matrix are shown above.

For an upper triangular matrix $A_{ij} = 0$ whenever $i > j$; for a lower triangular matrix $A_{ij} = 0$ whenever $i < j$.

Special Properties

$$\text{eig}(A) = \text{diag}(A) \quad (67)$$

$$\det(A) = \prod_i \text{diag}(A)_i \quad (68)$$

The product of two upper (lower) triangular matrices is still upper (lower) triangular.

The inverse of a nonsingular upper (lower) triangular matrix is still upper (lower) triangular.

5.17 Vandermonde Matrix

$$V = \begin{bmatrix} 1 & \alpha_1 & \alpha_1^2 & \dots & \alpha_1^{n-1} \\ 1 & \alpha_2 & \alpha_2^2 & \dots & \alpha_2^{n-1} \\ 1 & \alpha_3 & \alpha_3^2 & \dots & \alpha_3^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \alpha_m & \alpha_m^2 & \dots & \alpha_m^{n-1} \end{bmatrix} \quad (69)$$

Alternatively,

$$V_{i,j} = \alpha_i^{j-1} \quad (70)$$

Uses


Polynomial interpolation of data.

Special Properties

- $\det(V) = \prod_{1 \leq i < j \leq n} (x_j - x_i)$

6 | Matrix Decompositions

6.1 LLT/UTU: Cholesky Decomposition




If \mathbf{A} is symmetric, positive definite, square, then

$$\mathbf{A} = \mathbf{U}^T \mathbf{U} = \mathbf{L} \mathbf{L}^T \quad (71)$$

where \mathbf{U} is a unique upper triangular matrix and \mathbf{L} is a unique lower-triangular matrix.

6.2 LDLT



If \mathbf{A} is a non-singular symmetric definite square matrix, then

$$\mathbf{A} = \mathbf{L} \mathbf{D} \mathbf{L}^T = \mathbf{L}^T \mathbf{D} \mathbf{L} \quad (72)$$

where \mathbf{L} is a unit lower triangular matrix and \mathbf{D} is a diagonal matrix. If $\mathbf{A} \succ 0$, then $\mathbf{D}_{ii} > 0$.


6.3 PCA: Principle Components Analysis

Find normalized directions in data space such that the variance of the projections of the centered data points is maximal. For centered data $\tilde{\mathbf{X}}$, the mean-square variation of data along a vector \mathbf{x} is $\mathbf{x}^T \tilde{\mathbf{X}} \tilde{\mathbf{X}}^T \mathbf{x}$.

$$\max_{\mathbf{x} \in \mathbb{R}^n, \|\mathbf{x}\|_2=1} \mathbf{x}^T \tilde{\mathbf{X}} \tilde{\mathbf{X}}^T \mathbf{x} \quad (73)$$

Taking an SVD of $\tilde{\mathbf{X}} \tilde{\mathbf{X}}^T$ gives $H = \mathbf{U}_r \mathbf{D}^2 \mathbf{U}^T$, which is maximized by taking $\mathbf{x} = \mathbf{u}_1$. By repeatedly removing the first principal components and recalculating, all the principal axes can be found.

6.4 QR: Orthogonal-triangular




For $\mathbf{A} \in \mathbb{R}^{n,n}$, $\mathbf{A} = \mathbf{QR}$ where \mathbf{Q} is orthogonal and \mathbf{R} is an upper triangular matrix. If \mathbf{A} is non-singular, then \mathbf{Q} and \mathbf{R} are uniquely defined if $\text{diag}(\mathbf{R})$ are imposed to be positive.

Algorithms

Gram-Schmidt.

6.5 SVD: Singular Value Decomposition



Any matrix $\mathbf{A} \in \mathbb{R}^{m,n}$ can be written as

$$\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T = \sum_{i=1}^r \sigma_i u_i v_i^T \quad (74)$$

where

$$U = \text{eigenvectors of } \mathbf{A}\mathbf{A}^T \quad \mathbb{R}^{m,m} \quad (75)$$

$$D = \text{diag}(\sigma_i) = \sqrt{\text{diag}(\text{eig}(\mathbf{A}\mathbf{A}^T))} \quad \mathbb{R}^{n,n} \quad (76)$$

$$V = \text{eigenvectors of } \mathbf{A}^T \mathbf{A} \quad \mathbb{R}^{n,n} \quad (77)$$

Let σ_i be the non-zero singular values for $i = 1, \dots, r$ where r is the rank of \mathbf{A} ; $\sigma_1 \geq \dots \geq \sigma_r$.

We also have that

$$\mathbf{A}\mathbf{v}_i = \sigma_i \mathbf{u}_i \quad (78)$$

$$\mathbf{A}^T \mathbf{u}_i = \sigma_i \mathbf{v}_i \quad (79)$$

$$\mathbf{U}^T \mathbf{U} = \mathbf{I} \quad (80)$$

$$\mathbf{V}^T \mathbf{V} = \mathbf{I} \quad (81)$$

\mathbf{D} can be written in an expanded form:

$$\tilde{\mathbf{D}} = \begin{bmatrix} \mathbf{D} & 0_{r,n-r} \\ 0_{m-r,r} & 0_{m-r,n-r} \end{bmatrix} \quad (82)$$

The final $n - r$ columns of \mathbf{V} give an orthonormal basis spanning $\mathcal{N}(\mathbf{A})$. An orthonormal basis spanning the range of \mathbf{A} is given by the first r columns of \mathbf{U} .

$$\|\mathbf{A}\|_F^2 = \text{Frobenius norm} = \text{tr}(\mathbf{A}^T \mathbf{A}) = \sum_{i=1}^r \sigma_i^2 \quad (83)$$

$$\|\mathbf{A}\|_2^2 = \sigma_1^2 \quad (84)$$

$$\|\mathbf{A}\|_* = \text{nuclear norm} = \sum_{i=1}^r \sigma_i \quad (85)$$

The **condition number** κ of an invertible matrix $\mathbf{A} \in \mathbb{R}^{n,n}$ is the ratio of the largest and smallest singular value. Matrices with large condition numbers are closer to being singular and more sensitive to changes.

$$\kappa(\mathbf{A}) = \frac{\sigma_1}{\sigma_n} = \|\mathbf{A}\|_2 \cdot \|\mathbf{A}^{-1}\|_2 \quad (86)$$

Low-Rank Approximation

Approximating $\mathbf{A} \in \mathbb{R}^{m,n}$ by a matrix \mathbf{A}_k of rank $k > 0$ can be formulated as the optimization problem

$$\min_{\mathbf{A}_k \in \mathbb{R}^{m,n}} \|\mathbf{A} - \mathbf{A}_k\|_F^2 : \text{rank } \mathbf{A}_k = k, 1 \leq k \leq \text{rank}(\mathbf{A}) \quad (87)$$

The optimal solution of this problem is given by

$$\mathbf{A}_k = \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T \quad (88)$$

where

$$\frac{\|\mathbf{A}_k\|_F^2}{\|\mathbf{A}\|_F^2} = \frac{\sigma_1^2 + \dots + \sigma_k^2}{\sigma_1^2 + \dots + \sigma_r^2} \quad (89)$$

$$1 - \frac{\|\mathbf{A}_k\|_F^2}{\|\mathbf{A}\|_F^2} = \frac{\sigma_{k+1}^2 + \dots + \sigma_r^2}{\sigma_1^2 + \dots + \sigma_r^2} \quad (90)$$

is the fraction of the total variance in \mathbf{A} explained by the approximation \mathbf{A}_k .

Range and Nullspace

$$\mathcal{N}(\mathbf{A}) = \mathcal{R}(\mathbf{V}_{nr}) \quad (91)$$

$$\mathcal{N}(\mathbf{A})^\perp \equiv \mathcal{R}(\mathbf{A}^T) = \mathcal{R}(\mathbf{V}_r) \quad (92)$$

$$\mathcal{R}(\mathbf{A}) = \mathcal{R}(\mathbf{U}_r) \quad (93)$$

$$\mathcal{R}(\mathbf{A})^\perp \equiv \mathcal{N}(\mathbf{A}^T) = \mathcal{R}(\mathbf{U}_{nr}) \quad (94)$$

where \mathbf{V}_r is the first r columns of \mathbf{V} and \mathbf{V}_{nr} are the last $[r+1, n]$ columns; similarly for \mathbf{U} .

Projectors

The projection of \mathbf{x} onto $\mathcal{N}(\mathbf{A})$ is $(\mathbf{V}_{nr}\mathbf{V}_{nr}^T)\mathbf{x}$. Since $\mathbf{I}_n = \mathbf{V}_r\mathbf{V}_r^T + \mathbf{V}_{nr}\mathbf{V}_{nr}^T$, $(\mathbf{I}_n - \mathbf{V}_r\mathbf{V}_r^T)\mathbf{x}$ also works. The projection of \mathbf{x} onto $\mathcal{R}(\mathbf{A})$ is $(\mathbf{U}_r\mathbf{U}_r^T)\mathbf{x}$.

If $\mathbf{A} \in \mathbb{R}^{m,n}$ is full row rank ($\mathbf{A}\mathbf{A}^T \succ 0$), then the minimum distance to an affine set $\{x : \mathbf{A}\mathbf{x} = \mathbf{b}\}$, $\mathbf{b} \in \mathbb{R}^m$ is given by $\mathbf{x}^* = \mathbf{A}^T(\mathbf{A}\mathbf{A}^T)^{-1}\mathbf{b}$.

If $\mathbf{A} \in \mathbb{R}^{m,n}$ is full column rank ($\mathbf{A}^T\mathbf{A} \succ 0$), then the minimum distance to an affine set $\{x : \mathbf{A}\mathbf{x} = \mathbf{b}\}$, $\mathbf{b} \in \mathbb{R}^m$ is given by $\mathbf{x}^* = \mathbf{A}(\mathbf{A}^T\mathbf{A})^{-1}\mathbf{A}^T\mathbf{b}$.

Computational Notes

Since $\sigma \approx 0$, a *numerical rank* can be estimated for the matrix as the largest k such that $\sigma_k > \epsilon\sigma_1$ for $\epsilon \geq 0$.

6.6 Eigenvalue Decomposition for Diagonalizable Matrices

For a square, diagonalizable matrix $\mathbf{A} \in \mathbb{R}^{n,n}$

$$\mathbf{A} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^{-1} \quad (95)$$

where $\mathbf{U} \in \mathbb{C}^{n,n}$ is an invertible matrix whose columns are the eigenvectors of \mathbf{A} and $\mathbf{\Lambda}$ is a diagonal matrix containing the eigenvalues $\lambda_1, \dots, \lambda_n$ of \mathbf{A} in the diagonal.

The columns $\mathbf{u}_1, \dots, \mathbf{u}_n$ satisfy

$$\mathbf{A}\mathbf{u}_i = \lambda_i\mathbf{u}_i \quad i = 1, \dots, n \quad (96)$$

6.7 Eigenvalue (Spectral) Decomposition for Symmetric Matrices

A symmetric matrix $\mathbf{A} \in \mathbb{R}^{n,n}$ can be factored as

$$\mathbf{A} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T = \sum_i^n \lambda_i \mathbf{u}_i \mathbf{u}_i^T \quad (97)$$

where $\mathbf{U} \in \mathbb{R}^{n,n}$ is an orthogonal matrix whose columns \mathbf{u}_i are the eigenvectors of \mathbf{A} and $\mathbf{\Lambda}$ is a diagonal matrix containing the eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$ of \mathbf{A} in the diagonal. These eigenvalues are always real. The eigenvectors can always be chosen to be real and to form an orthonormal basis.

The columns $\mathbf{u}_1, \dots, \mathbf{u}_n$ satisfy

$$\mathbf{A}\mathbf{u}_i = \lambda_i \mathbf{u}_i \quad i = 1, \dots, n \quad (98)$$

6.8 Schur Complements

For $\mathbf{A} \in \mathbb{S}^n$, $\mathbf{B} \in \mathbb{S}^n$, $\mathbf{X} \in \mathbb{R}^{n,m}$ with $\mathbf{B} \succ 0$ and the block matrix

$$\mathbf{M} = \begin{bmatrix} \mathbf{A} & \mathbf{X} \\ \mathbf{X}^T & \mathbf{B} \end{bmatrix} \quad (99)$$

and the Schur complement of \mathbf{A} in \mathbf{M}

$$S = \mathbf{A} - \mathbf{X}\mathbf{B}^{-1}\mathbf{X}^T \quad (100)$$

Then

$$\mathbf{M} \succeq 0 \iff S \succeq 0 \quad (101)$$

$$\mathbf{M} \succ 0 \iff S \succ 0 \quad (102)$$

7 | Transpose Properties

$$(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T \tag{103}$$

$$(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T \tag{104}$$

$$(\mathbf{A}^{-1})^T = (\mathbf{A}^T)^{-1} \tag{105}$$

8 | Determinant Properties

Geometrically, if a unit volume is acted on by \mathbf{A} , then $|\det(\mathbf{A})|$ indicates the volume after the transformation.

$$\det(I_n) = 1 \quad (106)$$

$$\det(\mathbf{A}^T) = \det(\mathbf{A}) \quad (107)$$

$$\det(\mathbf{A}^{-1}) = \frac{1}{\det(\mathbf{A})} = \det(\mathbf{A})^{-1} \quad (108)$$

$$\det(AB) = \det(BA) \quad (109)$$

$$\det(AB) = \det(A) \det(B) \quad \mathbf{A}, \mathbf{B} \in \mathbb{R}^{n,n} \quad (110)$$

$$\det(c\mathbf{A}) = c^n \det(\mathbf{A}) \quad \mathbf{A} \in \mathbb{R}^{n,n} \quad (111)$$

$$\det(\mathbf{A}) = \prod \text{eig}(\mathbf{A}) \quad (112)$$

9 | Trace Properties

For $\mathbf{A} \in \mathbb{R}^{n,n}$

$$\text{tr}(\mathbf{A}) = \sum_{i=1}^n \mathbf{A}_{ii} \quad (113)$$

$$\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}(\mathbf{A}) + \text{tr}(\mathbf{B}) \quad (114)$$

$$\text{tr}(c\mathbf{A}) = c \text{tr}(\mathbf{A}) \quad (115)$$

$$\text{tr}(\mathbf{A}) = \text{tr}(\mathbf{A}^T) \quad (116)$$

For $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$ of compatible dimensions,

$$\text{tr}(\mathbf{A}^T \mathbf{B}) = \text{tr}(\mathbf{A} \mathbf{B}^T) = \text{tr}(\mathbf{B}^T \mathbf{A}) = \text{tr}(\mathbf{B} \mathbf{A}^T) \quad (117)$$

$$\text{tr}(\mathbf{ABCD}) = \text{tr}(\mathbf{BCDA}) = \text{tr}(\mathbf{CDAB}) = \text{tr}(\mathbf{DABC}) \quad (118)$$

(Invariant under cyclic permutations)

10 | Inverse Properties

The inverse of $\mathbf{A} \in \mathbb{C}^{n,n}$ is denoted \mathbf{A}^{-1} and defined such that

$$\mathbf{A}\mathbf{A}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}_n \quad (119)$$

where \mathbf{I}_n is the $n \times n$ identity matrix. \mathbf{A} is nonsingular if \mathbf{A}^{-1} exists; otherwise, \mathbf{A} is singular.

If individual inverses exist

$$(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1} \quad (120)$$

more generally

$$(\mathbf{A}_1\mathbf{A}_2 \dots \mathbf{A}_n)^{-1} = \mathbf{A}_n^{-1} \dots \mathbf{A}_2^{-1}\mathbf{A}_1^{-1} \quad (121)$$

$$(\mathbf{A}^{-1})^T = (\mathbf{A}^T)^{-1} \quad (122)$$

11 | Pseudo-Inverse Properties

For $\mathbf{A} \in \mathbb{R}^{m,n}$, a pseudoinverse satisfies:

$$\mathbf{A}\mathbf{A}^+\mathbf{A} = \mathbf{A} \quad (123)$$

$$\mathbf{A}^+\mathbf{A}\mathbf{A}^+ = \mathbf{A}^+ \quad (124)$$

$$(\mathbf{A}\mathbf{A}^+)^T = \mathbf{A}\mathbf{A}^+ \quad (125)$$

$$(\mathbf{A}^+\mathbf{A})^T = \mathbf{A}^+\mathbf{A} \quad (126)$$

11.1 Moore-Penrose Pseudoinverse

$$\mathbf{A}^+ = \mathbf{V}\mathbf{D}^{-1}\mathbf{U}^T \quad (127)$$

where the foregoing comes from a singular-value decomposition and $\mathbf{D}^{-1} = \text{diag}(\frac{1}{\sigma_1}, \dots, \frac{1}{\sigma_r})$

Special Properties

- $\mathbf{A}^+ = \mathbf{A}^{-1}$ if $\mathbf{A} \in \mathbb{R}^{n,n}$ and \mathbf{A} is square and nonsingular.
- $\mathbf{A}^+ = (\mathbf{A}^T\mathbf{A})^{-1}\mathbf{A}^T$, if $\mathbf{A} \in \mathbb{R}^{m,n}$ is full column rank ($r = n \leq m$). \mathbf{A}^+ is a left inverse of \mathbf{A} , so $\mathbf{A}^+\mathbf{A} = \mathbf{V}_r\mathbf{V}_r^T = \mathbf{V}\mathbf{V}^T = \mathbf{I}_n$.
- $\mathbf{A}^+ = \mathbf{A}^T(\mathbf{A}\mathbf{A}^T)^{-1}$, if $\mathbf{A} \in \mathbb{R}^{m,n}$ is full row rank ($r = m \leq n$). \mathbf{A}^+ is a right inverse of \mathbf{A} , so $\mathbf{A}\mathbf{A}^+ = \mathbf{U}_r\mathbf{U}_r^T = \mathbf{U}\mathbf{U}^T = \mathbf{I}_m$.

12 | Hadamard Identities

$$(\mathbf{A} \circ \mathbf{B})_{ij} = A_{ij}B_{ij} \quad \forall i, j \quad (128)$$

$$\mathbf{A} \circ \mathbf{B} = \mathbf{B} \circ \mathbf{A} \quad (129) \quad [14]$$

$$\mathbf{A} \circ (\mathbf{B} \circ \mathbf{C}) = (\mathbf{A} \circ \mathbf{B}) \circ \mathbf{C} \quad (130)$$

$$\mathbf{A} \circ (\mathbf{B} + \mathbf{C}) = \mathbf{A} \circ \mathbf{B} + \mathbf{A} \circ \mathbf{C} \quad (131) \quad [14]$$

$$a(\mathbf{A} \circ \mathbf{B}) = (a\mathbf{A}) \circ \mathbf{B} = \mathbf{A} \circ (a\mathbf{B}) \quad (132) \quad [14]$$

$$(\mathbf{A}^T \circ \mathbf{B}^T) = (\mathbf{A} \circ \mathbf{B})^T \quad (133)$$

$$(\mathbf{A}^T \circ \mathbf{B}^T) = (\mathbf{A} \circ \mathbf{B})^T \quad (134)$$

$$(\mathbf{x}^T \mathbf{A} \mathbf{x}) = \sum_{i,j} ((\mathbf{x} \mathbf{x}^T) \circ \mathbf{A}) \quad (135)$$

$$\mathbf{x}^T (\mathbf{A} \circ \mathbf{B}) \mathbf{y} = \text{tr}((\text{diag}(\mathbf{x}) \mathbf{A})^T \mathbf{B} \text{diag}(\mathbf{y})) \quad \mathbf{A}, \mathbf{B} \in \mathbb{R}^{m,n} \quad (136) \quad [15]$$

$$\text{tr}(\mathbf{A}^T \mathbf{B}) = \mathbf{1}^T (\mathbf{A} \circ \mathbf{B}) \mathbf{1} \quad (137)$$

13 | Eigenvalue Properties

$\lambda \in \mathbb{C}$ is an eigenvalue of $\mathbf{A} \in \mathbb{R}^{n,n}$ and $u \in \mathbb{C}^n$ is a corresponding eigenvector if $\mathbf{A}\mathbf{u} = \lambda\mathbf{u}$ and $\mathbf{u} \neq 0$. Equivalently, $(\lambda\mathbf{I}_n - \mathbf{A})\mathbf{u} = 0$ and $\mathbf{u} \neq 0$. Eigenvalues satisfy the equation $\det(\lambda\mathbf{I}_n - \mathbf{A}) = 0$.

Any matrix $\mathbf{A} \in \mathbb{R}^{n,n}$ has n eigenvalues, though some may be repeated. λ_1 is the largest eigenvalue and λ_n the smallest.

$$\text{eig}(\mathbf{A}\mathbf{A}^T) = \text{eig}(\mathbf{A}^T\mathbf{A}) \quad (138)$$

(Note that the number of entries in $\mathbf{A}\mathbf{A}^T$ and $\mathbf{A}^T\mathbf{A}$ may differ significantly leading to different compute times.)

$$\text{eig}(\mathbf{A}^T\mathbf{A}) \geq 0 \quad (139)$$

Computation

TODO: eigsh, small eigen value extraction, top-k

14 | Norms

14.1 Matrices

Matrix norms satisfy some properties:

$$f(\mathbf{A}) \geq 0 \quad (140)$$

$$f(\mathbf{A}) = 0 \iff \mathbf{A} = 0 \quad (141)$$

$$f(c\mathbf{A}) = |c|f(\mathbf{A}) \quad (142)$$

$$f(\mathbf{A} + \mathbf{B}) \leq f(\mathbf{A}) + f(\mathbf{B}) \quad (143)$$

Many popular matrix norms also satisfy “sub-multiplicativity”: $f(\mathbf{AB}) \leq f(\mathbf{A})f(\mathbf{B})$.

14.1.1 Frobenius norm

$$\|\mathbf{A}\|_F = \sqrt{\text{tr } \mathbf{A}\mathbf{A}^H} \quad (144)$$

$$= \sqrt{\sum_{i=1}^m \sum_{j=1}^n |\mathbf{A}_{ij}|^2} \quad (145)$$

$$= \sqrt{\sum_{i=1}^m \text{eig}(\mathbf{A}^H \mathbf{A})_i} \quad (146)$$

Special Properties

$$\|\mathbf{Ax}\|_2 \leq \|\mathbf{A}\|_F \|\mathbf{x}\|_2 \quad \mathbf{x} \in \mathbb{R}^n \quad (147)$$

$$\|\mathbf{AB}\|_F \leq \|\mathbf{A}\|_F \|\mathbf{B}\|_F \quad (148)$$

$$\left\| \mathbf{C} - \mathbf{xx}^T \right\|_F^2 = \|\mathbf{C}\|_F^2 + \|\mathbf{x}\|_2^4 - 2\mathbf{x}^T \mathbf{Cx} \quad (149)$$

14.1.2 Operator Norms

For $p = 1, 2, \infty$ or other values, an operator norm indicates the maximum input-output gain of the matrix.

$$\|\mathbf{A}\|_p = \max_{\|\mathbf{u}\|_p=1} \|\mathbf{Au}\|_p \quad (150)$$

$$\|\mathbf{A}\|_1 = \max_{\|\mathbf{u}\|_1=1} \|\mathbf{Au}\|_1 \quad (151)$$

$$= \max_{j=1, \dots, n} \sum_{i=1}^m |\mathbf{A}_{ij}| \quad (152)$$

$$= \text{Largest absolute column sum} \quad (153)$$

$$\|\mathbf{A}\|_\infty = \max_{\|\mathbf{u}\|_\infty=1} \|\mathbf{A}\mathbf{u}\|_\infty \quad (154)$$

$$= \max_{j=1,\dots,m} \sum_{i=1}^n |\mathbf{A}_{ij}| \quad (155)$$

$$= \text{Largest absolute row sum} \quad (156)$$

$$\|\mathbf{A}\|_2 = \text{"spectral norm"} \quad (157)$$

$$= \max_{\|\mathbf{u}\|_2=1} \|\mathbf{A}\mathbf{u}\|_2 \quad (158)$$

$$= \sqrt{\max(\text{eig}(\mathbf{A}^T \mathbf{A}))} \quad (159)$$

$$= \text{Square root of largest eigenvalue of } \mathbf{A}^T \mathbf{A} \quad (160)$$

Special Properties

$$\|\mathbf{A}\mathbf{u}\|_p \leq \|\mathbf{A}\|_p \|\mathbf{u}\|_p \quad (161)$$

$$\|\mathbf{A}\mathbf{B}\|_p \leq \|\mathbf{A}\|_p \|\mathbf{B}\|_p \quad (162)$$

$$(163)$$

14.1.3 Spectral Radius

Not a proper norm.

$$\rho(\mathbf{A}) = \text{spectral radius}(\mathbf{A}) = \max_{i=1,\dots,n} |\text{eig}(\mathbf{A})_i| \quad (164)$$

Special Properties

$$\rho(\mathbf{A}) \leq \|\mathbf{A}\|_p \quad (165)$$

$$\rho(\mathbf{A}) \leq \min(\|\mathbf{A}\|_1, \|\mathbf{A}\|_\infty) \quad (166)$$

$$(167)$$

14.2 Vectors

$$\|\mathbf{x}\|_1 = \sum_i |\mathbf{x}_i| \quad (\text{L1-norm}) \quad (168)$$

$$\|\mathbf{x}\|_p = \left(\sum_i |\mathbf{x}_i|^p \right)^{1/p} \quad (\text{P-norm}) \quad (169)$$

$$\|\mathbf{x}\|_\infty = \max_i |\mathbf{x}_i| \quad (\text{L}\infty\text{-norm, L-infinity norm}) \quad (170)$$

15 | Bounds

15.1 Matrix Gain

$$\lambda_{\min}(\mathbf{A}^T \mathbf{A}) \leq \frac{\|\mathbf{Ax}\|_2^2}{\|\mathbf{x}\|_2^2} \leq \lambda_{\max}(\mathbf{A}^T \mathbf{A}) \quad (171)$$

$$\max_{\mathbf{x} \neq 0} \frac{\|\mathbf{Ax}\|_2}{\|\mathbf{x}\|_2} = \|\mathbf{A}\|_2 = \sqrt{\lambda_{\max}(\mathbf{A}^T \mathbf{A})} \implies \mathbf{x} = u_1 \quad (172)$$

$$\min_{\mathbf{x} \neq 0} \frac{\|\mathbf{Ax}\|_2}{\|\mathbf{x}\|_2} = \sqrt{\lambda_{\min}(\mathbf{A}^T \mathbf{A})} \implies \mathbf{x} = u_n \quad (173)$$

15.2 Norms

For $\mathbf{x} \in \mathbb{R}^n$

$$\frac{1}{\sqrt{n}} \|\mathbf{x}\|_2 \leq \|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_2 \leq \|\mathbf{x}\|_1 \leq \sqrt{\text{card}(\mathbf{x})} \|\mathbf{x}\|_2 \leq \sqrt{n} \|\mathbf{x}\|_2 \leq n \|\mathbf{x}\|_\infty \quad (174)$$

For any $0 < p < q$ we have that $\|\mathbf{x}\|_q \leq \|\mathbf{x}\|_p$.

15.3 Rayleigh quotients

The Rayleigh quotient of $\mathbf{A} \in \mathbb{S}^n$ is given by

$$\frac{\mathbf{x}^T \mathbf{Ax}}{\mathbf{x}^T \mathbf{x}} \quad \mathbf{x} \neq 0 \quad (175)$$

$$\lambda_{\min}(\mathbf{A}) \leq \frac{\mathbf{x}^T \mathbf{Ax}}{\mathbf{x}^T \mathbf{x}} \leq \lambda_{\max}(\mathbf{A}) \quad \mathbf{x} \neq 0 \quad (176)$$

$$\lambda_{\max}(A) = \max_{\mathbf{x}: \|\mathbf{x}\|_2=1} \mathbf{x}^T \mathbf{Ax} = u_1 \quad (177)$$

$$\lambda_{\min}(A) = \min_{\mathbf{x}: \|\mathbf{x}\|_2=1} \mathbf{x}^T \mathbf{Ax} = u_n \quad (178)$$

where u_1 and u_n are the eigenvectors associated with λ_{\max} and λ_{\min} , respectively.

16 | Linear Equations

The linear equation $\mathbf{Ax} = \mathbf{y}$ with $\mathbf{A} \in \mathbb{R}^{m,n}$ admits a solution iff $\text{rank}([\mathbf{Ay}]) = \text{rank}(\mathbf{A})$. If this is satisfied, the set of all solutions is an affine set $\mathcal{S} = \{\mathbf{x} = \bar{\mathbf{x}} + \mathbf{z} : \mathbf{z} \in \mathcal{N}(\mathbf{A})\}$ where $\bar{\mathbf{x}}$ is any vector such that $\mathbf{A}\bar{\mathbf{x}} = \mathbf{y}$. The solution is unique if $\mathcal{N}(\mathbf{A}) = \{0\}$.

$\mathbf{Ax} = \mathbf{y}$ is *overdetermined* if it is tall/skinny ($m > n$); that is, if there are more equations than unknowns. If $\text{rank}(\mathbf{A}) = n$ then $\dim \mathcal{N}(\mathbf{A}) = 0$, so there is either no solution or one solution. Overdetermined systems often have no solution ($\mathbf{y} \notin \mathcal{R}(\mathbf{A})$), so an approximate solution is necessary. See section 16.1.

$\mathbf{Ax} = \mathbf{y}$ is *underdetermined* if it is short/wide ($n > m$); that is, if it has more unknowns than equations. If $\text{rank}(\mathbf{A}) = m$ then $\mathcal{R}(\mathbf{A}) = \mathbb{R}^m$, so $\dim \mathcal{N}(\mathbf{A}) = n - m > 0$, so the set of solutions is infinite. Therefore, finding a single solution that optimizes some quantity is of interest.

$\mathbf{Ax} = \mathbf{y}$ is *square* if $n = m$. If \mathbf{A} is invertible, then the equations have the unique solution $\mathbf{x} = \mathbf{A}^{-1}\mathbf{y}$. See section 16.2.

16.1 Least-Squares

For an overdetermined system we wish to find:

$$\min_{\mathbf{x}} \|\mathbf{Ax} - \mathbf{y}\|_2^2 \quad (179)$$

Since $\mathbf{Ax} \in \mathcal{R}(\mathbf{A})$, we need a point $\tilde{\mathbf{y}} = \mathbf{Ax}^* \in \mathcal{R}(\mathbf{A})$ closest to \mathbf{y} . This point lies in the nullspace of \mathbf{A}^T , so we have $\mathbf{A}^T(\mathbf{y} - \mathbf{Ax}^*) = 0$. There is always a solution to this problem and, if $\text{rank}(\mathbf{A}) = n$, it is unique [16, p. 161]

$$\mathbf{x}^* = (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{y} \quad (180)$$

16.1.1 Regularized least-squares with low-rank data

For $\mathbf{A} \in \mathbb{R}^{m,n}$, $\mathbf{y} \in \mathbb{R}^m$, $\lambda \geq 0$, the regularized least-squares problem

$$\text{argmin}_{\mathbf{x}} \|\mathbf{Ax} - \mathbf{y}\|_2^2 + \lambda \|\mathbf{x}\|_2^2 \quad (181)$$

has a closed form solution

$$\mathbf{x} = (\mathbf{A}^T \mathbf{A} + \lambda \mathbf{I})^{-1} \mathbf{A}^T \mathbf{y} \quad (182)$$

However, if \mathbf{A} has a rank $r \ll \min(n, m)$ and a known low-rank decomposition $\mathbf{A} = \mathbf{L}\mathbf{R}^T$ with $\mathbf{L} \in \mathbb{R}^{m,r}$ and $\mathbf{R} \in \mathbb{R}^{n,r}$, then we can rewrite Equation 182 as

$$\mathbf{x} = (\mathbf{R}^T \mathbf{R} \mathbf{L}^T \mathbf{L} + \lambda \mathbf{I})^{-1} \mathbf{L}^T \mathbf{y} \quad (183)$$

This decreases the time complexity from $O(mn^2 + n^3)$ to $O(nr^2 + mr^2)$.

16.2 Minimum Norm Solutions

For undertermined systems in which $\mathbf{A} \in \mathbb{R}^{m,n}$ with $m < n$. We wish to find

$$\min_{\mathbf{x}: \mathbf{Ax}=\mathbf{y}} \|\mathbf{x}\|_2 \quad (184)$$

The solution \mathbf{x}^* must be orthogonal to $\mathcal{N}(\mathbf{A})$, so $\mathbf{x}^* \in \mathcal{R}(\mathbf{A}^T)$, so $\mathbf{x}^* = \mathbf{A}^T c$ for some c . Substituting into $\mathbf{Ax} = \mathbf{y}$ gives $\mathbf{AA}^T c = \mathbf{y}$, therefore [16, p. 162]:

$$\mathbf{x}^* = \mathbf{A}^T (\mathbf{AA}^T)^{-1} \mathbf{y} \quad (185)$$

17 | Updates

17.1 Removing a row from $\mathbf{A}^T \mathbf{A}$ ($\mathbf{A}^T \mathbf{A} \rightarrow \mathbf{A}_{\setminus i}^T \mathbf{A}_{\setminus i}$)

Plain English: Matrix times its transpose after eliminating row i from the matrix

Inputs: $\mathbf{A} \in \mathbb{R}^{k,m}$, $\mathbf{u} \in \mathbb{R}^m$, $\mathbf{v} \in \mathbb{R}^n$ and i , the row to remove from \mathbf{A}

Reduces to: $\mathbf{C} \in \mathbb{R}^{k,l}$

Algorithm:

Recall that

$$(\mathbf{AB})_{kl} = \sum_m \mathbf{A}_{km} \mathbf{B}_{ml} \quad \mathbf{A} \in \mathbb{R}^{k,m}, \mathbf{B} \in \mathbb{R}^{m,l} \quad (186)$$

then we have that

$$(\mathbf{A}^T \mathbf{A})_{kl} = \sum_m \mathbf{A}_{mk} \mathbf{A}_{ml} = \sum_{m \neq i} \mathbf{A}_{mk} \mathbf{A}_{ml} + \mathbf{A}_{jk} \mathbf{A}_{jl} = \sum_{m \neq i} \mathbf{A}_{mk} \mathbf{A}_{ml} + (\mathbf{A}_{k*})_j (\mathbf{A}_{l*})_j \quad (187)$$

where $(\mathbf{A}_{k*})_j$ is the j th element of the k th column of \mathbf{A} .

Thus,

$$\mathbf{A}_{\setminus i}^T \mathbf{A}_{\setminus i} = \mathbf{A}^T \mathbf{A} - \mathbf{A}_{*j} \mathbf{A}_{*j}^T \quad (188)$$

17.2 $\mathbf{1}_r^T \mathbf{A} \mathbf{1}_c$

Plain English: The sum of the elements of the matrix.

Reduces to: Scalar

Notation: For $\mathbf{A} \in \mathbb{R}^{r \times c}$, $\mathbf{1}_r$ is in $\mathbb{R}^{r \times 1}$ and $\mathbf{1}_c$ is in $\mathbb{R}^{c \times 1}$.

Algorithm: Traverse all the element of the matrix in order keeping track of the sum. For applications where accuracy is important and the matrices have a large dynamic range, Kahan summation or a similar technique should be used.

Update Algorithm: If an entry changes, subtract its old value from the sum and add its new value to the sum.

17.3 $\mathbf{x}^T \mathbf{A} \mathbf{x}$

Plain English: TODO

Reduces to: Scalar

Notation: \mathbf{A} must be in $\mathbb{R}^{i \times i}$. \mathbf{x} is in $\mathbb{R}^{i \times 1}$.

Algorithm: TODO

Update Algorithm: We make use of the identity $(\mathbf{x}^T \mathbf{A} \mathbf{x}) = \sum_{i,j} ((\mathbf{x} \mathbf{x}^T) \circ \mathbf{A})$. If an entry $\mathbf{A}_{i,j}$ in the matrix changes subtract its old value $\mathbf{x}_i \mathbf{x}_j \mathbf{A}_{i,j}$ and add the new value $\mathbf{x}_i \mathbf{x}_j \mathbf{A}'_{i,j}$. If an entry \mathbf{x}_i changes TODO.

18 | Optimization

18.1 Standard Forms

Least Squares

$$\min_{\mathbf{x} \in \mathbb{R}^n} \|\mathbf{y} - \mathbf{A}\mathbf{x}\|_2 \quad (189)$$

LASSO

$$\min_{\mathbf{b} \in \mathbb{R}^n} \left(\frac{1}{N} \|\mathbf{y} - \mathbf{X}\mathbf{b}\|_2^2 + \lambda \|\mathbf{b}\|_1 \right) \quad (190)$$

LP: Linear program

$$\min_{\mathbf{x}} \mathbf{c}^T \mathbf{x} \quad (191)$$

$$\text{s.t. } \mathbf{A}_{\text{eq}} \mathbf{x} = \mathbf{b}_{\text{eq}} \quad (192)$$

$$\mathbf{A}\mathbf{x} \leq \mathbf{b} \quad (193)$$

Linear Fractional Program

$$\underset{\mathbf{x}}{\text{maximize}} \quad \frac{\mathbf{c}^T \mathbf{x} + a}{\mathbf{d}^T \mathbf{x} + b} \quad (194a)$$

$$\text{subject to} \quad \mathbf{A}\mathbf{x} \leq \mathbf{b} \quad (194b)$$

Additional constraints must ensure $\mathbf{d}^T \mathbf{x} + b$ has the same sign throughout the entire feasible region.

$$\underset{\mathbf{x}}{\text{max}} \quad \frac{\mathbf{c}^T \mathbf{x} + a}{\mathbf{d}^T \mathbf{x} + b} \quad (195)$$

$$\text{s.t.} \quad (196)$$

QCQP: Quadratic Constrained Quadratic Programs

$$\min_{\mathbf{x}} \mathbf{x}^T \mathbf{H}_0 \mathbf{x} + 2\mathbf{c}_0^T \mathbf{x} + \mathbf{d}_0 \quad (197)$$

$$\text{s.t. } \mathbf{x}^T \mathbf{H}_i \mathbf{x} + 2\mathbf{c}_i^T \mathbf{x} + \mathbf{d}_i \leq 0, \quad i \in \mathcal{I} \quad (198)$$

$$\mathbf{x}^T \mathbf{H}_j \mathbf{x} + 2\mathbf{c}_j^T \mathbf{x} + \mathbf{d}_j = 0, \quad j \in \mathcal{E} \quad (199)$$

Note that, in general, QCQPs are NP-Hard.

QP: Quadratic Program

$$\min_{\mathbf{x}} \quad \frac{1}{2} \mathbf{x}^T \mathbf{H}_0 \mathbf{x} + \mathbf{c}_0^T \mathbf{x} \quad (200)$$

$$\text{s.t. } \mathbf{A}_{\text{eq}} \mathbf{x} = \mathbf{b}_{\text{eq}} \quad (201)$$

$$\mathbf{A}\mathbf{x} \leq \mathbf{b} \quad (202)$$

SOCP: Second Order Cone Program (Standard Form)

$$\min_{\mathbf{x}} \mathbf{c}^T \mathbf{x} \quad (203)$$

$$\text{s.t. } \|\mathbf{A}_i \mathbf{x} + \mathbf{b}_i\|_2 \leq \mathbf{c}_i^T \mathbf{x} + \mathbf{d}_i, \quad i = 1, \dots, m \quad (204)$$

SOCP: Second Order Cone Program (Conic Standard Form)

$$\min_{\mathbf{x}} \mathbf{c}^T \mathbf{x} \quad (205)$$

$$\text{s.t. } (\mathbf{A}_i \mathbf{x} + \mathbf{b}_i, \mathbf{c}_i^T \mathbf{x} + \mathbf{d}_i) \in \mathcal{K}_{m_i} \quad i = 1, \dots, m \quad (206)$$

18.2 Transformations

18.2.1 Linear-Fractional to Linear

We transform a Linear-Fractional Program

$$\begin{aligned} & \underset{\mathbf{x}}{\text{maximize}} && \frac{\mathbf{c}^T \mathbf{x} + a}{\mathbf{d}^T \mathbf{x} + b} \end{aligned} \quad (207a)$$

$$\text{subject to} \quad \mathbf{A} \mathbf{x} \leq \mathbf{b} \quad (207b)$$

where $\mathbf{d}^T \mathbf{x} + b$ has the same sign throughout the entire feasible region to a linear program using the Charnes–Cooper transformation [17] by defining

$$\mathbf{y} = \frac{1}{\mathbf{d}^T \mathbf{x} + b} \cdot \mathbf{x} \quad (208)$$

$$t = \frac{1}{\mathbf{d}^T \mathbf{x} + b} \quad (209)$$

to form the equivalent program

$$\underset{\mathbf{y}, t}{\text{maximize}} \quad \mathbf{c}^T \mathbf{y} + at \quad (210a)$$

$$\text{subject to} \quad \mathbf{A} \mathbf{y} \leq \mathbf{b}t, \quad (210b)$$

$$\mathbf{d}^T \mathbf{y} + bt = 1, \quad (210c)$$

$$t \geq 0 \quad (210d)$$

We then have $\mathbf{x}^* = \frac{1}{t} \mathbf{y}$.

19 | Algorithms

19.1 Gram-Schmidt

TODO

Bibliography

- [1] Volker Strassen. Gaussian elimination is not optimal. *Numerische mathematik*, 13(4):354–356, 1969.
- [2] Günter Rote. Division-free algorithms for the determinant and the pfaffian: algebraic and combinatorial approaches. In *Computational discrete mathematics*, pages 119–135. Springer, 2001.
- [3] Erwin H. Bareiss. Sylvester’s identity and multistep integer-preserving gaussian elimination. *Mathematics of Computation*, 22(103):565–578, 1968. ISSN 00255718, 10886842. doi: 10.2307/2004533. URL <http://www.jstor.org/stable/2004533>.
- [4] V Ya Pan. Strassen’s algorithm is not optimal trilinear technique of aggregating, uniting and canceling for constructing fast algorithms for matrix operations. In *Foundations of Computer Science, 1978., 19th Annual Symposium on*, pages 166–176. IEEE, 1978. doi: 10.1109/SFCS.1978.34.
- [5] DARIO ANDREA Bini, Milvio Capovani, Francesco Romani, and Grazia Lotti. $o(n^{2.7799})$ complexity for n by n approximate matrix multiplication. *Information processing letters*, 8(5): 234–235, 1979.
- [6] Arnold Schönhage. Partial and total matrix multiplication. *SIAM Journal on Computing*, 10(3):434–455, 1981.
- [7] Francesco Romani. Some properties of disjoint sums of tensors related to matrix multiplication. *SIAM Journal on Computing*, 11(2):263–267, 1982.
- [8] Don Coppersmith and Shmuel Winograd. On the asymptotic complexity of matrix multiplication. *SIAM Journal on Computing*, 11(3):472–492, 1982.
- [9] Volker Strassen. The asymptotic spectrum of tensors and the exponent of matrix multiplication. In *Foundations of Computer Science, 1986., 27th Annual Symposium on*, pages 49–54. IEEE, 1986.
- [10] Don Coppersmith and Shmuel Winograd. Matrix multiplication via arithmetic progressions. *Journal of Symbolic Computation*, 9(3):251 – 280, 1990. ISSN 0747-7171. doi: 10.1016/S0747-7171(08)80013-2. URL <http://www.sciencedirect.com/science/article/pii/S0747717108800132>. Computational algebraic complexity editorial.
- [11] Virginia Vassilevska Williams. Multiplying matrices faster than coppersmith-winograd. In *Proceedings of the Forty-fourth Annual ACM Symposium on Theory of Computing, STOC ’12*, pages 887–898, New York, NY, USA, 2012. ACM. ISBN 978-1-4503-1245-5. doi: 10.1145/2213977.2214056. URL <http://doi.acm.org/10.1145/2213977.2214056>.
- [12] François Le Gall. Powers of tensors and fast matrix multiplication. In *Proceedings of the 39th International Symposium on Symbolic and Algebraic Computation, ISSAC ’14*, pages 296–303, New York, NY, USA, 2014. ACM. ISBN 978-1-4503-2501-1. doi: 10.1145/2608628.2608664. URL <http://doi.acm.org/10.1145/2608628.2608664>.
- [13] Gilbert Strang. *Introduction to Linear Algebra*. 2016.
- [14] Elizabeth Million. The hadamard product. <http://buzzard.ups.edu/courses/2007spring/projects/million-paper.pdf>, 2007.

- [15] Thomas P Minka. Old and new matrix algebra useful for statistics. <https://tminka.github.io/papers/matrix/minka-matrix.pdf>, 2000.
- [16] Giuseppe C Calafiore and Laurent El Ghaoui. *Optimization models*. Cambridge University Press, 2014. ISBN 978-1-107-05087-7.
- [17] A. Charnes and W. W. Cooper. Programming with linear fractional functionals. *Naval Research Logistics Quarterly*, 9(3-4):181–186, 1962. doi: 10.1002/nav.3800090303. URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/nav.3800090303>.

Index

L^∞ -norm, 34

L^1 -norm, 34

P-norm, 34