

The Setting

We work in a compact, riemanian, orientable, smooth Manifold (M, g) of dimension m together with a Morse-Smale funktion

$$f : M \rightarrow \mathbb{R} \quad (1)$$

We assume that q is a critical point of index $k+1$ and p is a critical point of index k . we define $f(q) = b$ and $f(p) = a$ and assume that there is no kritical point in $f^{-1}(a, b) \subseteq M$. We define the sub and superlevelsets

$$M^t := \{x \in M | f(x) \leq t\} \quad \text{and} \quad M_t := \{x \in M | f(x) \geq t\}, \quad (2)$$

and the constants

$$c \in (a, b) \quad , \quad \varepsilon > 0 \text{ small} \quad , \quad T > 0 \text{ big} . \quad (3)$$

With this we define the sets:

$$N_q := \{x \in M_c | f(\varphi_{-T}(x)) \leq b + \varepsilon\}, \quad (4)$$

$$L_q := \{x \in N_q | f(x) = c\}, \quad (5)$$

$$N_p := \{x \in M^c | f(\varphi_T(x)) \geq a - \varepsilon\}, \quad (6)$$

$$L_p := \{x \in N_p | f(\varphi_T(x)) = a - \varepsilon\}, \quad (7)$$

and finally:

$$C := N_p \cup N_q, \quad (8)$$

$$B := N_p \cup L_q, \quad (9)$$


$$A := L_p \cup (L_q - \overset{\circ}{N_p}). \quad (10)$$

Lemma 0.0.1. *We claim that*

1. (N_q, L_q) is a regular index pair for q .
2. (C, B) is an index pair for q .
3. (N_p, L_p) is a regular index pair for p .
4. (B, A) is an index pair for p .
5. N_p is a tubular neighbourhood of $W(\rightarrow p) \cap M^c$.

The statements 1-4 are easy to proof by the definitions. The regularity can be derived from a general fact for pairs (X, Y) in metric spaces: If Y is closed in X and

there is a neighbourhood of Y that is open in X such that Y is a strong deformation retract of U . So the only thing left to show is, that N_p is a tubular neighbourhood. in [MorseTheorySalmbo] in the proof of lemma 3.2 Salamon claims this to be true without a proof (page 119 in the attached source). Similar, in [banyaga2004lectures] Banyaga claims this (also without any argument). I find this difficult to proof, since we cannot use the flow for the map from the normal bundle, as points leave N_p along the flow: The Set N_p is sketched in the figure below.



So to proof this, we first formulate two lemmas. First we restate/ refine the λ -lemma:

Lemma 0.0.2 (The Technical λ -Lemma). *Let M be a smooth manifold and let p be a hyperbolic fixpoint of a diffeomorphism φ on M with $\dim W(p \rightarrow) = r$ for $0 < r < m = \dim M$. Let N be an injective immersed submanifold of M that is invariant under φ and has a point of transverse intersection q with $W(\rightarrow p)$. Then for any $\varepsilon > 0$, any r -cell neighbourhood D of q in N that also intersects $W(\rightarrow p)$ transversal and any r -cell neighbourhood B of p in $W(p \rightarrow)$ there is a smaller cell $\tilde{D} \subset D$, an $n \in \mathbb{N}$ such that the following is true:*

We can assume that B and $\varphi^n(\tilde{D})$ lives in a chart centered at p with a product structure $U \cong W(\rightarrow p) \cap B_\mu(0) \times W(p \rightarrow) \cap B_\mu(0)$. Here, the invars of the projection into $B \subseteq W(\rightarrow p)$:

$$h : B \rightarrow \varphi^n(\tilde{D})$$

satisfies:

$$1. \sup_{x \in B} \|x - h(x)\| < \varepsilon \text{ and}$$

$$2. \text{ For the inclusions } i_1 : B \rightarrow M \text{ and } i_2 : \varphi^n(\tilde{D}) \rightarrow M \text{ we have that}$$

$$\sup_{x \in B} \|d(i_1)_x - d(i_2 \circ h)_x\| < \varepsilon.$$

Proof. Let q be a point of transversal intersection of $W(\rightarrow p)$ with N . Since both N and $W(\rightarrow p)$ are invariant under φ , it follows that $\varphi^n(q) \in N \cap W(\rightarrow p)$ for all $n \in \mathbb{N}$. Moreover, the intersection remains transversal: the direct sum decomposition $T_q M = T_q W(\rightarrow p) \oplus T_q N$ is preserved under $d\varphi_q$, which implies transversality at $\varphi^n(q)$ as well.

Therefore, we may assume that q lies in a coordinate chart centered at the hyperbolic fixed point p , and identify a neighborhood of p with an open subset of $\mathbb{R}^m$, where $m = \dim M$. By hyperbolicity of p , the tangent space at p splits as

$$T_p M = T_p^s M \oplus T_p^u M,$$

and this splitting is preserved by $d\varphi_p$. As discussed earlier, there exists a chart in which $d\varphi_p$ is represented by a real diagonal matrix (in suitable coordinates). Hence, we may choose an orthonormal basis of coordinates $(x_s, x_u) \in \mathbb{R}^s \times \mathbb{R}^u$, adapted to the stable and unstable subspaces, such that

$$d\varphi_0 = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix},$$

where the eigenvalues of A have modulus less than 1, and those of B greater than 1.

Since we are working in $\mathbb{R}^m$, we identify the manifold and its tangent space locally around 0. We then expand φ in a Taylor series at the origin:

$$\varphi(x_s, x_u) = \varphi(0) + d\varphi_0(x_s, x_u) + f(x_s, x_u),$$

reference
einfügen

where $f(0) = 0$ and $df(0) = 0$. Writing this in coordinates gives:

$$\varphi(x_s, x_u) = (Ax_s + f_s(x_s, x_u), Bx_u + f_u(x_s, x_u)). \quad (11)$$

Furthermore, since $f = \varphi - d\varphi|_0$ we know that:

$$\frac{\partial f_s}{\partial x_u}|_{(0, x_u)} = \frac{\partial \varphi_s}{\partial x_u}|_{(0, x_u)} = 0 = \frac{\partial f_u}{\partial x_s}|_{(x_s, 0)} = \frac{\partial \varphi_u}{\partial x_s}|_{(x_s, 0)}.$$

Similar to Corollary ?? we can define an inner product such that for the operator norm we have

$$\|A\| \leq \alpha < 1 \quad \text{and} \quad \|B^{-1}\| \leq \frac{1}{\beta} < 1.$$

Since $1 = \|\text{id}\| = \|BB^{-1}\| \leq \|B\| \cdot \|B^{-1}\|$ we know that $\|B\| \geq \beta$. For a neighbourhood $U_1 \subseteq U$ of the origin we define

$$k := \max_{i,j=s,u} \left\{ \sup_{U_1} \left\| \frac{\partial f_i}{\partial x_j} \right\| \right\},$$

and choose U_1 small enough such that k satisfies:

$$\begin{aligned} 0 &< k < 1 \\ \beta_1 &:= \beta - k > 1 \\ \alpha_1 &:= \alpha + k < 1 \\ k &< \frac{1}{4}(b_1 - 1)^2. \end{aligned}$$

This can be achieved since $df|_0 = 0$. Let B be a cell neighbourhood of the origin in $W(p \rightarrow)$. For some $n \in \mathbb{N}$ we have that $\varphi^{-n}(B) \subseteq U_1$. If the λ -lemma holds for $\varphi^{-n}(B)$ for all ε it also holds for B , since the metric and φ are continuous and the r -cell is compact. Therefore, the change of distance after applying φ^{-n} is finite and we can choose the distance from $\varphi^{-n}(B)$ to D small enough such that B is ε C^1 -close to D . Hence, without restrictions $B \subseteq U_1$ and $q \in U_1$. Next we define the inclination of N with respect to the (un-)stable manifolds as follows:

Let $x \in N \cap U_1$ and $v^0 \in T_x N \subseteq T_0 M = T_o^s M \oplus T_o^u M$ and let $v^0 = (v_s^0, v_u^0)$ be the coordinate entries. Furthermore let v be a unit vector. Now we define the **inclination** of v^0 :

$$\lambda(v^0) = \frac{\|v_s^0\|}{\|v_u^0\|}.$$

In other words: The smaller the inclination, the more parallel we are to $W(p \rightarrow)$. Notice how we can only define the inclination if $v_u^0 \neq 0$. Furthermore define:

$$x^n := \varphi^n(x), \quad v^n := d\varphi_{x^{n-1}}(v^{n-1}) \quad \text{and} \quad \varphi(x) = (\varphi_s(x), \varphi_u(x)) \in W(\rightarrow 0) \times W(0 \rightarrow).$$

Now we want to calculate the derivative $d\varphi|_{q^n}$. But first inspect:

$$d\varphi|_x = \begin{pmatrix} \frac{\partial \varphi_s}{\partial x_s}|_x & \frac{\partial \varphi_s}{\partial x_u}|_x \\ \frac{\partial \varphi_u}{\partial x_s}|_x & \frac{\partial \varphi_u}{\partial x_u}|_x \end{pmatrix} \stackrel{(11)}{=} \begin{pmatrix} A + \frac{\partial f_s}{\partial x_s}|_x & \frac{\partial f_s}{\partial x_u}|_x \\ \frac{\partial f_u}{\partial x_s}|_x & B + \frac{\partial f_u}{\partial x_u}|_x \end{pmatrix}$$

Since $q^n \in W(\rightarrow 0)$ and $\frac{\partial f_u}{\partial x_s}(x_s, u)=0$ this gives us:

$$d\varphi|_{q^n} = \begin{pmatrix} A + \frac{\partial f_s}{\partial x_s}|_{q^n} & \frac{\partial f_s}{\partial x_u}|_{q^n} \\ 0 & B + \frac{\partial f_u}{\partial x_u}|_{q^n} \end{pmatrix}$$

Let $B_s(\delta)$ denote the ball of radius δ around the origin that lives in $W(\rightarrow p)$ and $U_2 := B_s(\delta) \times B$. Since $\frac{\partial f_s}{\partial x_u}|_{(0, x_u)} = 0$ and by the smoothness of φ , for any $\varepsilon > 0$ there exists a δ such that

$$\underbrace{\sup_{U_2} \left\| \frac{\partial f_s}{\partial x_u} \right\|}_{:=k_1} < \min\{\varepsilon, k\}.$$

Again, we assume $q \in D \subseteq U_2$. Now we want to get a hold of the inclination: Let $x \in N$ such that $x^j \in U_2$ for all $j \leq n$ and let $v^0 \in T_x N$. Denote $v^n = (v_s^n, v_u^n) \in T_0 W(\rightarrow 0) \times T_0 W(0 \rightarrow)$.

$$\begin{aligned} \lambda(v^n) &= \frac{\|d\varphi_s|_{x^{n-1}}(v^{n-1})\|}{\|d\varphi_u|_{x^{n-1}}(v^{n-1})\|} \\ &= \frac{\|Av_s^{n-1} + \frac{\partial f_s}{\partial x_s}|_{x^{n-1}}v_s^{n-1} + \frac{\partial f_s}{\partial x_u}|_{x^{n-1}}(v_u^{n-1})\|}{\|\frac{\partial f_u}{\partial x_s}|_{x^{n-1}}(v_s^{n-1}) + Bv_u^{n-1} + \frac{\partial f_u}{\partial x_u}|_{x^{n-1}}(v_u^{n-1})\|} \\ &\leq \frac{(\alpha + k)\|v_s^{n-1}\| + k_1\|(v_u^{n-1})\|}{k\|(v_s^{n-1})\| + (\beta + k)\|v_u^{n-1}\|} \\ &\leq \frac{(\alpha + k)\|v_s^{n-1}\| + k_1\|(v_u^{n-1})\|}{(\beta - k)\|v_u^{n-1}\| - k_1\|(v_s^{n-1})\|} && \text{since } \beta - k > 1 \text{ and } k_1 < k < 1 \\ &= \frac{(\alpha + k)\lambda(v^{n-1}) + k_1}{\beta_1 - k_1\lambda v^{n-1}} && \text{we divided by } \|v_u^{n-1}\| \\ &\leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1 - k_1\lambda(v^{n-1})} \end{aligned} \tag{12}$$

If $x \in W(\rightarrow 0)$ the blue part above disappears and we get:

$$\lambda(v^n) \leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1}.$$

By induction we get:

$$\begin{aligned}\lambda(v^n) &\leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1} = \frac{\frac{\lambda(v^{n-2}) + k_1}{\beta_1} + k_1}{\beta_1} = \frac{\frac{\lambda(v^{n-2}) + \frac{k_1\beta_1}{\beta_1} + k_1}{\beta_1}}{\beta_1} = \frac{\lambda(v^{n-2}) + k_1 + k_1\beta_1}{\beta_1^2} \leq \dots \\ &\leq \frac{\lambda(v^0) + \sum_{l=0}^{n-1} k_1\beta_1^l}{\beta_1^n}.\end{aligned}\tag{13}$$

By using the geometric series we see:

$$\begin{aligned}\lambda(v^n) &\leq \frac{\lambda(v^0)}{\beta_1^n} + k_1 \sum_{l=1}^n \frac{1}{\beta_1^l} \leq \frac{\lambda(v^0)}{\beta_1^n} + k_1 \left(\frac{1}{1 - \beta_1} - 1 \right) = \frac{\lambda(v^0)}{\beta_1^n} + k_1 \left(\frac{b}{\beta_1 - 1} - \frac{\beta_1 - 1}{\beta_1 - 1} \right) \\ &= \frac{\lambda(v^0)}{\beta_1^n} + k_1 \left(\frac{1}{\beta_1 - 1} \right).\end{aligned}$$

Since β_1 was greater then 1 we can conclude that $\lim_{n \rightarrow \infty} \frac{\lambda(v^0)}{\beta_1^n} = 0$ and since $k < \frac{1}{4}(\beta_1 - 1)^2$ we know that $\frac{k}{\beta_1 - 1} < \frac{b-1}{4}$. Because of that, a n_0 exists such that for all $n > n_0$:

$$\lambda(v^n) \leq \frac{\lambda(v^0)}{\beta_1^n} + \frac{k}{b-1} < \frac{\beta_1 - 1}{3}.$$

Lets collect what we have done so far: We take a point x from a transverse intersection of N and $W(\rightarrow p)$. Then, we inspect the inclination of a tangent vector v in $T_x N \subseteq T_0^s M \times T_0^u M$ and move the tangent vector via $d\varphi$ alongside of the stable set and realize that the inclination gets smaller, the more we move it.

For any $v \in T_{q^n} N$ there exists a $\tilde{v} \in T_q N$ such that $v = \tilde{v}^n$, since $d\varphi|_x : T_x M \rightarrow T_{\varphi(x)} M$ is bijective. If we take an arbitrary local section $f : M \rightarrow TM$ that denotes the choice of $v^0 \in T_x N$ we know that $\lambda \circ f$ is continuous, as it is the composition of continuous functions. In other words, the inclination smoothly depends on v^0 and x . This tells us that for a small enough cellular neighbourhood $D_0 \subseteq D \subseteq N \cap U_2$ such that for any $x \in D_0$ and $v_0 \in T_x N$ of length 1 we have

$$\lambda(v^0) \leq \frac{1}{2}(\beta_1 - 1).\tag{14}$$

(This is true as long as the invlination is well defined) Now we claim that if $\varphi^j(v^0) \in U_2$ for all $j = 0, \dots, n$ then $\lambda(v^n) \leq \frac{1}{2}(\beta_1 - 1)$. We prove this by induction: The case $n = 0$ is already done. For the induction step we use the estimate (12), $k_1 < k < 1$ and $k < \frac{1}{4}(\beta_1 - 1)^2$:

$$\lambda(v^{n+1}) \leq \frac{\lambda(v^n) + k_1}{\beta_1 - k_1\lambda(v^n)} \leq \frac{\frac{1}{2}(\beta_1 - 1) + \frac{1}{4}(\beta_1 - 1)^2}{\beta_1 - \frac{1}{2}(\beta_1 - 1)} = \frac{1}{2}(\beta_1 - 1).$$

Using the above estimate and (12) again we conclude that for all $x \in D_0$ such that $\varphi^j(x) \in U_2$ for all $j = 0, \dots, n$ and for any unit vector $v^0 \in T_x N$ we have:

$$\begin{aligned}\lambda(v^n) &\leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1 - k_1 \lambda(v^{n-1})} \\ &\leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1 - \frac{1}{2}(\beta_1 - 1)} \\ &\leq \frac{\lambda(v^{n-1}) + k_1}{(\frac{1}{2}(1 + \beta_1))}.\end{aligned}$$

Notice that for $\beta_2 := \frac{1}{2}(1 + \beta_1)$ we have $1 < \beta_2 < \beta_1$. Again like in the equation (13) we can use induction to conclude:

$$\lambda(v^n) \leq \frac{\lambda(v^0)}{\beta_2^n} + \frac{k_1}{\beta_2 - 1},$$

and since $k_1 \leq \varepsilon$ we have for any n large enough, any $x \in D_0$ such that $\varphi^j(x) \in U_2$ for $j = 0, \dots, n$ and any unit vector $v^0 \in T_x N$:

$$\lambda(v^n) \leq \varepsilon(1 + \frac{1}{\beta_2 - 1}).$$

Let $\pi_u : U \rightarrow W(0 \rightarrow)$ be the projection. Since φ is expanding on $W(0 \rightarrow)$ we know that a $n_2 \geq n_1$ exists such that $B \subseteq \pi_u(D_{n_2})$, where D_{n_2} denotes the connected component of $\varphi^{n_2}(D_0) \cap U$ that contains q^{n_2} . If $\dim N > \dim W(0 \rightarrow)$ we can choose a subset of $\dim W(0 \rightarrow)$ and then the projection restricted to D_{n_2} : $\pi_u|_{D_{n_2}}$ can be modified to be one to one. For this we use the fact that we are an embedded manifold. Hence, we are given by an implicit function where the derivatives in $W(\rightarrow p)$ -direction are invertible as a matrix. By the implicit function theorem we can take any point x from D_{n_2} and conclude that there is a neighbourhood $U(x) \subset W(p \rightarrow)$ around $\pi_u(x)$ in D_{n_2} and a smooth function $G : W(p \rightarrow) \rightarrow W(\rightarrow p)$, such that around x D_{n_2} looks like $(y, G(y))$ telling us that the projection onto the domain of G is one to one after restricting D_{n_2} to $U(x) \times \text{im}(G)$. Now we can define h to be the inverse of that projection (that is $G \times \text{id}$) and get the closeness of B to D_{n_2} . To see that we first check the distances and the difference after differentiating. For this we need the maps: $i_1 : B \rightarrow M, x \mapsto (0, x)$, and $i_2 : D_{n_2} \rightarrow M, x \mapsto x$ and $h : B \rightarrow D_{n_2}, x \mapsto \pi_u^{-1}(x) = (h_s(x), x) = (h_s(x), h_u(x))$:

$$\begin{aligned}
\sup_{x \in B} \|d(i_1)_x - d(i_2 \circ h)_x\| &= \sup_{x \in B} \left\| \begin{pmatrix} 0 \\ 1 \end{pmatrix} - \begin{pmatrix} d(h_s)_x \\ 1 \end{pmatrix} \right\| \\
&= \sup_{x \in B} \|d(h_s)_x\| \\
&= \sup_{x \in B} \sup_{\|v\| \leq 1} \left\| \frac{d(h_s)_x}{d(h_u)_x}(v) d(h_u)_x \right\| (v) \quad \text{here: } \begin{cases} h_u \equiv \text{id}_{W(\rightarrow p)} \times h_u \\ h_s \equiv h_s \times \text{id}_{W(p \rightarrow)} \end{cases} \\
&= \sup_{x \in B} \sup_{\|v\| \leq 1} \frac{\|(d(h)_x)_s(v)\|}{\|(d(h)_x)_u(v)\|} \|v\| \\
&= \sup_{x \in B} \sup_{\|v\|=1} \lambda(dh_x(v)) \quad \text{since } dh_x \text{ is linear} \\
&\leq \varepsilon \left(1 + \frac{1}{\beta_2 - 1}\right).
\end{aligned}$$

The last inequality follows from $dh_x(v)$ being an element in $T_x N$ and the existence of a $\tilde{v} \in D_0 : \tilde{v}^n = v$. Finally since β_2 is independent of ε we can make this construction for any ε . Furthermore since we can increase n (and decrease the size of $\tilde{D}$) such that D_{n^2} is arbitrary close to the origin, the statement that

$$\sup_{x \in B} \|x - h(x)\| < \varepsilon$$

is clear.

Therefore, by definition we have that for any $\mathcal{N}^r(f, (\varphi, U), (\psi, V), \varepsilon)$ that contains i_2 there is a r -cell D_{n^2} and a h such that $i_2 \circ h \in \mathcal{N}^r(f, (\varphi, U), (\psi, V), \varepsilon)$. This tells us that every neighbourhood of i_1 contains such an h and therefore, in the C^r -generating metric d_1 , an r -cell and a h exists such that $d_1(i_1, i_2 \circ h) < \varepsilon$. This concludes the proof of the lambda lemma.

we proofed the same with the only difference, that we defined the r -cell D arbitrary. We now want to show that this r -cell can be defined to be the image of a moved smaller cell of a given r -cell. So lets inspect the already given proof:

We started by moving the point of transverse intersection q closer to the fixpoint by applying the diffeomorphism to it. Then we defined an neighbourhood D_0 of that moved point in the stable manifold (compare ??). Here we were able to get a hold of the inclination. Then we moved D_0 via applying the isomorphism closer to the fixed point and restricted it to the dimension of the unstable manifold. This restriction we can refine as follows: First we define D_{n^2} to be the moved D_0 . Lets say we moved q n -times. In other words $q \in \varphi^{-n}(D_{n^2})$. Then there exist a small r -cell neighbourhood of q that lives in $D \cap \varphi^{-n}(D_{n^2})$. That is because $\varphi^{-n}(D_{n^2})$ is an in N open neighbourhood of y , hence $\varphi^{-n}(D_{n^2}) \cap D$ is open in D . Hence we can find a ball around y in D that is contained in the intersection. We call that r -cell $\tilde{D}$. Then $\varphi^n(\tilde{D})$ is an r -dimensional subset of D_{n^2} and the rest of the proof can be done with that subset. All arguments work, as that subset is transverse to $W(\rightarrow p)$. $\square$

Next we want to see, that locally the flow is monoton in the normal direction:

Lemma 0.0.3. *Let U be a normal neighbourhood of $W(\rightarrow p)$. For $x \in W(\rightarrow p)$ we define U_x to be the image of the normal fibre above x . Then for any $T > 0$ big enough and $\varepsilon > 0$ small enough, the function $f \circ \varphi_T : U_x \rightarrow R$ is monoton decreasing in a small ball of radius ε meaning that if $\lambda > 1, v \in U_x$ with $\|v\| < \|\lambda v\| < \varepsilon$ then $f \circ \varphi_T(v) < f \circ \varphi_T(\lambda v)$.*

Proof. This is an application of the technical λ -lemma. Because, we can define $N = M$ and U_x is our r -cell. Then for a smaller r -cell we know that $\varphi_n(U_x)$ is arbitrary close with the derivative being arbitrary close. Now Assume that there is a fibre where the map is **not** monotonly increasing. This means that there is a $x \in W(\rightarrow p)$ and a $v \in U_x$ such that the map

$$\begin{aligned} g : (0, \varepsilon) &\rightarrow \mathbb{R} \\ \lambda &\mapsto f \circ \varphi_T(\lambda \cdot v) \end{aligned}$$

is increasing in some μ and hence $g'(\mu) \geq 0$. □