

# A Morse-Theoretic Construction of the Atiyah–Hirzebruch Spectral Sequence

David Seez

October 7, 2026

## Contents

|          |                                                      |            |
|----------|------------------------------------------------------|------------|
| <b>1</b> | <b>Topological Preliminaries</b>                     | <b>2</b>   |
| 1.1      | Compactly Generated Weak Hausdorff Spaces . . . . .  | 2          |
| 1.2      | Topological Pairs . . . . .                          | 4          |
| 1.3      | Mapping Spaces . . . . .                             | 6          |
| <b>2</b> | <b>Vector Bundles and Riemannian Geometry</b>        | <b>7</b>   |
| <b>3</b> | <b>Complex K-Theory</b>                              | <b>24</b>  |
| 3.1      | Mathematical Goals . . . . .                         | 24         |
| 3.2      | Classification of Vector Bundles . . . . .           | 24         |
| 3.3      | K-Theory as a Cohomology Theory . . . . .            | 39         |
| 3.4      | Homotopy-Theoretic definition of $K$ . . . . .       | 50         |
| 3.5      | K-Theory as a Cohomology Theory . . . . .            | 59         |
| <b>4</b> | <b>Orientations</b>                                  | <b>75</b>  |
| 4.1      | Oriented Vector Spaces and Bundles . . . . .         | 75         |
| 4.2      | K-theoretic Orientation . . . . .                    | 80         |
| 4.3      | Local K-Theoretic Orientation . . . . .              | 84         |
| 4.4      | Orientations Calculated by a Jacobean Sign . . . . . | 87         |
| <b>5</b> | <b>Morse Theory</b>                                  | <b>91</b>  |
| 5.1      | Signed Flow Lines . . . . .                          | 103        |
| 5.2      | A Morse-Theoretic Künneth Formula . . . . .          | 105        |
| <b>6</b> | <b>A Morse-Theoretic Filtration</b>                  | <b>112</b> |
| 6.1      | Conley index . . . . .                               | 112        |
| 6.2      | Commuting Boundary Maps . . . . .                    | 117        |

## 1 Topological Preliminaries

This section aims to set the playing ground of the thesis. Since Morse theory needs the topological spaces to be rather restricted and the goal is to relate Morse Homology to topological K-Theory, there is an urge to just not worry about topological nuances. However, for proving things we still need the vocabulary of topological notions. This chapter serves to provide the notions and theorems with references whilst omitting the technical proofs. Steenrod himself writes in his article "A convenient category of topological spaces" [Ste67]:

"Our need is to be able to make a variety of constructions and to know that the results have good properties without the tedious spelling out at each step of lengthy hypotheses such as countably paracompact normal, completely regular, first axiom of countability, metrizable, and so forth. It may be good research technique and an enjoyable exercise to analyse the precise circumstances for which an argument works; but if a developing theory is to be handy for research workers and attractive to students, then simplicity of the fundamentals must be the goal."

Whilst Steenrod argues for the category of *compactly generated Hausdorff spaces* the standard nowadays is the category of *compactly generated weak Hausdorff spaces*. The latter has direct limits, whilst Steenrod's doesn't. One might wonder why we don't just use the category of CW-complexes, like Hatcher does. There are two reasons for this. The main goal is a Morse theoretic Spectral sequence, which relies on a Morse theoretic filtration. It would be a theoretical drawback to assume that all spaces already omit a CW filtration, as such a filtration naturally already leads to a Atiyah Hirzebruch Spectral sequence. The second reason is, that whilst a lot of pointset topological technicalities can be omitted using CW-complexes, not all of them vanish. This results in the necessity of a discussion of those technicalities, or ignoring the details. Hence we stay away from CW complexes. Let us start constructing this convenient category. All topological constructions that are taught in the compulsory section of a university are assumed to be known. Most proofs are omitted but can be found in [Str].

### 1.1 Compactly Generated Weak Hausdorff Spaces

**Remark 1.1.1** (Notation and Convention). The literature on topological spaces in geometric contexts is widely non-uniform. For this work, a topological space is *compact* if it is quasi-compact (every open cover has a finite subcover) and Hausdorff (every pair of points can be separated by disjoint open subsets).

**Definition 1.1.2** (Compactly Generated). A topological space  $X$  is **compactly generated**, if a subset  $A \subseteq X$  is closed, if and only if for all compact  $K$  and continuous  $f : K \rightarrow X$   $f^{-1}(A)$  is closed.

$\text{def: topPre-compactlyGenerated}$




Figure 1: The definition of compactly generated weak Hausdorff spaces.

def:topPre-weakHausdorff

**Definition 1.1.3** (Weak Hausdorff). A topological space  $X$  is called weak Hausdorff, if for all compact spaces  $K$  and continuous map  $f : K \rightarrow X$  the image  $f(K)$  is closed in  $X$ .

**Definition 1.1.4.** We call the category of compactly generated weak Hausdorff spaces together with continuous functions as morphisms **CGWH**.

prop:topPre-spacesThatAreCGWH

**Proposition 1.1.5.** 1. Every closed subset of a compactly generated weak Hausdorff space is compactly generated weak Hausdorff in the subspace topology.

2. Every locally compact Hausdorff space, and hence every compact space, is compactly generated weak Hausdorff.
3. Every manifold is compactly generated weak Hausdorff.
4. Every metric space is compactly generated weak Hausdorff.
5. Every disjoint union of compactly generated weak Hausdorff spaces is compactly generated weak Hausdorff.
6. If  $X$  is compactly generated weak Hausdorff and  $Y$  locally compact Hausdorff, then  $X \times Y$  is compactly generated weak Hausdorff in the product topology.
7. If  $X$  is compactly generated weak Hausdorff and  $Y \subseteq X$  a closed subset of  $X$ . Then the quotient topology on  $X/Y$  is compactly generated and weak Hausdorff.

There is another caveat and reason for the usage of **CGWH**. If we view two CW-complexes as topological spaces, their product might not be a CW-complex. However as objects in **CGWH** their product is a CW-complex. The reason behind this is that the general product doesn't produce the final topology; when it fails to be compactly generated it is strictly coarser than the CW-topology. Passing to the product in **CGWH**

(hence making the topology finer to become compactly generated) repairs this, and the product of CW-complexes is again a CW-complex.

**Remark 1.1.6.** From now on all topological spaces will be compactly generated weak Hausdorff. For such spaces the product  $X \times Y$  always means the product in **CGWH**. Proposition 1.1.5 (6) tells us: If one of the factors is also locally compact Hausdorff, the product in **CGWH** agrees with the classical product.

For the construction of topological K-Theory we will heavily rely on the two theorems from pointset topology:

**Theorem 1.1.7** (Tietze Extension Theorem). *Let  $X$  be a normal space (and hence the theorem is true in **CGWH**),  $Y \subseteq X$  a closed subspace,  $V$  a finite dimensional real vector space, and  $f : Y \rightarrow V$  a continuous map. Then there exists a continuous map  $g : X \rightarrow V$  such that  $g|_Y = f$ .*

**Theorem 1.1.8** (Existence of Partitions of Unity). *Let  $X$  be a compact space,  $\{U_i\}$  a finite open covering. Then there exist continuous maps  $f_i : X \rightarrow \mathbb{R}$  such that:*

1.  $f_i(x) \geq 0$  for all  $x \in X$ .
2.  $\text{supp}(f_i) \subseteq U_i$ .
3.  $\sum_i f_i(x) = 1$  for all  $x \in X$ .

*Such a collection  $\{f_i\}$  is called a **partition of unity**.*

## 1.2 Topological Pairs

**Definition 1.2.1** (Homotopy Extension Property). Let  $(X, A)$  be a topological pair, i.e.  $A \subseteq X$ . The pair satisfies the homotopy extension property if for any map  $f : X \rightarrow Y$  and any homotopy  $f_t : A \rightarrow Y$  with  $f_0 = f|_A$  there exists an extension  $\tilde{f}_t : X \rightarrow Y$  with  $\tilde{f}_t|_A = f_t$  and  $\tilde{f}_0 = f$ . The inclusion  $i : A \rightarrow X$  of such a pair is called a **cofibration**. We denote the category of pairs of compactly generated weak Hausdorff spaces where the inclusion is a cofibration by **CGWH**<sub>cof</sub><sup>2</sup>.

A more intuitive but (for  $A$  closed in  $X$ ) equivalent definition is:

**Definition 1.2.2.** If  $(X, A)$  is a topological pair,  $A$  is closed in  $X$  and there is a continuous function  $u : X \rightarrow I$  and a Homotopy  $H : X \times I \rightarrow X$  such that

1.  $H(x, 0) = x$  for all  $x \in X$ .
2.  $H(a, t) = a$  for all  $a \in A$  and  $t \in I$ .

Falsche  
Definition  
!!




Figure 2: The homotopy extension property for the pair  $(X, F)$ , where  $F$  gets rotated.

3.  $H(x, 1) \in A$  for all  $x \in X$  with  $u(x) < 1$ .
4.  $u^{-1}(0) = A$ .

Such a pair is called an **NDR**-pair which stands for neighbourhood deformation retract-pair. The name comes from the interpretation of  $u^{-1}[0, 1)$  being a neighbourhood that deforms onto  $A$  under the global homotopy  $H$ .

**Remark 1.2.3.** If we encounter an NDR pair in this thesis that lives in a manifold, we will often omit proving the property. A proof would mostly consist of applying the tubular neighbourhood theorem and the collar neighbourhood theorem.

lem:topPre-he  
pImpliescontr  
actingisHomot  
opicalInvaria  
nt

**Lemma 1.2.4.** *If the pair  $(X, A)$  satisfies the homotopy extension property and  $A$  is contractible, then the quotient map*

$$q : X \rightarrow X/A$$

*is a homotopy equivalence.*

*Proof.* Let  $f_t : X \rightarrow X$  be a homotopy extending the contraction of  $A$  with  $f_0 = \text{id}$ . Then the composition  $q \circ f_t$  maps  $A$  to a point and hence factors over  $X/A$  for all  $t$  giving us the commutative diagram

$$\begin{array}{ccc}
X & \xrightarrow{f_t} & X \\
\downarrow q & & \downarrow q \\
X/A & \xrightarrow{\overline{f_t}} & X/A
\end{array}$$

Here,  $\overline{f_t}$  denotes the induced map on the quotient. Now  $f_1(A) = pt$  where  $pt$  is a point in  $A$ . So  $f_1$  factors over  $X/A$  and denote the induced map by  $g: X/A \rightarrow X$ . Now  $g \circ q = f_1$  and hence

$$q \circ g(\overline{x}) = q \circ g \circ q(x) = q \circ f_1(x) = \overline{f_1}(\overline{x}).$$

In other words we have the commutative diagram

$$\begin{array}{ccc}
X & \xrightarrow{f_1} & X \\
\downarrow q & \nearrow g & \downarrow q \\
X/A & \xrightarrow{\overline{f_1}} & X/A
\end{array}$$

This concludes the proof, since  $g \circ q = f_1 \simeq f_0 = \text{id}$  and  $q \circ g = \overline{f_1} \simeq \overline{f_0} = \text{id}$ .  $\square$

**Definition 1.2.5.** We call a pointed topological space  $(X, x_0)$  well pointed, if the inclusion  $x_0 \rightarrow X$  is a cofibration. We denote the category of well pointed compactly generated weak Hausdorff spaces by  $\mathbf{CGWH}_{\text{cof}}^*$ .

### 1.3 Mapping Spaces

**Definition 1.3.1** (Compact-open topology). Let  $X, Y$  be topological spaces. Then the set of continuous maps  $C(X, Y)$  can be equipped with the topology that has

$$\{V(K, U) | K \subseteq X \text{ compact}, U \subseteq Y \text{ open}\} \text{ where } V(K, U) := \{f \in C(X, Y) | f(K) \subseteq U\}$$

as a subbasis. For real vector spaces  $V, W$  in the standard topology, this coincides with the topology of  $\text{Hom}(V, W)$  as a subset of  $\mathbb{R}^n$  after identification with the corresponding matrix.

**Proposition 1.3.2.** If  $X$  is compactly generated and  $Y$  compactly generated weak Hausdorff, then  $C(X, Y)$  is compactly generated weak Hausdorff.

**Proposition 1.3.3.** Let  $X, Y, Z$  be objects in  $\mathbf{CGWH}$ . The natural map

$$C(X, C(Y, Z)) \rightarrow C(X \times Y, Z)$$

is a homeomorphism.

prop:kTheo-cu  
rryingIsConti  
nuous

## 2 Vector Bundles and Riemannian Geometry

sec:background-geometry

Due to differential geometry and the theory of vector bundles being areas of mathematics used by pure mathematicians interested in analytical aspects like curvature, by topologists interested in invariants like Euler characteristics, physicists using the spaces as playgrounds for general relativity theories, and many others, there are numerous competing notational conventions. This thesis lies at the tripoint of differential geometry, dynamical systems, and algebraic topology. Hence, there is no a priori preferred notational convention. In this chapter, we establish the necessary background in the theory of vector bundles and Riemannian manifolds whilst not proving all the basics; we establish the notation used throughout this thesis. We establish manifolds and vector bundles in parallel, as they are interdependent on one another. We begin by defining a topological manifold. Afterwards, we equip that space with increasingly more structure and hence reduce the generality until we arrive at the playground of the thesis:

def:vb-topologicalManifold

**Definition 2.0.1** (Topological Manifold). A second countable Hausdorff space  $M$  is called a **topological manifold** of dimension  $m \in \mathbb{N}$ , if it is locally homeomorphic to  $\mathbb{R}^m$ . To be precise, for all  $p \in M$  there exists an open neighbourhood  $U \subseteq M$  of  $p$ , an open set  $V \subseteq \mathbb{R}^m$  and a map  $\varphi : U \rightarrow V$  that is a homeomorphism. We call the map  $\varphi : U \rightarrow V$  a **chart around  $p$  on  $M$** , and  $\varphi^{-1}$  a **local coordinate system around  $p$  on  $M$** .

def:vb-differentiableManifold

**Definition 2.0.2** (Differentiable Manifold). Let  $M$  be a topological manifold of dimension  $m$ .

1. A **differentiable atlas of class  $r \in \mathbb{N} \cup \{\infty\}$**  is a family of charts  $\mathfrak{A} = (\varphi_i : U_i \rightarrow V_i)_{i \in I}$  such that
  - a)  $\bigcup_{i \in I} U_i = M$ , meaning that  $(U_i)$  is an open covering of  $M$ .
  - b) For every pair  $(i, j) \in I^2$ , the **transition function**:

$$\begin{aligned} \varphi_{ij} : \varphi_j(U_i \cap U_j) &\rightarrow \varphi_i(U_i \cap U_j) \\ x &\mapsto (\varphi_i \circ \varphi_j^{-1})(x) \end{aligned}$$

is differentiable of class  $r$ .

We call such an atlas a  $C^r$ -atlas.

2. Two  $C^r$ -atlases  $\mathfrak{A}$  and  $\mathfrak{B}$  are called **equivalent** if the family  $\mathfrak{A} + \mathfrak{B} = (\varphi_i, \varphi_j)_{ij}$  is a  $C^r$ -atlas.

A **differentiable structure of class  $r$**  on  $M$  is an equivalence class  $c$  of  $C^r$ -atlases. For  $r = \infty$ , we call the pair  $(M, c)$  a smooth manifold.

**Remark 2.0.3.** The requirement of second countability together with Hausdorff ensures that we have partitions of unity subordinate to an open cover of the manifolds. In the smooth case we can assume the continuous maps of the partition to be smooth.

**Corollary 2.0.4.** Every transition function  $\varphi_{ij}$ ,  $i, j \in I^2$  of a differentiable atlas  $\mathfrak{A} = (\varphi_i)_{i \in I}$  is not merely a homeomorphism but also a diffeomorphism. This follows from the fact that  $\varphi_{ij}^{-1} = \varphi_{ji}$  which provides smooth inverses.

def:vb-smooth  
Functions

**Definition 2.0.5** (Smooth Functions). Let  $(M, c)$  be a differentiable manifold of class  $r$  and  $U \subseteq M$  open. We call a continuous function

$$f : U \rightarrow \mathbb{R}$$

**differentiable of class  $r$** , if for any chart (and hence for all)  $(\varphi_i)_{i \in I} = \mathfrak{A} \in c$  the compositions  $f \circ \varphi_i^{-1}$  are differentiable of class  $r$ . For  $r = \infty$  we define:

$$\mathcal{E}(U) = \{f : U \rightarrow \mathbb{R} \text{ continuous} \mid f \text{ is differentiable of class } \infty\}.$$

**Corollary 2.0.6.** Let  $(M, c)$  be a smooth manifold of dimension  $m$  and  $U \subseteq M$  be an open subset. With pointwise defined operations, the set  $(\mathcal{E}(U), +, \cdot, \circ)$  becomes an  $\mathbb{R}$ -algebra. Furthermore,  $\mathcal{E}$  becomes a sheaf of  $\mathbb{R}$ -algebras.

*Proof.* There is not really a need for a proof. However, it might help to work through the definition of a sheaf. First,  $\mathcal{E}$  is a presheaf, where the restriction in the domain of a function gives the needed restriction homomorphism:

$$\begin{aligned} \text{res}_V^U : \mathcal{E}(U) &\rightarrow \mathcal{E}(V) \\ f &\mapsto f|_V. \end{aligned}$$

The required properties of a presheaf are trivial. Furthermore, this gives a sheaf as the requirement of locality is trivial for functions and the property of gluing is also trivial for functions, since differentiability is a local property.  $\square$

**Definition 2.0.7.** If  $p \in M$  is fix,  $f \in \mathcal{E}(U)$  and  $g \in \mathcal{E}(U')$  such that  $p \in U \cap U'$  we say that  $f$  and  $g$  have the same **germ in  $p$** , if there is another open neighbourhood  $W \subseteq U \cap U'$  of  $p$  such that  $f|_W = g|_W$ . This defines an equivalence relation  $\sim_p$ . An equivalence class  $s$  of local functions around  $p$  is called a **germ in  $p$** . We write  $s = f_p$ , if  $s = [f]$  with  $f \in \mathcal{E}(U)$ . We write

$$\mathcal{E}_p(M) = \left( \sum_{U \text{ open}, p \in U} \mathcal{E}(U) \right) / \sim_p$$

for the set of germs, and call it the **stalk in  $p$** . Where  $\sum$  denotes the co-product (disjoint union) in **CGWH**.

**Corollary 2.0.8.** For a smooth manifold  $(M, c)$  the set  $\mathcal{E}_p(M)$  inherits an  $\mathbb{R}$ -algebra structure from the  $\mathcal{E}(U)$ . Furthermore, it carries a natural (evaluation-)homomorphism:

$$\begin{aligned} \text{eval}_p : \mathcal{E}_p(M) &\rightarrow \mathbb{R} \\ f_p &\mapsto f(p) =: f_p(p) \end{aligned}$$

The stalks are also local rings with maximal ideal  $\mathfrak{m}_p = \ker(\text{eval}_p)$ . Hence, the pair  $(M, \mathcal{E})$  gives us a locally ringed space.

*Proof.* We only need to verify the local ring structure of the stalks. An element  $f_p \in \mathcal{E}_p(M)$  is invertible if and only if  $f(p) \neq 0$ , equivalently if  $f_p \notin \ker(\text{eval}_p)$ . Thus  $\mathcal{E}_p(M)/\ker(\text{eval}_p)$  is a field, which proves that  $\ker(\text{eval}_p)$  is maximal.  $\square$

**Definition 2.0.9.** Let  $(M, c)$  be a smooth manifold of dimension  $m$  and  $p \in M$ . We call an  $\mathbb{R}$ -linear map  $\delta : \mathcal{E}_p(M) \rightarrow \mathbb{R}$  a **derivation**, if it satisfies the Leibniz rule:

$$\delta(f_p \cdot g_p) = \delta(f_p)g_p(p) + f_p(p)\delta(g_p) \quad \text{for all } f_p, g_p \in \mathcal{E}_p(M).$$

We call  $\text{Der}_{\mathbb{R}}(\mathcal{E}_p(M), \mathbb{R})$  the set of derivations and give it an  $\mathbb{R}$ -vector space structure by pointwise operations. We define the **tangent space of  $M$  at  $p$**  to be the vector space

$$T_p M := \text{Der}_{\mathbb{R}}(\mathcal{E}_p(M), \mathbb{R}).$$

**Corollary 2.0.10.** Let  $(M, c)$  be a smooth manifold of dimension  $m$  and  $\varphi : U \rightarrow V$  be a chart around  $p$  with  $x_0 = \varphi(p)$  (where  $\varphi \in \mathfrak{A} \in c$ ). Then we define the derivation

$$\begin{aligned} \partial_i(p) : \mathcal{E}_p(M) &\rightarrow \mathbb{R} \\ f_p &\mapsto \partial_i(p)(f_p) := \frac{\partial(f \circ \varphi^{-1})}{\partial x_i} \Big|_{x_0}. \end{aligned}$$

This is well-defined and hence gives a tangent vector. Moreover, the family

$$(\partial_1(p), \dots, \partial_m(p))$$

defines a basis of  $T_p M$ . Hence, the dimension of  $T_p M$  is  $m$ .

This is a good point to take a break from smooth manifolds and introduce the structure of vector bundles.

Analogously to the definition of a smooth manifold, we start by defining families of vector spaces and equip them with increasingly more structure:

def: vb-vector  
Bundles

**Definition 2.0.11** (Vector Bundles). Let  $\mathbb{K}$  be a field. A  $\mathbb{K}$  **vector bundle** of rank  $m$  is a surjection

$$\pi : E \rightarrow X$$

such that  $E$  and  $X$  are topological spaces and  $\pi$  is continuous. Furthermore  $\pi$  satisfies:

1. For all  $x \in X$ ,  $\pi^{-1}(x)$  carries the structure of an  $m$ -dimensional  $\mathbb{K}$ -vector space.
2. For all  $x \in X$ , there exist a neighbourhood  $U \subseteq X$  and a homeomorphism

$$\psi : E|_U := \pi^{-1}(U) \rightarrow U \times \mathbb{K}^m,$$

that is compatible in the following sense: Let  $\pi_1$  denote the projection  $U \times \mathbb{K}^m \rightarrow U$  onto the first factor. Then

$$\pi_1 \circ \psi = \pi$$

and

$$\psi_x : E_x := \pi^{-1}(x) \rightarrow \{x\} \times \mathbb{K}^m$$

is a vector space isomorphism.

We call  $E$  the **total space**,  $X$  the **base space**,  $E_x$  the **fibre over  $x$**  and  $\psi$  a **local trivialisation**.

**Remark 2.0.12.** In the context of differential geometry we will mostly consider  $\mathbb{R}$  vector bundles. However, the K-theory that we will develop will depend on  $\mathbb{C}$  vector bundles.

**Definition 2.0.13.** Let  $\pi : E \rightarrow X$  be a vector bundle. A continuous function  $s : X \rightarrow E$  such that  $\pi \circ s = \text{id}_X$  is called a (global) **section** (if  $s : U \subseteq X \rightarrow E$  we call it a local section). We denote the space of global sections by  $\Gamma(E)$  and by  $\Gamma(U)$  the space of local sections over an open subset  $U \subseteq X$ .

**Definition 2.0.14** (Homomorphism of Vector Bundles). A **homomorphism between two vector bundles**  $\pi : E \rightarrow X$  and  $\tilde{\pi} : F \rightarrow X$  over the same base space is a continuous function  $\varphi : E \rightarrow F$  such that:

$$1. \quad \begin{array}{ccc} E & \xrightarrow{\varphi} & F \\ \downarrow \pi & \swarrow \tilde{\pi} & \\ X & & \end{array} \quad \text{commutes.}$$

2. For all  $x \in X$  the function  $\varphi_x : E_x \rightarrow F_x$  is linear.

**Definition 2.0.15** (Pullback of Vector Bundles). Let  $\pi : E \rightarrow Y$  be a vector bundle,  $X$  be a topological space and  $f : X \rightarrow Y$  be continuous. Then we define the **pullback**  $f^*\pi : f^*(E) \rightarrow X$  as follows: The total space  $f^*(E)$  is defined as the subspace of  $(x, e) \in X \times E$  such that  $f(x) = \pi(e)$ , and the map  $f^*\pi$  is the projection to the second factor restricted to said subspace.

**Example 2.0.16.** Let  $\pi : E \rightarrow X$  be a vector bundle. For any  $Y \subseteq X$  we have the vector bundle

$$\pi|_Y : \pi^{-1}(Y) \rightarrow Y,$$

which is defined as the pullback along the inclusion of  $Y$  into  $X$ .

**Remark 2.0.17.** We will often omit the notation of the projection and denote a vector bundle by its total space. The restriction  $\pi|_Y : \pi^{-1}(Y) \rightarrow Y$  is often denoted by  $E|_Y$ . The restriction may equivalently be defined by restricting the domain and codomain of the projection.

**Remark 2.0.18.** Notice that this construction of the pullback  $f^*\pi : f^*(E) \rightarrow X$  is the simplest space to make the square commute:

$$\begin{array}{ccc} f^*(E) & \xrightarrow{\pi_2} & E \\ \downarrow f^*\pi & & \downarrow \pi \\ X & \xrightarrow{f} & Y \end{array}$$

The notion of simplicity is captured by the universal property, that for any other vector bundle  $\tilde{\pi} : \tilde{E} \rightarrow X$  and function  $\tilde{f} : \tilde{E} \rightarrow E$  there exists a unique  $h : \tilde{E} \rightarrow f^*(E)$  (which is given by  $h(y) := (\tilde{\pi}(y), \tilde{f}(y))$ ) such that the following diagram commutes:

$$\begin{array}{ccccc}
\tilde{E} & & \tilde{f} & & \\
\downarrow h & \searrow & & \searrow & \\
f^*(E) & \xrightarrow{\pi_2} & E & & \\
\downarrow f^*\pi & & \downarrow \pi & & \\
X & \xrightarrow{f} & Y & & 
\end{array}$$

(Note: There is also a curved arrow from  $\tilde{E}$  to  $X$  labeled  $\tilde{\pi}$ .)

**Definition 2.0.19** (Morphism of Vector Bundles over Different Base Spaces). Let  $\pi : E \rightarrow X$  and  $\pi' : E' \rightarrow X'$  be two vector bundles and  $f : X \rightarrow Y$  continuous. A **bundle morphism over  $f$**  is a map continuous map  $\Phi : E \rightarrow E'$  such that

$$f \circ \pi = \pi' \circ \Phi$$

and for all  $x$ , the restriction  $\varphi_x : E_x \rightarrow E'_{f(x)}$  is a linear map.

**Remark 2.0.20.** Notice how a bundle morphism  $\varphi$  always factors over the pullback  $f^*(E')$ . Furthermore, if  $\varphi_x$  is an isomorphism for all  $x$ , then  $E \cong f^*(E')$ .

**Remark 2.0.21.** In the context of K-Theory, bundle maps will always be homomorphisms between two vector bundles over the same base space. Hence, we omit the *over the same base space*. If a bundle map appears that is not over the same base space, we will note that explicitly.

**Remark 2.0.22.** Let  $\pi : E \rightarrow X$  be a vector bundle. Then the space of sections  $\Gamma(E)$  is a (in general, infinite-dimensional)  $\mathbb{R}$ -vector space. Furthermore, and more relevant to the purposes of this thesis, we view  $\Gamma(E)$  (and  $\Gamma(U)$  for open subsets) as a  $C(X)$  (or  $C(U)$ ) module. The latter denotes the ring of continuous functions from  $E$  (or  $U$ ) to  $\mathbb{R}$ .

**Example 2.0.23.** For any smooth manifold  $M$  of dimension  $m$ , we can equip the union

$$TM := \bigsqcup_{p \in M} T_p M$$

with a topology, such that  $\pi : TM \rightarrow M$  defines a vector bundle which we call **tangent bundle**. For this we first define the trivialisations and then equip  $TM$  with a topology to make them homeomorphisms: Let  $\varphi : U \rightarrow V$  be a chart on  $M$ . If we denote the  $i$ -th coordinate function of  $\varphi$  by  $\varphi_i$  we can define the bijection

$$\tilde{\varphi} : \pi^{-1}(U) \rightarrow V \times \mathbb{R}^m, \quad \xi_p \mapsto \left( \varphi(p), \begin{pmatrix} \xi_p(\varphi_1) \\ \vdots \\ \xi_p(\varphi_m) \end{pmatrix} \right).$$

Now we declare a subset  $W \subseteq TM$  to be open, if and only if for all charts  $\varphi$  the set  $\tilde{\varphi}(W \cap \pi^{-1}(U)) \subset V \times \mathbb{R}^m$  is open. We can now extend and give meaning to the notation of the local basis vectors  $\partial_i(p) \in T_p M$ : For any chart  $\varphi : U \rightarrow V$  we have the local sections

$$\partial_i : U \rightarrow TU, \quad p \mapsto \partial_i(p).$$

Furthermore,  $TM$  is a  $2m$ -dimensional manifold. Using a smooth atlas on  $M$  yields a smooth atlas on  $TM$  making it a smooth manifold. Notice that the family  $(\partial_i)_i$  is a basis of the  $C(U)$ -module  $\Gamma(U)$ . This follows by expressing any section in local coordinates via  $\tilde{\varphi}$ .

Notice that  $TM$  as a bundle satisfies more than just the bundle definition: Taking two different charts induces two different local trivialisations. Transitioning between them is not just fibrewise linear, but also smooth. This is captured in the definition:

Henceforth we suppress the differentiable structure from the notation.

**Definition 2.0.24** (The Derivative). Let  $M$  and  $M'$  be smooth manifolds. We call a continuous function  $f : M \rightarrow M'$  **smooth**, if for all  $\varphi \in \mathfrak{A} \in \mathcal{C}$  and  $\varphi' \in \mathfrak{A}' \in \mathcal{C}'$  the maps

$$\varphi' \circ f \circ \varphi^{-1} : V \rightarrow V'$$

are smooth. A given smooth function induces a smooth map between the tangent bundles as manifolds as follows:

$$df : TM \rightarrow TM', df_p(\xi_p)(g_p) = \xi((g \circ f)_p)$$

Here,  $\xi_p \in T_p M$ ,  $g_p \in \mathcal{E}_p(M')$ . The function  $df$  is called the **derivative** or **pushforward** of  $f$ .

**Remark 2.0.25.** Notice how the pushforward  $df$  is a bundle morphism over  $f$  with the base spaces being  $M$  and  $M'$ .

**Definition 2.0.26** (Smooth Vector Bundles). A **smooth vector bundle of rank  $r$**  is a surjection  $\pi : E \rightarrow M$  where the codomain (or base space) is a smooth manifold together with a family of continuous functions (or local trivialisations)

$$\Phi_i : \pi^{-1}(U_i) \rightarrow U_i \times \mathbb{R}^r$$

such that  $\bigcup_i U_i = M$  and for all overlapping  $U_i \cap U_j \neq \emptyset$ , the transition functions

$$\Phi_j \circ \Phi_i^{-1} : \pi^{-1}(U_i \cap U_j) \rightarrow (U_i \cap U_j) \times \mathbb{R}^r$$

are of the form

$$\Phi_j \circ \Phi_i^{-1}(p, v) = (p, g_{i,j}(v)), \quad \text{where} \quad g_{i,j} : (U_i \cap U_j) \rightarrow \text{GL}(r, \mathbb{R})$$

is smooth.

**Corollary 2.0.27.** A smooth vector bundle is a smooth manifold and also a vector bundle. The latter is ensured by the codomain of  $g_{i,j}$  which is also called the **structure group**. We can enforce structure on the bundle by reducing the structure group. As a preview: reducing it to  $\text{GL}^+(k, \mathbb{R})$  the linear maps with positive determinant means that we can induce the concept of orientations. Reducing it to  $\text{O}(k)$  lets us equip it with a Riemannian metric.




Figure 3: A smooth vector bundle (If we ignore the face and feet and assume the hedgehogs spines to be infinitely tight and long)

def:vb-frames  
FromTrivialis  
ations

**Definition 2.0.28.** Let  $\pi : E \rightarrow M$  be a smooth vector bundle of rank  $r$  and let

$$\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^r$$

be a local trivialisation. Then a section is called smooth if it is a smooth map of manifolds. A family of smooth local sections  $(s_1, \dots, s_r)$  with  $s_i \in \Gamma(U)$  is called a **local frame** if they form a basis of the  $C(U)$ -module  $\Gamma(U)$ . Such a family is obtained from  $\Phi$  by defining  $s_i(p) := \Phi^{-1}(p, e_i)$ . Similarly, from a local frame we can obtain a local trivialisation by defining:

$$\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^r, \quad v = \sum_{i=1}^r \lambda_i s_i(\pi(v)) \mapsto (\pi(v), \lambda_1, \dots, \lambda_r).$$

**Example 2.0.29.** The local trivialisations  $\partial_i$  of the tangent bundle of a smooth manifold  $M$  form a local frame and make the tangent bundle into a smooth vector bundle.

def:vb-homomo  
rphismBundle

**Definition 2.0.30** (Homomorphism Bundle). Let  $W, V$  be vector spaces and  $E = V \times X$ ,  $F = W \times X$  the product bundles over  $X$ . Now any bundle homomorphism  $\varphi : E \rightarrow F$  determines a map  $\Phi : X \rightarrow \text{hom}(V, W)$  by sending  $x \mapsto \varphi_x$ . With respect to the compact-open topology, this  $\Phi$  is continuous. Furthermore, a continuous map  $\Phi : X \rightarrow \text{hom}(V, W)$  determines a bundle homomorphism  $\varphi : E \rightarrow F$ .

*Proof.* Let  $\{e_i\}$  and  $\{f_i\}$  be bases of  $V$  and  $W$ . Then each  $\Phi(x)$  is a matrix  $\Phi_{i,j}(x)$ , such that:

$$\Phi(x)e_i = \sum_j \Phi_{i,j}(x)f_j.$$

Now  $\Phi$  is continuous if and only if  $\Phi_{i,j}$  is continuous. Since  $\varphi(x, v) = (x, \Phi(x)v)$  this is equivalent to  $\varphi$  being continuous.  $\square$

thm:vb-isomorphicVectorBundles

**Lemma 2.0.31** (Isomorphic Vector Bundles). *Let  $E$  and  $F$  be vector bundles and  $\varphi : E \rightarrow F$  be a homomorphism. Then  $\varphi$  is an isomorphism if and only if all  $\varphi_x$  are isomorphisms.*

*Proof.* To show that it is an isomorphism we need a continuous inverse, equivalently, we need to show that  $\varphi$  is a homeomorphism. First of all we can deduce, that  $\varphi$  is bijective: Assume it wasn't injective, then there would be a fibre in which  $\varphi_x$  is not injective. Assume that it wasn't surjective. Then there would be a  $w \in F$  that has no preimage which means that in the fibre containing  $w$  there is no isomorphism. Thus, there is a unique inverse and it remains to verify that it is continuous, which is a local condition. Therefore we can assume that  $E$  and  $F$  are product bundles. Then we have a continuous map  $\Phi : X \rightarrow \text{Hom}(V, W)$ . Furthermore, since all  $\varphi_x$  are isomorphisms, we can restrict the image to  $\text{Iso}(V, W)$  which is an open subset of  $\text{Hom}(V, W)$ . Now we can define the map  $x \mapsto \Phi(x)^{-1}$  which is continuous since taking inverses is continuous. By the corollary above, we obtain a continuous map  $\psi : F \rightarrow E$  by sending  $(x, w) \mapsto (x, \Phi(x)^{-1}(w))$ , which is the inverse.  $\square$

**Example 2.0.32.** Let  $M$  be a smooth manifold of dimension  $m$ ,  $\varphi : U \rightarrow V$  a chart and  $\tilde{\varphi} : TU \rightarrow V \times \mathbb{R}^m$  be the corresponding bundle chart. Then it follows from the lemma, that  $TU$  is isomorphic to  $V \times \mathbb{R}^m$ . This is verified by observing that  $\partial_i(p)$  is a basis of  $T_pU$  for all  $p \in U$ . In this case  $\tilde{\varphi}(\partial_i) = e_i$  giving us isomorphisms in each fibre.

def:vb-subBundles

**Definition 2.0.33** (Sub-bundles). Let  $\pi : E \rightarrow X$  be a vector bundle. A **sub-bundle** is a subset of  $E$  which is a vector bundle in the induced topology and fibrewise vector space structure.

def:vb-monoAndEpi

**Definition 2.0.34** (Mono- and Epimorphisms). A homomorphism  $\varphi : E \rightarrow F$  is a **monomorphism** if all  $\varphi_x$  are monomorphisms (i.e. injective). Respectively, it is called an **epimorphism** if all  $\varphi_x$  are epimorphisms (i.e. surjective).

**Corollary 2.0.35.** If  $F$  is a sub-bundle of  $E$ , then the inclusion  $\varphi : F \rightarrow E$  is a monomorphism.

lem:vb-monoInduceSubBundles

**Lemma 2.0.36.** *If  $\varphi : E \rightarrow F$  is a monomorphism, then  $\varphi(E)$  is a sub-bundle and  $\varphi : E \rightarrow \varphi(E)$  is an isomorphism.*

*Proof.* Once we know that  $\varphi(E)$  is a sub-bundle, the map  $\varphi : E \rightarrow \varphi(E)$  is a bijective homomorphism and fibrewise an isomorphism, hence an isomorphism by Lemma 2.0.31. So we need to check that  $\varphi(E)$  is a sub-bundle, which is a local property. Thereby we can assume that  $E$  and  $F$  are product bundles  $E = X \times V$  and  $F = X \times W$ . Let  $x \in X$  and let  $W_x \subseteq W$  be a subspace complementary to  $\varphi_x(V)$ . Then  $G := X \times W_x \subseteq F$  and we define

$$\begin{aligned} \theta : E \oplus G &\rightarrow F \\ (a, b) &\mapsto \varphi(a) + i(b) \quad \text{where } i \text{ denotes the inclusion } G \hookrightarrow F. \end{aligned}$$

In the fibre over  $x$ , the map  $\theta_x$  is an isomorphism and thereby there exists an open neighbourhood  $U$  of  $x$  such that  $\theta|_U$  is an isomorphism. That is because  $\text{Iso}(V \oplus W_x, W)$  is open in  $\text{Hom}(V \oplus W_x, W)$ . Since  $E$  is a sub-bundle of  $E \oplus G$  we have that  $\theta(E)|_U = \varphi(E)|_U$  is a sub-bundle of  $F|_U$ .  $\square$

cor:vb-subBundlesAreLocalSummands

**Corollary 2.0.37.** In the proof, we have shown more: locally a sub-bundle is a direct summand. Furthermore, we have shown that the set of points  $x$  where  $\varphi_x$  is a monomorphism, is open.

def:vb-quotientBundle

**Definition 2.0.38** (Quotient Bundle). If  $F$  is a sub-bundle of  $E$  we define the **quotient bundle**  $E/F$  as the union of the spaces  $E_x/F_x$  equipped with the quotient topology.

**Corollary 2.0.39.** The quotient bundle is a vector bundle, because locally  $F$  is a direct summand and thereby  $E/F$  is locally trivial.

**Lemma 2.0.40.** *Quotient bundles of smooth subbundles are smooth.*

lem:vb-quotientBundlesOfSmoothSubbundlesAreSmooth

*Proof.* Let  $F \subseteq E$  be a smooth subbundle of rank  $k$ , where  $E$  has rank  $n$  and  $M$  is the base manifold. We need to check that the transition functions consist of smooth maps into the structure group. First, we will show that there are smooth local frames of  $E$  that restricts to a smooth local frame of  $F$ :

Let  $p \in M$ . Since  $F$  is smooth there exists a local frame  $(s_1, \dots, s_k)$  of  $F$  around  $p$ . In particular, the  $s_i$  are smooth local sections into  $E$  around  $p$ . Let  $(t_1, \dots, t_k)$  be a smooth local frame of  $F$  near  $p$ . (without restrictions over the same domain as the sections  $s_i$ ) Now we make use of steinitz exchange lemma to choose  $n - k$  many  $t_i$  (Without restrictions  $t_{k+1}, \dots, t_n$ ) such that

$$(s_1(p), \dots, s_k(p), t_{k+1}(p), \dots, t_n(p))$$

is a basis of  $E_p$ . Being a basis over a point  $q \in M$  near  $p$  is equivalent to the invertability of the matrix where the columns are the sections at  $q$ , and since invertability (or rather nonzero determinant) is an open condition, there exists a neighbourhood, such that the family

$$(s_1, \dots, s_k, t_{k+1}, \dots, t_n)$$

is a local frame, whose first  $k$  sections span  $F$ . By Definition 2.0.28 it induces a smooth trivialisation  $\Phi_U$  of  $E$  with  $\Phi_U(F|_U) = U \times \mathbb{R}^k \times \{0\}$ , and fibrewise passing to the quotient  $\mathbb{R}^n/(\mathbb{R}^k \times \{0\}) \cong \mathbb{R}^{n-k}$  yields a trivialisation  $\bar{\Phi}_U$  of  $(E/F)|_U$ .

Now we can cover  $M$  by such adapted trivializations. Their transition functions  $g_{i,j} : U_i \cap U_j \rightarrow \text{GL}(n, \mathbb{R})$  are smooth and pointwise preserve  $\mathbb{R}^k \times \{0\}$ : for  $v \in \mathbb{R}^k \times \{0\}$  we have  $\Phi_i^{-1}(p, v) \in F_p$  since  $\Phi_i$  is adapted, and hence  $\Phi_j(\Phi_i^{-1}(p, v)) \in \{p\} \times \mathbb{R}^k \times \{0\}$  since  $\Phi_j$  is adapted. Hence they have block form

$$g_{i,j} = \begin{pmatrix} A_{i,j} & B_{i,j} \\ 0 & C_{i,j} \end{pmatrix},$$

where all coordinate entries and hence all blocks smoothly depend on the basepoint. Finally, the transition functions of the quotient with respect to the trivialization  $\bar{\Phi}_U$  of  $(E/F)|_U$  consist of the smooth maps  $p \mapsto C(p) = (C_{i,j})_{i,j}(p)$  into the structure group  $\text{GL}(n-k, \mathbb{R})$ .  $\square$

Now we want to construct new bundles from old ones. There are many operations to construct new vector spaces such as the tensor product, direct sum, Hom-Spaces and many more. Instead of individually proving that those operations applied fibrewise yield new bundles, we subsume these operations within the following definition:

**Definition 2.0.41** (Continuous Functor). Let  $T$  be a covariant endofunctor in the category of finite-dimensional vector spaces. We call  $T$  a **continuous functor**, if for all  $V, W$  the map  $T : \text{hom}(V, W) \rightarrow \text{hom}(T(V), T(W))$  is continuous. All vector spaces are given the topology induced by their isomorphism to real Euclidean space. If  $E$  is a vector bundle we define the set:

$$T(E) := \bigcup_{x \in X} T(E_x).$$

Now we want to equip this with a topology such that  $T(E)$  is a bundle over the same base space as  $E$ .

def: vb-constructingNewBundles

**Definition 2.0.42** (Constructing new bundles). We start by assuming that  $E = X \times V$ . In that case we define the topology on  $T(E)$  as the product topology  $X \times T(V)$ . Now suppose that  $F = X \times W$  is another product bundle and  $f : E \rightarrow F$  is a homomorphism. Let  $\Phi : X \rightarrow \text{hom}(V, W)$  be the corresponding map. Then  $T\Phi : X \rightarrow \text{hom}(T(V), T(W))$  is continuous by assumption. Thereby,  $T(\varphi) : X \times T(V) \rightarrow X \times T(W)$  is continuous. If  $\varphi$  is an isomorphism, then  $T(\varphi)$  is also an isomorphism. That follows because it is continuous and an isomorphism fibrewise.

Now assume that  $E$  is trivial. Then we can choose an isomorphism  $\alpha : E \rightarrow X \times V$  and induce a topology on  $T(E)$  by pulling it back via  $T(\alpha) : T(E) \rightarrow X \times T(V)$ . This topology does not depend on  $\alpha$  by the above. For general vector bundles we do this construction locally.

If  $f : X \rightarrow Y$  is continuous, there is a natural isomorphism

$$f^*T(E) \cong Tf^*(E)$$

This construction is carried out in detail by Atiyah in [Ati89, Chapter 1.2] and can be generalised for continuous functors with several variables both co- and contravariant. Hence this procedure gives us for vector bundles  $E$  and  $F$  over the same base space the new bundles:

- (i)  $E \oplus F$ , their direct sum
- (ii)  $E \otimes F$ , their tensor product
- (iii)  $\text{Hom}(E, F)$
- (iv)  $E^*$ , the dual bundle of  $E$
- (v)  $\Lambda^i(E)$  the  $i$ -th exterior power

and by the natural isomorphism we have natural isomorphisms:

- (i)  $E \oplus F \cong F \oplus E$
- (ii)  $E \otimes F \cong F \otimes E$
- (iii)  $E \otimes (F \oplus F') \cong E \otimes F \oplus E \otimes F'$
- (iv)  $\text{Hom}(E, F) \cong E^* \otimes F$

cor:vb-sectionsIntoHom

**Corollary 2.0.43.** Let  $\pi_E : E \rightarrow X$  and  $\pi_F : F \rightarrow X$  be two vector bundles over  $X$ . Then, the space  $\Gamma(\text{Hom}(E, F))$  can be identified with the space of bundle morphisms  $\text{hom}(E, F)$  since for any section  $s : X \rightarrow \text{Hom}(E, F)$  we can define the bundle homomorphism  $\tilde{s} : E \rightarrow F$  by sending  $v \mapsto s_{\pi_E(v)}(v)$ . This is a homomorphism since  $\pi_F(\tilde{s}(v)) = \pi_E(v)$ .

**Example 2.0.44** (Cotangent bundle). Let  $M$  be a smooth manifold of dimension  $m$ . Then we can define the **cotangent bundle**  $T^*M$  to be the dual of the tangent bundle. A local frame of said bundle is given by the fibrewise dual of the frame  $(\partial_i)$ . Sections into the tangent bundle are called **vector fields** and sections into the cotangent bundle are called **one-forms**.

def:vb-linearIdentification

**Definition 2.0.45.** Let  $V$  be a linear manifold (i.e. a finite-dimensional vector space). Then for any  $v \in V$  we have the canonical maps

$$R_v : V \rightarrow T_v V$$

$$x \mapsto \left( f_v \mapsto \frac{\partial}{\partial t} \Big|_{t=0} f(v + tx) \right)$$

Let  $L : V \rightarrow V$  be a linear map viewed as a smooth map between manifolds. Then we have the identity:

$$dL_v \circ R_v = R_{L(v)} \circ L.$$

This is proved by a short calculation:

$$\left( dL_v(R_v(x)) \right) (f_{L(v)}) = R_v(x) (f \circ L) = \frac{\partial}{\partial t} \Big|_{t=0} f \circ L(v + tx) = \underbrace{\frac{\partial}{\partial t} \Big|_{t=0} f(L(v) + tL(x))}_{=R_{L(v)} \circ L(x)}$$

Now let  $v, v' \in V$ . Then we have the translation

$$tr_{v,v'} : V \rightarrow V; \quad x \mapsto (x + (v' - v))$$

This allows us to relate  $R_v$  and  $R_{v'}$  as follows:

$$\left( dtr_{v,v'}|_v (R_v(x)) \right) (f_{v'}) = \frac{\partial}{\partial t} \Big|_{t=0} (f \circ tr_{v,v'}(v + tx)) = \frac{\partial}{\partial t} \Big|_{t=0} (f(v' + tx)) = R_{v'}(x)(f_{v'}).$$

Hence:

$$R_{v'} = dtr_{v,v'} \circ R_v.$$

The **linear identification** is the map  $Q$  given on the whole tangent bundle:

$$Q : TV \rightarrow V; \quad \xi_p \mapsto R_p^{-1}(\xi_p),$$

and satisfies for any linear  $L : V \rightarrow V$

$$Q \circ dL = L \circ Q.$$

cor: vb-vector  
FieldsTakeFun  
ctions

**Corollary 2.0.46.** Let  $f : M \rightarrow \mathbb{R}$  be smooth. Then we can interpret  $df$  as a one-form. To be precise, let  $Q : T\mathbb{R} \rightarrow \mathbb{R}$  denote the linear identification with respect to the identity on  $\mathbb{R}$ . For  $v \in \Gamma(TM)$  we have

$$(Q \circ df)(v) = v(f)$$

Here on the right side,  $f$  denotes a map  $p \mapsto f_p$  such that  $v(f)(p) := v_p(f_p)$  is well defined. We retain this notation so that vector fields can take functions as an input.

*Proof.* Let  $\varphi : U \rightarrow V$  be a chart of  $M$ . We prove this by showing it for the local sections  $\partial_i$  and the general case follows by  $C(U)$ -linearity, since those sections form a local frame. Now let  $g_p \in \mathcal{E}_p(\mathbb{R})$  and  $\varphi(p) = x_0$ . Then

$$df_p(\partial_i(p))(g_p) = \partial_i((g \circ f)_p) = \frac{\partial}{\partial x_i} \Big|_{x_0} (g \circ f \circ \varphi^{-1}) = \frac{\partial}{\partial x_{f_p(p)}} (g_p) \cdot \partial_i(p)(f_p)$$

Hence, applying  $Q$  to the calculation yields  $(Q \circ df)(v) = v(f)$ .  $\square$

**Definition 2.0.47** (A Metric). Let  $M$  be a smooth manifold. A section  $g \in \Gamma(T^*M \otimes T^*M)$  into the tensor product of the dual tangent bundle with itself is a **Riemannian metric** if the following are satisfied for all  $p \in M$ :

1.  $g$  is **non-degenerate**, meaning if for a  $v_p \in T_p M$  we have  $g_p(v_p, w_p) = 0$  for all  $w_p \in T_p M$  then  $v_p = 0$ .
2.  $g$  is **symmetric**, meaning  $g_p(v_p, w_p) = g_p(w_p, v_p)$  for all  $v_p, w_p \in T_p M$ .
3.  $g$  is **positive definite**, meaning that  $g_p(v_p, v_p) \geq 0$  for all  $v_p$  with equality only for  $v_p = 0$ .

We call a tuple  $(M, g)$  a **Riemannian manifold**.

**Remark 2.0.48.** Equipping a smooth manifold  $M$  of dimension  $m$  with a Riemannian metric is equivalent to reducing the structure group of the tangent bundle from  $GL(m)$  to  $O(m)$ . To be precise: A Riemannian metric  $g$  lets us canonically choose an atlas with structure group  $O(m)$  and an atlas with such a structure group lets us canonically choose a Riemannian metric.

*Proof.* We start by assuming that we have an atlas with structure group  $O(m)$ . In Definition 2.0.28 we saw that a covering of local trivialisations is equivalent to a covering of smooth frames. Now in a single trivialisation

$$\Phi : \pi^{-1}(U) \rightarrow TU \rightarrow U \times \mathbb{R}^m$$

we call the projection to the  $\mathbb{R}^m$ -part (the second factor)  $\pi_2$  and define

$$\begin{aligned} g : U &\rightarrow T^*U \otimes T^*U \\ p &\mapsto g_p^\Phi : T_p U \times T_p U \rightarrow \mathbb{R} \\ &\quad (\nu_p, \xi_p) \mapsto g_p^\Phi(\nu_p, \xi_p) := \pi_2(\Phi(\nu_p)) \cdot \pi_2(\Phi(\xi_p)), \end{aligned}$$

where  $\cdot$  denotes the standard scalar product. This gives us locally a Riemannian metric. If we do this for all trivialisations we have to make sure, that the result is well defined. So assume that two trivialisations  $\Phi$  and  $\Psi$  overlap in  $V$ . The structure group is  $O(m)$ ,  $\Phi = R\Psi$  for some function

$$R : V \rightarrow O(m).$$

Let  $p$  be in said overlapping and  $(\nu_p, \xi_p)$  be from  $T_p M \times T_p M$ . Then since  $R$  is orthogonal we can calculate the independence from the trivialisation:

$$g_p^\Phi(\nu_p, \xi_p) = \pi_2(R\Psi(\nu_p)) \cdot \pi_2(R\Psi(\xi_p)) = \pi_2(\Psi(\nu_p)) \cdot \pi_2(\Psi(\xi_p)) = g_p^\Psi(\nu_p, \xi_p).$$

To get an orthonormal atlas from a smooth one given a Riemannian metric, we start by building a covering of local frames from the trivialisations. We then apply Gram–Schmidt to the local frames to get orthonormal frames. Now a covering of orthonormal frames gives a covering of trivialisations, where the structure group consists of basis changes between orthonormal bases and hence is  $O(m)$ .  $\square$

**Definition 2.0.49** (The Gradient). Assume that  $(M, g)$  is a smooth manifold together with a Riemannian metric. Let  $f : M \rightarrow \mathbb{R}$  be a smooth function and  $Q : T\mathbb{R} \rightarrow \mathbb{R}$  the

linear identification. We define its **gradient** to be a vector field  $\text{grad}(f) \in \Gamma(TM)$  such that for any vector field  $V$ :

$$Q \circ df(V) = g(\text{grad}(f), V).$$

Notice how the non-degeneracy of  $g$  ensures existence and uniqueness.

def:vb-localD  
escriptionOfG  
radientGij

**Definition 2.0.50.** Let  $(M, g)$  be a Riemannian manifold and  $\varphi : U \rightarrow V$  be a chart. We define the smooth functions  $g_{ij} : U \rightarrow \mathbb{R}$  to be:

$$g_{ij}(p) = g_p(\partial_i(p), \partial_j(p)).$$

Then if two vector fields  $v, w$  over  $U$  are given with  $v = \sum v_i \partial_i$  and  $w = \sum w_i \partial_i$  we can calculate the metric locally:

$$g(v, w) = \sum_{i,j} g_{ij} v_i w_j : U \rightarrow \mathbb{R}$$

Furthermore, we can invert the matrices  $(g_{ij}(p))_{ij}$  for all  $p$  (which is a smooth procedure as can be seen by the construction via Cramer's rule) to get smooth functions

$$g^{ij} : U \rightarrow \mathbb{R}$$

$$p \mapsto g^{ij}(p) = \left( (g_{kl}(p))_{kl}^{-1} \right)_{ij}$$

**Remark 2.0.51.** Due to the rare appearance of big summations with vectors and co-vectors in this work, we omit the Einstein summation convention and assign all coordinate entries lower indices.

lem:vb-localD  
escriptionOfG  
radient

**Lemma 2.0.52** (Local Description of the Gradient). *Let  $(M, g)$  be a smooth Riemannian manifold of dimension  $m$  and  $\varphi : U \rightarrow V$  be a chart. Then the gradient has the local form:*

$$\text{grad}(f) = \sum_{i,j} g^{ij} (\partial_i(f)) \cdot \partial_j.$$

*Proof.* Assume that  $v, w \in \Gamma(TU)$  such that  $v = \sum_i v_i \partial_i$  and  $w = \sum_i w_i \partial_i$  (Here  $v_i$  and  $w_i$  are smooth functions  $U \rightarrow \mathbb{R}$ ). Then  $g(v, w) = \sum_{i,j} g_{ij} v_i w_j$ . Suppose that  $\text{grad}(f) = \sum_j G^j \partial_j$  and  $Q$  is the linear identification  $T\mathbb{R} \rightarrow \mathbb{R}$  in each tangent space. Then by definition of the gradient and corollary 2.0.46 we have:

$$\partial_j(f) = Q \circ df(\partial_j) = g(\text{grad}(f), \partial_j) = g\left(\sum_i G^i \partial_i, \partial_j\right) = \sum_i g_{ij} G^i$$

Reading this as a pointwise matrix product:

$$(G^1, \dots, G^m)(g_{ij}) = (\partial_1(f), \dots, \partial_m(f))$$

$$\text{which gives } (G^1, \dots, G^m) = (\partial_1(f), \dots, \partial_m(f)) (g^{ij}).$$

But then we can conclude that  $G^j = \sum_i g^{ij} \partial_i(f)$ .

□




Figure 4: A dynamical system with a critical point created by printing a flow of inc.

**Definition 2.0.53** (A Dynamical System). Let  $M$  be a smooth manifold. A **dynamical system on  $M$**  (sometimes also called a **flow on  $M$** ) is given by a smooth map  $\varphi : \Omega \rightarrow M$  such that:

1.  $\{0\} \times M \subseteq \Omega \subseteq \mathbb{R} \times M$  open, and for any  $p \in M$  the set  $I(p) := \pi_1(\Omega \cap \mathbb{R} \times \{p\})$  is connected. ( $\pi_1$  is the projection onto the first factor)
2. For all  $p \in M$  and  $t \in I(p)$  we have that  $s \in I(\varphi(t, p))$  if and only if  $s + t \in I(p)$ . Then we have:

$$\varphi(s, (\varphi(t, p))) = \varphi(s + t, p)$$

3. For all  $p$  we require  $\varphi(0, p) = p$ .

We will write  $\varphi_t(p) := \varphi(t, p)$  to keep the notation manageable. If for all  $p$   $I(p) = \mathbb{R}$  we call the system **global**.

**Corollary 2.0.54.** For a global dynamical system  $\varphi$  on a smooth manifold  $M$  the map

$$\rho : \mathbb{R} \rightarrow \text{Diff}(M), \quad t \mapsto \varphi_t$$

is a homomorphism. To verify well-definedness notice how for all  $t \in \mathbb{R}$ ,  $\varphi_{-t}$  is an inverse of  $\varphi_t$ . The map  $\rho$  is called a **one-parameter group of diffeomorphisms**.

**Definition 2.0.55** (Vector Field of a Dynamical System). Let  $\varphi : \Omega \rightarrow M$  be a dynamical system on a smooth manifold  $M$ . For a chart  $\psi : U \rightarrow V$  of  $M$  we have the chart  $\text{id} \times \psi$  of  $\Omega$ . For this chart we denote the associated local frame by  $(\partial_t, \partial_1, \dots, \partial_m)$ . Using this we define the mapping

$$\begin{aligned} M &\rightarrow TM \\ p &\mapsto X(p) := d\varphi(\partial_t(0, p)) \end{aligned}$$

and call it the **associated vector field**. This is in fact a smooth vector field.

**Definition 2.0.56** (Dynamical System of a Vector Field). Let  $X \in \Gamma(TM)$  be a global vector field on the smooth manifold  $M$ . Then we call a dynamical system  $\varphi$  on  $M$  the **associated dynamical system** to  $X$ , if for all  $t \in \mathbb{R}$  and  $x \in M$

$$d\varphi(\partial_t(t, x)) = X(\varphi_t(x)).$$

**Remark 2.0.57.** Regarding the left-hand side of the definition  $d\varphi(\partial_t(t, x))$  we could also differentiate the map

$$t \mapsto \varphi_t(x),$$

for fixed  $x$  at  $t$  and evaluate this at the constant vector field  $t \mapsto R_t(1) \in T_t\mathbb{R}$  from definition 2.0.45. The condition on the dynamical system reads then

$$\frac{d}{dt}\varphi_t(x)(R_t(1)) = X(\varphi_t(x))$$

**Definition 2.0.58.** Let  $X \in \Gamma(TM)$  be a global vector field on the smooth manifold  $M$  and denote the associated flow by  $\varphi$ . Then we call  $q \in M$  a **critical point of  $X$** , if  $X(q) = 0$ .

prop:vb-onceC  
riticalAlways  
Critical

**Proposition 2.0.59.** Let  $X \in \Gamma(TM)$  be a global vector field on the smooth manifold  $M$  and denote the associated flow by  $\varphi$ . Then the function

$$t \mapsto \varphi_t(x)$$

is constant if  $x$  is a critical point of  $X$ . Furthermore, if  $x$  is not critical, then for all  $t$ ,  $\varphi_t(x)$  is also not critical.

*Proof.* The first statement follows from Picard–Lindelöf, as the constant function  $\text{const}_x : t \mapsto x$  for all  $t$  satisfies the condition

$$\frac{d}{dt}\text{const}_x = 0 = X(\text{const}_x(t)).$$

Now assume that for some  $t_0$ ,  $y := \varphi_{t_0}(x)$  is critical. Then by the flow property  $x = \varphi_{-t_0}(y)$  and hence  $x$  was already critical.  $\square$

prop:vb-ffDec  
reasesAlongGr  
radientFlow

**Proposition 2.0.60** (The Flow of the Gradient). Let  $f : M \rightarrow \mathbb{R}$  be smooth function on a Riemannian manifold  $(M, g)$  and  $\varphi$  be the dynamical system associated to  $\text{grad}(f)$ . Then for any  $x \in M$ , the function

$$f \circ \varphi_t(x) : \mathbb{R} \rightarrow \mathbb{R}, \quad t \mapsto f \circ \varphi_t(x)$$

is strictly increasing if  $x$  is not a critical point of  $\text{grad}(f)$  and constant if  $x$  is a critical point of  $\text{grad}(f)$ . (If the function is only defined on an open subset of  $\mathbb{R}$ , the statement still holds.)

*Proof.* We just calculate the differential of  $\varphi_t(x)$  and make use of positive definiteness of  $g$ :

$$\frac{d}{dt} f \circ \varphi_t(x)(R_t(1)) = df_{\varphi_t(x)}(\text{grad}_{\varphi_t(x)}(f)) = g(\text{grad}_{\varphi_t(x)}(f), \text{grad}_{\varphi_t(x)}(f)) \geq 0 \quad \forall t \in \mathbb{R}$$

where equality holds only if and only if  $\text{grad}_{\varphi_t(x)}(f) = 0$ . Prop 2.0.59 tells us that the latter is equivalent to  $x$  being a critical point of our vector field.  $\square$

These two constructions are mutually inverse, giving the following theorem:

**Theorem 2.0.61.** *Let  $M$  be a smooth manifold. Then there is a one-to-one correspondence between global dynamical systems and vector fields given by the constructions above.*

## 3 Complex K-Theory

### 3.1 Mathematical Goals

**Theorem 3.1.1** (The Theorems and Mathematical Structures). • *Definition of The K-Functor*

- *Cohomology structures*
  - *Additivity (pointed sum)*
  - *Long exact sequences*
  - *Excision*

### 3.2 Classification of Vector Bundles

The first goal is to classify all vector bundles over sufficiently nice spaces. The latter refers to spaces admitting partitions of unity for which Tietze's extension theorem holds. For simplicity we just consider compact spaces. Along the way we will discover that the set of isomorphism classes of vector bundles is a homotopy invariant.

Before we go back to the analytical reals, we will further develop the theory of vector bundles:

**Definition 3.2.1.** For a topological space  $X$  we define  $\text{Vect}_n$  to be the set of isomorphism classes of complex vector bundles of rank  $n$  and  $\text{Vect}(X)$  to be the set of isomorphism classes of complex vector bundles of any rank. By declaring  $\text{Vect}(f) = f^*$  to be the pullback we get a contravariant functor  $\text{Vect}$  from **CGWH** to **Set**. We can equip the set  $\text{Vect}(X)$  with the direct sum and tensor product operation. It is then easy to see that these operations descend to isomorphism classes and hence we get an abelian semiring  $(\text{Vect}(X), \oplus, \otimes)$  (abelian semigroup with associative and distributive multiplication). In fact we even get a contravariant functor from **CGWH** to the category of abelian semirings.

lem:kTheo-sec  
tionExtension

**Lemma 3.2.2.** Let  $X$  be a compact space,  $Y \subseteq X$  a closed subspace and  $E$  a vector bundle over  $X$ . Then any section  $s : Y \rightarrow E|_Y$  can be extended to a global section of  $X$ .

*Proof.* For a detailed proof see Chapter 1.4 of [Ati89]. The general argument however is easy to follow: By Tietze's extension theorem we can extend the section to around every  $x \in X$ . By compactness we can pick finitely many of those extensions. Now using a partition of unity with respect to their domains we can build the sum of all the extensions to get a well defined global one.  $\square$

[Ati89]

cor:kTheo-iso  
morphismExten  
sion

**Corollary 3.2.3.** Let  $Y$  be a closed subspace of a compact space  $X$ , and let  $E, F$  be bundles over  $X$ . If  $f : E|_Y \rightarrow F|_Y$  is an isomorphism, then there is an open neighbourhood  $U \supset Y$  and an extension  $f : E|_U \rightarrow F|_U$  which is an isomorphism.

*Proof.* By corollary 2.0.43 we can view  $f$  as a section into the bundle  $\text{Hom}(E|_Y, F|_Y)$ , which can be extended to a global one in  $\text{Hom}(E, F)$ . Let  $U$  be the open set for which the section maps to isomorphisms. Since  $U$  contains  $Y$  this yields the required extension.  $\square$

thm:kTheo-hom  
otopyInvarian  
ce

**Theorem 3.2.4** (Homotopy Invariance of the Pullback). *Let  $X, Y$  be two compact spaces,  $I = [0, 1]$ , and*

$$f : Y \times I \rightarrow X$$

*be a homotopy. For a fixed  $t \in I$ , we define the function  $f_t : Y \rightarrow X$ ,  $f_t(y) = f(t, y)$ . Let  $\pi : E \rightarrow X$  be a vector bundle. Then*

$$f_0^*(E) \cong f_1^*(E).$$

*Proof.* Consider the following two vector bundles over  $Y \times I$ :

$$f^*E \quad \text{and} \quad \pi_1^* f_t^* E,$$

where  $\pi_1 : Y \times I \rightarrow Y$  denotes the projection and  $t \in I$ . Since the restriction is by definition the pullback along the inclusion  $i : Y \times \{t\} \rightarrow Y \times I$ . The equality of the two maps:

$$f \circ i = f_t \circ \pi_1 \circ i : Y \times \{t\} \rightarrow X$$

together with the functoriality gives us an isomorphism

$$\Phi : f^*E|_{Y \times \{t\}} \rightarrow \pi_1^* f_t^* E|_{Y \times \{t\}}.$$

By Corollary 3.2.3 we can extend the isomorphism for some open neighbourhood of  $Y \times \{t\}$  and by the product topology we can deduce that there exists a  $\delta_t$  such that  $U$  contains the thickened line  $Y \times (t - \delta_t, t + \delta_t)$ . The isomorphism class of  $f^*E|_{Y \times \{t\}}$  as a bundle over  $Y$  is locally constant in  $t$ : Let  $0 < \varepsilon < \delta$  and  $r \in I$ . Then define the inclusions

$$\begin{array}{ccc} i_t : Y & \rightarrow & Y \times \{t\} \subseteq Y \times I \\ y & \mapsto & (y, t) \end{array} \quad \text{and} \quad \begin{array}{ccc} i_{t+\varepsilon} : Y & \rightarrow & Y \times (t - \delta_t, t + \delta_t) \\ y & \mapsto & (y, t + \varepsilon) \end{array}$$

Then we have the identities  $f_t = f \circ i_t$  and  $\pi_1 \circ i_{t+\varepsilon} = \text{id}_Y$  and we can calculate:

$$f_{t+\varepsilon}^* E = i_{t+\varepsilon}^* f^* E = i_{t+\varepsilon}^* \pi_1^* f_t^* E = (f_t \circ \pi_1 \circ i_{t+\varepsilon})^* E = f_t^* E.$$

Finally, since  $I$  is connected, any locally constant map from  $I$  is globally constant and therefore  $f_0^* E \cong f_1^* E$ .  $\square$

lem:kTheo-con  
tractibleBund  
lesAreTivial

**Corollary 3.2.5.** 1. If  $f : X \rightarrow Y$  is a homotopy equivalence,  $f^* : (Y) \rightarrow (X)$  is a semiring isomorphism.

2. If  $X$  is contractible, every bundle over  $X$  is trivial and  $(X) \cong \mathbb{N}_0$

def:kTheo-tri  
vialisation

**Definition 3.2.6** (Trivialisation of a Vector Bundle). Let  $Y$  be a closed subspace of  $X$ ,  $E$  a vector bundle over  $X$  and  $\alpha : E|_Y \rightarrow Y \times V$  be an isomorphism. Then we call  $\alpha$  **trivialisation of  $E$  over  $Y$** .

lem:kTheo-tri  
vialisationDe  
finesABundle

**Lemma 3.2.7.** *A trivialisation  $\alpha$  of a bundle  $E$  over  $Y \subseteq X$  defines a bundle  $E/\alpha$  over  $X/Y$ . The isomorphism class of  $E/\alpha$  depends only on the homotopy class of  $\alpha$ .*

*Proof.* We define the bundle  $E/\alpha$  as follows: Let  $\pi_2 : Y \times V \rightarrow V$  denote the projection and define an equivalence relation on  $E|_Y$  by

$$e \sim e' \Leftrightarrow \pi_2 \alpha(e) = \pi_2 \alpha(e').$$

We extend this relation by the identity on  $E|_{X \setminus Y}$  and we define  $E/\alpha$  to denote the quotient space of  $E$  defined by that relation. To see why this is a bundle we need to check the local triviality only around the base point  $Y/Y$  of  $X/Y$ . This follows from the fact that by Corollary 3.2.3 we can extend  $\alpha$  to an isomorphism  $\tilde{\alpha} : E|_U \rightarrow U \times V$ . Then  $\tilde{\alpha}$  induces an isomorphism

$$E/\alpha|_{U/Y} \cong (U/Y) \times V$$

proving the local triviality. Now for the homotopy invariance of said bundle suppose that  $\alpha_0$  and  $\alpha_1$  are homotopic trivialisations of  $E$  over  $Y$ . This means that we have a trivialisation  $\beta$  of  $E \times I$  over  $Y \times I \subset X \times I$  that induces the  $\alpha_i$  at the endpoints. Let  $f : (X/Y) \times I \rightarrow (X \times I)/(Y \times I)$  be the natural map. Then  $f^*((E \times I)/\beta)$  is a bundle on  $(X/Y) \times I$  whose restriction to  $(X/Y) \times \{i\}$  is  $E/\alpha_i$  ( $i = 0, 1$ ). Hence by the homotopy invariance of the pullback (Theorem 3.2.4) we have

$$E/\alpha_0 \cong E/\alpha_1.$$

□

cor:kTheo-tri  
vializationIs  
RightInverse

**Corollary 3.2.8.** The trivialisation acts as a right inverse of the pullback along the collapse: If we call  $f : X \rightarrow X/Y$  the collapse, and  $fE$  can be trivialised over  $Y$  via  $\alpha$ , then

$$f^*(E/\alpha) \cong E.$$

*Proof.* Intuitively this is clear, as the pullback along the collapse builds a bundle, that is trivial over  $Y$  and agrees with  $E$  outside of  $Y$ . To prove the statement we will construct a bundle map, that is an isomorphism in each fibre: Let  $q : E \rightarrow E/\alpha$  denote the quotient map from the construction and let  $\pi_E : E \rightarrow X$ ,  $\pi_{E/\alpha} : E/\alpha \rightarrow X/Y$  be the bundle projections. By construction of  $E/\alpha$  we have  $\pi_{E/\alpha} \circ q = f \circ \pi_E$ , so

$$\Phi : E \rightarrow f^*(E/\alpha), \quad \Phi(e) = (\pi_E(e), q(e))$$

is a well-defined bundle morphism. We show  $\Phi$  is a fibrewise linear isomorphism, and hence a bundle isomorphism by Lemma 2.0.31.

*Case  $x \notin Y$ :* Here the equivalence relation defining  $E/\alpha$  restricts to the identity on  $E_x$ , so

$$q_x : E_x \xrightarrow{\cong} (E/\alpha)_{f(x)} = (E/\alpha)_x$$

is a linear isomorphism.

*Case  $x \in Y$ :* By definition of the equivalence relation, two points  $e, e' \in E_x$  are identified with  $\pi_2 \alpha(e) = \pi_2 \alpha(e') \in V$ , and all fibres  $E_x$  ( $x \in Y$ ) are glued to the single fibre  $(E/\alpha)_{y_0} \cong V$  over the basepoint  $y_0 = Y/Y$ . Hence

$$q_x = \pi_2 \circ \alpha_x : E_x \rightarrow V = (E/\alpha)_{y_0},$$

where  $\alpha_x : E_x \rightarrow V$  is the (linear) restriction of  $\alpha$  to the fibre over  $x$ . Since  $\alpha_x$  is a linear isomorphism, so is  $q_x$ .

Thus  $\Phi$  is a bundle morphism which is a linear isomorphism on every fibre.  $\square$

lem:kTheo-col  
lapsingOfCont  
ractibleSubsp  
ace

**Lemma 3.2.9.** *Let  $Y \subseteq X$  be a closed contractible subspace of a compact space  $X$ . Then*

$$f : X \rightarrow X/Y \quad \text{induces a semiring isomorphism} \quad f^* : \text{Vect}(X/Y) \rightarrow \text{Vect}(X).$$

*Proof.* Since  $Y$  is contractible,  $E|_Y$  is trivial and thereby a trivialisation  $\alpha : E|_Y \rightarrow Y \times V$  exists. Two such trivialisations only differ by the automorphism of  $Y \times V$  which in turn corresponds to a map  $Y \rightarrow \text{GL}(V) \cong \text{GL}_{\text{rank}(E)}(\mathbb{C})$ . Now since  $Y$  is contractible those two maps factor (up to homotopy) over any one point  $y \in Y$  and are thereby homotopically equivalent to the constant map. Finally since  $\text{GL}_{\text{rank}(E)}(\mathbb{C})$  is connected all constant maps are homotopically equivalent. Thereby  $\alpha$  is unique up to homotopy. By Lemma 3.2.7 the isomorphism class of  $E/\alpha$  only depends on the homotopy class of  $\alpha$  we have a well defined map

$$\text{Vect}(X) \rightarrow \text{Vect}(X/Y),$$

which in turn is an inverse of  $f^*$  concluding the proof.  $\square$

def:kTheo-glu  
ingAndClutchi  
ng

**Definition 3.2.10** (Gluing and Clutching of Vector Bundles). Let

$$X = X_1 \cup X_2, \quad A = X_1 \cap X_2,$$

and all be compact spaces. Assume that  $E_i$  is a vector bundle over  $X_i$  and  $\varphi : E_1|_A \rightarrow E_2|_A$  is an isomorphism. Then we define a vector bundle  $E_1 \cap_\varphi E_2$  on  $X$  as follows: We topologise the space  $E_1 \cup_\varphi E_2$  as the quotient of  $E_1 + E_2 / \sim$ , where  $\sim$  is the identification of points with their image under  $\varphi$ . Identifying  $X$  with  $X_1 + X_2$  in the quotient space we get the natural projection  $p : E_1 \cap_\varphi E_2 \rightarrow X$  and the fibres have a natural vector space structure. It remains to show that this is locally trivial. This is obvious outside of  $A$  so let  $a \in A$  and  $V_1$  be a closed neighbourhood of  $a$  in  $X_1$  such that  $E_1|_{V_1}$  is trivial giving us the isomorphisms

$$\begin{aligned} \theta_1 : E_1|_{V_1} &\rightarrow V_1 \times \mathbb{C}^n \\ \theta_1^A : E_1|_{V_1 \cap A} &\rightarrow (V_1 \cap A) \times \mathbb{C}^n \end{aligned}$$

Finally, define

$$\theta_2^A : E_2|_{V_1 \cap A} \rightarrow (V_1 \cap A) \times \mathbb{C}^n$$

by composition  $\theta_1^A \circ \varphi^{-1}$ . Let  $V_2$  be a neighbourhood of  $a$  in  $X_2$  such that

$$\theta_2 : E_2|_{V_2} \rightarrow V_2 \times \mathbb{C}^n$$

is an extension of  $\theta_2^A$ . This extension exists again by extending the section into the homomorphism bundle corresponding to the isomorphism  $\theta_2^A$ . Finally  $\theta_1$  and  $\theta_2$  define a well defined isomorphism

$$\theta_1 \cap_\varphi \theta_2 : E_1 \cap_\varphi E_2 \rightarrow (V_1 \cap V_2) \times \mathbb{C}^n.$$

This proves the local triviality.

We call such a triple  $(E_1, E_2, \varphi)$  a **clutching datum** and  $\varphi$  the **transition function**.

**Corollary 3.2.11** (Elementary Properties of the Gluing and Clutching Construction). Elementary properties of the gluing and clutching construction are:

1. If  $E$  is a bundle over  $X$ , then the identity defines an isomorphism and

$$E_1 \cup_{\text{id}_A} E_2 \cong E.$$

2. If  $\beta_i : E_i \rightarrow E'_i$  are isomorphisms on  $X_i$  and  $\varphi' \beta_1 = \beta_2 \varphi$ , then

$$E_1 \cup_\varphi E_2 \cong E'_1 \cup_{\varphi'} E'_2.$$

3. If  $(E_i, \varphi)$  and  $(E'_i, \varphi')$  are two “clutching data” on  $X_i$ , then

$$\begin{aligned} (E_1 \cup_\varphi E_2) \oplus (E'_1 \cup_{\varphi'} E'_2) &\cong E_1 \oplus E'_1 \bigcup_{\varphi \oplus \varphi'} E_2 \oplus E'_2, \\ (E_1 \cup_\varphi E_2) \otimes (E'_1 \cup_{\varphi'} E'_2) &\cong E_1 \otimes E'_1 \bigcup_{\varphi \otimes \varphi'} E_2 \otimes E'_2, \\ (E_1 \cup_\varphi E_2)^* &\cong E_1^* \bigcup_{(\varphi^*)^{-1}} E_2^*. \end{aligned}$$

$(-)^*$  denotes the dual here.

lem:kTheo-hom  
otopyInvariant  
ceClutchingFu  
nction

**Lemma 3.2.12.** *The isomorphism class of  $E_1 \cup_\varphi E_2$  depends only on the homotopy class of the isomorphism  $\varphi : E_1|_A \rightarrow E_2|_A$ .*

*Proof.* A homotopy of isomorphisms is a map

$$\Phi : E_1|_A \times I \rightarrow E_2|_A,$$

such that for all  $t$  the map

$$\Phi_t : E_1|_A \rightarrow E_2|_A \quad \Phi_t(\xi) := \Phi(\xi, t)$$

is a bundle isomorphism. We can encode this  $\Phi$  into a bundle isomorphism as follows: Let  $\pi : X \times I \rightarrow X$  be the projection. Then elements  $\xi \in \pi^* E_1$  are by definition 2.0.15 of the form  $\xi_{x,t} = ((x,t), \xi_x) \in (\pi^* E_1)_{x,t}$ . With this notation define the bundle isomorphism:

$$\Psi : \pi^*(E_1)|_{A \times I} \rightarrow \pi^*(E_2)|_{A \times I}, \quad \Psi(\xi_{x,t}) := ((x,t), \Phi_t(\xi_x))$$

This is a bundle isomorphism, since  $\Phi_t$  is one. Now denote the inclusion

$$i_t : X \rightarrow X \times I, \quad i_t(x) := (x, t),$$

and notice that the pullback along  $i_t$  is the same as restricting to  $X \times \{t\}$ .

With those definitions it is clear by the construction of the gluing, that

$$E_1 \cup_{\Phi_t} E_2 \cong i_t^*(\pi^*(E_1) \cup_{\Psi} \pi^*(E_2)). \quad (1)$$

{eq:kTheo-gluingHomotopyInvariant}

The isomorphism can be seen by part (2) of Corollary 3.2.11 as follows: First notice the isomorphism  $\pi^*(E_i)|_{A \times \{t\}} \cong E_i$  for  $i = 1, 2$  and any  $t$ . By construction of the gluing this leads to the isomorphism

$$E_1 \cup_{\Phi_t} E_2 \cong \pi^*(E_1)|_{A \times \{t\}} \cup_{(\Phi_t, \text{id})} \pi^*(E_2)|_{A \times \{t\}}.$$

Now again by construction we notice, that gluing restricted spaces is the same as gluing the whole space and then restricting. Furthermore the map  $(\Phi_t, \text{id})$  is the restriction of  $\Psi$  to  $X \times \{t\}$ . Therefore:

$$\pi^*(E_1)|_{A \times \{t\}} \cup_{(\Phi_t, \text{id})} \pi^*(E_2)|_{A \times \{t\}} \cong (\pi^*(E_1) \cup_{\Psi} \pi^*(E_2))|_{X \times \{t\}}$$

Finally, pulling back via  $i_t$  is the same as restricting  $X \times i$  to  $X \times \{t\}$  and identifying  $X$  with  $X \times \{t\}$ . This concludes the chain of arguments:

$$(\pi^*(E_1) \cup_{\Psi} \pi^*(E_2))|_{X \times \{t\}} \cong i_t^*(\pi^*(E_1) \cup_{\Psi} \pi^*(E_2)).$$

Since the pullback is homotopy invariant we can conclude by the isomorphism (1):

$$E_1 \cup_{\Phi_0} E_2 \cong E_1 \cup_{\Phi_1} E_2.$$

□

def:kTheo-homotopyClassesOfMaps

**Definition 3.2.13.** Let  $X, Y$  be topological spaces. By  $[X, Y]$  we denote the set of homotopy classes of maps from  $X$  to  $Y$ .

lem:kTheo-gluingAndHomotopyClassesToGLn

**Lemma 3.2.14.** Let  $X = X_1 \cup X_2$  with  $X_1$  and  $X_2$  closed contractible spaces. Then there is a canonical bijection

$$\text{Vect}_n(X) \rightarrow [A, \text{GL}_n(\mathbb{C})].$$

*Proof.* Let  $[E]$  be any vector bundle over  $X$  of rank  $n$ . Then  $E$  can be represented by a vector bundle obtained by gluing that is trivial on both  $X_i$ : Since  $X_i$  is contractible we can choose two trivialisations

$$\alpha_1 : E|_{X_1} \rightarrow X_1 \times \mathbb{C}^n, \quad \text{and} \quad \alpha_2 : E|_{X_2} \rightarrow X_2 \times \mathbb{C}^n.$$

With part (2) of Corollary 3.2.11 we can conclude:

$$E \cong E|_{X_1} \cup_{\text{id}_A} E|_{X_2} \cong X \times \mathbb{C}^n \cup_{\alpha_2 \circ \alpha_1^{-1}|_A} X \times \mathbb{C}^n$$

Since a map between the two trivial bundles  $A \times \mathbb{C}^n \rightarrow A \times \mathbb{C}^n$  can be identified with a map  $A \rightarrow \text{GL}_n(\mathbb{C})$  we get an assignment from bundles over  $X$  together with the trivialisations to maps  $A \rightarrow \text{GL}_n(\mathbb{C})$ . The next step is to show that two isomorphic vector bundles give homotopic maps: Assume that we are given two vector bundles  $E, F$  over  $X$  that are isomorphic via  $T : E \rightarrow F$  and without loss of generality we assume that they are given as the gluing of two trivial bundles:

$$E : X_1 \times \mathbb{C}^n \cup_{\varphi} X_2 \times \mathbb{C}^n, \quad \text{and} \quad F : X_1 \times \mathbb{C}^n \cup_{\psi} X_2 \times \mathbb{C}^n.$$

We now want to define a homotopy between  $\psi$  and  $\varphi$ . For this we start by splitting the isomorphism  $T$  into its two parts over  $X_1$  and  $X_2$ :

$$T|_{X_i} : X_i \times \mathbb{C}^n \rightarrow X_i \times \mathbb{C}^n.$$

These maps are given by maps

$$T_i : X_i \rightarrow \text{GL}_n(\mathbb{C}),$$

where the gluing compatibility of the  $T|_{X_i}$  yield for any  $a \in A$ :

$$\psi(a)T_1(a) = T_2(a)\varphi(a), \quad \text{equivalently} \quad \psi(a) = T_2(a)\varphi(a)T_1^{-1}(a).$$

By the contractibility of  $X_i$  we have for any fixed  $x_i \in X_i$  a homotopy  $h_t^i : X_i \rightarrow X_i$  such that  $h_0^i = c_{x_i}$  is the collapsing map whilst  $h_1^i = \text{id}_{X_i}$ . Furthermore, since  $\text{GL}_n(\mathbb{C})$  is connected we have a path  $\gamma^i : I \rightarrow \text{GL}_n(\mathbb{C})$  that connects  $T_i(x_i) = \gamma^i(1)$  to  $\mathbb{1}_n = \gamma^i(0)$ . With this we can define the homotopies

$$T_{i,t} : X_i \rightarrow \text{GL}_n(\mathbb{C})$$

$$x \mapsto \begin{cases} T_i \circ h_{2t}^i(x) & \text{if } t \leq \frac{1}{2}, \\ \gamma^i(2(t - \frac{1}{2})) & \text{if } t > \frac{1}{2}, \end{cases}$$

that satisfy  $T_{i,1} = T_i$  and  $T_{i,0}(x) = \mathbb{1}_n$  for all  $x$ . Finally we define the homotopy  $H_t : A \rightarrow \text{GL}_n(\mathbb{C})$  by

$$H_t(a) := T_{2,t}(a)\varphi(a)T_{1,t}^{-1}(a)$$

to get a homotopy from  $\varphi = H_0$  to  $\psi = H_1$ . Hence, we get a well defined map from isomorphism classes of bundles over  $X$  of rank  $n$  to the homotopy classes of maps from  $A$  to  $\text{GL}_n(\mathbb{C})$  that is the inverse of the gluing construction and hence is bijective.  $\square$

ex:kTheo-taut  
opogicalLineB  
undleOverCP1

**Example 3.2.15.** We now want to analyse the tautological line bundle  $l$  over  $\mathbb{CP}^1$ . For this we cut  $\mathbb{CP}^1$  into the two pieces which are contractible as they come with homeomorphisms into the unit disc in  $\mathbb{C}$ :

$$\begin{aligned} H^+ &:= \{[x : y] \in \mathbb{CP}^1 \mid |x| \geq |y|\} \rightarrow D \subset \mathbb{C}, \quad [x : y] \mapsto \frac{y}{x} \\ H^- &:= \{[x : y] \in \mathbb{CP}^1 \mid |x| \leq |y|\} \rightarrow D \subset \mathbb{C}, \quad [x : y] \mapsto \frac{x}{y} \end{aligned}$$

Those maps give us unique representations of elements in  $H^+$  and  $H^-$ . We now want to evaluate the bijection  $\text{Vect}_1(\mathbb{CP}^1) \rightarrow [H^+ \cap H^-, \text{GL}_1(\mathbb{C})]$  at  $[l]$ . The image of the intersection  $H^+ \cap H^-$  under the two maps is the boundary of the disc and hence a circle. Furthermore,  $\text{GL}_1(\mathbb{C}) \cong \mathbb{C}^*$ . Hence, we know that the image is isomorphic to  $[S^1, \mathbb{C}^*] \cong \pi_1(S^1) \cong \mathbb{Z}$ . This tells us, that there must be nontrivial vector bundles on  $\mathbb{CP}^1$ . By the proof of 3.2.14 we know that  $[l]$  gets mapped to the function gluing together trivialisations over  $H^+$  and  $H^-$  along their intersection. To define the trivialisations we need to define local frames over  $H^+$  and  $H^-$ . Those frames are given by a single section defined as:

$$\begin{aligned} s_+ : H^+ &\rightarrow l, \quad [x : y] \mapsto (1, \frac{y}{x}), \\ s_- : H^- &\rightarrow l, \quad [x : y] \mapsto (\frac{x}{y}, 1). \end{aligned}$$

With those we define the trivialisations:

$$\begin{aligned} t_+ : l|_{H^+} &\rightarrow H^+ \times \mathbb{C}, \quad ([x : y], \lambda \cdot s_+[x : y]) \mapsto ([x : y], \lambda) \\ t_- : l|_{H^-} &\rightarrow H^- \times \mathbb{C}, \quad ([x : y], \mu \cdot s_-[x : y]) \mapsto ([x : y], \mu) \end{aligned}$$

in each fibre over the intersection we have

$$\lambda \cdot s_+([x : y]) = \lambda(1, \frac{y}{x}) = \lambda \cdot \frac{y}{x}(\frac{x}{y}, 1) = \lambda \cdot \frac{y}{x}s_-([x : y])$$

Hence in each fibre over  $[x : y] \in H^+ \cap H^-$  the gluing function can be calculated:

$$\lambda \mapsto t_-|_{[x:y]} \circ t_+|_{[x:y]}^{-1}(\lambda) = t_-|_{[x:y]}(\lambda \cdot s_+([x : y])) = t_-|_{[x:y]}(\lambda \cdot \frac{y}{x} \cdot s_-([x : y])) = \lambda \cdot \frac{y}{x}$$

Hence, as a map from  $H^+ \cap H^- \rightarrow \text{GL}_1(\mathbb{C}) \cong \mathbb{C}^*$  this map reads:

$$H^+ \cap H^- \rightarrow \mathbb{C}^* \quad [x : y] \mapsto \frac{y}{x}$$

viewing this as an element of  $\pi(S^1)$  by identifying the intersection with a circle via the homeomorphism from  $H^+ \rightarrow D \subseteq \mathbb{C}$  we notice that the gluing map corresponds to the identity in  $\pi_1(S^1)$ . If the bundle  $l$  would be trivial, the gluing function would be homotopically equivalent to the constant one-function  $H^+ \cap H^- \rightarrow \mathbb{C}^*$  that maps all  $[x : y] \mapsto 1$ . This cannot be, because interpreted as elements in  $\pi(S)$  they are different.

def:kTheo-met-  
ricsOnBundles

**Definition 3.2.16** (Metrics on Bundles). A **metric** on a complex bundle  $E$  is a section  $h : X \rightarrow \text{Herm}(E)$  such that  $h(x)$  is positive definite for all  $x$ .  $\text{Herm}(E)$  denotes the bundle of hermitian forms (conjugate-symmetric sesquilinear) in each fibre. A bundle together with a metric is a **Hermitian bundle**.

def:kTheo-pro-  
jectionOperat-  
or

**Definition 3.2.17** (Projection Operator). A **projection operator**  $P : E \rightarrow E$  is a bundle homomorphism in  $E$  such that  $P^2 = P$ .

lem:kTheo-pro-  
jectionsGiveS-  
umdecompositi-  
on

**Lemma 3.2.18.** *Any projection operator  $P : E \rightarrow E$  determines a direct sum decomposition  $E = PE \oplus (1 - P)E$ .*

*Proof.* Observe that fibrewise  $1 - P$  maps each  $\xi$  to the kernel of  $P$ , and hence fibrewise  $P(E)$  and  $(1 - P)(E)$  are disjoint. Furthermore  $\text{im}(P) + \text{im}(1 - P) = E$ , because  $v = P(v) + (v - P(v))$ . Therefore in each fibre we have the direct sum  $E_x = P(E_x) \oplus (1 - P)(E_x)$ . Now the question is, if that gives a global splitting, meaning that  $P(E)$  and  $(1 - P)(E)$  are continuous as subbundles of  $E$ . For this to be true, we need local triviality, which is clear as long as the rank of  $P$  (and  $(1 - P)$ ) is locally constant. In each fibre we know that  $\text{rank}(P_x) + \text{rank}(1 - P) = \dim(E_x)$ . Furthermore,  $\text{rank}(P_x)$  and  $\text{rank}(1 - P)$  are locally constant. To see this first notice that they are lower semi-continuous, meaning that for  $y$  from a small enough neighbourhood of  $x$   $\text{rank}(P_x) \leq \text{rank}(P_y)$ . To see this, pick a  $\text{rank}(P_x)_x \times \text{rank}(P_x)_x$  minor with nonzero determinant. The latter is an open condition and hence stable under small perturbation. But now we have the sum of two lower semi-continuous functions that give something locally constant and hence, each summand is locally constant. Finally the above shows that  $\ker(P) = (1 - P)(E)$  we have the decomposition.  $\square$

lem:kTheo-met-  
ricInducedCom-  
plement

**Lemma 3.2.19** (Metric Induced Complement). *Suppose that  $F$  is a sub-bundle of  $E$ . Then a metric provides a canonical complementary sub-bundle.*

*Proof.* Let  $h$  denote the metric. For each  $x$  we have the orthonormal projection  $P_x : E_x \rightarrow F_x$  inducing a map  $P : E \rightarrow F$ . Let  $(f_1, \dots, f_n)$  be an orthonormal frame of  $F$ . Then fibrewise with  $\xi \in E_x$  the orthogonal projection is given by

$$P_x(\xi) = \sum_i h_x(\xi, f_i(x)) f_i(x).$$

And since  $h$  is continuous the whole map  $P$  is. Furthermore  $P^2 = P$  and hence we get a canonical sum decomposition.  $\square$

**Definition 3.2.20.** We call a sequence of bundle homomorphisms

$$\dots \longrightarrow E \longrightarrow F \longrightarrow \dots$$

exact, if it is fibrewise exact.

lem:kTheo-spl  
ittingPrinzip  
le

**Lemma 3.2.21** (Splitting Principle). *Every short exact sequence of bundles over the same base space splits.*

*Proof.* Let  $X$  be the base space and assume that the sequence is given as

$$0 \longrightarrow E' \xrightarrow{\varphi'} E \xrightarrow{\varphi''} E'' \longrightarrow 0$$

Now it is well known that any short exact sequence of vector spaces splits. We just need to ensure that this splitting is continuous. For this we give  $E$  a metric. Now since  $\varphi'$  is injective, it is a monomorphism and  $\varphi'(E')$  is a subbundle by Lemma 2.0.36. By Lemma 3.2.19 we can split  $E \cong \varphi'(E') \oplus L$ . However,  $\varphi''|_L$  is fibrewise (and hence as a bundle map) an isomorphism. This can be seen as  $\varphi''$  being surjective and trivial on  $\varphi'(E')$  hence the restriction is still surjective. The injectivity is also clear as the kernel by the homomorphism theorem of linear maps. But then we clearly have  $E = E' \oplus E''$  and the sequence is isomorphic as a short exact sequence to the splitting one:

$$0 \longrightarrow E' \xrightarrow{i_1} E' \oplus E'' \xrightarrow{\pi_2} E'' \longrightarrow 0$$

□

def:kTheo-amp  
leSubspace

**Definition 3.2.22.** A subspace  $V \subseteq \Gamma(E)$ , where  $\Gamma(E)$  is viewed as a  $\mathbb{C}$ -vector space is called **ample**, if

$$\varphi : X \times V \rightarrow E, \quad \varphi(x, s) := s(x)$$

is a surjection.

**Corollary 3.2.23.** If the subspace  $V$  is ample and  $\dim(V) = \text{rank}(E)$ , then  $E$  is trivialisable.

lem:kTheo-amp  
leSubspaceExi  
sts

**Lemma 3.2.24.** *For any vector bundle  $E$  over a compact space  $X$ , there exists a finite dimensional ample subspace of  $\Gamma(E)$ .*

*Proof.* Let  $\{U_\alpha\}$  be a finite open cover of  $X$  such that  $E|_{U_\alpha}$  is trivial. Let  $\{s_{\alpha,i}\}$  be the corresponding local frames, and define  $V_\alpha$  to be the finite dimensional ample subspace  $V_\alpha \subseteq \Gamma(E|_{U_\alpha})$ . Let  $\rho_\alpha$  be a partition of unity subordinate to the cover  $\{U_\alpha\}$ . Define

$$\theta_\alpha : V_\alpha \rightarrow \Gamma(E), \quad \theta_\alpha(s_\alpha)(x) = \begin{cases} s_\alpha(x) \cdot \rho_\alpha(x) & \text{if } x \in U_\alpha, \\ 0 & \text{else.} \end{cases}$$

Then the  $\theta_\alpha$  give rise to a homomorphism

$$\theta : \Pi_\alpha V_\alpha \rightarrow \Gamma(E),$$

and the image of  $\theta$  is finite dimensional and ample. The surjectivity is clear, as for any  $x$  there exists a  $\rho_\alpha(x) > 0$  making the map

$$\theta_\alpha(V_\alpha) \rightarrow E_x$$

a surjection.

□

cor:kTheo-epi  
FromTrivial

**Corollary 3.2.25.** If  $E$  is any bundle over  $X$ , there exists an epimorphism  $\varphi : X \times \mathbb{C}^m \rightarrow E$  for some integer  $m$ .

*Proof.* Just take a basis  $(s_1, \dots, s_m)$  of an ample subspace  $V$  and define  $\varphi(x, e_i) = s_i(x)$ , where  $e_i$  is the  $i$ -th standard basis vector of  $\mathbb{C}^m$ .  $\square$

cor:kTheo-spl  
ittingTrivial

**Corollary 3.2.26.** If  $E$  is a bundle over  $X$ , there exists a bundle  $F$  such that  $E \oplus F$  is trivial.

*Proof.* By Corollary 3.2.25 there exists an epimorphism  $\varphi : X \times \mathbb{C}^m \rightarrow E$ . The kernel of  $\varphi$  is a subbundle of  $X \times \mathbb{C}^m$  and hence by Lemma 3.2.21 we have a splitting of the short exact sequence

$$0 \longrightarrow \ker(\varphi) \longrightarrow X \times \mathbb{C}^m \xrightarrow{\varphi} E \longrightarrow 0$$

Hence, the middle term becomes isomorphic to the direct sum of the outer terms and we get  $X \times \mathbb{C}^m \cong E \oplus \ker(\varphi)$ .  $\square$

def:kTheo-gra  
ssmanManifold

**Definition 3.2.27** (Grassmann Manifold). Let  $V$  be a vector space and  $n$  be any integer. The **Grassmann manifold**  $\text{Gr}_n(V)$  is the set of all subspaces of  $V$  with **codimension**  $n$ . If  $V$  is given a Hermitian metric, we can topologise  $\text{Gr}_n(V)$  as a subspace of the set of all projection operators on  $V$  and this defines a map

$$P : \text{Gr}_n(V) \rightarrow \text{End}(V),$$

where  $\text{End}(V)$  is the space of all endomorphisms of  $V$  with the operator norm topology. We give  $\text{Gr}_n(V)$  the subspace topology of  $\text{End}(V)$ .

def:kTheo-cla  
ssifyingMap

**Definition 3.2.28.** Suppose that  $E$  is a bundle over  $X$ ,  $V$  is a vector space and  $\varphi : X \times V \rightarrow E$  is an epimorphism. Then we can define a map  $f : X \rightarrow \text{Gr}_{\dim(V)}(V)$  by sending  $x$  to the kernel of  $\varphi_x : V \rightarrow E_x$ . We call  $f$  **induced by**  $\varphi$ . This map is continuous for any metric on  $V$ . To see this notice that the topology on  $V$  is metric invariant, as  $V$  is finite dimensional. Now the map  $f$  is continuous if for the map

$$P : \text{Gr}_n(V) \rightarrow \text{End}(V)$$

the composition  $P \circ f$  is continuous. But  $P \circ f$  is given by sending  $x$  to the projection onto  $\ker(\varphi_x)$ , which is continuous as  $\varphi$  is.

def:kTheo-cla  
ssifyingBundl  
e

**Definition 3.2.29** (Classifying Bundle). Let  $V$  be a vector space and  $F \subseteq \text{Gr}_n(V) \times V$  be the subbundle consisting of all points  $(g, v)$  such that  $v \in g$ . We call  $F$  the **tautological bundle** over  $\text{Gr}_n(V)$ . Furthermore, if  $E = (\text{Gr}_n(V) \times V)/F$  is the quotient bundle, we call  $E$  the **classifying bundle** over  $\text{Gr}_n(V)$ .

**Corollary 3.2.30.** If  $E'$  is a bundle over  $X$ , and  $\varphi : X \times V \rightarrow E'$  is an epimorphism, then  $\varphi$  induces a map  $f : X \rightarrow \text{Gr}_m(V)$  such that  $E' \cong f^*E$ , where  $E$  is the classifying bundle over  $\text{Gr}_m(V)$ . The isomorphism follows from a fibrewise construction because  $E_x \cong V / \ker(\varphi_x)$ . The latter however is exactly the fibre over  $\ker(x)$  in the classifying bundle and hence the pullback via the  $\varphi$  induced map gives an isomorphism. That motivates the name “classifying bundle” for  $E$  and “classifying map” for  $f$ .

Begründung  
ergänzen,  
z.B.:  
Metriken  
auf  
 $X \times \mathbb{C}^m$   
und  $E$   
wählen;  
dann  
ist  $P :=$   
 $\varphi^*(\varphi\varphi^*)^{-1}\varphi$   
ein  
stetiger  
Projek-  
tionsop-  
erator  
mit  $(1 -$   
 $P)(X \times$   
 $\mathbb{C}^m) =$   
 $\ker(\varphi)$ ,  
und  
Lemma  
3.2.18  
liefert die  
Zerlegung.  
( $\varphi\varphi^*$  ist  
invertier-  
bar, da  $\varphi$   
Epimor-  
phismus.)

lem:kTheo-Gra  
ssmanTopology  
MetricInvaria  
nt

**Lemma 3.2.31.** *The topology on  $\text{Gr}_n(V)$  is independent of the choice of the metric on  $V$ .*

*Proof.* Let  $h$  and  $h'$  be two metrics on  $V$ . Then we denote the topological space  $\text{Gr}_n(V)$  with the topology induced by  $h$  by  $\text{Gr}(V_h)$  and the one induced by  $h'$  by  $\text{Gr}(V_{h'})$ . We need to show that the identity map  $\text{id} : \text{Gr}(V_h) \rightarrow \text{Gr}(V_{h'})$  is a homeomorphism. For this we need to show that  $\text{id}$  is continuous. The projection  $\text{Gr}_n(V_h) \times V_{h'} \rightarrow E$  (where  $E$  is the classifying bundle) is an epimorphism and hence induces a classifying map  $f : \text{Gr}_n(V_h) \rightarrow \text{Gr}_n(V_{h'})$  such that  $f^*E \cong E$ . However that map is given by the identity on  $\text{Gr}_n(V)$  and hence the identity is continuous by Definition 3.2.28.  $\square$

**Remark 3.2.32.** To continue further we need direct limits of directed systems to construct general Grassmannians. However we will also need said construction for topological spaces and groups. Hence, we will give the definitions for arbitrary categories.

**Definition 3.2.33** (Directed System). Let  $\mathcal{C}$  be a category and  $(I, \leq)$  be a partially ordered set where every finite subset is bounded from above. Let  $(A_i)_{i \in I}$  be a family of objects from  $\mathcal{C}$  and  $f_{ij} : A_i \rightarrow A_j$  be a morphism for all  $i \leq j$  that satisfies:

1.  $f_{ii} = \text{id}_{A_i}$  for all  $i \in I$ .
2.  $f_{kj} \circ f_{ik} = f_{ij}$  for all  $i \leq k \leq j$ .

The pair  $((A_i)_{i \in I}, (f_{ij})_{i \leq j})$  is called a **direct system**.

**Definition 3.2.34** (Direct Limit). Let  $((A_i)_{i \in I}, (f_{ij})_{i \leq j})$  be a direct system in a category  $\mathcal{C}$ . A pair  $(X, (\varphi_i)_{i \in I})$  where  $X$  is an object of  $\mathcal{C}$  and  $\varphi_i : A_i \rightarrow X$  are morphisms such that  $\varphi_j \circ f_{ij} = \varphi_i$  for all  $i \leq j$  is called a direct limit, if the following universal property holds: For any family of morphisms  $g_i : A_i \rightarrow Y$  that satisfies  $g_i \circ f_{ji} = g_j$  for all  $i \leq j$  there exists a unique morphism  $g : X \rightarrow Y$  such that  $g_i = g \circ \varphi_i$ . We denote the space  $X$  by  $\text{colim}_i A_i$  (or just  $\text{colim } A_i$ ), where the maps  $f_{ij}$  are omitted from the notation.

**Corollary 3.2.35.** As always, the direct limit is (if it exists) unique up to unique isomorphism.

**Remark 3.2.36** (Construction of Direct Limits). We will now see three constructions of direct limits. Let  $((A_i)_{i \in I}, (f_{ij})_{i \leq j})$  denote a directed system in the noted category.

*In the category of sets we define the direct limit by*

$$\text{colim}_i A_i := \bigsqcup A_i / \sim,$$

where  $x \sim y$  if there exists  $i, j, k \in I$  such that  $f_{ik}(x) = f_{jk}(y)$ . This gives an equivalence relation. The maps  $\varphi_i : A_i \rightarrow \text{colim}_i A_i$  are the composition of the inclusion into the disjoint union with the projection onto the quotient space. The universal property is easy to verify. For a family of suitable maps  $g_i : A_i \rightarrow B$  we define the map  $g : \text{colim}_i A_i \rightarrow B$

as  $g(x) = g_m(x)$  for  $x \in A_m$ . This is well defined since the  $g_i$  are compatible with the  $f_{ij}$  and unique by definition.

In the category of topological spaces we define the direct limit to be the set-theoretic direct limit equipped with the final topology of the inclusions, i.e. a subset  $U \subseteq \text{colim}_i A_i$  is open if and only if  $\varphi_i^{-1}(U)$  is open for all  $m$ . Again the universal property is easy to prove by using the construction of the universal map from the set-theoretic limit. To show the continuity we show that  $g^{-1}(U)$  is open for any open  $U$  which is checked by verifying openness of  $\varphi_i^{-1}(g^{-1}(U))$  for all  $i$ , which in turn equals  $g_i^{-1}(U)$  and is open as the  $g_i$  are continuous. Uniqueness follows from the uniqueness of  $g$  established already in set.

In an algebraic category like semigroups, groups, rings we define the direct limit again as the set-theoretic direct limit, where the algebraic operations are defined such that the  $\varphi_i$  are homomorphisms meaning for example (if  $+$  is an operation in the  $A_i$ )  $a \in A_i$  and  $b \in A_j$  we define  $a + b$  in  $\text{colim}_i A_i$  by including the sum  $f_{ik}(a) + f_{jk}(b)$  which is defined in  $A_k$ . Again the universal morphism  $g$  is the one from the set-theoretic limit which now is a homomorphism by construction.

rem:kTheo-sta  
irCaseMapsClo  
sedEmbedding

**Remark 3.2.37.** If we build the direct limit of sets and all the maps of the direct system are injective, then the inclusions into the direct limit are injective. If all the maps of the direct system are closed embeddings, then the inclusions into the direct limit are also closed embeddings.

*Proof.* Recall that the inclusions  $\varphi_i : A_i \rightarrow \text{colim}_i A_i$  are defined to be the composition of the inclusion into the disjoint union with the projection onto the quotient. Now assume that all maps  $f_{ij}$  in the direct system are injective and  $\varphi_i(x) = \varphi_i(y)$ . Then  $x \sim y$  in the quotient and hence, there exists  $i, j, k \in I$  such that  $f_{ik}(x) = f_{jk}(y)$  which implies that  $x = y$ .

Now assume that all  $f_{ij}$  are closed embeddings. Then, the inclusions are injective by the set-theoretic argument. Furthermore, the images of  $\varphi_i : A_i \rightarrow \text{colim}_i A_i$  are closed. That follows from subsets in the final topology being closed, if and only if all their preimages are closed. Hence we check if  $\varphi_j^{-1}(\varphi(A_i))$  is closed for all  $j \in I$ :

$$\begin{aligned} \text{If } j \leq i \text{ then: } & \varphi_j^{-1}(\varphi_i(A_i)) = (\varphi_i \circ f_{ji})^{-1}(\varphi_i(A_i)) = f_{ji}^{-1}(A_i) \text{ which is closed.} \\ \text{If } i < j \text{ then: } & \varphi_j^{-1}(\varphi_i(A_i)) = \underbrace{\varphi_j^{-1}(\varphi_j \circ f_{ij}(A_i))}_{\varphi \text{ is injective}} = f_{ij}(A_i) \text{ which is closed.} \end{aligned} \quad (2)$$

req:kTheo-sta  
irCaseMapsClo  
sedEmbedding

To show that  $\varphi_i^{-1} : \text{im}(\varphi_i) \rightarrow A_i$  is continuous we assume that  $B \subset A_i$  is open. Then  $\varphi_i(B)$  is open if and only if for any  $j$   $\varphi_j^{-1}(\varphi_i(B))$  is open. This follows from the same calculations as in equation (2) using that the  $f_{ij}$  are homeomorphisms onto their image.  $\square$

cor:kTheo-dir  
ectGrassmanSy  
stem

**Corollary 3.2.38.** The so-called inverse system given by the projections

$$\pi : \mathbb{C}^m \rightarrow \mathbb{C}^{m-1}, \quad (x_1, \dots, x_m) \mapsto (x_1, \dots, x_{m-1}),$$

induces a direct system of Grassmann manifolds with the induced continuous maps

$$i_{m-1} : \mathrm{Gr}_n(\mathbb{C}^{m-1}) \rightarrow \mathrm{Gr}_n(\mathbb{C}^m), \quad U \mapsto \pi^{-1}(U).$$

If we denote the classifying bundle over  $\mathrm{Gr}_n(\mathbb{C}^m)$  by  $E_m$ , then the pullback of  $E_m$  along  $i_{m-1}$  is isomorphic to  $E_{m-1}$ .

thm:kTheory-c  
lassifyingSpa  
ce

**Theorem 3.2.39.** *The map*

$$\mathrm{colim}_m[X, \mathrm{Gr}_n(\mathbb{C}^m)] \rightarrow \mathrm{Vect}_n(X),$$

*induced by  $f \mapsto f^*E_m$  is a bijection for all compact spaces  $X$ .*

*Proof.* We shall construct an inverse map. If  $E$  is a bundle over  $X$ , then by Corollary 3.2.25 there exists an epimorphism  $\varphi : X \times \mathbb{C}^m \rightarrow E$  for some integer  $m$ . Let  $f : X \rightarrow \mathrm{Gr}_n(\mathbb{C}^m)$  be the map induced by  $\varphi$ . If we manage to show that the homotopy class of  $f$  as a map to  $\mathrm{Gr}_n(\mathbb{C}^{m'})$  for some large enough  $m'$  is independent of the choice of  $\varphi$  and  $m$ , then we get a well defined map from  $\mathrm{Vect}_n(X)$  to  $\mathrm{colim}_m[X, \mathrm{Gr}_n(\mathbb{C}^m)]$  that is inverse to the classifying map.

Suppose that  $\varphi_i : X \times \mathbb{C}^{m_i} \rightarrow E$  for  $i = 1, 2$  are two epimorphisms. Let  $g_i : X \rightarrow \mathrm{Gr}_n(\mathbb{C}^{m_i})$  be the maps induced by  $\varphi_i$ . We want to show that  $g_1$  and  $g_2$  are homotopic as maps to  $\mathrm{Gr}_n(\mathbb{C}^{m_1+m_2})$ . For this we define the map

$$\psi_t : X \times \mathbb{C}^{m_1} \times \mathbb{C}^{m_2} \rightarrow E, \quad \psi_t(x, v_1, v_2) = t\varphi_1(x, v_1) + (1-t)\varphi_2(x, v_2).$$

Since the  $\varphi_i$  are epimorphisms, this map is an epimorphism for any  $t$ . Let

$$f_t : X \rightarrow \mathrm{Gr}_n(\mathbb{C}^{m_1+m_2})$$

be the maps induced by  $\psi_t$ . (where we identified  $\mathbb{C}^{m_1} \times \mathbb{C}^{m_2}$  with  $\mathbb{C}^{m_1+m_2}$  by mapping  $(v_1, v_2)$  to  $(v_1, v_2)$ ). Then

$$f_0 = j_0 \circ g_1 \quad \text{and} \quad f_1 = T \circ j_1 \circ g_2,$$

where  $j_i : \mathrm{Gr}_n(\mathbb{C}^{m_i}) \rightarrow \mathrm{Gr}_n(\mathbb{C}^{m_1+m_2})$  is the map induced by the inclusion  $\mathbb{C}^{m_i} \rightarrow \mathbb{C}^{m_1+m_2}$  and  $T$  is the map induced by the isomorphism  $\mathbb{C}^{m_1+m_2} \rightarrow \mathbb{C}^{m_1+m_2}$  that sends  $(v_1, v_2)$  to  $(v_2, v_1)$ . Since  $T$  is homotopic to the identity, we can conclude that  $j_0 \circ g_1$  and  $j_1 \circ g_2$  are homotopic as maps to  $\mathrm{Gr}_n(\mathbb{C}^{m_1+m_2})$ . Hence, the homotopy class of the map induced by an epimorphism is independent of the choice of the epimorphism and the integer  $m$ . This concludes the proof.  $\square$

lem:kTheo-col  
imUndHomVerta  
uschen

**Lemma 3.2.40.** *For any compact  $X$  and direct system  $((Y_m)_{m \in \mathbb{N}}, (i_m)_{m \in \mathbb{N}})$  where all  $Y_m$  are Hausdorff and all  $i_m$  are closed embeddings we have a canonical bijection between the set theoretic direct limit of homotopy classes of maps and the*

homotopy classes of maps to the topological direct limit:

$$\operatorname{colim}_m [X, X_m] \cong [X, \operatorname{colim}_m X_m].$$

*Proof.* We denote  $\operatorname{colim}_m Y_m = Y$  and the corresponding inclusion  $\varphi_m : Y_m \rightarrow Y$ . Since  $X$  is compact, any map  $f : X \rightarrow Y$  has image that lies in some  $\operatorname{im}(\varphi_m)$  for  $m$  big enough. Assume that this was not the case, then there would exist an increasing sequence  $(j_i)_{i \in \mathbb{N}}$  such that for all  $i$  there is an  $x_i$  with  $f(x_i) \in \operatorname{im} \varphi_{j_i} \setminus \operatorname{im} \varphi_{j_i-1}$ . With this we define the set

$$A := \{f(x_i) \mid i \in \mathbb{N}\} \subseteq \operatorname{im} f \subseteq Y.$$

Then  $A$  is closed, since for all  $m$  the intersection  $\operatorname{im} \varphi_m \cap A$  is finite and since  $\operatorname{im}(\varphi_m)$  is closed, said intersection is closed. By the final topology we can conclude that the whole  $A$  is closed and since it is contained in the image of  $f$  it is compact (Here we use that the  $Y_i$  are Hausdorff!). Furthermore, the same argument shows that every subset  $B \subset A$  is closed, because again  $B \cap \operatorname{im} \varphi_m$  is finite. This tells us that  $A$  as a subset of  $Y$  carries the discrete topology: Let  $C \subseteq A$  be an arbitrary subset. Then  $A \setminus C \subseteq A$  is closed in  $Y$  and hence its complement in  $Y$  is open and intersected with  $A$  yields  $C$ . Hence,  $C$  is open in the subspace topology. This however is a contradiction, as  $A$  cannot be infinite, compact and carry the discrete topology. Hence any map  $f : X \rightarrow Y$  has image that lies in some  $\operatorname{im}(\varphi_m)$  for  $m$  big enough. But then there exists a  $f_m$  such that  $f = \varphi_m \circ f_m$ . This assignment is well defined on homotopy classes because if  $f, s : X \rightarrow Y$  are homotopic, then there exists a homotopy  $H : X \times I \rightarrow Y$  between them: By the same argument as before, the image of  $H$  is compact and hence contained in some  $\operatorname{im} \varphi_m$ . Hence,  $f_m$  and  $s_m$  are homotopic via  $H_m$  as maps to  $\operatorname{im} \varphi_m$ . Hence, we get a well defined canonical map

$$\Phi : [X, \operatorname{colim}_m Y_m] \rightarrow \operatorname{colim}_m [X, Y_m].$$

This map is injective, as if  $f_m$  and  $s_m$  are homotopic as maps to  $\operatorname{im} \varphi_m$ , then they are also homotopic as maps to the direct limit. Finally  $\Phi$  is surjective, as any element of the direct limit is represented by some  $f_m : X \rightarrow Y_m$  and hence  $\Phi([\varphi_m \circ f_m]) = [f_m]$ .  $\square$

**Definition 3.2.41.** We denote  $\operatorname{colim}_m \operatorname{Gr}_n(\mathbb{C}^m)$  by  $\operatorname{BU}(n)$  and call it the **classifying space** for complex vector bundles of rank  $n$ . The name originates from the functor  $B$  that sends a topological group  $G$  to the classifying space of vector bundles with structure group  $G$ .

**Corollary 3.2.42.** For any compact space  $X$ , there is a canonical bijection between the homotopy classes of maps from  $X$  to  $\operatorname{BU}(n)$  and the isomorphism classes of vector bundles of rank  $n$  over  $X$ :

$$[X, \operatorname{BU}(n)] \cong \operatorname{Vect}_n(X).$$

*Proof.* This is a direct application of Lemma 3.2.40.  $\square$

### 3.3 K-Theory as a Cohomology Theory

**Definition 3.3.1.** We need the following categories of topological spaces:

- **CWH** is the category of compact spaces.
- $\mathbf{CWH}_{\text{cof}}^*$  is the category of well-pointed compact spaces.
- $\mathbf{CWH}_{\text{cof}}^2$  is the category of pairs of compact spaces, where the inclusions are cofibrations.

Here, we have the natural functors:

- $\mathbf{CWH}_{\text{cof}}^2 \rightarrow \mathbf{CWH}_{\text{cof}}^*$  by sending a pair  $(X, Y) \mapsto (X/Y, [Y])$  with obvious morphisms. If  $Y = \emptyset$  this is the disjoint union of  $X$  with a point.
- $\mathbf{CWH} \rightarrow \mathbf{CWH}_{\text{cof}}^2$  by sending  $X \mapsto (X, \emptyset)$  again with obvious morphisms. The inclusion of  $\emptyset$  into  $X$  is a cofibration, as the empty set itself is an open neighbourhood and hence the pair is a NDR pair.
- $\mathbf{CWH} \rightarrow \mathbf{CWH}_{\text{cof}}^*$  as the composition. Again, the inclusion of  $[\emptyset]$  into the disjoint union  $X \sqcup [\emptyset]$  is a cofibration, as  $\{[\emptyset]\}$  is an open neighbourhood of itself.

If we descend to the category where the morphisms are homotopy classes of maps, we get the categories  $\mathbf{hCWH}$ ,  $\mathbf{hCWH}_{\text{cof}}^*$  and  $\mathbf{hCWH}_{\text{cof}}^2$ . Furthermore, the construction gives a functor from the category of abelian semigroups to the category of abelian groups.

**Definition 3.3.2** (Grothendieck Group and Ring). Let  $A$  be an abelian semigroup. Then the **Grothendieck group** of  $A$  is the abelian group  $K(A)$  together with a homomorphism  $\varphi : A \rightarrow K(A)$  such that for any other group  $G$  together with a homomorphism  $\psi : A \rightarrow G$ , there exists a unique homomorphism  $\tilde{\psi} : K(A) \rightarrow G$  such that the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & K(A) \\ & \searrow \psi & \downarrow \tilde{\psi} \\ & & G \end{array}$$

If  $G$  is a semiring, then  $K(G)$  inherits a ring structure and we call  $K(G)$  the **Grothendieck ring** of  $G$ .

**Corollary 3.3.3.** The Grothendieck group exists for any abelian semigroup and is unique up to canonical isomorphism. If  $G$  is a semiring, then the Grothendieck ring exists and is unique up to canonical isomorphism.

*Proof.* The construction follows the following idea: We define tuples  $(a, b)$  that are to be interpreted as differences “ $a - b$ ”. For this construction to yield something useful we want to enforce the rule that “ $a - b = a + c - (b + c)$ ” or said differently  $(a + c, b + c) = (a, b)$ . Finally, to account for the universal property, we want to use the smallest relation

def:kTheo-gro  
thendiekGroup  
AndRing

cor:kTheo-Con  
structionOfGro  
thendiekGroup

that obeys said rule. This gives us the following construction: Let  $A$  be an abelian semigroup and  $\Delta : A \rightarrow A \times A$  be the diagonal homomorphism that sends  $a$  to  $(a, a)$ . Let  $K(A) := A \times A / \Delta(A)$ . By this notation we mean:  $\Delta(A)$  acts on  $A \times A$  via

$$\Delta(A) \times (A \times A) \rightarrow A \times A \quad ((a, a), (b, c)) \mapsto (a + b, a + c).$$

With this we have the set of orbits which gives rise to a relation:

$$\mathcal{R} := \{((a, b), (a + c, b + c)) | c \in A\}$$

Now we take the smallest equivalence relation on  $A \times A$  that contains  $\mathcal{R}$  (This relation is made explicit in Remark 3.3.4). We define  $K(A)$  to be  $A \times A$  modulo said relation. Then  $K(A)$  is a priori an abelian semigroup, but for  $a, b \in A$  we have

$$(a, b) + (b, a) = (a + b, b + a) \in \Delta(A),$$

and hence  $K(A)$  is an abelian group. Also we can see that

$$\overline{(a, b)} = \overline{(a, 0)} - \overline{(b, 0)} = \alpha_A(a) - \alpha_A(b)$$

Together with the homomorphism  $\alpha_A : A \rightarrow K(A)$  that sends  $a$  to the class of  $(a, 0)$ , we get the universal property of the Grothendieck group:

Let  $G$  be another abelian group and  $\psi : A \rightarrow G$  a semigroup homomorphism. From this define

$$\tilde{\psi} : K(A) \rightarrow G, \quad \overline{(a, b)} \mapsto \psi(a) - \psi(b)$$

The well definition of this map can be verified by a calculation. By calculating it also follows that  $\tilde{\psi}$  is a group homomorphism and that  $\tilde{\psi} \circ \alpha_A = \psi$ . Hence, for the universal property we only need to verify uniqueness. So assume that  $\theta : K(A) \rightarrow G$  is another group homomorphism. Then:

$$\theta(\overline{(a, b)}) = \underbrace{\theta(\alpha_A(a))}_{\psi(a)} - \underbrace{\theta(\alpha_A(b))}_{\psi(b)} = \tilde{\psi}(\overline{(a, b)})$$

This concludes the construction of the Grothendieck group. To see the functoriality we first need to define what  $K(\beta)$  means, when  $\beta : A \rightarrow B$  defines a semigroup homomorphism between two abelian semigroups. For this we use the universal property and define  $K(\beta)$  to be the unique group homomorphism coming from  $\alpha_B \circ \beta : A \rightarrow K(B)$ . Now the uniqueness of said definition shows the equalities  $K(\text{id}) = \text{id}$  and  $K(f \circ g) = K(f) \circ K(g)$ .  $\square$

rem:kTheo-gro  
thendieckEqui  
valence

**Remark 3.3.4.** In  $K(A)$  we get the following equivalence for any two pairs  $\overline{(a, b)}$  and  $\overline{(a', b')}$ :

$$\overline{(a, b)} = \overline{(a', b')} \Leftrightarrow \text{There exists a } c \in A \text{ such that } (a + b' + c) = (a' + b + c).$$

*Proof.* We start by proving “ $\Leftarrow$ ”. This direction follows from the properties of an equivalence relation, as then, every element of an equivalence class is also a representative:

$$\overline{(a, b)} = \overline{(a + (a' + b + c), b + (a + b' + c))} = \overline{(a' + (a + b + c), b' + (a + b + c))} = \overline{(a', b')}. \quad (3)$$

The other direction follows from the minimality of the equivalence relation: To see this notice how the argument above works in every equivalence relation containing the orbits of  $\Delta(A)$  and hence works in the smallest one. Furthermore,

$$(a, b) \sim (a', b') \Leftrightarrow \text{There exists a } c \in A \text{ such that } (a + b' + c) = (a' + b + c)$$

defines an equivalence relation that contains the orbits and hence this is the smallest one.  $\square$

**Remark 3.3.5** (A simpler Construction of the Grothendieck Group?). One might be wondering, why we defined the relation on the pair  $A \times A$  as the smallest equivalence relation starting by identifying the orbits of the operation of the diagonal, when Remark 3.3.4 might look and feel different. The reason for this detour is that I wanted to understand what Atiyah meant when he defined in Chapter II.1 [Ati89] word for word:

*A slightly different construction of  $K(A)$  which is sometimes convenient is the following: Let  $\Delta : A \rightarrow A$  be the diagonal homomorphism of semi-groups, and let  $K(A)$  denote the set of cosets of  $\Delta(A)$  in  $A \times A$ . It is a quotient semi-group, but the interchange of factors in  $A \times A$  introduces an inverse in  $K(A)$  so that  $K(A)$  is a group.*

Whilst this definition feels easy to the reader, it carries a lot of baggage. First of all, if one is not familiar with commutative monoid or abelian semigroup theory, one might want to generalise the notion of coset from group theory that states that for a subgroup  $A \subseteq G$  of an abelian group  $G$  the cosets are of the form  $g + A$ . Whilst this notion can be generalised to semigroups, it would yield something useless. To see this just take  $\mathbb{N}$ . In the semigroup  $\mathbb{N} \times \mathbb{N} / \Delta(\mathbb{N})$  the cosets  $(2, 0) + \mathbb{N}$  and  $(3, 1) + \mathbb{N}$  are different, as  $(2, 0)$  is not contained in the latter. To give Atiyah the benefit of the doubt we could generalise the definition of cosets as orbits of the right translation, which is what we did in our construction. However, the literature on semigroups from that time defines the notion of quotients in the setting of semigroups with the help of *congruence* instead of cosets. The latter refers to an equivalence relation that behaves well with the operation of said semigroup. Compare Chapter 1.5 [CP61]. It was even shown that the notion of congruence and coset are equivalent for groups (compare for example: Chapter 6.1 [Gre65]). Regardless of what Atiyah meant by his notion of coset, the construction became known and cited. ([Bro96], [Lan05], [Cou04]) None of those references define coset in semigroups and hence this little detour into the history should function as a reminder, that we need to be careful when generalising things from the known categories. We are so used to cosets giving equivalence relations, that it's easy to overlook the needed care.

**Remark 3.3.6.** We will omit the inclusion maps  $\alpha$  from now on and interpret (if not said different) the equivalence class  $[E]$  of a vector bundle as an element in  $K(X)$ .

{eq:kTheo-the  
GrothendieckRe  
lation}

**Corollary 3.3.7.** In the proof above we have shown that the assignment  $K$  is a functor from the category of abelian semigroups to the category of abelian groups. Alternatively, we can view  $K$  as a functor from the category of semirings to the category of rings.

**Definition 3.3.8.** Let  $X$  be a compact Hausdorff space. We define the **K-group** of  $X$  to be the Grothendieck ring of the semiring  $\text{Vect}(X)$  of isomorphism classes of vector bundles over  $X$  with the direct sum as addition and the tensor product as multiplication. We denote the K-group of  $X$  by  $K(X)$ . As a whole we get a contravariant functor from the category of compact Hausdorff spaces to the category of rings

$$\mathbf{CWH} \rightarrow \mathbf{Ring} . \quad (4)$$

**Remark 3.3.9.** For a continuous map  $f : X \rightarrow Y$  we denote the induced map  $K(f^*) : K(Y) \rightarrow K(X)$  by just  $f^*$ .

**Theorem 3.3.10.** *The functor  $K : \mathbf{CWH} \rightarrow \mathbf{Ring}$  factors through the homotopy category  $\mathbf{hCWH}$  of compact Hausdorff spaces together with homotopy classes of maps.*

*Proof.* This directly follows from Theorem 3.2.4. □

def:kTheo-reducedKTheory

**Definition 3.3.11** (Reduced K-Theory). Let  $X$  be a compact Hausdorff space and  $x_0 \in X$  be a base point. We define the **reduced K-group**  $\tilde{K}(X)$  of  $X$  to be the kernel of the map  $K(X) \rightarrow K(\{x_0\})$  induced by the inclusion  $\{x_0\} \rightarrow X$ .

cor:kTheo-splittingReducedGroupNatural

**Corollary 3.3.12.** If  $c : X \rightarrow x_0$  is the collapsing map sending each point to the base-point, then we get a natural splitting

$$K(X) \cong \tilde{K}(X) \oplus K(x_0) .$$

Naturality here means, that for any homomorphism of pointed spaces  $f : (X, x_0) \rightarrow (Y, y_0)$  we have a commuting diagram:

$$\begin{array}{ccc} K(X) & \longrightarrow & \tilde{K}(X) \oplus K(x_0) \\ \downarrow f^* & & \downarrow \tilde{K}(f) \oplus (\{x_0\} \rightarrow \{y_0\})^* \\ K(Y) & \longrightarrow & \tilde{K}(Y) \oplus K(y_0) \end{array}$$

*Proof.* We have the exact sequence of rings:

$$0 \longrightarrow \tilde{K}(X) \xrightarrow{\text{inclusion}} K(X) \xrightarrow{i_X^*} K(x_0) \longrightarrow 0$$

where  $c_X^* : K(x_0) \rightarrow K(X)$  is a right inverse of  $i_X^*$ , and hence the sequence splits and  $K(X) \cong \tilde{K}(X) \oplus K(x_0)$ . The isomorphism is explicitly given by

$$K(X) \rightarrow \tilde{K}(X) \oplus K(x_0), \quad \xi \mapsto (\xi - c_X^* \circ i_X^*(\xi), i_X^*(\xi))$$

and its inverse maps  $(a, b) \mapsto a + c_X^*(b)$ . The splitting is natural with respect to maps in  $\mathbf{CWH}_{\text{cof}}^*$ , because for a map of pointed spaces  $f : X \rightarrow Y$  we get commutativity:

diag:kTheo-NaturalSplitting

$$\begin{array}{ccccccc}
0 & \longrightarrow & \tilde{K}(X) & \hookrightarrow & K(X) & \xrightarrow{i_X^*} & K(x_0) \longrightarrow 0 \\
& & \uparrow f^*|_{\tilde{K}(Y)} & & \uparrow f^* & & \uparrow (f|_{x_0})^* \\
0 & \longrightarrow & \tilde{K}(Y) & \hookrightarrow & K(Y) & \xrightarrow{i_Y^*} & K(y_0) \longrightarrow 0
\end{array}$$

$\xleftarrow{c_X^*}$ 
 $\xleftarrow{c_Y^*}$

The well definition of  $f^*|_{\tilde{K}(Y)}$  follows immediately from the commutativity. The commutativity itself follows from  $K$  being a contravariant functor and all topological maps commute. Finally, define  $\tilde{K}(f) := f^*|_{\tilde{K}(Y)}$  and we are done, since the naturality of the splittings, i.e. commutativity of the diagram in the corollary, then follows from commutativity of each square separately: the left square controls the  $\tilde{K}$ -summand and the right square (together with the sections  $c_X^*, c_Y^*$ ) controls the  $K(x_0)$ -summand.  $\square$

**Remark 3.3.13.** For a map  $f : (X, x_0) \rightarrow (Y, y_0)$  we denote  $\tilde{K}(f)$  and  $K(f)$  by  $f^*$ . The context will make clear if we work in the reduced theory or not.

**Corollary 3.3.14.**  $\tilde{K}$  is a functor from  $\mathbf{CWH}_{\text{cof}}^*$  to  $\mathbf{Ring}$ .

*Proof.* The proof follows immediately from the naturality of the splitting in Corollary 3.3.12.  $\square$

cor:kTheo-unpointedReducedIsNonReduced

**Corollary 3.3.15.** Let  $X$  be a compact Hausdorff space, and let

$$F : \mathbf{CWH} \rightarrow \mathbf{CWH}_{\text{cof}}^*$$

be the functor that sends  $X$  to  $(X, [\emptyset])$  with obvious homomorphisms. Then there is a natural isomorphism between the functors  $\tilde{K} \circ F$  and  $K$ .

*Proof.* This follows from the fact that disjoint unions give rise to direct sums of semirings:  $\text{Vect}(X \sqcup \{\emptyset\}) \cong \text{Vect}(X) \oplus \text{Vect}(\{\emptyset\})$  and the isomorphism is given by gluing the bundles and hence natural. Furthermore, the Grothendieck ring of a direct sum gives the direct sum of rings (again in a natural manner). (This can be verified using the universal property). Hence we get the commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \tilde{K}(F(X)) & \xrightarrow{\text{inclusion}} & K(F(X)) & \xrightarrow{i^*} & K([\emptyset]) \longrightarrow 0 \\
\parallel & & & & \downarrow \Phi & & \parallel \\
0 & \longrightarrow & K(X) & \xrightarrow{i_1} & K(X) \oplus K([\emptyset]) & \xrightarrow{\pi_2} & K([\emptyset]) \longrightarrow 0
\end{array}$$

But then we can define the map  $\Phi|_{\tilde{K}(F(X))}$  and diagram yoga shows that this is an isomorphism. Furthermore, since  $\Phi$  is natural, the new map is also natural and defines a natural transformation between the two functors.  $\square$

**Definition 3.3.16.** Let  $F : \mathbf{CWH}_{\text{cof}}^2 \rightarrow \mathbf{CWH}_{\text{cof}}^*$  be the functor that collapses the second space. Then the composition  $\tilde{K} \circ F$  defines a contravariant functor.

**Definition 3.3.17.** Let  $(X, x_0)$  and  $(Y, y_0)$  be two pointed spaces. Then we define the **pointed sum** as:

$$X \vee Y := (X \sqcup Y) / (x_0 \sim y_0).$$

With this, we can define the **smash product** as:

$$X \wedge Y := (X \times Y) / (X \vee Y)$$

**Definition 3.3.18.** For the sake of intuition, I want to give a bit of insight into tensor products without being too technical and opening Pandora's box of monoidal categories. We define a tensor product in a category simply as the space over which bilinear maps factor uniquely. To capture bilinearity in such a category we define the following. Let  $\pi_i : X_0 \times X_1 \rightarrow X_i$  be the canonical projection for  $i = 0, 1$ . Any reasonable product should come with such surjective maps. Since those are surjective, the  $\pi_i$  admit a family of right inverses  $r_i : X_i \rightarrow X_0 \times X_1$ . We call such a right inverse “product preserving” if the compositions  $\pi_2 \circ r_1$  and  $\pi_1 \circ r_2$  are constant. Hence the right inverse maps into just one fibre. Now bilinearity in this general setting states, that for any product preserving right inverse of the projections  $r_i$ , the composition  $f \circ r_i$  is structure preserving.

**Remark 3.3.19** (What is the Smash Product). We can think of the smash product as the tensor product in the pointed topological spaces as follows. Let  $f : (X, x_0) \times (Y, y_0) \rightarrow (Z, z_0)$  be bilinear. For any  $y \in Y$  we have the product preserving right inverse of  $\pi_1$  given by

$$r_1^y : (X, x_0) \rightarrow (X, x_0) \times (Y, y_0), \quad x \mapsto (x, y).$$

Analogously we define  $r_2^x$  for any  $x \in X$  as

$$r_2^x : (Y, y_0) \rightarrow (X, x_0) \times (Y, y_0), \quad y \mapsto (x, y).$$

Now for any  $x \in X$  we have  $f(x, y_0) = f \circ i_Y^x(y_0) = z_0$  as the composition is said to be basepoint preserving. Similarly we see that  $f(x_0, y) = z_0$  for any  $y \in Y$ . Hence  $f$  factors through the quotient space

$$X \times Y / (X \times \{y_0\} \cup (Y \times \{x_0\})),$$

where the numerator is the canonically embedded  $X \vee Y$  into the product. Furthermore by the universal property of quotient spaces, this factorisation is unique.

**Example 3.3.20.** Let  $S^n$  be the  $n$ -dimensional sphere given by all points of norm one in  $\mathbb{R}^{n+1}$ . Then a general fact is that  $S^n \cong I^n / \partial I^n$  and hence for all  $n, m > 0$  we have  $S^n \wedge S^m \cong S^{n+m}$ . Later we will make this homeomorphism canonical.

ref

def:orient-fr  
omRelaSpaceTo  
TheSphere

**Definition 3.3.21** (From Real Space to the Sphere). We now want to construct a canonical homeomorphism between  $S^m$  and  $\overline{\mathbb{R}^m}$ . First we use the stereographic projection from the South Pole: If  $m = 0$  we define this to be

$$\pi_s : S^0 = \{-1, 1\} \setminus \{-1\} \rightarrow \mathbb{R}^0, \quad 1 \mapsto 0.$$

For  $m > 0$  we define:

$$\begin{aligned} \pi_S : S^m \setminus \{S\} \subseteq \mathbb{R}^{m+1} &\rightarrow \mathbb{R}^m \\ (x_1, \dots, x_m, x_{m+1}) &\mapsto \left( \frac{x_1}{1+x_{m+1}}, \dots, \frac{x_m}{1+x_{m+1}} \right) \end{aligned}$$

with its inverse:

$$\begin{aligned} \pi_S^{-1} : \mathbb{R}^m &\rightarrow S^m \setminus \{S\} \subseteq \mathbb{R}^{m+1} \\ (y_1, \dots, y_m) &\mapsto \left( \frac{2y_1}{1+\|y\|^2}, \dots, \frac{2y_m}{1+\|y\|^2}, \frac{1-\|y\|}{1+\|y\|} \right) \end{aligned}$$

Those are inverse maps and they give rise to maps between their one-point-compactifications. Hence  $\overline{\mathbb{R}^m} \cong S^m$ . Lets calculate  $\pi_S^{-1} \circ \pi_S$ , since we will later need a little bit of insight in those calculations:

$$\begin{aligned} \pi_S^{-1} \circ \pi_S(x_1, \dots, x_{m+1}) &= \pi_S^{-1} \left( \frac{x_1}{1+x_{m+1}}, \dots, \frac{x_m}{1+x_{m+1}} \right) \\ &= \left( \frac{2 \frac{x_1}{1+x_{m+1}}}{1 + \sum_{i=1}^m \left( \frac{x_i^2}{(1+x_{m+1})^2} \right)}, \dots, \frac{2 \frac{x_m}{1+x_{m+1}}}{1 + \sum_{i=1}^m \left( \frac{x_i^2}{(1+x_{m+1})^2} \right)}, \frac{1 - \sum_{i=1}^m \left( \frac{x_i^2}{(1+x_{m+1})^2} \right)}{1 + \sum_{i=1}^m \left( \frac{x_i^2}{(1+x_{m+1})^2} \right)} \right) \end{aligned}$$

Before we keep calculating, we notice the denominator can be rewritten as:

$$1 + \sum_{i=1}^m \left( \frac{x_i^2}{(1+x_{m+1})^2} \right) = \frac{(1+x_{m+1})^2 + \sum_{i=1}^m x_i^2}{(1+x_{m+1})^2}$$

And hence:

$$\frac{2 \frac{x_j}{1+x_{m+1}}}{1 + \sum_{i=1}^m \left( \frac{x_i^2}{(1+x_{m+1})^2} \right)} = \frac{2 \frac{x_j}{1+x_{m+1}}}{\frac{(1+x_{m+1})^2 + \sum_{i=1}^m x_i^2}{(1+x_{m+1})^2}} \quad (5)$$

$$= \frac{2x_j(1+x_{m+1})}{(1+x_{m+1})^2 + \sum_{i=1}^m x_i^2} \quad (6)$$

$$= \frac{2x_j(1+x_{m+1})}{(1+x_{m+1})^2 + \sum_{i=1}^m x_i^2} \quad (7)$$

$$= \frac{2x_j(1+x_{m+1})}{1 + 2x_{m+1} + \underbrace{\sum_{i=1}^{m+1} x_i^2}_{=1}} \quad (8)$$

$$= x_j \quad (9)$$

And the calculation of the last term is:

$$\begin{aligned}
 \frac{1 - \sum_{i=1}^m \left( \frac{x_i^2}{(1+x_{m+1})^2} \right)}{1 + \sum_{i=1}^m \left( \frac{x_i^2}{(1+x_{m+1})^2} \right)} &= \frac{\frac{(1+x_{m+1})^2 - \sum_{i=1}^m x_i^2}{(1+x_{m+1})^2}}{\frac{(1+x_{m+1})^2 + \sum_{i=1}^m x_i^2}{(1+x_{m+1})^2}} \\
 &= \frac{(1+x_{m+1})^2 - \sum_{i=1}^m x_i^2}{(1+x_{m+1})^2 + \sum_{i=1}^m x_i^2} \\
 &= \frac{1 + 2x_{m+1} + x_{m+1}^2 - \sum_{i=1}^m x_i^2}{1 + 2x_{m+1} + \underbrace{x_{m+1}^2 + \sum_{i=1}^m x_i^2}_{=1}} \\
 &= \frac{1 + 2x_{m+1} + 2x_{m+1}^2 - \sum_{i=1}^{m+1} x_i^2}{2 + 2x_{m+1}} \\
 &= \frac{x_{m+1}(2 + 2x_{m+1})}{2 + 2x_{m+1}} = x_{m+1}
 \end{aligned}$$

This concludes the calculation of  $\pi_S^{-1} \circ \pi_s = \text{id}$  and  $\pi_s \circ \pi_S^{-1} = \text{id}$  can be calculated in a similar manner.

**Definition 3.3.22** (One-Point Compactification). For a topological space  $X$  we define  $\bar{X} := X \sqcup \{\infty\}$ , where open sets are either open subsets of  $X$ , or sets of the form  $\{\infty\} \sqcup X \setminus C$  where  $C$  is compact and closed in  $X$ . For non compact Hausdorff spaces this is the one-point compactification that is characterised by being the unique (up to homeomorphism) compactification that contains  $X$  and one more point. If  $X$  is already compact, then  $\{\infty\}$  is an open subset of  $\bar{X}$  (just choose  $C = K$ ) and thereby isolated. Hence  $\bar{X}$  is the pointing of  $X$  given by  $(X, [\emptyset])$  from Corollary 3.3.15.

lem:orient-quotientIsCompactification

**Lemma 3.3.23** (Open subspaces of compact spaces). *Let  $Z$  be a compact Hausdorff space and  $U \subseteq Z$  open. Then the map*

$$Z/(Z \setminus U) \rightarrow \bar{U}, \quad [z] \mapsto z \text{ for } z \in U, \quad [Z \setminus U] \mapsto \infty$$

*is a homeomorphism.*

*Proof.* The map is the quotient of the function  $\varphi : Z \rightarrow \bar{U}$  that is the identity on  $U$  and sends  $Z \setminus U$  constant to  $\infty$ . Hence, we only need to prove continuity for  $\varphi$ . Nor  $O \subseteq \bar{U}$  is open, if it is open in  $U$ , or the complement of a compact set  $C \subset U$  in  $\bar{U}$ . In both cases,  $\varphi^{-1}(U)$  is open and hence  $\varphi$  is continuous. Since the quotient map is now a continuous bijection from a compact space to a Hausdorff space, it is a homeomorphism.  $\square$

lem:orient-compactificationOfProduct

**Lemma 3.3.24** (Compactification of a product). *Let  $X$  and  $Y$  be locally compact Hausdorff spaces, and point  $\bar{X}$  and  $\bar{Y}$  at their respective points at infinity. Then*

there is a homeomorphism

$$\Xi_{X,Y} : \overline{X \times Y} \xrightarrow{\cong} \overline{X} \wedge \overline{Y}$$

which is the identity on  $X \times Y$  and sends  $\infty$  to the base point. It is natural in the sense that for proper maps  $f : X \rightarrow X'$  and  $g : Y \rightarrow Y'$  the square

$$\begin{array}{ccc} \overline{X \times Y} & \xrightarrow{\Xi_{X,Y}} & \overline{X} \wedge \overline{Y} \\ f \times g \downarrow & & \downarrow \bar{f} \wedge \bar{g} \\ \overline{X' \times Y'} & \xrightarrow{\Xi_{X',Y'}} & \overline{X'} \wedge \overline{Y'} \end{array}$$

commutes.

*Proof.* Let  $Z := \overline{X \times Y}$ . Then  $Z$  is a compact Hausdorff space that contains  $U := X \times Y$  as an open subset. The complement of  $U$  consists of those points where one of the coordinates is infinit and hence

$$Z \setminus U = (\{\infty\} \times \overline{Y}) \cup (\overline{X} \times \{\infty\}) = \overline{X} \vee \overline{Y}.$$

Therefore, by definition of the smash product,  $Z/(Z \setminus U) = \overline{X} \wedge \overline{Y}$  and by Lemma 3.3.23 we can conclude that

$$\overline{X \times Y} \cong (Z/(Z \setminus U)) = \overline{X} \wedge \overline{Y}.$$

For the naturality notice that the product of proper maps is proper and hence all maps are continuous. The commutation then comes down to checking where infinity gets mapped which is always the basepoint of the wedge product.  $\square$

lem:kTheo-Sma  
schIsAssociat  
ive

**Lemma 3.3.25** (Associativity and Kommutativität). *For pointed compact spaces  $(A, a_0), (B, b_0), (C, c_0)$  there is a natural homeomorphism*

$$(A \wedge B) \wedge C \rightarrow A \wedge (B \wedge C)$$

*That restricts to the identity over  $(A \setminus \{a_0\}) \times (B \setminus \{b_0\}) \times (C \setminus \{c_0\})$ . Furthermore, there is a natural homeomorphism*

$$A \wedge B \rightarrow B \wedge A$$

*that restricts to the change of coordinates over*

$$(A \setminus \{a_0\} \times B \setminus \{b_0\}) \rightarrow (B \setminus \{b_0\} \times A \setminus \{a_0\})$$

For the kommutativity we can proceed in the same way:

$$A \wedge B \cong \overline{(A \wedge B)^\circ} \cong \overline{(B \wedge A)^\circ} \cong B \wedge A \quad (10)$$

*Proof.* Notice how for all compact pointed spaces  $(X, x_0)$  the identity  $(x_0 \mapsto \infty)$  is a natural homeomorphism  $\overline{X \setminus \{x_0\}} \cong X$  and we denote  $X^\circ := X \setminus \{x_0\}$ . Furthermore, the smash product of compact spaces is again compact (as it is a quotient of a closed space). Hence we can just calculate

$$\begin{aligned} (A \wedge B) \wedge C &\cong \overline{(A \wedge B)^\circ} \wedge \overline{C^\circ} \\ &\cong \overline{(A^\circ \times B^\circ)} \wedge \overline{C^\circ} \\ &\cong \overline{(A^\circ \times B^\circ) \times C^\circ} \\ &\cong \overline{A^\circ \times (B^\circ \times C^\circ)} \\ &\cong A \wedge (B \wedge C). \end{aligned}$$

□

**Remark 3.3.26.** The statement about associativity can be extended to the category of  $\mathbf{CGWH}_{\text{cof}}^*$ , but not all pointed topological spaces.

ref  
einfügen

def:ktheo-wed  
geOfShoperesI  
sSphere

**Definition 3.3.27** (Convention). Let  $\pi_s^n$  denote the compactified stereographical projection  $\overline{S^n} \rightarrow \overline{\mathbb{R}^n}$ . With those methods we can now define the homeomorphism

$$(\pi_s^{n+m})^{-1} \circ \Xi^{-1} \circ (\pi_s^n) \wedge (\pi_s^m) : S^n \wedge S^m \rightarrow S^{n+m}.$$

Often we will identify smash product of spheres and for those identification we will always use this homeomorphism. For the composition to make sense we identify  $\mathbb{R}^n \times \mathbb{R}^m$  with  $\mathbb{R}^{n+m}$  (also for  $n$  or  $m$  equal zero.)

**Definition 3.3.28.** Let  $(X, x_0)$  be a pointed topological space. We define the **reduced suspension** of  $X$  to be

$$\Sigma X := S^1 \wedge X.$$

Furthermore, we define inductively  $\Sigma^n X := \Sigma(\Sigma^{n-1} X) \cong S^n \wedge X$  and using the principle of permanence we extend this definition to  $\Sigma^0 X := S^0 \wedge X \cong X$ . The last homeomorphism is canonical.

**Definition 3.3.29.** Let  $(X, x_0)$  be a pointed topological space. We define the **suspension** of  $X$  to be the space

$$S(X) := X \times I / \sim \quad (11)$$

where  $(x, t) \sim (x', t')$  if  $t = 1$  or  $t = 0$ . Here we identify the top to a point and the bottom to a different point.  $((x, 1) \sim (0, 0))$ . We view  $(S(X), (x, 1))$  as a pointed space.

In general the reduced and unreduced suspension differ. However, under the following condition there will be a natural homotopy equivalence:

lem:kTheo-wel  
lpointedImpli  
esCylinderCof  
ibration

**Lemma 3.3.30.** *If  $(X, x_0)$  is well-pointed, then the inclusion  $\{x_0\} \times I \hookrightarrow S(X)$  is a cofibration.*

*Proof.* This proof is omitted but is Proposition 5.1.16 in [tom08].  $\square$

cor:kTheo-red  
ucedSuspensio  
nisSuspension

**Corollary 3.3.31.** *If  $(X, x_0)$  is well pointed, then  $S(X)$  is homotopically equivalent to  $\Sigma(X)$ .*

*Proof.* By lemma 3.3.30 we conclude that  $(S(X), x_0 \times I)$  satisfies the homotopy extension property. Hence, by Lemma 1.2.4 we know that  $S(X) \simeq S(X)/x_0 \times I$  and the homeomorphism

$$S(X)/x_0 \times I \cong \Sigma(X)$$

concludes the proof.  $\square$

def:kTheo-low  
erKGroub

**Definition 3.3.32** (Lower K-groups). For  $n \geq 0$  we define:

$$\begin{aligned} \tilde{K}^{-n}(X, x_0) &:= \tilde{K}(\Sigma^n X) && \text{for } (X, x_0) \in \text{Ob}(\mathbf{CWH}_{\text{cof}}^*), \\ K^{-n}(X, Y) &:= \tilde{K}^{-n}(X/Y, [Y]) && \text{for } (X, Y) \in \text{Ob}(\mathbf{CWH}_{\text{cof}}^2), \\ K^{-n}(X) &:= K^{-n}(X, \emptyset) && \text{for } X \in \text{Ob}(\mathbf{CWH}). \end{aligned}$$

All those definitions give contravariant and homotopically invariant functors into the category of rings.

cor:kTheo-K(X)  
)isK(X,nix)

**Corollary 3.3.33.** Let  $X$  be a topological space. Then we have a natural isomorphism between the functors  $K^0(X)$  and  $K(X)$ . Furthermore, if  $(X, x_0)$  is a pointed topological space, we have a natural isomorphism between  $\tilde{K}$  and  $K^0$ .

*Proof.* When disentangling the definitions, we see that

$$K^0(X) = \tilde{K}(\Sigma^0(X, [\emptyset])) \cong \tilde{K}(X, [\emptyset]) \cong K(X)$$

where the first isomorphism comes from  $\Sigma^0$  being canonically isomorphic to the identity and the second isomorphism is given in Corollary 3.3.15. The natural isomorphism between  $K^0$  and  $\tilde{K}$  is given by definition and the canonical isomorphism  $X \cong S^0 \wedge (X)$ .  $\square$

**Remark 3.3.34.** If we have pointed topological spaces  $(X, x_0)$  or pairs  $(X, A)$  and want to denote  $K^0(X, pt)$  or  $K^0(X, A)$  we will often omit the exponent and just write  $K(X, x_0)$  and  $K(X, A)$ .

**Theorem 3.3.35** (Excision). *For any pair  $(X, A)$  and  $U \subseteq A$  open such that  $\bar{U} \subseteq \mathring{A}$  the inclusion*

$$i : (X \setminus U, A \setminus U) \rightarrow (X, A)$$

induces an isomorphism

$$i^* : K^{-n}(X, A) \rightarrow K^{-n}(X \setminus U, A \setminus U)$$

*Proof.* Since  $U$  is open and  $A$  is closed,  $(X \setminus U, A \setminus U)$  is a compact pair. The inclusion  $i$  induces a continuous bijection

$$\varphi : (X \setminus U) / (A \setminus U) \rightarrow X / A$$

which is a homeomorphism as a continuous bijection from a compact space to a Hausdorff space. Hence  $i^*$  is an isomorphism.  $\square$

### 3.4 Homotopy-Theoretic definition of $K$

**Definition 3.4.1.** For any natural number  $n \in \mathbb{N}$  we define  $\underline{n}$  to be the  $n$ -dimensional trivial bundle over a base space  $X$ . With this we define the inclusions

$$i_n : \text{Vect}_n(X) \rightarrow \text{Vect}_{n+1}(X), \quad \xi \mapsto \xi \oplus [\underline{1}],$$

and this gives rise to the direct limit  $\text{colim}_n \text{Vect}_n(X)$ .

**Definition 3.4.2** (Stable Equivalence). Two vector bundles  $E, F$  over the same base space are called **stably equivalent** if there exists an  $n$  such that  $E$  and  $F$  become isomorphic after adding the trivial bundle:

$$E \oplus \underline{n} \cong F \oplus \underline{n}.$$

**Lemma 3.4.3.** In  $K(X)$  two equivalence classes  $[E]$  and  $[F]$  are the same, if and only if  $E$  and  $F$  are stably equivalent.

*Proof.* Assume that two vector bundles satisfy

$$E \oplus \underline{n} \cong F \oplus \underline{n},$$

for some  $n \in \mathbb{N}$ . By abuse of notation we let  $E$  (and  $\underline{n}$ ) denote a vector bundle and its isomorphism class simultaneously. Then by Remark 3.3.4

$$[E] = \overline{(E, 0)} = \overline{(E + \underline{n}, \underline{n})} = \overline{(F + \underline{n}, \underline{n})} = \overline{(F, 0)} = [F].$$

On the other hand, if  $[E] = [F]$ , then by the same Remark we can conclude:

$$(E, 0) \sim (F, 0) \Leftrightarrow \text{there exists } G \text{ such that } E \oplus G = F \oplus G.$$

Now by Corollary 3.2.26 we can find a vector bundle  $G'$  such that  $G \oplus G' = \underline{n}$  for some  $n$ . Then

$$E \oplus G \oplus G' = F \oplus G \oplus G' \Rightarrow E \oplus \underline{n} = F \oplus \underline{n}$$

$\square$

def:kTheo-tri  
vialBundleAnd  
LimitInclusin  
os

**Definition 3.4.4.** We call two vector bundles  $E, F$  over the same base space **eventually equivalent**, if there exist  $n, m \in \mathbb{N}$  such that  $E \oplus \underline{n} \cong F \oplus \underline{m}$ .

**Lemma 3.4.5.** *The set of equivalence classes using the notion of eventual equivalence is the limit*

$$\operatorname{colim}_n \operatorname{Vect}_n(X)$$

*Proof.* Let  $\operatorname{EVect}(X)$  be the set of equivalence classes using the notion of eventual equivalence. This satisfies the universal property of the direct limit. To see this let  $f_m : \operatorname{Vect}_m(X) \rightarrow Y$  be a family of maps, that commute with the inclusions  $i_n$  from Definition 3.4.1. From those we define the map

$$f : \operatorname{EVect}(X) \rightarrow Y, \quad [\xi] \mapsto f_{\operatorname{rank}(\xi)}(\xi).$$

This map is well defined, because  $f$  commuting with the inclusions  $i_n$  is equivalent to  $f$  being invariant under eventual equivalence. Uniqueness follows from every element of  $\operatorname{EVect}(X)$  being represented by a vector bundle.  $\square$

**Lemma 3.4.6.** *The set of equivalence classes using the notion of eventual equivalence together with the direct sum is an abelian group*

*Proof.* Everything except inverses is clear. The neutral element is the class of  $0$ . For inverses we take a bundle  $E$  and by Corollary 3.2.26 there exists a  $G$  such that  $E \oplus G \cong \underline{n}$ . Now  $[E] + [G] = [\underline{n}] = [0]$ .  $\square$

lem:kTheo-  
ucedKisEventu-  
ally

**Lemma 3.4.7.** *For a compact connected space  $X$  there is a natural group isomorphism.*

$$K(X) \cong \mathbb{Z} \times \operatorname{EVect}(X) \cong \mathbb{Z} \times \operatorname{colim}_n \operatorname{Vect}_n(X).$$

*Furthermore if  $(X, p)$  is pointed, there is a natural group isomorphism*

$$\tilde{K}(X) \cong \operatorname{EVect}(X) \cong \operatorname{colim}_n \operatorname{Vect}_n(X).$$

*Proof.* Let  $K$  be an element of  $K(X)$ . Then by Corollary 3.3.3 we can write  $K$  as the difference of two classes of vector bundles  $K = [E] - [F]$ . We abuse the notation of equivalence classes in this proof meaning we won't differentiate between isomorphism classes of vector bundles in  $\operatorname{Vect}_n$ , eventually or stable isomorphism classes or equivalence classes of naturally included vector bundles in  $K(X)$ . The context should always be clear. With this we define the function

$$\operatorname{rank}(K) : K(X) \rightarrow \mathbb{Z}, \quad K = [E] - [F] \mapsto \operatorname{rank}(E) - \operatorname{rank}(F).$$

This is well defined as if  $K = [E] - [F] = [E'] - [F']$ , then by the relation in  $K(X)$  we know that there is an  $n$  such that

$$E \oplus F' \oplus \underline{n} \cong E' \oplus F \oplus \underline{n}.$$

and hence:

$$\text{rank}(E) - \text{rank}(F) = \text{rank}(E') - \text{rank}(F')$$

Furthermore, rank has a natural right inverse given by

$$r : \mathbb{Z} \rightarrow K(X), \quad m \mapsto \begin{cases} [m] - [0] & \text{if } m \geq 0 \\ [0] - [-m] & \text{if } m < 0. \end{cases}$$

Now we claim that the kernel  $\ker(\text{rank}) \cong \text{EVect}(X)$ . For this we define the map

$$\varphi : \text{EVect}(X) \rightarrow K(X), \quad [E] \mapsto [E] - [\underline{\text{rank}(E)}]$$

To see the well-definedness we assume that  $E$  and  $E'$  are eventually isomorphic and hence there are  $n, n' \in \mathbb{N}$  such that  $E \oplus \underline{n} \cong E' \oplus \underline{n'}$ . But then in  $K(X)$  we have:

$$[E] - [\underline{\text{rank}(E)}] = [E \oplus \underline{n}] - [\underbrace{\underline{\text{rank}(E) \oplus n}}_{=\underline{\text{rank}(F) \oplus n'}}] = [E' \oplus \underline{n'}] - [\underline{\text{rank}(F) \oplus n'}] = [F] - [\underline{\text{rank}(F)}].$$

Obviously  $\text{rank} \circ \varphi \equiv 0$ . To get the other inclusion we first notice the following: Let  $K \in K(X) = [E] - [F]$  be arbitrary. By Corollary 3.2.26 there exists a  $G$  such that  $F \oplus G \cong \underline{n}$  for some  $n \in \mathbb{N}$ . Hence if  $K$  is in the kernel of rank we have:

$$K = [E] - [F] = [E \oplus G] - [\underline{n}] = [E \oplus G] - [\underline{\text{rank}(E \oplus G)}] = \varphi([E \oplus G])$$

This leads to the naturally split short exact sequence

$$0 \longrightarrow \text{EVect}(X) \xrightarrow{\varphi} K(X) \xrightleftharpoons[\text{rank}]{r} \mathbb{Z} \longrightarrow 0$$

This already proves the isomorphisms

$$K(X) \cong \mathbb{Z} \times \text{EVect}(X) \cong \mathbb{Z} \times \text{colim}_n \text{Vect}_n(X).$$

Now we define the group isomorphism

$$l : \mathbb{Z} \rightarrow K(p), \quad m \mapsto \begin{cases} [m] - [0] & \text{if } m \geq 0 \\ [0] - [-m] & \text{if } m < 0. \end{cases}$$

This gives the commutative diagram:

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathrm{SVect}(X) & \xrightarrow{\varphi} & K(X) & \xrightarrow[\mathrm{rank}]{r} & \mathbb{Z} \longrightarrow 0 \\
& & & & \downarrow \mathrm{id} & & \downarrow l \\
0 & \longrightarrow & \tilde{K}(X) & \xrightarrow{\text{inclusion}} & K(X) & \xrightarrow{i^*} & K(p) \longrightarrow 0 \\
& & & & & \nwarrow c_p^* & \\
& & & & & & 
\end{array}$$

Where  $i$  is the inclusion  $\{p\} \rightarrow X$  and  $c_p$  is the collapse  $X \rightarrow \{p\}$ . Then by some diagram chasing the map

$$\tilde{\varphi} : \mathrm{SVect}(X) \rightarrow \tilde{K}(X)$$

is well defined and by the five lemma (add zeros to the short sequences on the left) it is an isomorphism.  $\square$

rem:kTheo-  
elementsInReduce  
dKtheoryAreGi  
venByEMionusN

**Remark 3.4.8.** In the proof above we have seen that every element  $k \in K(X)$  is given as the difference of a vector bundle and a trivial bundle  $k = [\xi] - [n]$  for some vector bundle  $\xi$  and  $n \in \mathbb{N}$ . We also saw that  $\tilde{K}(X)$  is in the kernel of rank and hence any  $k \in \tilde{K}(X)$  is given by  $k = [\xi] - [\mathrm{rank}(\xi)]$  where  $\xi$  is a vector bundle.

**Definition 3.4.9.** We define the **unitary group**  $U$  to be the direct limit of the unitary group of rank  $n$  under the standard inclusions

$$U(n) \rightarrow U(n+1), \quad A \mapsto A \oplus \mathbb{1}_1 = \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix}.$$

**Definition 3.4.10.** We now want to make  $BU(n)$  into a direct system. Notationally, this proof will have many maps with two indices. The convention is that lower indices show for which colimit the maps form a directed system. As an example, the maps  $i_m^n$  are used for the colimit over  $m$ . We start with the inclusion

$$j_m : \mathbb{C}^m \rightarrow \mathbb{C}^{m+1}, \quad (x) \mapsto (x, 0),$$

which gives rise to the maps

$$f_n^m : \mathrm{Gr}_n(\mathbb{C}^m) \rightarrow \mathrm{Gr}_{n+1}(\mathbb{C}^{m+1}), \quad V \mapsto j_m(V).$$

Those inclusions commute with the maps

$$i_m^n : \mathrm{Gr}_n(\mathbb{C}^m) \rightarrow \mathrm{Gr}_n(\mathbb{C}^{m+1}), \quad V \mapsto V \oplus \langle e_{m+1} \rangle$$

from the direct system from Corollary 3.2.38. If we denote the inclusion by  $\varphi_m^{n+1} : \mathrm{Gr}_{n+1}(\mathbb{C}^m) \rightarrow \mathrm{colim}_m(\mathrm{Gr}_{n+1}(\mathbb{C}^m))$  we get the following commuting diagram, for all  $n \leq m$ :

$$\begin{array}{ccccc}
\mathrm{Gr}_n(\mathbb{C}^m) & \xrightarrow{f_n^m} & \mathrm{Gr}_{n+1}(\mathbb{C}^{m+1}) & \xrightarrow{\varphi_{m+1}^{n+1}} & \mathrm{colim}_m(\mathrm{Gr}_{n+1}(\mathbb{C}^m)) = BU(n+1) \\
\downarrow i_m^n & & \downarrow i_m^n & \nearrow \varphi_{m+2}^{n+1} & \\
\mathrm{Gr}_n(\mathbb{C}^{m+1}) & \xrightarrow{f_n^{m+1}} & \mathrm{Gr}_{n+1}(\mathbb{C}^{m+2}) & & 
\end{array}$$

Now for all  $n$  we use the universal property of the direct limit  $\operatorname{colim}_m(\operatorname{Gr}_n(\mathbb{C}^m))$  on the family

$$\left( \varphi_{m+1}^{n+1} \circ f_n^m : \operatorname{Gr}_n(\mathbb{C}^m) \rightarrow BU(n+1) \right)_m$$

to induce maps  $f_n : BU(n) \rightarrow BU(n+1)$ , which in turn give us a directed system and let us define:

$$BU := \operatorname{colim}_n BU(n) = \operatorname{colim}_n \left( \operatorname{colim}_m \operatorname{Gr}_n(\mathbb{C}^m) \right).$$

**Lemma 3.4.11.** *The maps  $f_n : BU(n) \rightarrow BU(n+1)$  are closed embeddings.*

*Proof.* Since  $\operatorname{Gr}_n(\mathbb{C}^m)$  is compact and Hausdorff for any  $n \leq m$  the maps  $f_n^m$  are closed embeddings. Furthermore, the natural maps  $\varphi_m^n : \operatorname{Gr}_n(\mathbb{C}^m) \rightarrow BU(n)$  are also closed embeddings. Now assume that  $f_n(x) = f_n(x')$ . Then for  $r$  big enough (omitting the inclusions)  $f_n^r(x) = f_n^r(x')$  which contradicts the injectivity of the  $f_n^r$ . Now we want to show that  $f_n$  has a closed image. For this we let  $C := \operatorname{im} f_n \subset BU(n+1)$  and call the natural inclusions  $\varphi_m^{n+1} : \operatorname{Gr}_{n+1}(\mathbb{C}^m) \rightarrow BU(n+1)$ . Then  $C$  is closed, if and only if  $C \cap \operatorname{im} \varphi_m^{n+1}$  is closed for all  $m > n$ . To show this we first show that  $C \cap \operatorname{im} \varphi_m^{n+1} = \operatorname{im} \varphi_m^{n+1} \circ f_n^{m-1}$ . The first inclusion  $\subseteq$  is clear. The other inclusion follows directly from the construction of  $f_n$  that satisfies  $f_n \circ \varphi_{m-1}^n = \varphi_m^{n+1} \circ f_n^{m-1}$ . By Remark 3.2.37 we know that  $\varphi_m^{n+1} \circ f_n^{m-1}$  is a closed embedding hence  $C \cap \operatorname{im} \varphi_m^{n+1}$  is closed for all  $m > n$  and hence  $C$  is closed. Now finally we need to show that  $f_n$  are homeomorphisms onto their image. For this we show that they are closed maps, meaning they map closed sets to closed sets. This property together with injectivity implies the closed embedding property. Let  $U \subset BU(n)$  be closed. By construction of the direct limit we can write  $U = \bigcup_m (\operatorname{im}(\varphi_m^n) \cap U) = \bigcup_m (\varphi_m^n(U_m))$ , where  $U_m := (\varphi_m^n)^{-1}(U)$ . Now by the same argument as before we have the equality  $f_n(U) \cap \operatorname{im} \varphi_{m+1}^{n+1} = \varphi_{m+1}^{n+1} \circ f_n^m(U_m)$ . This is closed for all  $m$  because the maps  $\varphi_{m+1}^{n+1}$  and  $f_n^m$  are closed embeddings and hence closed. But then  $f_n(U)$  is closed in the final topology and hence  $f_n$  is a closed embedding.  $\square$

thm:kTheo-red  
ucedKgroupAsH  
omotopy

**Theorem 3.4.12.** *For a connected pointed space  $X$  we have a natural homomorphism*

$$\tilde{K}(X) \cong [X, BU]$$

*Proof.* We can now interchange the homotopy classes of maps and the limit by Lemma 3.2.40:

$$\begin{aligned} [X, BU] &= [X, \operatorname{colim}_n (BU(n))] \\ &\cong \operatorname{colim}_n \operatorname{Vect}_n(X) && \text{by Corollary 3.2.42,} \\ &\cong \tilde{K}(X) && \text{by Lemma 3.4.7.} \end{aligned}$$

$\square$

lemma:kTheo-  
generalGroupCo  
ntractsToUnit  
ary

**Lemma 3.4.13.** *The general linear group  $\mathrm{GL}_n(\mathbb{C})$  retracts onto the unitary group  $U(n)$  and the retraction commutes with the inclusions  $A \mapsto \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix}$ .*

*Proof.* Let  $A \in \mathrm{GL}_n(\mathbb{C})$ . Then  $A^*A$  is a positive-definite Hermitian matrix, since for  $v \in \mathbb{C}^n$ :

$$v^* A^* A v = (Av)^* (Av) = \|Av\|^2 \geq 0.$$

Hence, it is diagonalisable with real non-negative eigenvalues  $\lambda_1, \dots, \lambda_n > 0$ . We define  $P_A := \sqrt{A^*A}$  to be the matrix, that scales all eigenspaces by  $\sqrt{\lambda_i}$ . Then  $P_A^2 = A^*A$ , and  $\sqrt{A^*A}$  is also positive definite Hermitian, as all eigenvalues are positive and hence, it is invertible. Notice, that

$$\sqrt{\begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix}^* \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix}} = \begin{pmatrix} \sqrt{A^*A} & 0 \\ 0 & 1 \end{pmatrix}.$$

We then define  $U_A := A(P_A)^{-1}$ . By definition  $A = U_A P_A$ , and since  $P_A$  is Hermitian,  $P_A^{-1}$  is also Hermitian letting us conclude that  $U_A$  is unitary.

$$U_A^* U_A = ((P_A)^{-1})^* \underbrace{A^* A}_{P_A^2} (P_A)^{-1} = \mathbb{1}.$$

Notice again, that for  $\tilde{A} = A \oplus \mathbb{1}_1 \in \mathrm{GL}_{n+1}(\mathbb{C})$ , we have  $U_{\tilde{A}} = U_A \oplus \mathbb{1}_1 \in U(n+1)$ . Now the space of positive-definite Hermitian matrices is convex. This is just a calculation of all the axioms for the matrices on the connecting line of two positive-definite Hermitian matrices. But as a convex space, it is contractible.  $\square$

lem:kTheo-red  
ucedSuspended  
KGroupHomotop  
ically

**Lemma 3.4.14.** *For a connected pointed space  $X$  with well sitting basepoint, we have a natural homomorphism*

$$\tilde{K}(\Sigma X) \cong [X, U]$$

*The group structure in  $[X, U]$  comes from viewing  $[X, U]$  as the direct limit of abelian groups  $\mathrm{colim}[X, \mathrm{GL}_n(\mathbb{C})]$ , where the group structure is derived from the one in  $\mathrm{GL}_n(\mathbb{C})$ .*

*Proof.* If the base point is well sitting, we can apply Corollary 3.3.31, which gives us the homotopy equivalence

$$\Sigma X \simeq S(X).$$

Now  $S(X)$  itself can be cut into two contractible cones

$$\begin{aligned} C_+(X) &:= \{(x, t) \in S(X) \mid t \geq \tfrac{1}{2}\} \\ C_-(X) &:= \{(x, t) \in S(X) \mid t \leq \tfrac{1}{2}\}. \end{aligned}$$

Those cones intersect in  $C_+ \cap C_- = X \times \{\tfrac{1}{2}\} \cong X$ . Now denote the trivial  $n$  bundle over  $C_+(X)$  by  $\underline{n}_+$  and over  $C_-(X)$  by  $\underline{n}_-$ . With this notation we know by the proof of Lemma 3.2.14 that the map

$$\begin{aligned} \varphi_n : [X, \mathrm{GL}_n(\mathbb{C})] &\rightarrow \mathrm{Vect}_n(\Sigma X) \\ f &\mapsto \underline{n}_+ \bigcup_f \underline{n}_- \end{aligned} \tag{12}$$

$\{\text{eq:kTheo-SXt}$   
 $\text{oHomotoy}\}$

is a bijection for all  $n$ . Furthermore, the inclusion of  $\mathrm{GL}_n(\mathbb{C})$  into  $\mathrm{GL}_{n+1}(\mathbb{C})$  gives an inclusion  $[X, \mathrm{GL}_n(\mathbb{C})]$  into  $[X, \mathrm{GL}_{n+1}(\mathbb{C})]$  which commutes with the inclusion  $\mathrm{Vect}_n(\Sigma X)$  into  $\mathrm{Vect}_{n+1}(\Sigma X)$  under the bijection (12) meaning that for any  $n \leq m$  the diagram

$$\begin{array}{ccc} [X, \mathrm{GL}_n(\mathbb{C})] & \xrightarrow{\varphi_n} & \mathrm{Vect}_n(\Sigma X) \\ \downarrow i_{nm} & & \downarrow i_{nm} \\ [X, \mathrm{GL}_m(\mathbb{C})] & \xrightarrow{\varphi_m} & \mathrm{Vect}_m(\Sigma X) \end{array}$$

commutes. Lemma 3.4.7 gives us

$$\tilde{K}(X) \cong \mathrm{colim}_n [X, \mathrm{GL}_n(\mathbb{C})] \cong [X, \mathrm{colim}_n \mathrm{GL}_n(\mathbb{C})].$$

The latter bijection follows from Lemma 3.2.40. Finally, Lemma 3.4.13 gives us a retraction  $\mathrm{GL}_n(\mathbb{C}) \rightarrow U(n)$  that commutes with the inclusions and hence we get the same direct limit.

$$[X, \mathrm{colim}_n \mathrm{GL}_n(\mathbb{C})] \cong [X, \mathrm{colim}_n U(n)] = [X, U]$$

concluding that we have a natural bijection. To get the group structure we need to look into said mapping. For this we inspect the mapping (12) closer. Let

$$\Phi_n : [X, \mathrm{GL}_n(\mathbb{C})] \rightarrow \mathrm{colim} \mathrm{Vect}_n(\Sigma X)$$

be the composition of  $\varphi_n$  with the inclusion  $\mathrm{Vect}_n(\Sigma X) \rightarrow \mathrm{colim} \mathrm{Vect}_n(\Sigma X)$ . By the commutative diagram above, the maps  $\varphi_n$  induce a map from  $\mathrm{colim}[X, \mathrm{GL}_n(\mathbb{C})]$  which is a group homomorphism, if all  $\Phi_n$  are one. Hence take two functions  $f, g : X \rightarrow \mathrm{GL}_n(\mathbb{C})$ . Then by definition

$$f \oplus g : X \rightarrow \mathrm{GL}_n(\mathbb{C}), \quad x \mapsto f(x) \times g(x)$$

Now  $f \oplus g$  can be included into  $[X, \mathrm{GL}_{2n}(\mathbb{C})]$ :

$$f \oplus g : X \rightarrow \mathrm{GL}_{2n}(\mathbb{C}), \quad x \mapsto \begin{pmatrix} f(x) \times g(x) & 0 \\ 0 & \mathbb{1}_n \end{pmatrix}.$$

Now we have that  $f \oplus g$  is homotopic to

$$\widetilde{f \oplus g} : X \rightarrow GL_{2n}(\mathbb{C}), \quad x \mapsto \begin{pmatrix} f(x) & 0 \\ 0 & g(x) \end{pmatrix}$$

via the homotopy  $\rho_t(x)$  for  $t \in (0, \frac{\pi}{2})$ :

$$\rho_t(x) = \begin{pmatrix} f(x) & 0 \\ 0 & \mathbb{1}_n \end{pmatrix} \begin{pmatrix} \cos(t) & \sin(t) \\ \sin(t) & \cos(t) \end{pmatrix} \begin{pmatrix} \mathbb{1}_n & 0 \\ 0 & g(x) \end{pmatrix} \begin{pmatrix} \cos(t) & \sin(t) \\ \sin(t) & \cos(t) \end{pmatrix} \quad (13)$$

{eq:kTheo-hom  
otopyBetween  
irectSumAndPr  
oduct}

Finally we can look at the image of  $[f \oplus g] = [\widetilde{f \oplus g}]$  under  $\Phi_{2n}$  which is given by inclusion of

$$\varphi_n([\widetilde{f \oplus g}]) = \underline{n}_+ \oplus \underline{n}_+ \bigcup_{\widetilde{f \oplus g}} \underline{n}_- \oplus \underline{n}_- = [\varphi_n(f) \oplus \varphi_n(g)]$$

into the colimit, where the equation

$$[\varphi_n(f) \oplus \varphi_n(g)] = [\varphi_n(f)] \oplus [\varphi_n(g)]$$

holds. The second last equality follows from the properties of the gluing construction given in Corollary 3.2.11.  $\square$

**Remark 3.4.15.** The proof of Lemma 3.4.14 gives a slightly more general statement: For a connected well pointed topological space  $(X, x_0)$  with a splitting into two contractible parts  $X^+$  and  $X^-$  with  $x_0 \in A := X^+ \cap X^-$  and  $X^+ \cup X^- = X$  and the inclusion of  $A$  into the two contractible parts are cofibrations, there is a natural group isomorphism

$$[A, U] \cong \tilde{K}(X)$$

**Corollary 3.4.16.** Let  $Y$  be a pointed space. Define

$$T : S^1 \rightarrow S^1 \\ t \mapsto 1 - t$$

and let  $T \wedge 1 : SY \rightarrow SY$  be the map induced by  $T$  on  $S^1$  and the identity on  $Y$ . Then:

$$(T \wedge 1)^* : \tilde{K}(SY) \rightarrow \tilde{K}(SY) \\ y \mapsto -y$$

*Proof.* We have the following commutative diagram:

$$\begin{array}{ccc} \tilde{K}(SX) & \longleftarrow & \text{colim}[Y, GL(n, \mathbb{C})] \\ (\Sigma T)^* \downarrow & & \downarrow f \mapsto f^{-1}(\text{pointwise}) \\ \tilde{K}(SX) & \longleftarrow & \text{colim}[Y, GL(n, \mathbb{C})] \end{array}$$

lem:kTheo-inv  
ersedInKGiven  
ByHomotopical  
lyMethods

To see why this commutes, notice how we defined the map by gluing via  $f$ , where  $\Sigma T$  changes the place of the upper and lower vector bundle. Hence the gluing function needed is now the inverse. The horizontal maps are natural group homomorphisms and on the right side the inverse is taken. Thereby the commutativity tells us that the left side is also the same as taking the inverse.  $\square$

ex:kTheo-Line  
BundlesSatlqu  
adequalslPlus  
1

**Example 3.4.17.** Let  $l$  and  $l'$  be line bundles over  $S^2$ . Then the clutching function for the vector bundle  $l \otimes l'$  is given by the tensor product of clutching functions. So let  $f$  and  $f'$  be two clutching functions

$$f, f' : S^1 \rightarrow \mathrm{GL}_1(\mathbb{C})$$

Then the product bundle  $l \otimes l'$  is given by the clutching function

$$f \otimes f' : S^1 \rightarrow \mathrm{GL}(\mathbb{C} \otimes \mathbb{C}) \cong \mathrm{GL}_1(\mathbb{C}), \quad x \mapsto f(x)f'(x)$$

Hence, the product in  $\tilde{K}(S^2)$  is given by  $f \otimes f' = f \cdot f'$ . Now, the homotopy (13) of the proof above shows that  $f \cdot f' \oplus \mathrm{id}_1 \simeq f \oplus f'$  and hence  $l \otimes l' \oplus \underline{1} \cong l \oplus l'$ .

**Proposition 3.4.18.** For compactly generated pointed Hausdorff spaces  $(X, x_0), (Y, y_0)$  there is a natural bijection

$$[SX, Y] \rightarrow [X, \Omega Y],$$

where  $\Omega$  denotes the loop space

$$\Omega Y := \{f : S^1 \rightarrow Y \mid f \text{ continuous and } f(1) = y_0\}.$$

together with the compact open topology.

*Proof.* For any three spaces  $(X, x_0), (Y, y_0), (Z, z_0)$  we have that a basepoint preserving map  $f : X \wedge Y \rightarrow Z$  can be lifted to a map  $\tilde{f} : X \times Y \rightarrow Z; \tilde{f}(z) = \tilde{f}(\bar{z})$  that maps the whole  $X \vee Y$  to  $z_0$ . This map is continuous by the universal property of the quotient topology. Hence we get a bijection

$$C(X \wedge Y, Z) \rightarrow \{f \in C(X \times Y, Z) \mid f|_{X \vee Y} \equiv z_0\}$$

its inverse is given by the quotient induced maps. We can compose this map with the natural currying map  $C(X \times Y, Z) \rightarrow C(X, C(Y, Z))$ . For  $Y = S^1$  we gain a map

$$C(SX, Z) \rightarrow C(X, \Omega Z)$$

that is again a bijection, since any  $f : X \rightarrow \Omega Z$  gives a map  $\tilde{f} : X \times S \rightarrow Z; \tilde{f}(x, s) \mapsto f(x)(s)$ . If  $(x, s) \in X \vee S \subseteq X \times S$ , then either  $x = x_0$  or  $s = 1$ . In the first case,  $f(x_0)$  is the constant loop and hence  $f(x_0)(s) = z_0$  and in the second case we evaluate a based loop at its base and hence  $f(x)(1) = z_0$ . Hence, the  $\tilde{f}$  factors through  $X \vee S$  giving a well defined element in  $C(SX, Z)$ . This is its inverse. The continuity part of the argument follows from Proposition 1.3.3.  $\square$

**Remark 3.4.19.** By Theorem 3.4.12, and the Proposition above we know that for a connected pointed compact space we have

$$\tilde{K}(SX) \cong [SX, BU] \cong [X, \Omega BU]$$

On the other hand we have Lemma 3.4.14 which tells us that

$$\tilde{K}(SX) \cong [X, U].$$

This hints at a connection that states that  $\Omega BU$  is homotopically equivalent to  $U$ :

### 3.5 K-Theory as a Cohomology Theory

This chapter aims to give the definitions and theorems that make K theory an axiomatic cohomology theory as defined by Eilenberg and Steenrod. For this we start trying to find a long exact sequence and on the way we will prove the additivity:

lem:kTheo-exa  
ctShortPartFo  
rThePair

**Lemma 3.5.1.** *Let  $(X, Y) \in \text{ob}(\mathbf{CWH}_{\text{cof}}^2)$  and define  $i : (Y, \emptyset) \rightarrow (X, \emptyset)$  and  $j : (X, \emptyset) \rightarrow (X, Y)$  to be the inclusions. Then we have an exact sequence:*

$$K(X, Y) \xrightarrow{j^*} K(X) \xrightarrow{i^*} K(Y)$$

*Here, we identify  $K(X) = K(X, \emptyset)$  and the same with  $K(Y) = K(Y, \emptyset)$  such that we have well defined induced functions from the same functor in the sequence. This is just a formal note, as  $K(X, \emptyset) \cong K(X)$  by Corollary 3.3.33.*

*Proof.* First, we show that  $\text{im}(i^*) \subseteq \ker(j^*)$ . To see this, notice that  $i^* \circ j^* = (j \circ i)^*$ , and  $j \circ i$  factors through  $(Y, Y)$ :

$$\begin{array}{ccc} (Y, \emptyset) & \xrightarrow{i} & (X, \emptyset) \\ \downarrow & & \downarrow j \\ (Y, Y) & \longrightarrow & (X, Y) \end{array}$$

Hence,  $j^* \circ i^*$  factors through the trivial ring  $K(Y, Y)$  and thereby is trivial. Now suppose  $k \in \ker(i^*)$ . The intuition now is that if we have a vector bundle that gets mapped to 0 this means that it is trivial over  $Y$  which lets us define a vector bundle over the quotient  $X/Y$  by a trivialisation: We write  $k = [\xi] - [\underline{n}]$  for  $\xi$  being a vector bundle over  $X$ . (Compare Remark 3.4.8) Since  $i^* \xi = 0$  it follows that  $i^*([\xi] - [\underline{n}]) = [\xi|_Y] - [\underline{n}|_Y] = 0$  and hence  $\xi$  and  $\underline{n}$  are stably equivalent over  $Y$  giving us an  $m$  such that

$$(\xi \oplus \underline{m})|_Y \cong \xi|_Y \oplus \underline{m} \cong \underline{n} \oplus \underline{m}$$

Hence, there is a trivialisation  $\alpha : (\xi \oplus \underline{m})|_Y \rightarrow Y \times \mathbb{C}^{\text{rank}(\xi)+m}$  that defines a bundle  $(\xi \oplus \underline{m})|_Y / \alpha$  as shown in the proof of Lemma 3.2.7 over  $X/Y$ . Now we define

$$\eta = [(\xi \oplus \underline{m})|_Y / \alpha] - [\underline{n} + \underline{m}] \in \tilde{K}(X/Y) = K(X, Y)$$

and since the trivialisation acts as a right inverse of the pullback of the restriction (compare Corollary 3.2.8)

$$j^*(\eta) = [\xi \oplus \underline{m}] - [\underline{n} + \underline{m}] = k.$$

□

lem:kTheo-poi  
ntedUnionExac  
tSequence

**Lemma 3.5.2.** *Let  $X, Y$  be pointed topological spaces and define the maps*

*$i : X \rightarrow X \vee Y$  the inclusion, and  $j : X \vee Y \rightarrow Y$  the collapse of  $X$ .*

*Then the sequence*

$$0 \longrightarrow \tilde{K}(X) \xrightarrow{j^*} \tilde{K}(X \vee Y) \xrightarrow{i^*} \tilde{K}(Y) \longrightarrow 0$$

*is exact and naturally splits via the map  $q^*$ , that is a right inverse of  $j^*$ , where  $q : X \vee Y \rightarrow Y$  is the collapse.*

*Proof.* Notice how there is a natural homeomorphism  $X \cong (X \vee Y)/Y$ . Hence, Lemma 3.5.1 provides the exactness at  $\tilde{K}(X \vee Y)$ . Now the map  $q$  is a left inverse of  $j$  as continuous maps, and hence  $q^*$  is a right inverse of  $j^*$  in the K group, proving the injectivity of  $j^*$ . Now only the surjectivity of  $i^*$  is missing. However the inclusion  $l : Y \rightarrow X \vee Y$  is a topological right inverse of  $i$  and thereby  $l^*$  is a left inverse of  $i^*$  giving us the injectivity and hence proving the exactness and splitting of the short exact sequence. □

thm:kTheo-unp  
ointedAdditiv  
ity

**Theorem 3.5.3** (Pointed Additivity). *Let  $X, Y$  be pointed topological spaces and denote the inclusions of  $X$  (resp.  $Y$ ) into  $X \vee Y$  by  $i_X$  (resp.  $i_Y$ ). Then the map*

$$i_X^* \oplus i_Y^* : \tilde{K}(X \vee Y) \rightarrow \tilde{K}(X) \oplus \tilde{K}(Y)$$

*is a natural isomorphism with inverse*

$$\psi : \tilde{K}(X) \oplus \tilde{K}(Y) \rightarrow \tilde{K}(X \vee Y), \quad (x, y) \mapsto c_X^*(x) + c_Y^*(y),$$

*where  $c_X$  (resp.  $c_Y$ ) denotes the collapse of  $X$  (hence onto  $Y$ ) respectively the collapse of  $Y$  (hence onto  $X$ ).*

*Proof.* This is a direct consequence of Lemma 3.5.2 and the algebraic machinery of splitting short exact sequences. □

thm:kTheo-unp  
ointedAdditiv  
ity

**Corollary 3.5.4** (Unpointed Additivity). *Let  $X, Y$  be topological spaces, then there is a natural isomorphism*

$$K(X \sqcup Y) \rightarrow K(X) \oplus K(Y)$$

*Proof.* This follows from  $K(A) \cong \tilde{K}(A, [\emptyset])$  for all topological spaces. Furthermore,  $(X \sqcup Y, [\emptyset]) \cong ((X, [\emptyset]) \vee (Y, [\emptyset]))$ . Hence, using those isomorphisms, the statement is a direct consequence of the pointed additivity.  $\square$

**Corollary 3.5.5.** If  $(X, Y) \in ob(\mathbf{CWH}_{\text{cof}}^2)$  and  $Y \in ob(\mathbf{CWH}_{\text{cof}}^*)$  such that  $X$  is pointed by using the base point  $y_0$  from  $Y$ . Then the sequence

$$K(X, Y) \xrightarrow{j^*} \tilde{K}(X) \xrightarrow{i^*} \tilde{K}(Y)$$

is exact.

*Proof.* We have the commutative diagram with exact top row:

$$\begin{array}{ccccc} K(X, Y) & \longrightarrow & K(X) & \longrightarrow & K(Y) \\ & \searrow & \downarrow \cong & & \downarrow \cong \\ & & \tilde{K}(X) \oplus K(y_0) & \longrightarrow & \tilde{K}(Y) \oplus K(y_0) \\ & & \downarrow \Downarrow & & \downarrow \\ & & \tilde{K}(X) & \longrightarrow & \tilde{K}(Y) \end{array}$$

The middle horizontal map is a map that respects the grading, since we have the same base point. With this we can do a bit of diagram yoga concluding the statement.  $\square$

**Definition 3.5.6.** Let  $X$  be a topological space and  $I = [0, 1]$  be the unit interval. We define the cone over  $X$  to be the space

$$CX := (X \times I) / X \times \{0\}.$$

Let  $(X, Y)$  be a pair of compact topological spaces. We define the cone over the subspace  $Y$  to be

$$(X \sqcup CY) / Y \sim Y \times \{1\}$$

and denote that with  $(X \cup CY)$ .

**Proposition 3.5.7.** If we define for  $f : X \rightarrow Y$  the induced maps on the cones  $Cf$  by

$$Cf : CX \rightarrow CY \quad \overline{(x, t)} \mapsto \overline{(f(x), t)},$$

the cone defines a functor  $\mathbf{CGWH} \rightarrow \mathbf{CGWH}^*$ . The cone over  $Y \subseteq X$  defines in similar fashion a functor from  $\mathbf{CGWH}_{\text{cof}}^2$  to  $\mathbf{CGWH}^*$ . There is a natural isomorphism:

$$T : (X \cup CY) / X \rightarrow CY / Y$$

Here, the left and right hand side both define functors from  $\mathbf{CGWH}_{\text{cof}}^2$  to  $\mathbf{CGWH}^*$  with the obvious morphisms. Furthermore, we have that  $CY/Y$  is homotopically equivalent to  $\Sigma Y$  (via the map induced from  $\overline{(x, t)} \mapsto \overline{(t, x)}$ ) which is homotopically equivalent to  $SY$  as long as  $(Y, y_0)$  is well pointed. All the equivalences are natural up to homotopy.

cor:kTheo-exa-  
ctnesOfTheRed-  
ucedInclusion  
Sequence

def:kTheo-con-  
eOverSubspace

prop:kTheo-co-  
neIsFunctorAn-  
dCYoverYIsRed-  
Suspension

**Corollary 3.5.8.** From the considerations above we get a natural isomorphism

$$\theta : K(X \cup CY, Y) \rightarrow \tilde{K}(SY) = \tilde{K}^{-1}(Y).$$

def:kTheo-connectingHomomorphism

**Definition 3.5.9** (Connecting Homomorphism). Let  $(X, Y)$  be a topological pair of compact spaces and  $Y$  be pointed. Define the two maps

$$m : (X \cup CY, \emptyset) \rightarrow (X \cup CY, X) \quad (14)$$

$$\pi : (X \cup CY, \emptyset) \rightarrow X/Y \quad (15)$$

where  $m$  is the inclusion in the second space, and  $\pi$  is the collapse of  $CY$ . Since  $CY$  is contractible, the induced map  $\pi^* : \tilde{K}(X/Y) \rightarrow K(X \cup CY, \emptyset)$  is an isomorphism by Lemma 3.2.9. Together with the isomorphism

$$\theta : K(X \cup CY, X) \rightarrow \tilde{K}(SY) = \tilde{K}^{-1}(Y).$$

We define the connecting homomorphism

$$\delta := (\pi^*)^{-1} \circ m^* \circ \theta^{-1} : \tilde{K}^{-1}(Y) \rightarrow K(X, Y)$$

This map defines also the higher degrees connecting homomorphisms by replacing the spaces with their suspensions.

**Remark 3.5.10.** The map  $\delta : K^{-n}(A) \rightarrow K^{-n+1}(X, A)$  is a natural transformation between the functors

- $K^{-n} \circ R$  from  $\mathbf{CWH}_{\text{cof}}^2$  to  $\mathbf{Ab}$ , where  $R$  is an endofunctor in  $\mathbf{CWH}_{\text{cof}}^2$  sending  $(X, A) \mapsto (A, \emptyset)$ .
- $K^{-n+1}$  from  $\mathbf{CWH}_{\text{cof}}^2$  to  $\mathbf{Ab}$ .

*Proof.* Let  $f : (X, A) \rightarrow (Y, B)$  be a map of pairs. Then we need to check if the diagram

$$\begin{array}{ccccc} (X, A) & & K^{-n}(A, \emptyset) & \xrightarrow{\delta} & K^{-n+1}(X, A) \\ \downarrow f & & (f|_A)^* \uparrow & & f^* \uparrow \\ (Y, B) & & K^{-n}(B, \emptyset) & \xrightarrow{\delta} & K^{-n+1}(Y, B) \end{array}$$

commutes. This however is the case since  $\delta$  is defined via the pullback of the inclusion  $(m^*)$  together with the natural isomorphisms  $\theta$  and  $\pi^*$ .  $\square$

thm:kTheo-longExactSequence

**Theorem 3.5.11** (Long Exact Sequence). *For  $(X, Y) \in \text{ob}(\mathbf{CWH}_{\text{cof}}^2)$  we have a natural exact sequence that is infinite to the left:*

$$\begin{array}{ccccccc} \cdots & \longrightarrow & K^{-2}(Y) & \xrightarrow{\delta} & K^{-1}(X, Y) & \xrightarrow{j^*} & K^{-1}(X) \xrightarrow{i^*} K^{-1}(Y) \\ & & & & & & \\ & & & & \xrightarrow{\delta} & K^0(X, Y) & \xrightarrow{j^*} K^0(X) \xrightarrow{i^*} K^0(Y) \end{array}$$

The maps  $i^*$  are induced by the inclusion  $i : Y \rightarrow X$ ,  $j^*$  are induced by the inclusion  $j : (X, \emptyset) \rightarrow (X, Y)$  and  $\delta$  is the connecting homomorphism. For the lower degrees,  $\delta$  is the connecting homomorphism where  $X$  and  $Y$  are replaced by the reduced suspensions of  $X$  and  $Y$ .

*Proof.* We will prove the pointed version of the theorem, that states that for a compact pair  $(X, Y)$  such that  $Y$  is itself compact pointed (and  $X$  is pointed with the same basepoint) we have the exact sequence (infinite to the left):

$$\begin{aligned} \dots \longrightarrow \tilde{K}^{-2}(Y) &\xrightarrow{\delta} K^{-1}(X, Y) \xrightarrow{j^*} \tilde{K}^{-1}(X) \xrightarrow{i^*} \tilde{K}^{-1}(Y) \\ &\xrightarrow{\delta} K^0(X, Y) \xrightarrow{j^*} \tilde{K}^0(X) \xrightarrow{i^*} \tilde{K}^0(Y) \end{aligned}$$

The pointed sequence induced the statement for unpointed spaces because for any topological space  $A$  we have a natural isomorphism  $\tilde{K}(A, [\emptyset]) \cong K(A)$ .

We will furthermore just prove the exactness of the segment:

$$\tilde{K}^{-1}(X) \xrightarrow{i^*} \tilde{K}^{-1}(Y) \xrightarrow{\delta} K^0(X, Y) \xrightarrow{j^*} \tilde{K}^0(X) \xrightarrow{i^*} \tilde{K}^0(Y) \quad (16)$$

{eq:kTheo-pro  
ofSesShortAnd  
reducedSequen  
ce}

From this we regain the long sequence by replacing  $(X)$  and  $Y$  with  $S^n(X)$  and  $S^n Y$ .

In  $\tilde{K}^0(X)$ , the exactness was shown in Corollary 3.5.5. To get the exactness at the other spots we apply the same corollary to the pairs:

- $(X \cup CY, X)$
- $((X \cup CY) \cup CX, X \cup CY)$

For the middle part we have the commutative diagram where the naming of the maps is the same as in the definition of the connecting homomorphisms (Definition 3.5.9).

$$\begin{array}{ccccc} K(X \cup CY, X) & \xrightarrow{m^*} & \tilde{K}(X \cup CY) & \xrightarrow{k^*} & \tilde{K}(X) \\ \downarrow \theta & & \uparrow \pi^* & & \downarrow \text{id} \\ K^{-1}(Y) & \xrightarrow{\delta} & K(X, Y) & \xrightarrow{j^*} & \tilde{K}(X) \end{array}$$

The commutativity in the left square is the definition of  $\delta$  and in the right square follows from  $\pi \circ k = j : X \rightarrow X/Y$ . Since all vertical maps are isomorphisms, Corollary 3.5.5 implies exactness in the bottom row  $K(X, Y)$ . Now only the exactness in the first three terms is missing. For this we look at the pair  $((X \cup C_1 Y) \cup C_2 X, X \cup C_1 Y)$ , where the  $C_i$  are cones and denoted with the index to distinguish the cones. First we get again an exact sequence:

$$K((X \cup C_1 Y) \cup C_2 X, X \cup C_1 Y) \xrightarrow{l^*} \tilde{K}((X \cup C_1 Y) \cup C_2 X) \xrightarrow{s^*} \tilde{K}(X \cup C_1 Y)$$

Our goal is now to show that we commute with the start of the exact sequence (16), hence to find vertical maps between the top rows (denoted with  $\psi$  in the diagram below) such that the diagram commutes:

diag:kTheo-pr  
oofConnecting  
Homo

$$\begin{array}{ccccc}
 K((X \cup C_1 Y) \cup C_2 X, X \cup C_1 Y) & \xrightarrow{l^*} & \tilde{K}((X \cup C_1 Y) \cup C_2 X) & \xrightarrow{s^*} & \tilde{K}(X \cup C_1 Y) \\
 \downarrow \psi_1 & & \downarrow \psi_2 & & \downarrow \psi_3 \\
 \tilde{K}^{-1}(X) & \xrightarrow{i^*} & \tilde{K}^{-1}(Y) & \xrightarrow{\delta} & K(X, Y)
 \end{array}$$

Diagram 17

We start with the right square and notice that by definition we get a commutative diagram:

$$\begin{array}{ccc}
 K(X \cup C_1 Y \cup C_2 X) & \xrightarrow{s^*} & \tilde{K}(X \cup C_1 Y) \\
 \downarrow (\tilde{\pi}^*)^{-1} & & \downarrow \text{id} \\
 K(X \cup C_1 Y, X) & \xrightarrow{m^*} & \tilde{K}(X \cup C_1 Y) \\
 \downarrow \theta & & \downarrow (\pi^*)^{-1} \\
 K^{-1}(Y) & \xrightarrow{\delta} & K(X, Y)
 \end{array}$$

The bottom row is added for clarification on what inclusion induces  $i^*$ .. besser erklären

The bottom square commutes by definition. In the top square, the map  $\tilde{\pi}$  denotes the collapsing map  $X \cup C_1 Y \cup C_2 X \rightarrow (X \cup C_1 Y)/X$  which is invertible, as the quotiented out space  $C_2 X$  is contractible. To see the commutativity in the top square notice how  $\tilde{\pi}^* \circ s^* = (s \circ \tilde{\pi})^*$  and the maps  $m$  and  $s \circ \tilde{\pi}$  coincide and hence the top square commutes. With  $\psi_2 := \theta \circ (\tilde{\pi}^*)^{-1}$  and  $\psi_3 := (\pi^*)^{-1} \circ \text{id}$ . For the left square, a natural choice is the homotopical equivalence given by quotienting the second space of the pair  $((X \cup C_1 Y) \cup C_2 X, X \cup C_1 Y)$  to get:

$$((X \cup C_1 Y) \cup C_2 X / X \cup C_1 Y) \cong (C_2 X / X) \simeq S(X).$$

We now have all the unknown maps, but we run into a problem, since the map  $l^*$  induces a map  $\lambda$  by the “wrong” inclusions (compared to  $i^*$  which is induced by the inclusion  $C_2 Y \rightarrow C_2 X$ , whereas we have  $C_1 Y$ ). This is depicted in the non-commutative diagram:

$$\begin{array}{ccc}
 K((X \cup C_1 Y) \cup C_2 X, X \cup C_1 Y) & \xrightarrow{l^*} & \tilde{K}((X \cup C_1 Y) \cup C_2 X) \\
 \downarrow \psi_1 & & \downarrow \psi_2 \\
 \tilde{K}(C_2 X / X) \cong \tilde{K}^{-1}(X) & \xrightarrow{\lambda} & \tilde{K}(C_1 Y / Y) \cong \tilde{K}^{-1}(Y) \\
 \uparrow & & \uparrow \\
 K(C_2 X, X) & \xrightarrow{i^*} & K(C_2 Y, Y)
 \end{array}$$

To resolve that problem we introduce the double cone on  $Y$  namely  $C_1Y \cup C_2Y$ , that fits into the commutative diagram where all unnamed maps are the natural ones:

$$\begin{array}{ccccc}
 X \cup C_1Y \cup C_2X & \xrightarrow{\quad\quad\quad} & C_1Y/Y & \xRightarrow{\quad} & \Sigma Y \\
 \downarrow & \swarrow & \uparrow & \searrow & \vdots \Sigma T \\
 & C_1Y \cup C_2Y & & & \\
 & \swarrow & \searrow & & \\
 C_2X/X & \xleftarrow{\quad\quad\quad} & C_2Y/Y & \xRightarrow{\quad} & \Sigma Y \\
 & \xleftarrow{i} & & & 
 \end{array}$$

The inclusion induced map  $i$  is the one, that induces the bottom left map in Diagram 17. The double arrows always induce isomorphisms in the  $K$ -rings. To see that all those maps induce isomorphism we note that we always, with one exception, collapse something contractible. The exception is the map  $C_1Y \cup C_2Y \rightarrow X \cup C_1Y \cup C_2X$ . Here, we get that the induced map is invertible, by noticing the commutative triangle on the top, where two maps induce isomorphisms and thereby also the third. The map  $\Sigma T$  is one we want to construct below. For said construction we formulate the following Lemma:

lem:kTheo-doubleConeFlip

**Lemma 3.5.12.** *Let  $Y$  be a well-pointed space and denote the points of  $CY$  and  $\Sigma Y$  by  $(x, t)$  such that  $CY$  has the apex  $[\{0\} \times Y]$  and base  $\{1\} \times Y \cong Y$ . Let*

$$D := C_1Y \cup C_2Y = (C_1Y \sqcup C_2Y) / ((x, 1)_1 \sim (x, 1)_2)$$

*denote the double cone glued along a common base. Now define the maps*

- $c_i : D \rightarrow C_iY/Y$  the collapse of the complementary cone  $C_jY$   $j \neq i$ .
- $\overline{\pi}_1 : C_iY/Y \rightarrow \Sigma Y$   $[(x, t)] \mapsto (x, t)$ .
- $p_i := \overline{\pi}_1 \circ c_i : D \rightarrow \Sigma Y$ .
- $R : D \rightarrow D$   $(x, t)_i \mapsto (x, t)_j$  where  $j \neq i$ , the cone swap.
- $\Sigma T : \Sigma Y \rightarrow \Sigma Y$ ,  $(x, t) \mapsto (x, 1 - t)$  the suspension flip.
- $\Phi : D \rightarrow \Sigma Y$ ,  $(x, t)_1 \mapsto (x, 1 - \frac{t}{2})$ ,  $(x, t)_2 \mapsto (x, \frac{t}{2})$ .

*Then we have that*

$$p_2 = p_1 \circ R \quad \text{and} \quad \Phi \circ R = \Sigma T \circ \Phi.$$

*Proof of Lemma 3.5.12.* The first statement  $p_2 = p_1 \circ R$  is clear, as:

$$\begin{aligned}
 p_2(x, t)_2 &= [(x, t)] = p_1(x, t)_1 = p_1 \circ R(x, t)_2, \quad \text{and} \\
 p_2(x, t)_1 &= [\{1\} \times Y] = p_1(x, t)_2 = p_1 \circ R(x, t)_1.
 \end{aligned}$$

Similarly, the second statement is also just a calculation:

$$\begin{aligned}\Sigma T \circ \Phi(x, t)_1 &= \Sigma T(x, 1 - \frac{t}{2}) = (x, \frac{t}{2}) = \Phi(x, t)_2 = \Phi \circ R(x, t)_1 \\ \Sigma T \circ \Phi(x, t)_2 &= \Sigma T(x, \frac{t}{2}) = (x, 1 - \frac{t}{2}) = \Phi(x, t)_1 = \Phi \circ R(x, t)_2.\end{aligned}$$

□

By the lemma above, we know that  $\Sigma T$  makes the diagram commute and by Lemma 3.4.16 we know that

$$(\Sigma T)^* : \tilde{K}(SY) \rightarrow \tilde{K}(SY)$$

is multiplication by minus one. Hence, the diagram where all maps are either topologically induced or topologically induced and inverted:

$$\begin{array}{ccccc}\tilde{K}(X \cup C_1 Y \cup C_2 X) & \xrightarrow{\psi_2} & \tilde{K}(C_1 Y / Y) & \longrightarrow & \tilde{K}^{-1}(Y) \\ & \searrow & & & \uparrow \\ & & \tilde{K}(C_1 Y \cup C_2 Y) & & (-1)^* \\ \uparrow \rho & & \searrow & & \parallel \\ \tilde{K}^{-1}(X) & \xrightarrow{i^*} & \tilde{K}(C_2 Y / Y) & \longrightarrow & \tilde{K}^{-1}(Y)\end{array}$$

is commutative. By construction (just composition of collapsing maps) we have that

$$l^* = \rho \circ \psi_1, \quad \text{and by the diagram} \quad \psi_2 \circ \rho = -i^*.$$

Which gives us in total:

$$i^* \circ \psi_1 = -\psi_2 \circ \rho \circ \psi_1 = -\psi_2 \circ l^*$$


and thereby the diagram commutes up to sign:

$$\begin{array}{ccccc}K((X \cup C_1 Y) \cup C_2 X, X \cup C_1 Y) & \xrightarrow{l^*} & \tilde{K}((X \cup C_1 Y) \cup C_2 X) & \xrightarrow{s^*} & \tilde{K}(X \cup C_1 Y) \\ \downarrow \psi_1 & & \downarrow \psi_2 & & \downarrow \psi_3 \\ \tilde{K}^{-1}(X) & \xrightarrow{i^*} & \tilde{K}^{-1}(Y) & \xrightarrow{\delta} & K(X, Y)\end{array}$$

The exactness in the top row now proves exactness in the bottom. □

ktheo:cor-tri  
pleSequence

**Corollary 3.5.13** (Triple Sequence). Let  $(X, Y, Z)$  be a topological triple, meaning that all inclusions are cofibrations. Then we can inspect the three long exact sequences from the pairs  $(X, Y)$ ,  $(Y, Z)$  and  $(X, Z)$  that are in the diagram below. The map  $\delta_{triple}$  is defined to be the composition  $\delta \circ j^*$ ,  $q^*$  is defined to be induced by the inclusion  $(X, Z) \rightarrow (X, Y)$  and  $s^*$  is also given by the inclusion  $(Y, Z) \rightarrow (X, Z)$ . Then, the whole diagram becomes commutative by the naturality of the maps appearing in the long exact sequences.



Furthermore, the yellow sequence is a chain complex which we need to verify at three locations:

1. At  $K^{-n-1}(X, Z)$  we can see that the composition  $s^* \circ q^*$  is the pullback of the inclusion  $(Y, Z) \rightarrow (X, Y)$  which factors through  $(Y, Y)$  and thereby the pullback factors through the trivial ring  $K^{-n-1}(Y, Y)$ .
2. At  $K^{-n-1}(Y, Z)$  we get that the composition is zero as it factors through two consecutive maps in the sequence of  $(X, Y)$ :  $\delta_{triple} \circ s^* = \delta \circ i^* \circ j^* = 0$ .
3. At  $K^{-n}(X, Y)$  we get  $q^* \circ \delta = 0$  as it factors through  $i^* \circ j^*$ .

Then, by the braid lemma (which is typical diagram yoga), we know that the orange sequence is exact. We call this sequence the **triple exact sequence** and it is natural with respect to maps of triples, meaning that for two such triples and a map of triples, we get two sequences and an induced map of sequences.

**Corollary 3.5.14.** If  $(X, Y)$  is a compact pair such that  $Y \subseteq X$  is a retract, i.e. the inclusion has a continuous up to homotopy left inverse. Then for all  $n \geq 0$ , the sequence

$$K^{-n}(X, Y) \xrightarrow{j^*} K^{-n}(X) \xrightarrow{i^*} K^{-n}(Y)$$

is a splitting short exact sequence giving us an isomorphism

$$K^{-n}(X) \cong K^{-n}(X, Y) \oplus K^{-n}(Y).$$

This splitting is naturally dependent on the retraction map.

**Corollary 3.5.15.** Let  $X, Y$  be objects in  $\mathbf{CWH}_{\text{cof}}^*$ . Then we have the projection maps

$$\pi_X : X \times Y \rightarrow X, \quad \pi_Y : X \times Y \rightarrow Y.$$

Those maps induce isomorphisms for all  $n \geq 0$ :

$$\tilde{K}^{-n}(X \times Y) \cong \tilde{K}^{-n}(X \wedge Y) \oplus \tilde{K}^{-n}(X) \oplus \tilde{K}^{-n}(Y)$$

cor:kTheo-ret  
ractionsInduc  
eSplittingSeq  
uences

referenz  
einfügen:  
Munkres,  
p. 148  
of Ele-  
ments of  
algebraic  
topology?

*Proof.*  $X$  is a retract of  $X \times Y$  as the projection yields a continuous left inverse of the natural inclusion, and  $Y$  is a retract of  $(X \times Y)/X$  (for the quotient to be natural, we include  $X$  at the base point of  $Y$ ). With this we apply corollary 3.5.14 twice:

$$\begin{aligned} \tilde{K}^{-n}(X \times Y) &\cong \tilde{K}^{-n}(X \times Y/X) \oplus \tilde{K}^{-n}(X) \\ &\cong \tilde{K}^{-n}(Y) \oplus \underbrace{\tilde{K}^{-n}((X \times Y/X)/Y)}_{\tilde{K}^{-n}(X \wedge Y)} \oplus \tilde{K}^{-n}(X) \end{aligned}$$

□

**Corollary 3.5.16.** Let's inspect the isomorphism induced by the splitting: Since we apply the splitting twice, we know that the final isomorphism

$$\Phi : \tilde{K}^{-n}(X \times Y) \rightarrow \underbrace{\tilde{K}^{-n}((X \times Y/X)/Y)}_{\tilde{K}^{-n}(X \wedge Y)} \oplus \tilde{K}^{-n}(X) \oplus \tilde{K}^{-n}(Y)$$

is of the form

$$\Phi = (g \circ l, \tilde{i}_Y^* \circ l, i_X^*),$$

where:

- $i_Y : Y \rightarrow X \times Y$
- $\tilde{i}_Y : Y \rightarrow (X \times Y/X)$ .
- $j : (X \times Y) \rightarrow (X \times Y/X)$ .
- $l : \tilde{K}^{-n}(X \times Y) \rightarrow \tilde{K}^{-n}(X \times Y/X)$  is the unique left inverse of  $j^*$  such that  $j^* \circ l = \text{id} - \pi_X^* \circ i_X^*$ .
- $g : \tilde{K}^{-n}(X \times Y/X) \rightarrow \tilde{K}^{-n}(X \times Y/X, Y)$  is a left inverse of the map  $K(X \times Y/X, Y) \rightarrow \tilde{K}^{-n}(X \times Y/X)$ .

We now want to show that  $\tilde{i}_Y^* \circ l$  is induced by the inclusion of  $Y$  into  $X \times Y$ .

We have that  $\tilde{i}_Y = j \circ i_Y$  and with this we can deduce:

$$\tilde{i}_Y^* \circ l = i_Y^* \circ j^* \circ l = i_Y^* (\text{id} - \pi_X^* \circ i_X^*) = i_Y^* - i_Y^* \circ \pi_X^* \circ i_X^* = i_Y^* - \underbrace{(\pi_X \circ i_Y)^*}_{\text{constant}} \circ i_X^* = i_Y^*$$

Hence, in the proof above the isomorphism has the form:

$$\begin{aligned} \Phi : \tilde{K}^{-n}(X \times Y) &\rightarrow \tilde{K}^{-n}(X \wedge Y) \oplus \tilde{K}^{-n}(Y) \oplus \tilde{K}^{-n}(X) \\ \Phi &= g \oplus i_Y^* \oplus i_X^* \end{aligned}$$

or said differently, the short exact sequence that splits looks like:

$$0 \longrightarrow \tilde{K}^{-n}(X \wedge Y) \xrightarrow{q^*} \tilde{K}^{-n}(X \times Y) \xrightarrow{(i_X)^* \oplus (i_Y)^*} \tilde{K}^{-n}(X) \oplus \tilde{K}^{-n}(Y) \longrightarrow 0$$

Before we start constructing a cohomology theory, we extend the K groups to be defined not only over the negatives, but over the whole  $\mathbb{Z}$  which is done via an extension. The extension relies on periodic behaviour. So-called Bott periodicity.

**Definition 3.5.17** (External Product). The **external product**  $\mu : K(X) \otimes K(Y) \rightarrow K(X \times Y)$  is defined via the two projections  $\pi_X : X \times Y \rightarrow X$  and  $\pi_Y : X \times Y \rightarrow Y$  by:

$$\begin{aligned} \mu : K(X) \otimes K(Y) &\rightarrow K(X \times Y) \\ (a \otimes b) &\mapsto (\pi_X)^*(a) \cdot (\pi_Y)^*(b), \end{aligned}$$

where  $\cdot$  denotes the multiplication in the K-group, which was induced by the tensor product. This  $\mu$  is a ring homomorphism, since:

$$\begin{aligned} \mu((a \otimes b)(c \otimes d)) &= \mu(ac \otimes bd) \\ &= \pi_X^*(ac) \pi_Y^*(bd) \\ &= \pi_X^*(a) \pi_X^*(c) \pi_Y^*(b) \pi_Y^*(d) \\ &= \pi_X^*(a) \pi_Y^*(b) \pi_X^*(c) \pi_Y^*(d) \\ &= \mu(a \otimes b) \mu(c \otimes d) \end{aligned}$$

prop:kTheo-re  
ducedExternal  
Product

**Proposition 3.5.18.** Let  $(X, x_0)$  and  $(Y, y_0)$  be well-pointed compact spaces and let  $q : X \times Y \rightarrow X \wedge Y$  denote the quotient map that appears in the exact sequence of Corollary 3.5.16. Then:

1. For  $a \in \tilde{K}(X)$  and  $b \in \tilde{K}(Y)$  we have  $\mu(a \otimes b) \in q^* \tilde{K}(X \wedge Y)$ . Since  $q^*$  is injective, this defines the **reduced external product**

$$\tilde{\mu} := (q^*)^{-1} \circ \mu|_{\tilde{K}(X) \otimes \tilde{K}(Y)} : \tilde{K}(X) \otimes \tilde{K}(Y) \rightarrow \tilde{K}(X \wedge Y).$$

2. With respect to the decompositions

$$\begin{aligned} K(X) \otimes K(Y) &\cong (\tilde{K}(X) \otimes \tilde{K}(Y)) \oplus \tilde{K}(X) \oplus \tilde{K}(Y) \oplus \mathbb{Z}, \\ K(X \times Y) &\cong q^* \tilde{K}(X \wedge Y) \oplus \pi_X^* \tilde{K}(X) \oplus \pi_Y^* \tilde{K}(Y) \oplus \mathbb{Z} \cdot 1_{X \times Y} \end{aligned}$$

the external product is block diagonal:

$$\mu = (q^* \circ \tilde{\mu}) \oplus \pi_X^*|_{\tilde{K}(X)} \oplus \pi_Y^*|_{\tilde{K}(Y)} \oplus \text{id}_{\mathbb{Z}}.$$

In particular  $\mu$  is an isomorphism if and only if  $\tilde{\mu}$  is.

*Proof.* (i): Let  $i_X : X \rightarrow X \times Y$ ,  $x \mapsto (x, y_0)$ , and  $i_Y$  analogously. Then

$$i_X^* \mu(a \otimes b) = i_X^* (\pi_X^*(a) \cdot \pi_Y^*(b)) = \underbrace{(\pi_X \circ i_X)^*}_{=\text{id}_X}(a) \cdot \underbrace{(\pi_Y \circ i_X)^*}_{=\text{constant}}(b) = 0,$$

since  $b \in \ker(i_{y_0}^* : K(Y) \rightarrow K(\{y_0\}))$  and the constant map  $\pi_Y \circ i_X$  factors through  $i_{y_0} : \{y_0\} \hookrightarrow Y$ . Analogously  $i_Y^* \mu(a \otimes b) = 0$ , so by exactness of the sequence from Corollary 3.5.16

$$\mu(a \otimes b) \in \ker((i_X^*, i_Y^*)) = \text{im}(q^*) = q^* \tilde{K}(X \wedge Y).$$

(ii): The first decomposition arises by tensoring the splittings  $K(X) \cong \tilde{K}(X) \oplus \mathbb{Z} \cdot 1_X$ , the second by combining the analogous splitting of  $K(X \times Y)$  with the splitting  $\pi_X^* \oplus \pi_Y^*$  of the exact sequence from Corollary 3.5.16. On the first summand  $\mu$  equals  $q^* \circ \tilde{\mu}$  by (i) and the definition of  $\tilde{\mu}$ . Further  $\mu(a \otimes 1_Y) = \pi_X^*(a) \cdot 1_{X \times Y} = \pi_X^*(a)$ , and  $\pi_X^*|_{\tilde{K}(X)}$  is an isomorphism onto  $\pi_X^* \tilde{K}(X)$ . The third summand is symmetric, and  $\mu(1_X \otimes 1_Y) = 1_{X \times Y}$ . Finally, a direct sum of homomorphisms is an isomorphism if and only if each block is one, and  $q^* \circ \tilde{\mu}$  is an isomorphism onto  $q^* \tilde{K}(X \wedge Y)$  if and only if  $\tilde{\mu}$  is one.  $\square$

By example 3.4.17 we know that for any line bundle  $l$ ,  $[l \otimes l \oplus \underline{1}] = [l]^2 + [\underline{1}] = [l] + [l]$  which is equivalent to  $([l] - 1)^2 = [l]^2 - [l] + [l] + [\underline{1}] = 0$ . Now  $\{[\underline{1}], [l]\}$  is an additive basis of  $\mathbb{Z}/([l]^2 - [\underline{1}])$ . The mapping that is defined on a basis of the tensor product as follows

$$\begin{aligned} \left( \mathbb{Z}([l]) / ([l]^2 - [\underline{1}]) \right) \otimes K(X) &\rightarrow K(S^2 \times X) \\ [\underline{1}] \otimes [\xi] &\mapsto \mu([\underline{1}], [\xi]) \\ [l] \otimes [\xi] &\mapsto \mu([l], [\xi]) \end{aligned}$$

is well defined.

**Definition 3.5.19.** We denote the external product of two elements by

$$\zeta \boxtimes \xi := \tilde{\mu}(\zeta \otimes \xi).$$

**Proposition 3.5.20** (Naturality). Let  $f : X \rightarrow X'$  and  $g : Y \rightarrow Y'$  be pointed maps, and  $a \in \tilde{K}(X')$ ,  $b \in \tilde{K}(Y')$ . Then

$$f^*(a) \boxtimes g^*(b) = (f \wedge g)^*(a \boxtimes b).$$

*Proof.* We start unreduced. Let  $f : X \rightarrow X'$  and  $g : Y \rightarrow Y'$  be continuous maps. then  $\pi_{X'} \circ (f \times g) = f \circ \pi_X$  and similar  $\pi_{Y'} \circ (f \times g) = g \circ \pi_Y$ . Using that  $(f \times g)^*$  is a ring homomorphism we calculate

$$\begin{aligned} (f \times g)^* \mu(a \otimes b) &= (f \times g)^* \left( (\pi_{X'})^*(a) \cdot (\pi_{Y'})^*(b) \right) \\ &= ((f \times g)^* (\pi_{X'})^*(a)) \cdot ((f \times g)^* (\pi_{Y'})^*(b)) \\ &= (\pi_X^* \circ f^*(a)) \cdot (\pi_Y^* \circ g^*(b)) \\ &= \mu(f^*(a) \otimes g^*(b)). \end{aligned}$$

Now assume that  $f$  and  $g$  are pointed and denote the projections  $q' : X' \times Y' \rightarrow X' \wedge Y'$  and  $q : X \times Y \rightarrow X \wedge Y$ . Then  $q' \circ (f \times g) = (f \wedge g) \circ q$  by definition of the wedge product of maps. But then we can calculate

$$q^*(f \wedge g)^*(a \boxtimes b) = (f \times g)^* q'^*(a \boxtimes b) = (f \times g)^* \mu(a \otimes b) = \mu(f^* a \otimes g^* b) = q^*(f^* a \boxtimes g^* b).$$

finally since  $q^*$  is injective we are done.  $\square$

prop:kTheo-associativityExternalProduct

**Proposition 3.5.21** (Associativity of the Reduced External Product). Let  $A, B, C$  be pointed spaces and  $a \in \tilde{K}(A)$ ,  $b \in \tilde{K}(B)$  and  $c \in \tilde{K}(C)$ . Let  $\psi : A \wedge (B \wedge C) \rightarrow (A \wedge B) \wedge C$  be the homeomorphism from Lemma 3.3.25. Then

$$\psi^*((a \boxtimes b) \boxtimes c) = a \boxtimes (b \boxtimes c).$$

*Proof.* Write  $q_{X,Y} : X \times Y \rightarrow X \wedge Y$  for the quotient maps and  $\pi_A, \pi_B, \pi_C$  for the projections of  $A \times B \times C$ . Let

$$Q := q_{A \wedge B, C} \circ (q_{A,B} \times \text{id}_C) : A \times B \times C \rightarrow (A \wedge B) \wedge C$$

and analogously  $Q' := q_{A, B \wedge C} \circ (\text{id}_A \times q_{B,C})$ . Both are quotient maps, and by the construction of  $\psi$  in Lemma 3.3.25 we have

$$\psi \circ Q' = Q, \tag{18}$$

{eq:kTheo-assocQuotients}

since both sides are induced by the identity on  $A \times B \times C$ . Furthermore  $Q^*$  is injective, being the composite of the injective maps  $q_{A \wedge B, C}^*$  and  $(q_{A,B} \times \text{id}_C)^*$ .

We first compute  $Q^*$  of the left hand side. By definition of  $\boxtimes$  we have the equalities

$$q_{A \wedge B, C}^*((a \boxtimes b) \boxtimes c) = \mu((a \boxtimes b) \otimes c) \in \tilde{K}((A \wedge B) \times C) \tag{19}$$

{eq:kTheo-assocExPro1}

$$\pi_C \circ (q_{A,B} \times \text{id}_C) = \pi_C : (A \times B) \times C \rightarrow C \tag{20}$$

{eq:kTheo-assocExPro2}

$$\pi_{A \wedge B} \circ (q_{A,B} \times \text{id}_C) = q_{A,B} \circ \pi_{A \times B} : (A \times B) \times C \rightarrow C \tag{21}$$

{eq:kTheo-assocExPro3}

hence

$$\begin{aligned} Q^*((a \boxtimes b) \boxtimes c) &= (q_{A,B} \times \text{id}_C)^*(\pi_{A \wedge B}^*(a \boxtimes b) \cdot \pi_C^*(c)) && \text{using (19)} \\ &= (q_{A,B} \times \text{id}_C)^*\pi_{A \wedge B}^*(a \boxtimes b) \cdot \pi_C^*(c) && \text{using (20) and (21)} \\ &= (q_{A,B} \times \text{id}_C)^*(\pi_A^*(a) \cdot \pi_B^*(b)) \cdot \pi_C^*(c) && \text{similar to (19)} \\ &= (\pi_A^*(a) \cdot \pi_B^*(b)) \cdot \pi_C^*(c) && \pi_A \circ \pi_{A \times B} = \pi_A \text{ and for } \pi_B \end{aligned}$$

The same computation on the other side gives

$$Q'^*(a \boxtimes (b \boxtimes c)) = \pi_A^*(a) \cdot \pi_B^*(b) \cdot \pi_C^*(c),$$

using associativity of the multiplication in  $K(A \times B \times C)$ . Combining this with (18),

$$Q'^*\psi^*((a \boxtimes b) \boxtimes c) = Q^*((a \boxtimes b) \boxtimes c) = Q'^*(a \boxtimes (b \boxtimes c)),$$

and by corollary 3.5.16 we know that  $q_{A,B}$  is injective. Hence,  $Q'^*$  is injective, which proves the claim.  $\square$

rem:kTheo-neutralElementExternalProduct

**Remark 3.5.22.** Let  $\zeta$  be the vector bundle over  $S^0 = \{-1, +1\}$  that is  $\underline{1}$  over  $-1$  and  $\underline{0}$  over  $+1$ . Then  $[\xi]$  is a generator of  $\tilde{K}(S^0)$ , as any bundle that is in the kernel of the inclusion of  $+1$  is  $\underline{0}$  over  $+1$ , and trivial over  $\{-1\}$ . Let  $\xi \in \tilde{K}(Y)$  be represented by

$\xi = [\xi_1] - [\xi_2]$  such that  $\xi_i$  are vector bundles ( $i = 1, 2$ ) over a pointed topological space  $Y$ . Then for  $i = 1, 2$ :

$$[\zeta] \boxtimes [\xi_i] = (q^*)^{-1}(\mu([\zeta] \otimes [\xi_i])) = (q^*)^{-1}\pi_{S^0}^*(\zeta) \cdot \pi_Y^*([\xi_i]) = (q^*)^{-1}([\pi_{S^0}^*(\zeta) \otimes \pi_Y^*(\xi_i)])$$

The bundle  $\pi_{S^0}^*(\zeta)$  is  $\underline{1}$  over  $\{-1\} \times Y$  and  $\underline{0}$  over  $\{1\} \times Y$ .  $\pi_Y^*(\xi_i)$  on the otherhand is the pullback of  $\xi_i$  via the homeomorphism  $h_- : \{-1\} \times Y \rightarrow Y$  over  $\{-1\} \times Y$  and similar over  $\{+1\} \times Y$ . Hence, the tensor product  $\pi_{S^0}^*(\zeta) \otimes \pi_Y^*(\xi_i)$  is given by  $h_-^*(Y)$  over

$$\{-1\} \times Y$$

and  $\underline{0}$  over  $\{+1\} \times Y$ . This bundles equivalence class in  $\tilde{K}$  is given by the bundle  $h_-^*$  viewed over the pointed  $S^0 \wedge Y = \{-1\} \times Y$  with basepoint  $[(-1, 1)]$ .

thm:kTheo-fundamentalProductTheorem

**Theorem 3.5.23** (The Fundamental Product Theorem). *For any compact  $X$ , and the tautological line bundle  $H$  on  $\mathbb{CP}^1 \cong S^2$ , the external product map*

$$\left(\mathbb{Z}([H]) / ([H] - [\underline{1}])^2\right) \otimes K(X) \rightarrow K(S^2 \times X)$$

*is an isomorphism of rings.*

*Proof.* The proof is omitted here. There are elementary proofs given in Chapter 2.1 [Hatcher2002algebraic] or Chapter 2.3 [Atiyah1989k].  $\square$

**Corollary 3.5.24** (The Easy Spaces). ?? We already know that  $\text{Vect}(pt) \cong \mathbb{N}$  as a semiring, where  $n \mapsto [\underline{n}]$ . Hence, as a Ring  $K(\{pt\}) \cong \mathbb{Z}$ . Since pointing the unpointed space  $\{pt\}$  results in  $S^0$  we know by Corollary 3.3.33 that  $K^0(S^0) \cong \mathbb{Z}$  and generated by the trivial bundle of degree one over  $\{-1\}$  and degree 0 over  $\{+1\}$ . We call this generator  $\beta_0$ . With the fundamental product Theorem, this tells us, that  $K(S^2) \cong \left(\mathbb{Z}([H]) / ([H] - [\underline{1}])^2\right)$  as a ring. Hence, we are isomorphic to the ring  $\mathbb{Z} \times \mathbb{Z}$  with componentwise multiplication.  $K(S^2)$  is generated by  $\{[\underline{1}], [H]\}$ . Furthermore, together with the map

$$\text{rank} : \left(\mathbb{Z}([H]) / ([H] - [\underline{1}])^2\right) \rightarrow \mathbb{Z}$$

that maps the generators to their rank, and

$$\nu : \mathbb{Z} \rightarrow K(pt); \quad n \mapsto [\underline{n}]$$

we get the commutative diagram:

$$\begin{array}{ccc} \mathbb{Z}([H]) / ([H]^2 - [\underline{1}]) & \xrightarrow{\text{rank}} & \mathbb{Z} \\ \downarrow & & \downarrow \nu \\ K(S^2) & \xrightarrow{i_{pt}^*} & K(pt) \end{array}$$

By the commutativity, we get an isomorphism induced by the map from the product theorem:

$$\tilde{K}(S^2) = \ker(i_{pt}^*) \cong \ker(\text{rank}) = \text{span}_{\mathbb{Z}}([H] - [\underline{1}])$$

Hence the Abelian group  $\tilde{K}(S^2) \cong \mathbb{Z}$ .

def:kTheo-BottGenerator

**Definition 3.5.25.** We call the generator  $([H] - [\underline{1}])$  that generates  $\tilde{K}(S^2)$  the **Bott generator** denoted by  $\beta$ .

thm:Ktheo-BottPeriodicity

**Theorem 3.5.26** (Bott Periodicity). *Let  $X$  be a well-pointed compact space. The isomorphism from the fundamental product theorem applied in its reduced form gives a natural isomorphism:*

$$\beta \boxtimes (-) : K^{-n}(X) \rightarrow K^{-n}(S^2 \wedge X) \cong K^{-n-2}(X)$$

*Proof.* We start with  $n = 0$ . By Proposition 3.5.18 we know that the reduced external product applied to the spaces  $\tilde{K}(S^2) \cong \text{span}_{\mathbb{Z}}([H] - [\underline{1}])$  and  $\tilde{K}(X)$  is an isomorphism, as applying the non-reduced external product gives an isomorphism by the fundamental product theorem. Hence, we have that

$$\mathbb{Z}([H] - [\underline{1}]) \otimes \tilde{K}(X) \rightarrow K^{-2}(X)$$

is an isomorphism. And since  $\tilde{K}(X) \cong \mathbb{Z}(\beta) \otimes \tilde{K}(X)$  via  $[\xi] \mapsto (\beta) \otimes [\xi]$  we have the isomorphism

$$\beta \boxtimes (-) : K^{-2}(X) \otimes \tilde{K}(X), \quad [\xi] \mapsto \tilde{\mu}(\beta \otimes [\xi]).$$

Naturality follows from  $\mu$  being natural,  $q^*$  being natural (and thereby its inverse) and  $\tilde{\mu}$  being the composition of both of them. Furthermore, for Replacing  $X$  with  $S^n X$  proves the lower degree.  $\square$

cor:kTheo-kGroupsOfTheSpheres

**Corollary 3.5.27** (Spheres). Let  $S^n$  be pointed spheres. We already saw, that  $K^0(S^2) = \tilde{K}(S^2) \cong \mathbb{Z}$ . Hence, the Bott periodicity together with the homotopy equivalence  $S^2 \wedge S^{2k-1} \simeq S^{2k}$  lets us deduce that  $K^0(S^2) \cong K^0(S^4) \cong \dots \cong K^0(S^{2k}) \cong \mathbb{Z}$ , and a generator of  $K^0(S^{2k})$  is given by  $\underbrace{\beta \boxtimes \dots \boxtimes \beta}_{k\text{-times}}$ .

On the other hand we can inspect  $\tilde{K}(S^1)$ . By Lemma 3.4.14 we know that

$$\tilde{K}(S^1) \cong \tilde{K}(\Sigma\{pt\}) \cong [\{pt\}, U] = \pi_0(U).$$

However  $U$  is pathconnected and thereby the reduced K group of the one sphere is trivial. The path connectedness of  $U$  follows from all  $U(n)$  being path connected which again can be deduced from  $\text{GL}_n(\mathbb{C})$  being path connected and contracting onto  $U(n)$ . Then, by the Bott periodicity we can conclude that  $\tilde{K}(S^{2k+1}) = 0$  for all  $k \in \mathbb{N}$ .

natürlich  
im  
zweiten  
faktor mit  
festem er-  
sten

def:ktheo-higherBottGenerators

**Definition 3.5.28** (Convention). Let  $\beta_0 \in \tilde{K}^0(S^0)$  be the canonical generator from Corollary ??,  $\beta_1 \in \tilde{K}^{-1}(S^1)$  be the Bott generator of  $\tilde{K}(S^2)$ , viewed in  $\tilde{K}^{-1}(S^1) = \tilde{K}^0(S^1 \wedge S^1)$  via Definition 3.3.27. Writing

$$\Upsilon_{n,m} := (\overline{\pi_S^{n+m}})^{-1} \circ \Xi_{\mathbb{R}^n, \mathbb{R}^m}^{-1} \circ (\overline{\pi_S^n} \wedge \overline{\pi_S^m}) : S^n \wedge S^m \rightarrow S^{n+m}$$

for the homeomorphism of Definition 3.3.27, we define inductively

$$\beta_n := (\Upsilon_{1,n-1}^*)^{-1} (\beta_1 \boxtimes \beta_{n-1}) \in \tilde{K}(S^{2n}) \cong \tilde{K}^{-n}(S^n).$$

By Theorem 3.5.26, the external product with  $\beta_1$  is an isomorphism, so each  $\beta_n$  is a generator.

lem:ktheo-bottGeneratorsAreMultiplicative

**Lemma 3.5.29.** For all  $n, m \geq 0$  we have  $\Upsilon_{n,m}^*(\beta_{n+m}) = \beta_n \boxtimes \beta_m$ .

*Proof.* Set  $b_i := (\overline{\pi_S^i})^*(\beta_i) \in \tilde{K}^{-i}(\mathbb{R}^i)$ .

*Reduction.* Since  $\overline{\pi_S^n} \wedge \overline{\pi_S^m}$  is a homeomorphism, naturality of  $\boxtimes$  gives  $(\overline{\pi_S^n} \wedge \overline{\pi_S^m})^*(\beta_n \boxtimes \beta_m) = b_n \boxtimes b_m$ . Applying  $(\overline{\pi_S^n} \wedge \overline{\pi_S^m})^*$  to the claim and using  $(\overline{\pi_S^n} \wedge \overline{\pi_S^m})^* \Upsilon_{n,m}^* = (\Upsilon_{n,m} \circ (\overline{\pi_S^n} \wedge \overline{\pi_S^m}))^* = ((\overline{\pi_S^{n+m}})^{-1} \circ \Xi_{n,m}^{-1})^*$ , the claim becomes

$$\Xi_{n,m}^*(b_n \boxtimes b_m) = b_{n+m}, \quad (22)$$

{eq:kTheo-multiplicative}

and by definition  $b_n = \Xi_{1,n-1}^*(b_1 \boxtimes b_{n-1})$ .

*Base case*  $n = 0$ . Here  $\Xi_{0,m}$  is the canonical homeomorphism  $\mathbb{R}^m \rightarrow \mathbb{R}^0 \wedge \mathbb{R}^m$  and (22) is Remark 3.5.22.

*Induction step.* By definition of  $\psi$  in Lemma 3.3.25 the diagram above commutes, i.e.

$$(\Xi_{1,n-1} \wedge \text{id}) \circ \Xi_{n,m} = \psi \circ (\text{id} \wedge \Xi_{n-1,m}) \circ \Xi_{1,(n-1)+m}. \quad (23)$$

{eq:kTheo-coherence}

Now

$$\begin{aligned} \Xi_{n,m}^*(b_n \boxtimes b_m) &= \Xi_{n,m}^*(\Xi_{1,n-1}^*(b_1 \boxtimes b_{n-1}) \boxtimes \text{id}^*(b_m)) && \text{definition of } b_n \\ &= \Xi_{n,m}^*(\Xi_{1,n-1} \wedge \text{id})^*((b_1 \boxtimes b_{n-1}) \boxtimes b_m) && \text{naturality of } \boxtimes \\ &= ((\Xi_{1,n-1} \wedge \text{id}) \circ \Xi_{n,m})^*((b_1 \boxtimes b_{n-1}) \boxtimes b_m) && \text{functoriality} \\ &= \Xi_{1,(n-1)+m}^*(\text{id} \wedge \Xi_{n-1,m})^* \psi^*((b_1 \boxtimes b_{n-1}) \boxtimes b_m) && \text{using (23)} \\ &= \Xi_{1,(n-1)+m}^*(\text{id} \wedge \Xi_{n-1,m})^*(b_1 \boxtimes (b_{n-1} \boxtimes b_m)) && \text{Proposition 3.5.21} \\ &= \Xi_{1,(n-1)+m}^*(\text{id}^*(b_1) \boxtimes \Xi_{n-1,m}^*(b_{n-1} \boxtimes b_m)) && \text{naturality of } \boxtimes \\ &= \Xi_{1,(n-1)+m}^*(b_1 \boxtimes b_{(n-1)+m}) && \text{induction hypothesis} \\ &= b_{n+m} && \text{definition of } b_{n+m} \end{aligned}$$

□

## 4 Orientations

The idea of Morse homology is now to find an invariant under manipulations of the Morse–Smale function. Obviously neither the number of critical points nor the number of flow lines stays invariant. However certain linear combinations stay invariant, when one establishes a sophisticated way to count flow lines. For this count we want to introduce signs to the flow lines between critical points with index one apart where the signs rely on orientations of over the flow lines.

### 4.1 Oriented Vector Spaces and Bundles

**Definition 4.1.1** (Orientation). Let  $V$  be a finite dimensional  $\mathbb{R}$  vector space and  $\det(V) := \Lambda^{\dim(V)}$  the maximal exterior algebra (or Grassmann algebra). By the principle of permanence we define  $\Lambda^0(V) = \mathbb{R}$ . An orientation of  $V$  is a choice of one of the two connected components of  $\det(V) \setminus \{0\}$ . we call the tuple  $(V, [\mathfrak{o}])$  where  $\mathfrak{o}$  is a non-zero element of  $\det(V)$  and hence  $[\mathfrak{o}]$  is an orientation, an **oriented vector space**. Often we will just write  $V$  oriented, and omitt the orientation.

**Definition 4.1.2** (Convention). If we use  $\mathbb{R}^n$  as an oriented vector space we fix  $e := e_1 \wedge \cdots \wedge e_n$  to induce the orientation.

**Remark 4.1.3.** Classically, an orientation of a vector space  $V$  of dimension  $m$  is given by an ordered basis  $\mathfrak{b} = (b_1, \dots, b_m)$  of  $V$  and hence the product  $b_1 \wedge \cdots \wedge b_m$  is an element of the maximal exterior product  $\det(V) = \Lambda^m V$ . Two ordered bases define different elements in  $\det(V)$  and their difference is given by the determinant of the base change matrix. (This is just a calculation using multilinearity and the alternating property). Hence a choice of equivalence class of bases modulo base change matrix of positive determinant is the same as a choice of one connected component of  $\det(V) \setminus \{0\}$ .

**Definition 4.1.4** (Determinant of an endomorphism). Let  $f : V \rightarrow V$  be an endomorphism. Then we define

$$\Lambda^{\dim(V)}(f) : \det(V) \rightarrow \det(V) \quad (24)$$

$$v_1 \wedge \cdots \wedge v_{\dim(V)} \mapsto f(v_1) \wedge \cdots \wedge f(v_{\dim(V)}). \quad (25)$$

Since  $\dim(\det(V)) = 1$ , the function  $\Lambda^{\dim(V)}(f)$  is given by a scalar multiplication with a nonzero element  $\lambda \in \mathbb{R}$ . We define the determinant  $\det(f) := \lambda$  to be said scalar. Obviously, if  $\det(V)$  is positive, then  $f$  is orientation preserving in the sence, that  $f$  applied to the ordered basis gives again an ordered basis that is compatible.

**Remark 4.1.5.** From linear algebra we know, that we can calculate  $\det(f) = \det(M_{\mathfrak{b}}^{\mathfrak{b}}(f))$ , where  $\mathfrak{b}$  is an ordered basis of  $V$ . Since the determinant of a matrix commutes with taking products, we have that the determinant is invariant under conjugation of linear endomorphisms.

rem:orient-calculateDeterminantViaMatrixDeterminant

```
def:orient-Si
gnOfAAbstract
LinearMap
```

**Definition 4.1.6.** Let  $(V, [\mathfrak{o}])$  and  $(W, [\mathfrak{p}])$  be oriented vector spaces. We call a linear isomorphism  $f : V \rightarrow W$  between two oriented vector spaces **orientation preserving**, if  $[\Lambda^{\dim(V)}(f)(\mathfrak{o})] = [\mathfrak{p}]$ . Let  $o$  and  $p$  denote the basis induced from  $\mathfrak{o}$  resp.  $\mathfrak{p}$ . Then by the Remark 4.1.5 the map  $f$  is orientation preserving, if and only if  $\det(M_p^o(f))$  is positive. We define the sing of  $f$  (and sometimes even the seign of  $\det(f)$ ) to be the sign of  $\det(M_p^o(f))$ .

**Definition 4.1.7** (Orientation and Exact Sequences). A short exact sequence of vector spaces

FIX

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

induces a canonical isomorphism

$$\det(A) \otimes \det(C) \rightarrow \det(B)$$

$$(a_1 \wedge \cdots \wedge a_l) \otimes (c_1 \wedge \cdots \wedge c_k) \mapsto (f(a_1) \wedge \cdots \wedge f(a_l) \wedge \tilde{c}_1 \wedge \cdots \wedge \tilde{c}_k),$$

where  $\tilde{c}_j \in g^{-1}(c_j)$ . The well-definedness of this map (invariance under chosen lift of  $c$ ) can be calculated, since two different choices differ in an element in the kernel of  $g$  which equals the image of  $f$  and hence the different choices vanish in the wedge product. By this isomorphism we get an orientation induced in the middle from the outside. Notice the convention, that the basis from  $A$  is placed in front of the basis from  $C$ . Notice how the definition is consistent with the edge cases of trivial vector spaces  $A$  or  $C$ . If  $A = 0$  we get an isomorphism  $B \cong C$  and a basis from  $C$  induced to  $B$ , that is either the induced one from  $g$ , or exactly the opposite one, if the trivial vector space  $A$  is negatively oriented.

**Remark 4.1.8.** Let

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

be a short exact sequence of vector spaces and  $\mathfrak{a}$  be a basis of  $A$ ,  $\mathfrak{c}$  be a basis of  $C$  and  $\tilde{\mathfrak{c}}$  be a lift of  $\mathfrak{c}$  via  $g$ . Replacing the basis  $\mathfrak{a}$  with  $P\mathfrak{a}$  and  $\mathfrak{c}$  with  $Q\mathfrak{c}$  where  $P, Q$  are invertible matrices, gives us a new induced basis on  $B$ , which differs from the original

one by a matrix  $\begin{pmatrix} P & * \\ 0 & Q \end{pmatrix}$ , where  $*$  depends on the chosen lift. Hence the corresponding elements in  $\det(B)$  differ by the product  $\det(P)\det(Q)$ .

```
def:orient-or
ientedVectorB
undle
```

**Definition 4.1.9.** For a rank  $r$  vector bundle  $E \rightarrow X$  we set  $\det(E) := \Lambda^r(E)$ . An orientation is an equivalence class of a non-vanishing section into the determinant bundle, where two sections are equivalent if they differ by multiplication of a positive scalar valued function.  $E$  is orientable if such a section exists, i.e. if  $\det(E)$  is trivial. An orientation of a bundle restricts to an orientation in each fibre.

lem:morse-Ori  
entationsInAP  
ointInduceOri  
entationsOfBu  
ndle

**Lemma 4.1.10.** *Let  $E \rightarrow X$  be an orientable vector bundle and  $X$  be connected. For each  $x \in X$  we get the bijection induced from the restriction to the fibre:*

$$\{\text{Orientation on } E\} \rightarrow \{\text{Orientation on } E_x\}.$$

*Hence, an orientation in one fibre can introduce an orientation in every other fibre.*

*Proof.* Two orientations over  $E$  are given as nowhere-vanishing sections  $s, s'$ . The quotient  $x \mapsto \frac{s}{s'}$  is continuous and its image is contained in  $\mathbb{R} \setminus \{0\}$ . Hence the sign of the quotient is a continuous function into  $\{-1, 1\}$ . Since  $X$  is connected we conclude that the sign of said quotient is a constant function. Thereby, if the sections induce the same orientation, they differ by multiplication of the positive number  $\frac{s(x)}{s'(x)}$  and then the sign of  $\frac{s}{s'}$  is globally one. The surjectivity is clear by mapping the sections  $s$  and  $-s$ .  $\square$

**Definition 4.1.11.** Let  $\pi : E \rightarrow X$  and  $\pi' : E' \rightarrow X'$  be two oriented vector bundles,  $f : X \rightarrow X'$  continuous, and  $\Phi : E \rightarrow E'$  a morphism of vector bundles over  $f$  such that for all  $x$ ,  $\Phi_x$  is an isomorphism of vector bundles. Then we call  $\Phi$  orientation preserving, if and only if  $\Phi_x : E_x \rightarrow E'_{f(x)}$  is an orientation preserving map between oriented vector spaces, that inherit their orientation from the bundles via Lemma 4.1.9 for all  $x \in X$ .

def:orient-or  
ientedMorphis  
mOfBundlesEqu  
ivalentDef

**Proposition 4.1.12** (Equivalent Definition). Let  $s$  and  $t$  be non-vanishing sections into the determinant bundle that represent orientations of the oriented vector bundles  $\pi : E \rightarrow X$  and  $\pi' : E' \rightarrow X'$ ,  $f : X \rightarrow Y$  be continuous and  $\Phi : E \rightarrow E'$  be a bundle morphism over  $f$ . Then for all  $x \in X$  there exists a  $\lambda(x) \in \mathbb{R} \setminus \{0\}$  such that  $\Lambda^{\text{rank}(E)}(\Phi)(s(x)) = \lambda(x)t(f(x))$ . The map  $x \mapsto \lambda(x)$  is continuous, as  $\lambda = \frac{\Lambda^{\text{rank}(E)}(\Phi) \circ s}{t \circ f}$  is continuous (one can check this in a local frame of  $\det(E')$ ) and non-vanishing. Thereby its sign is locally constant. Hence, if  $X$  is connected, then  $\Phi$  is oriented, if for **any**  $x \in X$   $\Phi_x$  is an orientation preserving map between oriented vector spaces, that inherit their orientation from the bundles via Lemma 4.1.9

**Proposition 4.1.13.** Let

$$0 \longrightarrow E' \xrightarrow{f} E \xrightarrow{g} E'' \longrightarrow 0$$

be a short exact sequence of orientable vector bundles. Then the fibrewise isomorphisms  $\det(E'_p) \otimes \det(E''_p) \rightarrow \det(E_p)$  from the vector space case assemble into an isomorphism of line bundles

$$\det(E') \otimes \det(E'') \xrightarrow{\cong} \det(E).$$

In particular, orientations of  $E'$  and  $E''$  induce a canonical orientation of  $E$ .

*Proof.* Fibrewise the map is the canonical isomorphism from the vector space case: a lift  $\tilde{c} \in g_p^{-1}(c)$  exists since  $g_p$  is surjective, and the resulting element  $f(a_1) \wedge \cdots \wedge f(a_l) \wedge \tilde{c}_1 \wedge \cdots \wedge \tilde{c}_k$  is independent of the choice of lift.

It remains to check that this bundle map is continuous (resp. smooth) to ensure that it maps sections smoothly into sections, which is a local statement: locally choose a frame  $(a_1, \dots, a_l)$  of  $E'$  and a frame  $(c_1, \dots, c_k)$  of  $E''$ ; since  $g$  is an epimorphism it admits local sections, so we may choose local lifts  $\tilde{c}_j$  of the  $c_j$  which are (smooth) local sections of  $E$ . Then the map is given in the induced local frames of the determinant bundles by wedging these sections, hence continuous (resp. smooth). By this, two non-vanishing sections get mapped to a non-vanishing section and hence, orientations on  $E'$  and  $E''$  which are represented by sections give a orientation over  $E$ .  $\square$

**Definition 4.1.14.** We call a manifold orientable, if its tangent bundle is **orientable**. We call a tuple  $(M, [\omega])$  where  $[\omega]$  denotes an orientation of  $TM$  an **oriented manifold**. We will often omit the orientation and just call  $M$  oriented.

**Definition 4.1.15** (Orientation Preserving Charts and Maps). Let  $(M, [\omega])$  be an oriented  $n$ -dimensional manifold and let  $\varphi : U \rightarrow \mathbb{R}^n$  be a chart. We call  $\varphi$  **orientation preserving** if for every  $x \in U$   $\partial_1(x) \wedge \dots \wedge \partial_n(x)$  and  $\omega(x)$  lie in the same connected component of  $\det(T_x M) \setminus \{0\}$ . More generally, let  $(N, [\eta])$  be another oriented  $n$ -dimensional manifold and let  $h : M \rightarrow N$  be differentiable at  $x$  with  $dh_x$  invertible. We define

$$\text{sign det}(dh_x) := \begin{cases} +1 & \text{if } \Lambda^n(dh_x)(\omega(x)) \text{ and } \eta(x) \text{ induce the same orientation in } T_h(x)N \\ -1 & \text{otherwise.} \end{cases}$$

**Remark 4.1.16.** One easily checks that  $\text{sign det}(dh_x)$  is well-defined and reversing both orientations doesn't affect the sign. In particular, for  $N = M$  the number  $\text{sign det}(dh_x)$  does not depend on the chosen orientation.

lem:orient-computingSignInCharts

**Lemma 4.1.17** (Computing the Sign in Charts). *Let  $(M, [\omega])$  and  $(N, [\eta])$  be oriented  $n$ -dimensional manifolds and  $\varphi$  be an orientation preserving chart around  $x$  in  $M$  and  $\psi$  one around  $h(x)$ . Then*

$$\text{sign det}(dh_x) = \text{sign det} \left( d(\psi \circ h \circ \varphi^{-1})|_{\varphi(x)} \right),$$

where the right hand side calculates the sign of the determinant of a matrix.

*Proof.* Define  $A := d(\psi \circ h \circ \varphi^{-1})|_{\varphi(x)} = d\psi|_{h(x)} \circ dh_x \circ (d\varphi|_x)^{-1}$  and denote the standard orientation of  $\mathbb{R}^n$  by  $e$ . Since  $\varphi$  is orientation preserving and  $\Lambda^n(d\varphi|_x)(\partial_1(x) \wedge \dots \wedge \partial_n(x)) = e$ , we have by linearity  $\Lambda^n(d\varphi|_x)(\omega(x)) = a \cdot e$  where  $a > 0$ . Similar  $\Lambda^n(d\psi|_{h(x)})(\eta(h(x))) = b \cdot e$  with  $b > 0$ . Hence

$$\Lambda^n(d\psi|_{h(x)}) \circ \Lambda^n(dh_x)(\omega(x)) = \Lambda^n(A) \circ \underbrace{\Lambda^n(d\varphi|_x)(\omega(x))}_{=a \cdot e} = a \det(A) \cdot e$$

Now by definition we need to check if  $\Lambda^n(dh_x)(\omega(x)) = c \cdot \eta(x)$  for  $c > 0$ , which is equivalent to

$$\underbrace{\Lambda^n(d\psi|_{h(x)}) \circ \Lambda^n(dh|_x)(\omega(x))}_{=a \det(A) \cdot e} = c \cdot \underbrace{\Lambda^n(d\psi|_{h(x)})(\eta(x))}_{=b \cdot e}.$$

which is given if and only if  $\det(A) > 0$ .  $\square$

cor:orient-existsOrientationPreservingAtlas

**Corollary 4.1.18.** A smooth manifold  $M$  of dimension  $n$  is orientable, if and only if it has an atlas such that the determinants of the Jacobians of the transition functions are positive.

*Proof.* Let  $[\omega]$  be an orientation on  $M$  and let  $U_\alpha$  be an open cover of  $M$  with all  $U_\alpha$  connected and the charts  $\varphi_\alpha : U_\alpha \rightarrow \mathbb{R}^n$ . Let  $\partial_1^\alpha \wedge \cdots \wedge \partial_n^\alpha$  be the induced local section into the determinant bundle. Then

$$\omega|_{U_\alpha} = g_\alpha \partial_1^\alpha \wedge \cdots \wedge \partial_n^\alpha$$

with  $g_\alpha$  non vanishing and  $U_\alpha$  connected. If  $g_\alpha$  is negative, replace  $\varphi_\alpha$  with  $\tilde{\varphi}_\alpha(x) = (-\varphi_\alpha(x)_1, \dots, \varphi_\alpha(x)_n)$ , which changes the sign of  $g_\alpha$ . Doing this in all charts gives an orientation preserving atlas. Hence, by definition  $\text{sign det}(d\text{id}_M|_x) = 1$  and by Lemma 4.1.17 we can calculate  $\text{sign det}(d\text{id}_M|_x)$  in this atlas by the determinants of the Jacobians of the transition functions. Hence those determinants are positive.

For the other direction we want to glue together the local sections  $\partial_1^\alpha \wedge \cdots \wedge \partial_n^\alpha$  using a smooth partition of unity subordinate to the open cover  $\{U_i\}$ . This yields something non-vanishing, as all local sections are pointwise positive multiples of each other where they intersect.  $\square$

cor:orient-transitionFunctionPositiveDeterminantAlsoOrientationPreserving

**Corollary 4.1.19.** The proof of Corollary 4.1.18 also shows that if one chart is orientation preserving, then also another chart that intersects with the first and has positive determinant of the Jacobian of the transition function.

**Lemma 4.1.20.** Let  $M$  be a smooth manifold that can be covered with two charts such that their intersection is connected. Then  $M$  is orientable.

*Proof.* Just ensure that the two charts are compatible such that they build an atlas such that the determinants of the Jacobians of the transition functions are positive. To achieve this one just manipulates one of the charts.  $\square$

**Corollary 4.1.21.**  $S^n$  for  $n \geq 2$  is orientable, as it is covered by the two stereographic projections.  $S^1$  and  $S^0$  are also orientable which can be seen by picking charts such that they fit into corollary 4.1.18.

## 4.2 K-theoretic Orientation

K theoretic orientation relies on bases induced generators of the K group of the sphere. Hence, we want to fix a canonical map from  $R^n$  to  $S^n$ :

lem:reflectio  
n-is-minus-on  
e

**Lemma 4.2.1** (Reflection in one variable). *Let  $R_1 : \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto -x$ , and let  $Z$  be a pointed compact Hausdorff space. Then*

$$(\overline{R_1} \wedge \text{id}_Z)^* = -\text{id} \quad \text{on } \tilde{K}(\overline{\mathbb{R}} \wedge Z).$$

*Proof.* Realise  $S^1$  as  $\mathbb{R}/\mathbb{Z}$  with base point  $[0]$ , so that the map  $T$  of Corollary 3.4.16 is  $T([t]) = [-t]$ . Define

$$h : \overline{\mathbb{R}} \rightarrow \mathbb{R}/\mathbb{Z}, \quad h(x) = \left[ \frac{1}{2} + \frac{1}{\pi} \arctan(x) \right], \quad h(\infty) = [0].$$

Since  $x \mapsto \frac{1}{2} + \frac{1}{\pi} \arctan(x)$  is a strictly increasing bijection onto  $(0, 1)$ , the map  $h$  is bijective; it is continuous on  $\mathbb{R}$  and also at  $\infty$ , because  $\arctan(x) \rightarrow \pm \frac{\pi}{2}$  gives  $h(x) \rightarrow [1] = [0]$  resp.  $h(x) \rightarrow [0]$ . Being a continuous bijection from a compact space to a Hausdorff space,  $h$  is a pointed homeomorphism. It intertwines the two reflections:

$$h(-x) = \left[ \frac{1}{2} - \frac{1}{\pi} \arctan(x) \right] = \left[ -\frac{1}{2} - \frac{1}{\pi} \arctan(x) \right] = T(h(x)),$$

using  $[-\frac{1}{2}] = [\frac{1}{2}]$ , and  $h(\overline{R_1}(\infty)) = [0] = T(h(\infty))$ . Hence  $h \wedge \text{id}_Z$  is a pointed homeomorphism with

$$(h \wedge \text{id}_Z) \circ (\overline{R_1} \wedge \text{id}_Z) = (T \wedge 1) \circ (h \wedge \text{id}_Z).$$

Applying  $\tilde{K}$  and Corollary 3.4.16 gives

$$(\overline{R_1} \wedge \text{id}_Z)^* = (h \wedge \text{id}_Z)^* \circ (T \wedge 1)^* \circ ((h \wedge \text{id}_Z)^*)^{-1} = -\text{id},$$

as all maps are ring homomorphisms and thereby commute with multiplication by minus one.  $\square$

**Definition 4.2.2** (K-theoretic orientations). We define a **K-theoretic orientation** of a finite dimensional real vector space  $V$  to be a generator of the group  $\tilde{K}^{-m}(\overline{V})$ . As a sanity check we can confirm: since  $\overline{V} \cong S^m$  we are looking at a generator of a group isomorphic to  $\tilde{K}(S^{2m}) \cong \mathbb{Z}$ , which has two possible generators.

lem:orient-li  
nearMapsOnSma  
shProducts

**Lemma 4.2.3** (Linear Maps on Smash Products). *Let  $A \in \text{GL}(V)$  and let  $Z$  be a pointed compact space. Then*

$$(\overline{A} \wedge \text{id}_Z)^* = \text{sign det}(A) \cdot \text{id} \quad \text{on } \tilde{K}(\overline{V} \wedge Z).$$

*In particular, if  $\overline{\mathbb{R}^{m_1}} \wedge \cdots \wedge \overline{\mathbb{R}^{m_r}}$  is identified with  $\overline{R^m}$  by Lemma 3.3.24, a permutation of the smash factors acts on  $\tilde{K}(\overline{\mathbb{R}^{m_1}} \wedge \cdots \wedge \overline{\mathbb{R}^{m_r}} \wedge Z)$  by the sign of the*

induced permutation of the  $m = m_1 + \dots + m_r$  coordinates.

*Proof.* Since the determinant of an endomorphism is invariant under conjugation we can choose a basis and assume that  $V = \mathbb{R}^m$ . The group  $\mathrm{GL}_m(\mathbb{R})$  has exactly two path components, separated by the sign of the determinant. Hence, there is a continuous path  $A_\bullet : [0, 1] \rightarrow \mathrm{GL}_m(\mathbb{R})$  with  $A_0 = A$  and  $A_1 = \mathrm{id}$  if  $\det(A) > 0$  and  $R$  if  $\det(A) < 0$  with  $R$  being the reflection  $(x_1, \dots, x_m) \mapsto (-x_1, x_2, \dots, x_m)$ . The map

$$H : \mathbb{R}^m \times [0, 1] \rightarrow \mathbb{R}^m, \quad H(v, t) := A_t v$$

is a proper homomorphism into a locally compact space: If  $K \subseteq \mathbb{R}^m$  is compact and  $A_t v \in K$ , then by definition of the operator norm we have that  $\|v\| \leq (\sup_t \|A_t^{-1}\|) \cdot \max_{k \in K} \|k\|$ . Hence  $H^{-1}(K)$  is bounded and closed by continuity. Thereby the extension  $\bar{H} : \bar{V} \times \bar{I} \rightarrow \bar{V}$  is continuous. Let  $q : \bar{V} \times \bar{I} \rightarrow \bar{V} \times \bar{I}$  be the collapse of  $\{\infty\} \times I$  to  $\infty$ . Then we have that  $\bar{H} = \tilde{H} \circ q$  and hence  $\bar{H}$  is continuous. If  $\det(A) > 0$ , we are done. Let  $\bar{R}$  denote the extension of  $R$  to the compactifications. What is left to show is that  $(\bar{R} \wedge \mathrm{id}_Z)^*$  is given by multiplication with minus one.

For this we write  $R = R_1 \times \mathrm{id}_{\mathbb{R}^{n-1}}$  where  $R_1$  denotes the reflection  $x \mapsto -x$  on  $\mathbb{R}$ . Let  $\Xi : \mathbb{R}^n \rightarrow \mathbb{R} \wedge \mathbb{R}^{n-1}$  be the natural homeomorphism from Lemma 3.3.24. By the naturality for proper maps we have that  $\Xi \circ (\bar{R}) = (\bar{R}_1 \wedge \mathrm{id}_{\mathbb{R}^{n-1}}) \circ \Xi$ , giving us:

$$(\Xi \wedge \mathrm{id}_Z)(\bar{R} \wedge \mathrm{id}_Z) = (\Xi \circ \bar{R}) \wedge \mathrm{id}_Z = ((\bar{R}_1 \wedge \mathrm{id}_{\mathbb{R}^{n-1}}) \circ \Xi) \wedge \mathrm{id}_Z$$

If we now define  $L := \overline{\mathbb{R}^{m-1}} \wedge Z$  and let  $AS : (\overline{\mathbb{R}^1} \wedge \overline{\mathbb{R}^{m-1}}) \wedge Z \rightarrow \overline{\mathbb{R}^1} \wedge (L)$  be the associativity of the wedge product. Then we have

$$AS \circ (\Xi \wedge \mathrm{id}_Z)(\bar{R} \wedge \mathrm{id}_Z) = (\bar{R}_1 \wedge \mathrm{id}_L) \circ AS \circ \Xi \wedge \mathrm{id}_Z$$

Hence, if we define  $\Theta := AS \circ \Xi \wedge \mathrm{id}_Z$  we get

$$\bar{R} \wedge \mathrm{id}_Z = \Theta^{-1} \circ (\bar{R}_1 \wedge \mathrm{id}_L) \circ \Theta$$

or as maps of K-groups:

$$(\bar{R} \wedge \mathrm{id}_Z)^* = \Theta^* \circ \underbrace{(\bar{R}_1 \wedge \mathrm{id}_L)^*}_{= -\mathrm{id} \text{ by Lemma 4.2.1}} \circ (\Theta^{-1})^* = -\mathrm{id}.$$

□

prop:linearityTheDeterminandDertermines

**Proposition 4.2.4.** Let  $V$  be an  $n$ -dimensional real vector space and  $A \in \mathrm{GL}(V)$ . Then

$$\bar{A}^* = \mathrm{sign} \det(A) \cdot \mathrm{id} \quad \text{on } \tilde{K}^{-n}(\bar{V}).$$

*Proof.* The statement is almost a corollary of Lemma 4.2.3: We care for the map  $\bar{A}^*$  which is defined as the map

$$(\mathrm{id}_{S^n} \wedge A)^* : \tilde{K}^{-n}(V) = \tilde{K}(S^n \wedge V) \rightarrow \tilde{K}(S^n \wedge V) = \tilde{K}^{-n}(V)$$

Let  $\Theta : S^n \wedge V \rightarrow V \wedge S^n$  be the reordering from Lemma 3.3.25. Then we have that

$$\text{id}_{S^n} \wedge A = \Theta^{-1} \circ (A \wedge \text{id}_{S^n}) \circ \Theta.$$

Hence, lemma 4.2.3 gives us:

$$\overline{A}^* = (\Theta^{-1} \circ (A \wedge \text{id}_{S^n}) \circ \Theta)^* = \Theta^* \circ \underbrace{(A \wedge \text{id}_{S^n})^*}_{=\text{sign det}(A) \cdot \text{id}} \circ (\Theta^{-1})^* = \text{sign det}(A) \cdot \text{id}.$$

□

**Corollary 4.2.5.** Any finite dimensional real vector space  $V$  is homeomorphic to  $\mathbb{R}^n$  via the coordinate map  $\kappa_v$  that maps each vector to its coordinate representation. With this we can introduce a K-theoretic orientation from an ordered basis via the composition:

$$\tilde{K}^{-n}(S^n) \xrightarrow{(\pi_S^{-1})^*} \tilde{K}^{-n}(\mathbb{R}^n) \xrightarrow{\overline{\kappa}_v^*} \tilde{K}^{-n}(\overline{V})$$

With this we can map the bott generator  $\beta$  from Definition 3.5.25 to a generator of  $\tilde{K}^{-n}(\overline{V})$  and hence induce a K-Theoretic orientation. This orientation only depends on the orientation of  $V$  and two different orientations induce different K-theoretic orientations

*Proof.* The statement follows immediately from Proposition 4.2.4. □

**Corollary 4.2.6.** Let  $f : V \rightarrow W$  be a linear isomorphism between oriented vector spaces  $(V, \mathfrak{a})$  and  $(W, \mathfrak{b})$ . Let  $\beta_V$  be the K-Theoretic orientation of  $V$  and  $\beta_W$  of  $W$ . Then

$$\overline{f}^*(\beta_W) = \text{sign det}(f) \cdot (\beta_V),$$

where  $\text{sign det}(f)$  denotes the sign of  $f$  as in Definition 4.1.6.

*Proof.* Let  $\kappa_V : V \rightarrow \mathbb{R}^n$  and  $\kappa_W : W \rightarrow \mathbb{R}^n$  be positive coordinate maps. By Proposition 4.2.4 we know that

$$(\overline{\kappa}_W \circ \overline{f} \circ \overline{\kappa}_V^{-1})^* = \text{sign det}(f) \cdot \text{id}$$

Now  $\beta_V = \overline{\kappa}_V^*(\beta_{\mathbb{R}^n})$  and  $\beta_W = \overline{\kappa}_W^*(\beta_{\mathbb{R}^n})$ . This concludes the proof, as

$$\begin{aligned} \overline{f}^*(\beta_W) &= (\overline{\kappa}_V^* \circ (\overline{\kappa}_V^*)^{-1}) \circ \overline{f}^* \circ (\overline{\kappa}_W^*(\beta_{\mathbb{R}^n})) \\ &= \overline{\kappa}_V^* \circ (\overline{\kappa}_W \circ \overline{f} \circ \overline{\kappa}_V^{-1})^* \beta_{\mathbb{R}^n} \\ &= \text{sign det}(f) \overline{\kappa}_V^* \beta_{\mathbb{R}^n} = \text{sign det}(f) \beta_V. \end{aligned}$$

□

lem:orient-mu  
ltiplication

**Lemma 4.2.7 (Multiplication).** Let  $V, W$  be oriented real vector spaces of dimension  $n, m$  and give  $V \oplus W$  the orientation by placing the orientation of  $V$  in front of the one of  $W$ . Let  $\beta_{\mathfrak{o}}$  be the K Theoretic orientation on  $V$  induced by the orientation of  $V$  and similar  $\beta_{\mathfrak{p}}$  be the one on  $W$ , and finally  $\beta_{\mathfrak{o} \oplus \mathfrak{p}}$  be the K-theoretic

cor:orient-ve  
ctorspaceOrie  
ntationInduce  
sKtheoreticOr  
ientation

cor:orient-ma  
psbetweenabst  
ractVectorSpa  
cesOnKOrienta  
tions

orientation of  $V \oplus W$ . Then under the homeomorphism

$$\Xi : \overline{V \oplus W} \rightarrow \overline{V} \wedge \overline{W}$$

from Lemma 3.3.24 we have

$$\Xi^* (\beta_{\mathbf{o}} \boxtimes \beta_{\mathbf{p}}) = \beta_{\mathbf{o} \oplus \mathbf{p}}.$$

*Proof.* Let  $\kappa_v : V \rightarrow \mathbb{R}^n$  and  $\kappa_w : \mathbb{R}^n \rightarrow W$  be two basis maps from ordered basis compatible with the orientations of  $V$  and  $W$ . By definition and naturality of the reduced external product:

$$\begin{aligned} \Xi^* (\beta_{\mathbf{o}} \boxtimes \beta_{\mathbf{p}}) &= \Xi^* \circ \left( \left( (\overline{\pi_S^n}^{-1} \circ \overline{\kappa_v})^* (\beta_n) \right) \boxtimes \left( (\overline{\pi_S^m}^{-1} \circ \overline{\kappa_w})^* (\beta_m) \right) \right) \\ &= \Xi^* \circ \left( \left( (\overline{\pi_S^n}^{-1} \circ \overline{\kappa_v}) \wedge (\overline{\pi_S^m}^{-1} \circ \overline{\kappa_w}) \right)^* (\beta_n \boxtimes \beta_m) \right) \end{aligned}$$

Now, the function  $\left( (\overline{\pi_S^n}^{-1} \circ \overline{\kappa_v}) \wedge (\overline{\pi_S^m}^{-1} \circ \overline{\kappa_w}) \right) : \overline{V} \rightarrow S^n$  is the compactification of  $(\pi_S^n)^{-1} \circ \kappa_v$ . Hence, by naturality of  $\Xi$  we conclude:

$$\begin{aligned} \Xi^* \circ \left( \left( (\overline{\pi_S^n}^{-1} \circ \overline{\kappa_v}) \wedge (\overline{\pi_S^m}^{-1} \circ \overline{\kappa_w}) \right)^* \right) &= \overline{((\pi_S^n)^{-1} \circ \kappa_v) \times ((\pi_S^m)^{-1} \circ \kappa_w)}^* \circ \Xi^* \\ &= \overline{\kappa_v \times \kappa_w}^* \circ \overline{((\pi_S^n)^{-1} \times (\pi_S^m)^{-1})}^* \circ \Xi^* \end{aligned}$$

Next we will apply Lemma to replace 3.5.29  $(\beta_n \boxtimes \beta_m)$  with  $\Upsilon_{n-m}^*(\beta_{n+m})$  where

$$\Upsilon_{n,m} := (\overline{\pi_S^{n+m}})^{-1} \circ I \circ \Xi^{-1} \circ (\overline{\pi_S^n} \wedge \overline{\pi_S^m}).$$

In the definition of  $\Upsilon$  in Definition 3.5.28 we suppressed the map  $I$  that identifies  $\overline{\mathbb{R}^n \times \mathbb{R}^m}$  with  $\overline{\mathbb{R}^{n+m}}$ . For this proof we will write it out. This leads to

$$\begin{aligned} \Xi^* (\beta_n \boxtimes \beta_m) &= \left( (\overline{\pi_S^{n+m}})^{-1} \circ I \circ \Xi^{-1} \circ (\overline{\pi_S^n} \wedge \overline{\pi_S^m}) \circ \Xi \right)^* (\beta_{n+m}) \\ &= \left( (\overline{\pi_S^{n+m}})^{-1} \circ I \circ (\overline{\pi_S^n} \times \overline{\pi_S^m}) \right)^* (\beta_{n+m}) \end{aligned}$$

Now pulling everything together we get:

$$\begin{aligned} \Xi^* (\beta_{\mathbf{o}} \boxtimes \beta_{\mathbf{p}}) &= \overline{\kappa_v \times \kappa_w}^* \circ \overline{((\pi_S^n)^{-1} \times (\pi_S^m)^{-1})}^* \circ \Xi^* (\beta_n \boxtimes \beta_m) \\ &= \overline{\kappa_v \times \kappa_w}^* \circ \overline{((\pi_S^n)^{-1} \times (\pi_S^m)^{-1})}^* \circ \left( (\overline{\pi_S^{n+m}})^{-1} \circ I \circ (\overline{\pi_S^n} \times \overline{\pi_S^m}) \right)^* (\beta_{n+m}) \\ &= \left( \overline{((\pi_S^{n+m})^{-1})} \circ \underbrace{I \circ \overline{\kappa_v \times \kappa_w}}_{\kappa_{v \oplus w}} \right)^* (\beta_{n+m}) \\ &= \beta_{\mathbf{o} \oplus \mathbf{p}} \end{aligned}$$

□

## 4.3 Local K-Theoric Orientation

lem:orient-MinkowskyGauge

**Lemma 4.3.1** (Minkowsky Gauge). *Let  $V$  be a finite dimensional real vector space with a norm  $\|\cdot\|$  and let  $U \subseteq V$  be open, bounded, convex and contain the origin of  $V$ . Choose  $\varepsilon, r > 0$  such that  $\{v \in V \mid \|v\| < \varepsilon\} \subseteq U \subseteq \{v \in V \mid \|v\| < r\}$  and define the Minkowsky gauge*

$$\mu : V \rightarrow [0, \infty], \quad \mu(v) := \inf\{s > 0 \mid s^{-1}v \in U\}.$$

Then:

- (a)  $\lambda U \subseteq U$  for  $\lambda \in [0, 1]$ , and  $s^{-1}v \in U$  for every  $s > \mu(v)$ .
- (b)  $\mu(\lambda v) = \lambda(\mu(v))$  for all  $\lambda > 0$ .
- (c)  $\frac{\|v\|}{r} \leq \mu(v) \leq \frac{\|v\|}{\varepsilon}$ . In particular  $\mu(v) > 0$  for  $v \neq 0$ .
- (d)  $\mu(v + w) \leq \mu(v) + \mu(w)$ .
- (e)  $|\mu(v) - \mu(w)| \leq \frac{\|v-w\|}{\varepsilon}$ , so  $\mu$  is Lipschitz and hence continuous;
- (f)  $U = \{v \in V \mid \mu(v) < 1\}$  and  $\overline{U} = \{v \in V \mid \mu(v) \leq 1\}$ .

*Proof.* (a) For  $x \in U$  and  $\lambda \in [0, 1]$  we have  $\lambda x = \lambda x + (1 - \lambda)0 \in U$  by convexity and since  $0 \in U$ . Now let  $s > \mu(v)$ . By definition of the infimum there exists a  $s_0 \in (\mu(v), s]$  with  $s_0^{-1}v \in U$  and  $s^{-1}v = \frac{s_0}{s}s_0^{-1}v \in U$  by the first property (since  $\frac{s_0}{s} \in [0, 1]$ ).

(b) Follows directly from the definition if one substitutes  $s \mapsto \lambda s$ .

(c) If  $s^{-1}v \in U \subseteq \{v \in V \mid \|v\| \leq r\}$  then  $\|v\| < sr$ , i.e.  $s > \|v\| r^{-1}$ . After the infimum the reads  $\mu(v) \geq \|v\| r^{-1}$ . Conversely, if  $s > \|v\| \varepsilon^{-1}$  then  $\|s^{-1}v\| < \varepsilon$ , and hence  $s^{-1}v \in U$ . Thereby,  $\mu(v) \leq \|v\| \varepsilon^{-1}$ .

(d) Let  $s > \mu(v)$  and  $t > \mu(w)$ . Then by (a) both  $s^{-1}v$  and  $t^{-1}w$  lie in  $U$ , hence by convexity

$$\frac{v+w}{s+t} = \frac{s}{s+t} \cdot \underbrace{\frac{v}{s}}_{\in U} + \underbrace{\frac{w}{t}}_{\in U} \in U.$$

So  $\mu(v+w) \leq s+t$  and letting  $s$  approach  $\mu(v)$  and  $t$  approach  $\mu(w)$  gives the claim.

(e) By (b) and (c), for all  $v, w \in V$  we have:  $\mu(v) \leq \mu(w) + \mu(v-w) \leq \mu(w) + \|v-w\| \varepsilon^{-1}$ . Hence,  $\mu(v) - \mu(w) \leq \|v-w\| \varepsilon^{-1}$ . Interchangeing  $v$  and  $w$  gives the symmetric statement for  $-(\mu(v) - \mu(w)) = \mu(w) - \mu(v) \leq \|v-w\| \varepsilon^{-1}$  proving the statement.

(f) If  $v \in U$ , then  $s^{-1}v \in U$  for some  $s < 1$ , because  $U$  is open and  $s \mapsto s^{-1}v$  is continuous at  $s = 1$ . Hence  $\mu(v) < 1$ . The other inclusion follows directly from (a): if  $\mu(v) < 1$  then  $1^{-1}(v) \in U$ . Thus  $U = \{v \in V \mid \mu(v) < 1\}$ . Now, by continuity of  $\mu$  the set  $\{v \in V \mid \mu(v) \leq 1\}$  is closed and contains  $U$  and thereby also contains  $\overline{U}$ . conversely if  $\mu(v) \leq 1$  then  $\mu(tv) = t\mu(v)$  for  $t \in (0, 1)$  by (b), so  $tv \in U$  and if  $t$  approaches one,  $tv$  approaches  $v$ . Hence  $v$  is in the closure of  $U$ .  $\square$

lem:orient-ad-  
missableNeigh-  
bourhoodsGive  
CompactPairs

**Lemma 4.3.2.** *Let  $V$  be a finite dimensional real vector space and let  $U \subseteq V$  be open, bounded, convex and non empty. Then  $\overline{V} \setminus U$  is compact and homeomorphic to the closure  $\overline{U} \subseteq V$ . In particular  $\overline{V} \setminus U$  is contractible and contains  $\infty$ .*

*Proof.* The compactness follows from  $U$  being open and  $\overline{V}$  being compact. After a translation we may assume that  $U$  is open, bounded convex neighbourhood of the origin and hence we can define the Minkowsky gauge with respect to  $U$  and after fixing a norm, we may use all properties from Lemma 4.3.1 (and denote them with just (a)-(f)). Now we define

$$\iota: \overline{V} \setminus U \rightarrow \overline{U}, \quad \iota(v) := \frac{v}{\mu(v)^2} (v \neq \infty), \quad \iota(\infty) := 0.$$

The well-definition as a map follows from  $\mu(v) \geq 1$  if  $v \in V \setminus U$  and thereby  $\mu(\iota(v)) = \mu(v)\mu(v)^2 = \mu(v)^{-1} \leq 1$  which concludes the well definition by (f). The map  $w \mapsto w\mu(w)^{-2}$  ( $0 \mapsto \infty$ ) is a two sided inverse of  $\iota$  which can be seen by computation and realizing that  $\iota$  maps  $\infty$  to zero (lem:orient-MinkowskyGauge (c)). Finally, continuity of  $\iota$  in  $V \setminus U$  follows from  $\mu$  being continuous and in  $\infty$  the continuity follows from (c):

$$\|\iota(v)\| = \frac{\|v\|}{\mu(v)^2} \xrightarrow{\|v\| \rightarrow \infty} 0.$$

The same argument gives us continuity of our inverse function. Alternatively we can argue that  $\iota$  is a continuous bijection from a compact space to a Hausdorff space and thereby a homeomorphism. Finally,  $\overline{U}$  is convex and compact and thereby contractible.  $\square$

**Definition 4.3.3** (Admissible Neighbourhoods). Let  $V$  be a real finite dimensional vector space. For  $x \in V$  let

$$\mathcal{U}(x) := \{U \subset V \mid U \text{ open, bounded, convex neighbourhood of } x\},$$

and for  $U \in \mathcal{U}(x)$  put  $L(U) := \overline{V} \setminus U$ . By Lemma 4.3.2  $(\overline{V}, L(U))$  is a compact pair with  $L(U)$  contractible. Furthermore,  $(\overline{V}, L(U))$  is an NDR pair. To see this, note that in terms of the gauge  $\mu$  of  $U$  centered at  $x$ , the space subspace  $L(U)$  consists of the points of "radius" greater than or equal to one, together with  $\infty$ . A radial push towards  $L(U)$ , damped across a color so as to match the identity on  $\{v \in V \mid \mu(v) \leq 0.5\}$  and on  $\{\infty\}$ , gives the required homotopy. The function  $u$  from the definition of an NDR pair is adapted to said damping. Notice, how  $\mathcal{U}$  is a directed system with respect to inclusion.

def:orient-lo-  
calKtheoretic  
Orientation

**Definition 4.3.4** (Local K-theoretic orientation). For  $U \in \mathcal{U}(x)$  we call  $K^{-n}(\overline{V}, L(U))$  the local group of  $V$  at  $x$  of size  $U$ . A **local K-theoretic orientation of  $V$  at  $x$**  is a generator of this group.

Next we want to make sure that the naming of said generator is deserved, i.e. the generator is induced and detects orientations, and that the size  $U$  is irrelevant.

lem:orient-comparisonOfSizes

**Lemma 4.3.5** (Comparison of Sizes). *Let  $U \subseteq U'$  in  $\mathcal{U}(x)$ . Then  $\text{id}_{\bar{V}}$  gives a map of pairs  $(\bar{V}, L(U')) \rightarrow (\bar{V}, L(U))$  and the induced map  $\lambda_{U'U} : K^{-n}(\bar{V}, L(U)) \rightarrow K^{-n}(\bar{V}, L(U'))$  is an isomorphism. Furthermore let  $j$  and  $j'$  denote the inclusion of the pair  $(\bar{V}, \emptyset)$  into  $(\bar{V}, L(U))$  and  $(\bar{V}, L(U'))$ . Then, the triangle*

$$\begin{array}{ccc} K^{-n}(\bar{V}, L(U)) & \xrightarrow{\lambda_{U'U}} & K^{-n}(\bar{V}, L(U')) \\ & \searrow j^* & \swarrow j'^* \\ & \tilde{K}^{-n}(\bar{V}) & \end{array}$$

*commutes, and all maps are isomorphisms and all groups are isomorphic to  $\mathbb{Z}$ .*

*Proof.* The commutation follows from the topological commutation of the maps that induce the diagram. The maps  $j$  and  $j^*$  induce isomorphisms by Lemma 3.2.9, as they are induced by collapsing contractible spaces where contractibility follows from Lemma 4.3.2. Hence  $\lambda_{U'U}$  is also an isomorphism. Finally  $K^{-\dim(V)}(\bar{V}) \cong \mathbb{Z}$  gets a generator as a K-theoretic orientation.  $\square$

**Definition 4.3.6.** For  $x \in V$  and  $U \in \mathcal{U}(x)$  define the isomorphism

$$\Theta_{x,U} : \tilde{K}^{-n}(\bar{V}) \rightarrow K^{-n}(\bar{V}, L(U))$$

to be the inverse of the map induced by the pair inclusion  $(\bar{V}, \emptyset) \rightarrow (\bar{V}, L(U))$ .

**Proposition 4.3.7** (A K-theoretic Orientation Induces a Local Orientation). Let  $\beta$  be a K-theoretic orientation of  $V$ , i.e. a generator of  $\tilde{K}^{-n}(\bar{V})$ . Then for every  $x \in V$  and every  $U \in \mathcal{U}(x)$  the element  $\Theta_{x,U}(\beta)$  is a local K-theoretic orientation of  $V$  at  $x$  and these are compatible:  $\lambda_{U'U} \Theta_{x,U}(\beta) = \Theta_{x,U'}(\beta)$  for  $U' \subseteq U$ .

*Proof.* All statements follow from Lemma 4.3.5, as  $\Theta_{x,U}$  is an isomorphism and thereby maps generators to generators and the compatibility statement follows from the commutative triangle.  $\square$

**Corollary 4.3.8.** An orientation of  $V$  (i.e. an equivalence class of ordered bases) induce via Corollary 4.2.5 a K-theoretic orientation which in turn induces a local K-Theoretic orientation at every  $x \in V$ .

prop:orient-localKOrientationIsNatural

**Proposition 4.3.9** (Naturality). Let  $A : V \rightarrow W$  be an isomorphism of  $n$ -dimensional real vector spaces,  $x \in V$  and  $U \in \mathcal{U}(x)$ . Then  $A(U) \in \mathcal{U}(Ax)$ , the extension  $\bar{A} : \bar{V} \rightarrow \bar{W}$  is a homeomorphism with  $\bar{A}(LU) = L(AU)$ , and the square

$$\begin{array}{ccc} \tilde{K}^{-n}(\bar{W}) & \xrightarrow{\bar{A}^*} & \tilde{K}^{-n}(\bar{V}) \\ \downarrow \Theta_{Ax,AU} & & \downarrow \Theta_{x,U} \\ K^{-n}(\bar{W}, L(AU)) & \xrightarrow{\bar{A}^*} & K^{-n}(\bar{V}, L(U)) \end{array}$$

commutes. The same holds for affine isomorphisms.

*Proof.* A linear isomorphism carries open bounded convex sets to open bounded convex sets, so  $A(U) \in \mathcal{U}(AX)$ .  $A$  is proper, so  $\bar{A}$  is a homeomorphism that maps  $L(U)$  onto  $L(AU)$ . If we replace the maps  $\Theta$  with the inverses, the corresponding square commutes, as its inducing topological square commutes, and thereby our square also commutes.  $\square$

**Corollary 4.3.10.** For  $A \in \text{GL}(V)$ ,  $x \in V$  and  $U \in \mathcal{U}(x)$  the induced map  $\bar{A}^* : K^{-n}(\bar{V}, L(AU)) \rightarrow K^{-n}(\bar{V}, L(Q))$  equals  $\text{sign det}(A)$  after the canonical identification  $\Theta_{x,U} \circ \Theta_{Ax,AU}^{-1}$ .

*Proof.* This follows directly from Propositions 4.2.4 and 4.3.9.  $\square$

#### 4.4 Orientations Calculated by a Jacobean Sign

We now write  $B_r := \{v \in \mathbb{R}^n \mid \|v\| < r\}$  and  $D_r := \bar{B}_r$  for the euclidean balls around 0, and  $L_r := \bar{R}^n \setminus B_r = L(B_r)$ .

lem:orient-Li  
nearisation

**Lemma 4.4.1** (Linearisation). *Let  $\Omega \subseteq \mathbb{R}^n$  be open with  $0 \in \Omega$  and let  $g : \Omega \rightarrow \mathbb{R}^n$  be  $C^1$  with*

$$g(0) = 0, \quad g(w) \neq 0 \text{ for } w \in \omega \setminus \{0\}, \quad A := dg_0 \in \text{GL}_n(\mathbb{R}).$$

*Choose  $\delta > 0$  with  $D_\delta \in \Omega$  and let  $\rho \in (0, \delta)$ . Then there are  $c > 0$  and  $R > c$  such that both  $g|_{D_\delta}$  and  $A|_{D_\delta}$  are maps of pairs*

$$(D_\delta, D_\delta \setminus B_\rho) \rightarrow (D_R, D_R \setminus B_c)$$

*are homotopic as such. In particular  $(g|_{D_\delta})^* = (A|_{D_\delta})^*$  on  $K^{-n}(D_R, D_R \setminus B_c)$ . Moreover  $c$  may be taken arbitrarily small and  $R$  arbitrarily large.*

*Proof.* First we define the map

$$M : D_\delta \rightarrow \text{Mat}_n(\mathbb{R}), \quad M(w) := \int_0^1 dg|_{s \cdot w} ds.$$

Then  $M$  is continuous,  $M(0) = A$  and

$$g(w) = g(w) - g(0) = \int_0^1 \frac{d}{ds}(g(s \cdot w)) ds = M(w)w$$

With this we define the mapping

$$H : D_\delta \times [0, 1] \rightarrow \mathbb{R}^n, \quad H(v, t) := M(tv)v,$$

which is continuous, with  $H(\cdot, 1) = g|_{D_\delta}$  and  $H(\cdot, 0) = A|_{D_\delta}$ . For  $v \neq 0$  and  $t \in (0, 1]$  we can calculate

$$H(v, t) = \frac{M(tv)(tv)}{t} = \frac{g(tv)}{t} \neq 0$$

because  $g$  only vanishes at the origin. At  $t = 0$  we have  $H(v, 0) = Av \neq 0$ . Hence, the set  $\{H(v, t) \mid \rho \leq \|v\| \leq \delta, t \in [0, 1]\}$  is compact and does not contain 0. Thereby the number

$$c_0 := \min\{\|H(v, t)\| \mid \rho \leq \|v\| \leq \delta, t \in [0, 1]\}$$

is greater than zero. Now let  $c \in (0, c_0]$ . Since  $H(D_\delta \times [0, 1])$  is bounded we may choose  $R$  big enough such that  $H(D_\delta \times [0, 1]) \subseteq B_R$ . With those constants we can conclude that  $H(v, t) \in D_R$  for all  $(v, t)$  and if  $\|v\| \geq \rho$  we have  $\|H(v, t)\| \geq c_0 \geq c$  meaning that  $H(v, t) \in D_r \setminus B_c$ . Thereby we have our homotopy of pairs.  $\square$

prop:orient-1  
ocalIndex

**Proposition 4.4.2** (Local Index). Let  $n \geq 1$  and let  $\Psi : \overline{\mathbb{R}^n} \rightarrow \overline{\mathbb{R}^n}$  be continuous with

1.  $\Psi^{-1}(0) = \{0\}$ .
2. There is an open  $\Omega \subseteq \mathbb{R}^n$  with  $0 \in \Omega$ ,  $\Psi(\Omega) \in \mathbb{R}^n$  and  $\Psi|_\Omega$  is of class  $C^1$ .
3.  $A := d(\Psi|_\omega)_0$  is invertible.

Then

$$\Psi^* = \text{sign det}(A) \cdot \text{id}, \quad \text{on } \tilde{K}^{-n}(\overline{\mathbb{R}^n}).$$

*Proof.* Choose  $\delta > 0$  such that  $D_\delta \subseteq \Omega$  and fix  $\rho \in (0, \delta)$ . Choose  $c, R$  as in Lemma 4.4.1 applied to  $g|_\omega$  and assume  $c$  to be small enough such that

$$\Psi(\overline{R^n} \setminus B_\rho) \cap B_c = \emptyset, \quad (26)$$

which is possible, as the left side of the intersection doesn't contain the origin and is closed. Hence, there exist a small ball around the origin that isn't contained. Furthermore, define  $\sigma := \min\{\|Au\| \mid \|u\| = 1\} > 0$ . Now assume  $c \leq \sigma \cdot \rho$ . Then we get two maps of pairs

$$\begin{aligned} \Psi : (\overline{R^n}, L_\rho) &\rightarrow (\overline{R^n}, L_c) \\ \overline{A} : (\overline{R^n}, L_\rho) &\rightarrow (\overline{R^n}, L_c) \end{aligned}$$

For the well definition of the second map assume  $\|v\| \geq \rho$ . Then:

$$\|Av\| \geq \sigma \|v\| \geq \sigma \rho \geq c.$$

Now we apply Lemma 4.3.2 to get the two isomorphisms induced by inclusions:

$$j_\rho^* : K^{-n}(\overline{\mathbb{R}^n}, L_\rho) \rightarrow K^{-n}(D_\delta, D_\delta \setminus B_\rho), \quad j_c^* : K^{-n}(\overline{\mathbb{R}^n}, L_c) \rightarrow K^{-n}(D_\delta, D_\delta \setminus B_c).$$

Because  $\Psi(D_\delta) \subseteq B_R$  and  $\overline{A}(D_\delta) \subseteq B_R$  (by choice of  $R$ ) we get the commuting square

$$\begin{array}{ccc} K^{-n}(\overline{R^n}, L_c) & \xrightarrow{F^*} & K^{-n}(\overline{R^n}, L_\rho) \\ \downarrow j_c^* & & \downarrow j_\rho^* \\ K^{-n}(D_R, D_R \setminus B_c) & \xrightarrow{F|_{D_\delta}^*} & K^{-n}(D_\delta, D_\delta \setminus B_\rho) \end{array}$$

for both  $F = \Psi$  and  $F = \bar{A}$ . (Commutativity follows from the commutativity of the topological square that induces). Now we can apply Lemma 4.4.1 to deduce that both maps in the bottom row agree. As the vertical maps are isomorphisms we can deduce, that also the top two maps agree, namely  $\Psi^* = \bar{A}^*$ . Finally by lemma 3.3.23 we now that  $L_\rho$  and  $L_c$  are contractible and contain  $\infty$ . Hence the collaps  $c_i : \bar{R}^n \rightarrow \bar{\mathbb{R}}^n / L_i$  for  $i = \rho, c$  induces isomorphism that become the vertical maps in the commutative square:

$$\begin{array}{ccc} K^{-n}(\bar{R}^n, L_c) & \xrightarrow{F^*} & K^{-n}(\bar{R}^n, L_\rho) \\ \downarrow c_c^* & & \downarrow c_\rho^* \\ \tilde{K}^{-n}(\bar{R}^n) & \xrightarrow{F^*} & \tilde{K}^{-n}(\bar{R}^n) \end{array}$$

Again for  $F = \Psi$  and  $F = \bar{A}$ . Hence, again we can conclude that  $\Psi^* = \bar{A}^*$  now as maps in  $\tilde{K}^{-n}(\bar{\mathbb{R}}^n)$ , where Proposition 4.2.4 tells us that  $\bar{A}^* = \text{sign det}(A) \cdot \text{id}$ .  $\square$

conv:orient-orientationOfTheSphere

**Remark 4.4.3** (Orientation convention). Let  $n \geq 1$ . We orient  $S^n$  such that the stereographic projection  $\pi_S$  from Definition 3.3.21 is orientation preserving.

lem:orient-rotationsPreserveOrientation

**Lemma 4.4.4.** *Every  $C \in \text{SO}(n+1)$  acts on  $S^n$  as an orientation preserving diffeomorphism, and  $\pi_S \circ C$  is an orientation preserving chart on  $C^{-1}(S^n \setminus \{S\})$ .*

*Proof.* Choose a path  $\gamma : [0, 1] \rightarrow \text{SO}(n+1)$  with  $\gamma(0) = \text{id}$  and  $\gamma(1) = C$ , which exists since  $\text{SO}(n+1)$  is path-connected. Fix  $x \in S^n$ . The map  $t \mapsto \text{sign det}(\text{d}\gamma(t)|_x)$  takes values in  $\{-1, 1\}$  and is continuous, because in charts it is the sign of the determinant of a matrix depending continuously on  $t$ , which never vanishes. Hence it is constant, and at  $t = 0$  it equals  $+1$ . The second claim follows from the chain rule together with Corollary 4.1.19.  $\square$

lem:orient-singleRegularPreimage

**Lemma 4.4.5** (Single Regular Preimage). *Let  $n \geq 1$  and let  $\xi : S^n \rightarrow S^n$  be continuous. Suppose that there are  $y, z \in S^n$  with*

$$\xi^{-1}(z) = \{y\},$$

*such that  $\xi$  is  $C^1$  near  $y$  and  $\text{d}\xi|_y$  is invertible. Then*

$$\xi^* = \text{sign det}(\text{d}\xi_y) \cdot \text{id}, \quad \text{on } \tilde{K}^{-n}(S^n).$$

*Proof.* Let  $S$  be the south pole,  $N$  be the north pole and

$$\pi_S : S^n \rightarrow \bar{\mathbb{R}}^n$$

be the compactifications of  $\pi_S$  to the one-point-compactifications, meaning  $\overline{\pi_S}(S) = \infty$ . Remember that  $\pi_S(N) = 0$  is. Now choose two rotations  $B, C \in \text{SO}(n+1)$  such that  $B(z) = N$  and  $C(y) = N$ . Then define the two homeomorphisms

$$\overline{\varphi} := \overline{\pi_S} \circ C, \quad \overline{\psi} := \overline{\pi_S} \circ B$$

that restrict charts over  $S^n \setminus C^{-1}(S)$  respectively over  $S^n \setminus B^{-1}(S)$ . Those charts are orientation preserving by Lemma 4.4.4. Now define

$$\Psi : \overline{\psi} \circ \xi \circ \overline{\varphi}^{-1} : \overline{\mathbb{R}^n} \rightarrow \overline{\mathbb{R}^n},$$

Then  $\Psi$  satisfies the hypotheses of Proposition 4.4.2:

1.  $\Psi(w) = 0$  if and only if  $\xi \circ \overline{\varphi}^{-1}(w) = z$  if and only if  $\overline{\varphi}^{-1}(w) = y$  which is only the case for  $w = 0$ .
2. Near 0 all maps that compose to  $\Psi$  are diffeomorphisms and hence  $\Psi$  is a diffeomorphism near 0.
3. By constructioun  $d\Psi|_0$  is the coordinate representative of  $d\xi|_y$  in the smooth charts  $\varphi$  and  $\psi$ . Invertibility of the latter in the hypothesus exactly states the invertibility of the coordinate representation.

Hence, by Proposition 4.4.2 we have that  $\psi^* = \text{sign det}(d\Psi|_0) \cdot \text{id}$  on  $\tilde{K}^{-n}(\overline{\mathbb{R}^n})$ . Finally, since  $\varphi$  and  $\psi$  are orientation preserving, this sign equals  $\text{sign det}(d\xi_y)$  by Lemma 4.1.17. Finally, since  $\text{SO}(n+1)$  is path connected there is a path  $\gamma : [0, 1] \rightarrow \text{SO}(n+1)$  with  $\gamma(0) = C$  and  $\gamma(1) = B$ . Then the map

$$S^n \times [0, 1] \rightarrow \overline{\mathbb{R}^n}, \quad (x, t) \mapsto \overline{\pi_S}(\gamma(t)x)$$

is a homotopy from  $\overline{\varphi}$  to  $\overline{\psi}$  givin us that  $\overline{\varphi}^* = \overline{\psi}^*$  and both maps are isomorphisms. Hence:

$$\xi^* = (\overline{\varphi}^*) \circ \Psi^* \circ (\overline{\psi}^*)^{-1} = \text{sign det}(d\xi_y) \circ \underbrace{(\overline{\varphi}^*) \circ (\overline{\psi}^*)^{-1}}_{=\text{id}}.$$

□

## 5 Morse Theory

For everything concerning Morse theory our main reference will be the wonderful book [BH04]. In Morse theory, we will always deal with a smooth function  $f : M \rightarrow \mathbb{R}$  from a smooth Riemannian manifold  $(M, g)$  that is also compact. We will denote its dimension by  $m = \dim(M)$ .

**Definition 5.0.1** (Critical Points). A point  $p \in M$  at which the smooth map  $df_p = 0$  is called **critical**. Equivalently, a point is critical, if it is a critical point of the vectorfield  $\text{grad}(f)$ . By  $\text{Crit}(f)$  we denote the **set of critical points**.

def:morse-the  
Hessian

**Definition 5.0.2** (The Hessian). Let  $f : M \rightarrow \mathbb{R}$  be a smooth function. The **Hessian** of  $f$  at  $p \in \text{Crit}(f)$  is a symmetric bilinear form

$$\text{Hess}_p(f) : T_p M \times T_p M \rightarrow \mathbb{R}$$

that is defined as follows. Let  $\xi_p, \zeta_p \in T_p M$  be two tangent vectors. Now choose two vector fields  $\Xi$  and  $Z$  in a neighbourhood of  $p$  such that  $\Xi(p) = \xi_p$  and  $Z(p) = \zeta_p$ . We then define the Hessian to be:

$$\text{Hess}_p(f)(\xi_p, \zeta_p) = \Xi(Z(f))(p)$$

See 2.0.46 for the notation of applying a function to a vector field. In corollary 5.0.5 we will give a local description telling us that this definition is well defined, meaning independent of the extensions  $\Xi$  and  $Z$ .

**Definition 5.0.3.** A smooth function  $f : M \rightarrow \mathbb{R}$  is called **Morse**, if for any critical point, the Hessian is a symmetric bilinear and non-degenerate form.

For such a form there exists the following theorem:

thm:morse-syl  
vestersLawOfI  
nertia

**Theorem 5.0.4** (Sylvester's Law of Inertia). *Let  $A \in \text{Mat}(n, \mathbb{R})$  be symmetric and let  $T, T' \in \text{GL}(n, \mathbb{R})$  and  $k, k', l, l' \in \mathbb{N}$  such that:*

$$T^\top \circ A \circ T = \begin{pmatrix} \mathbb{1}_k & 0 & 0 \\ 0 & -\mathbb{1}_l & 0 \\ 0 & 0 & 0_{n-k-l} \end{pmatrix} \text{ and } T'^\top \circ A \circ T' = \begin{pmatrix} \mathbb{1}_{k'} & 0 & 0 \\ 0 & -\mathbb{1}_{l'} & 0 \\ 0 & 0 & 0_{n-k'-l'} \end{pmatrix}$$

*then  $k = k'$ ,  $l = l'$  and  $\text{rank}(A) = k + l$ .*

*Proof.* Since  $T, T'$  are invertible we have that

$$k + l = \text{rank}(T^\top \circ A \circ T) = \text{rank}(A) = \text{rank}(T'^\top \circ A \circ T') = k' + l'. \quad (27)$$

Hence it suffices to show that  $k = k'$ . Which we will do by proving the claim:

$$k = \max \{ \dim(U) \mid U \subseteq \mathbb{R}^n \text{ subspace such that } x^\top A x > 0 \ \forall x \in U \setminus \{0\} \}$$

So we start by showing “ $\leq$ ”: Denote the first  $k$  columns of  $T$  with  $x_1, \dots, x_k$ . They form a basis of the subspace  $\mathbb{R}^k \subseteq \mathbb{R}^n$  and with  $0 \neq x = \sum_{i=1}^k \lambda_i x_i$  we have that by bilinearity

$$x^\top A x = \sum_{i=1}^k \lambda_i x_i^\top A x = \sum_{i,j=1}^k \lambda_i \lambda_j x_i^\top A x_j = \sum_{i=1}^k (\lambda_i)^2 \geq 0. \quad (28)$$

This concludes the first inequality.

Now let  $U$  be any  $k$ -dimensional subspace such that for all non zero  $x \in U$ ,  $x^\top A x > 0$ . By a calculation analogous to the one above we have that for any  $x \in W := \text{span}(x_{k+1}, \dots, x_n)$  the number  $x^\top A x$  is less than or equal to zero. Hence,  $W \cap U = \{0\}$  and with this we conclude:

$$\dim(U) = \dim(U + W) - \dim(W) + \dim(U \cap W) \leq (k + (n - k)) - (n - k) + 0 = k. \quad (29)$$

This completes the proof.  $\square$

cor:morse-loc  
alHessian

**Corollary 5.0.5.** Let  $\varphi : U \rightarrow V$  be a chart around a critical point  $p \in M$ . Then in said chart the matrix corresponding to the Hessian has the form:

$$M_p(f) = \left( \frac{\partial^2 (f \circ \varphi^{-1})}{\partial x_i \partial x_j} \right)_{i,j}$$

*Proof.* This follows by direct calculation of  $\text{Hess}_p(f)(\partial_i(p), \partial_j(p))$ . Denote  $x := \varphi(p)$ , then:

$$\begin{aligned} (\partial_i(p) (\partial_j(f)))_p &= \left( \partial_i(p) \left( p \mapsto \frac{\partial f \circ \varphi^{-1}}{\partial x_j} \Big|_{\varphi(p)} \right) \right)_p \\ &= \frac{\partial}{\partial x_i} \left( y \mapsto \frac{\partial f \circ \varphi^{-1}}{\partial x_j} \Big|_y \right) \Big|_x \\ &= \frac{\partial^2 (f \circ \varphi^{-1})}{\partial x_i \partial x_j}. \end{aligned}$$

$\square$

**Remark 5.0.6.** If we have two different charts, then the matrix  $M_p(f)$  of the Hessian transforms from one basis to the other via conjugation of the differential of the transition function  $\varphi_{12} := \varphi_1 \circ \varphi_2^{-1}$ :

$$M_p^1(f) = (d\varphi_{12})^\top \cdot M_p^2(f) \cdot d\varphi_{12}$$

Hence we can define the following:

def:morse-mor  
seIndex

**Definition 5.0.7.** By Sylvester’s Law of inertia, the number  $l$  that denotes the dimension of the maximal subspace on which a matrix is negative definite, is an invariant under conjugation. We call that number the index of said matrix. Now in the setting of Morse theory we assign to each critical point  $p$  the index of  $M_p(f)$  and call that the **(Morse-)index** of  $p$ , denoted by  $\lambda_p$ . If omitting the function would cause ambiguity, we write  $\lambda_p^f$ . We denote the set of all critical points of index  $k$  by  $\text{Crit}_k(f)$ .

thm:morse-morseLemma

**Theorem 5.0.8** (Morse Lemma). *Let  $p$  be a critical point of index  $k$  of the Morse function  $f : M \rightarrow \mathbb{R}$ . There exists a centred chart  $\varphi : U \rightarrow \mathbb{R}^m$  around  $p$  such that*

$$(f \circ \varphi^{-1})(x_1, \dots, x_m) = f(p) + \sum_{i=1}^k -x_i^2 + \sum_{i=k+1}^m x_i^2.$$

*Proof.* We refer to Lemma 3.11 in [banyaga2004lectures](#) [BH04] for a proof.  $\square$

def:morse-gradientFlowLine

**Definition 5.0.9.** Let  $\varphi_t : M \rightarrow M$  be the one-parameter group of diffeomorphisms generated by  $-\text{grad}(f)$ , i.e. if  $R_t(1) : \mathbb{R} \rightarrow T\mathbb{R}$  denotes the canonical unit vector field that comes from the identity as the global chart on  $\mathbb{R}$  (compare definition 2.0.45), we have that

$$\begin{aligned} \frac{d}{dt} \varphi_t(x)(R_t(1)) &= -\text{grad}(f)_{(\varphi_t(x))}, \quad \text{and} \\ \varphi_0(x) &= x. \end{aligned}$$

We call that group the **negative gradient flow** of  $f$  with respect to the metric  $g$ . For compact  $M$  we get that the flow  $\varphi_t$  is global. (also we would hit the boundary (of which there is non), or leave every compact set, which is impossible). The integral curve  $\gamma_x : (a, b) \rightarrow M$  given by

$$\gamma_x(t) = \varphi_t(x)$$

is called a **gradient flow line**.

**Proposition 5.0.10.** Let  $(M, g)$  be a smooth compact Riemannian manifold without boundary, let  $f : M \rightarrow \mathbb{R}$  be a Morse function and let  $\varphi_t : M \rightarrow M$  be the flow of  $-\text{grad}(f)$ . Then the limits

$$\lim_{t \rightarrow \pm\infty} \gamma_x(t)$$

exist and are critical points.

*Proof.* We will show that the flow line has a limit as  $t$  tends to  $+\infty$  and that said limit is a critical point. (The statement for  $-\infty$  is proven analogously.) We define the functions:

$$\begin{aligned} h : \mathbb{R} &\rightarrow \mathbb{R}, & t &\mapsto f \circ \gamma_x(t) \\ \psi : \mathbb{R} &\rightarrow \mathbb{R}, & t &\mapsto \|\text{grad}_{\gamma_x(t)}(f)\|^2 \\ G : \mathbb{R} \times M &\rightarrow \mathbb{R}, & (t, x) &\mapsto \|\text{grad}_{\varphi_t(x)}(f)\|^2 \end{aligned}$$

Then we have that  $\lim_{t \rightarrow \infty} h(t)$  exists and is finite, due to its monotonicity (compare Proposition 2.0.60) and boundedness in  $\mathbb{R}$ . Furthermore,

$$-\frac{d}{dt} h = df_{\gamma_x(t)}(\text{grad}_{\gamma_x(t)}(f)) = g(\text{grad}_{\gamma_x(t)}(f), \text{grad}_{\gamma_x(t)}(f)) = \psi,$$

and for any  $t, \tau \in \mathbb{R}$ :

$$\psi(t + \tau) = G(\tau, \varphi_t(x)).$$

Let  $t_n \rightarrow \infty$ . Then  $\gamma_x(t_n)$  has an accumulation point (as  $M$  is compact) and after passing to a subsequence we assume that  $\gamma_x(t_n) \rightarrow y$ . Assume that  $y$  is not critical and hence there exists an  $\varepsilon > 0$  such that  $G(0, y) = \|\text{grad}_y(f)\|^2 \geq \varepsilon$ . Now continuity of  $G$  in  $(0, y)$  gives us a  $\delta > 0$  and a neighbourhood  $W \subseteq M$  of  $y$  such that  $G(s, z) \geq \frac{\varepsilon}{2}$  for all  $(s, z) \in [0, \delta] \times W$ . Now we can calculate for all  $n$  big enough, such that  $\gamma_x(t_n) \in W$ :

$$h(t_n) - h(t_n + \delta) = \int_0^\delta \psi(t_n + \tau) d\tau = \int_0^\delta \underbrace{G(\tau, \gamma_x(t_n))}_{\in [0, \delta] \times W} d\tau \geq \delta \frac{\varepsilon}{2}$$

This contradicts the convergence of  $h$ , if we apply the limit  $n \rightarrow \infty$  to the calculation. By this argument we see, that any accumulation point of  $(\gamma_x(t_n))$  is critical.

We now want to show that the limit  $\lim_{t \rightarrow \infty} \gamma_x(t)$  exists, meaning that all sequences approach the same critical point. For the sake of a contradiction we assume that  $\lim_{t \rightarrow \infty} \gamma_x(t)$  does not exist. However, by the compactness of  $M$  we can assume that there is a sequence  $t_n \rightarrow \infty$  such that  $\gamma_x(t_n) \rightarrow q$  and by the argument above,  $q$  has to be critical. The Morse lemma gives us a chart (and hence a neighbourhood  $U$ ) around  $q$ , where  $q$  is the only critical point.

Since  $\gamma_x(t)$  does not converge to  $q$  for  $t \rightarrow \infty$  we may assume that there exists an open neighbourhood  $V$  around  $q$  with  $\bar{V} \subseteq U$  and another sequence  $t'_n \rightarrow \infty$  such that  $\gamma_x(t'_n) \notin V$ . Assume on the other hand, that  $\gamma_x(t_n) \in V$  for all  $n$  (again, pass to a subsequence, if this is only possible for large  $n$ ). Now passing to subsequences we may assume that  $t_n < t'_n$  for all  $n$ . Then, we define  $\tau_n := \inf\{t \geq t_n \mid \gamma_x(t) \notin V\} \leq t'_n$ . By continuity of  $\gamma_x(t)$  in  $t$  we know that  $\tau_n > t_n$ . This tells us two things: Firstly,  $\tau_n \rightarrow \infty$  and secondly since  $\gamma_x(t) \in V$  for  $t \in [t_n, \tau_n)$  and  $\gamma_x(\tau_n) \notin V$  we know that  $\gamma_x(\tau_n) \in \partial V$ . Since  $\partial V$  is compact,  $\gamma_x(\tau_n)$  has an accumulation point which again is a critical point in  $U \setminus V$  (by the same argument as above) giving us a contradiction, as  $q$  was the only critical point in  $U$ . Thereby,  $\gamma_x(t) \rightarrow q$  as  $t \rightarrow \infty$  where  $q$  is a critical point.  $\square$

Next we will deepen our **local** understanding of the negative gradient. Notice how the Morse lemma is a statement about smooth manifolds, no metric needed nor mentioned. In fact there are many Morse charts and we can find one where the Riemannian metric in  $p$  is diagonal with non-zero entries. To prove such a statement we first need a somewhat obvious lemma:

lem:morse-linearChartChangeAndTangentBasisChange

**Lemma 5.0.11.** Assume that  $\psi : U \rightarrow V$  is a chart and  $L : \mathbb{R}^m \rightarrow \mathbb{R}^m$  a linear function. Denote the local coordinate maps of the tangent bundle in a point  $p$  that come from a basis in said point induced from a chart  $\chi$  by  $q_\chi$ . Then the diagram

$$\begin{array}{ccc} T_p M & \xrightarrow{q_\psi|_p} & \mathbb{R}^m \\ & \searrow q_{L \circ \psi}|_p & \downarrow L \\ & & \mathbb{R}^m \end{array}$$

commutes for all  $p \in U$ .

*Proof.* We show this by showing the inverse: For any  $v \in \mathbb{R}^m$  we have the two  $(q_\psi|_p)^{-1} \circ L^{-1}(v)$  and  $(q_{\psi \circ L}|_p)^{-1}$  and claim that they are the same. Assume that  $L^{-1} = (\lambda_{ij})_{ij}$ ,  $\psi(p) = x_0$ ,  $L(x_0) = \tilde{x}_0$ ,  $(v_1, \dots, v_m) = v \in \mathbb{R}^m$  and denote the basis of  $T_p M$  coming from the chart  $\psi$  by  $(\frac{\partial}{\partial x_i}|_p)_i$  and the one coming from  $L^{-1} \circ \psi$  by  $(\frac{\partial}{\partial \tilde{x}_i}|_p)_i$ . We now calculate for  $f_p \in \mathcal{E}_p M$ :

$$\begin{aligned}
 \left[ (q_{\psi \circ L}|_p)^{-1}(v) \right] (f_p) &= \sum_i v_i \frac{\partial}{\partial \tilde{x}_i} \Big|_p (f_p) \\
 &= \sum_i v_i \frac{\partial}{\partial x_i} \Big|_{\tilde{x}_0} (f \circ \psi^{-1} \circ L^{-1}) \\
 &= \sum_i v_i \sum_j \frac{\partial f \circ \psi^{-1}}{\partial x_j} \Big|_{x_0} \cdot \frac{\partial L_j^{-1}}{\partial x_i} \Big|_{\tilde{x}_0} \\
 &= \sum_{i,j} v_i \frac{\partial}{\partial x_j} \Big|_p (f_p) \cdot \lambda_{ji} \\
 &= \sum_{i,j} v_i q_\psi|_p^{-1}(e_j)(f_p) \cdot \lambda_{ji} \\
 &= \left[ q_\psi|_p^{-1} \left( \sum_{i,j} v_i \lambda_{ji} e^j \right) \right] (f_p) \\
 &= \left[ q_\psi|_p^{-1} (L^{-1}(v)) \right] (f_p)
 \end{aligned}$$

This concludes the proof. □

Now we can prove that a Morse chart exists where the gradient is of a particularly easy form:

thm:morse-simultaneousDiagonalisationOfQuadraticForms

**Theorem 5.0.12** (Simultaneous Diagonalisation of Quadratic Forms). *Let  $p$  be a critical point of a Morse function  $f$  on a manifold  $M$  and let  $g(p)$  be the Riemannian metric on  $T_p M$ . Then there exists a Morse chart  $\psi$  around  $p$  such that the representation of  $g(p)$  in the basis induced by the Morse chart is given by a diagonal matrix  $\text{diag}(\mu_1, \dots, \mu_m)$  with  $\mu_i > 0$  for all  $i$ .*

*Proof.* The quadratic form of  $f \circ \psi^{-1} - f(p)$  in the coordinates of any Morse chart  $\psi$  is  $q_H(v) = v^T H v$ . The quadratic form induced by the Riemannian metric  $g(p)$  is  $q_G(v) = v^T G v$ , where  $v \in \mathbb{R}^m$  are the coordinate vectors with respect to the Morse chart. To be precise this means for  $v, w \in \mathbb{R}^m$ :

$$g \circ (q_\psi|_p \oplus q_\psi|_p)(v, w) = v^T G w \text{ and } f \circ \psi^{-1}(v) = f(p) + v^T H v.$$

We now aim to manipulate  $\psi$  such that  $G$  becomes diagonal: Since  $g(p)$  is positive definite,  $G$  is also positive definite, and  $H$  is symmetric.

Consider the generalised eigenvalue problem  $Hv = \lambda Gv$ . Since  $H$  and  $G$  are real symmetric matrices and  $G$  is positive definite, this problem has  $m$  real eigenvalues  $\lambda_1, \dots, \lambda_m$  and corresponding eigenvectors  $v_1, \dots, v_m$ , which can be chosen to be orthogonal with respect to the bilinear form defined by  $G$ , such that  $v_i^T G v_j = \delta_{ij}$ , since if  $v_i$  and  $v_j$  are different eigenvectors with respect to different eigenvalues, we can calculate:

$$v_j^T H v_i = (H v_j)^T v_i = \lambda_j (G v_j)^T v_i = \lambda_j v_j^T G(v_i) \quad \text{and} \quad v_j^T H v_i = v_j^T \lambda_i G(v_i) = \lambda_i v_j^T G(v_i).$$

Hence we have the equality

$$\lambda_j (G v_j)^T v_i = \lambda_i v_j^T G(v_i) \Leftrightarrow \underbrace{(\lambda_j - \lambda_i)}_{\neq 0} v_j^T G(v_i) = 0$$

Hence all eigenspaces are orthogonal and we can use Gram–Schmidt to make the bases of the eigenspaces orthogonal and, by rescaling, orthonormal.

Let  $L = [v_1 | \dots | v_m]$  be the matrix whose columns are these  $G$ -orthonormal eigenvectors. The linear change of coordinates  $y = Lz$  leads to new coordinates  $z$ . In these new coordinates, the quadratic forms transform as follows:

$$\begin{aligned} q_G(y) &= y^T G y = (Lz)^T G (Lz) = z^T L^T G L z = z^T I z = \sum_{i=1}^m z_i^2 \\ q_H(y) &= y^T H y = (Lz)^T H (Lz) = z^T L^T H L z \end{aligned}$$

Since  $H v_i = \lambda_i G v_i$ , we have  $L^T H L = \text{diag}(\lambda_1, \dots, \lambda_m)$ . The eigenvalues  $\lambda_i$  are real and have the same signature as the eigenvalues of  $H$  ( $l$  negative,  $m-l$  positive). This is true by Sylvester's law of inertia. By a further scaling of the  $v_i$  and a reordering the matrix  $L^T H L$  can be brought into the form  $\text{diag}(-1, \dots, -1, 1, \dots, 1)$ . (by doing this however, the matrix  $L^T G L$  becomes  $L^T G L = \text{diag}(\mu_1, \dots, \mu_m)$  with  $\mu_j > 0 \ \forall j$ ). If  $\psi$  is the Morse chart we started with, then  $L^{-1} \circ \psi$  is the new Morse chart:

$$f \circ (\psi^{-1} \circ L)(y) = f \circ \psi^{-1}(L(y)) = q_H(L(y)) = \sum_{i=1}^l -y_i^2 + \sum_{i=l+1}^m y_i^2$$

Furthermore, we want to inspect the metric  $g|_p : T_p M^2 \rightarrow \mathbb{R}$  with respect to the chart  $L^{-1} \circ \psi$ -induced basis  $\left(\frac{\partial}{\partial x_1}|_p, \dots, \frac{\partial}{\partial x_m}|_p\right)$ . By the lemma 5.0.11

$$g(q_{L^{-1} \circ \psi}|_p^{-1}, q_{L^{-1} \circ \psi}|_p^{-1})(v, w) = g(q_\psi|_p^{-1}, q_\psi|_p^{-1})(Lv, Lw) = (Lv)^T G L w = v^T L^T G L w$$

□

**Definition 5.0.13.** We call a Morse chart where the matrix corresponding to the gradient in the critical point is of diagonal form a **refined Morse chart**.

Using such a chart, we can start looking at the flow itself and try to find easy local formulas around critical points. For this we start by looking at the Jacobian of the gradient in a critical point:

def:morse-jac  
obianOfTheGra  
dient

**Definition 5.0.14** (Jacobian of the Gradient). The gradient is a section into the tangent bundle,  $\text{grad}(f) : M \rightarrow TM$ . Let  $\psi : M \supseteq U \rightarrow V \subseteq \mathbb{R}^m$  be a chart. The coordinate map  $q_\psi$  on  $TU$  assigns to a vector  $v \in T_p M$  (where  $p \in U$  with  $\psi(p) = x$ ) its components with respect to the basis  $\left(\frac{\partial}{\partial x_i}\big|_p\right)$ , i.e.,  $q_\psi(v) = (v_1, \dots, v_m)$  if  $v = \sum_{i=1}^m v_i \frac{\partial}{\partial x_i}\big|_p$ . We define the Jacobian of the gradient with respect to the chart  $\psi$  as the Jacobian matrix of the coordinate representation of the gradient in this chart:

$$J(\text{grad}(f))_\psi(x) := J(q_\psi \circ \text{grad}(f) \circ \psi^{-1})(x) = \left( \frac{\partial(q_\psi \circ \text{grad}(f) \circ \psi^{-1})_i}{\partial x_j}(x) \right)_{ij}.$$

lem:morse-jab  
obianOfTheGra  
dient

**Lemma 5.0.15.** Let  $\psi$  be a coordinate system around a critical point  $p$ . I.e.  $\psi : p \in U \rightarrow V$  with  $\psi(p) = 0$  and let  $\left(\frac{\partial}{\partial x_1}\big|_x, \dots, \frac{\partial}{\partial x_m}\big|_x\right)$  be the induced basis of  $T_x M$ . Then the Jacobian of the gradient reads

$$J(\text{grad}(f))_\psi(0) = \left( \sum_k g^{ki}(0) \frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} \right)_{ij}, \quad (30)$$

where  $g^{ij}$  is defined as in definition 2.0.50.

*Proof.* Let  $q_\psi$  denote the coordinate function  $TU \rightarrow \mathbb{R}^m$  induced from the basis  $\left(\frac{\partial}{\partial x_1}\big|_x, \dots, \frac{\partial}{\partial x_m}\big|_x\right)$ . Then the gradient in  $\psi$  reads:

$$q_\psi \circ \text{grad}(f) = \left( \sum_k g^{k1} \frac{\partial f}{\partial x_k}, \dots, \sum_k g^{km} \frac{\partial f}{\partial x_k} \right)$$

Hence, we can differentiate:

$$\begin{aligned} \frac{\partial}{\partial x} q_\psi \circ \text{grad}(f) \circ \psi^{-1} &= \left( \frac{\partial}{\partial x_j} \sum_k g^{ki} \frac{\partial f \circ \psi^{-1}}{\partial x_k} \right)_{ij} \\ &= \left( \sum_k \left[ \frac{\partial g^{ki}}{\partial x_j} \frac{\partial f \circ \psi^{-1}}{\partial x_k} + g^{ki} \frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} \right] \right)_{ij}. \end{aligned}$$

and now in  $p = \psi^{-1}(0)$  we have

$$\frac{\partial}{\partial x} q_\psi \circ \text{grad}(f) \circ \psi^{-1}\big|_0 = \left( \sum_k \left[ g^{ki}(0) \frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} \right] \right)_{ij}.$$

□

Now that we know how the gradient locally looks, it is reasonable to assume that the flow locally looks like its exponential:

lem:morse-differentialOfTheFlowInCriticalPoint

**Lemma 5.0.16.** *Let  $t \in \mathbb{R}$  be small enough and  $\varphi_t : M \rightarrow M$  be the flow map corresponding to the negative gradient. Assume that  $U$  is the domain of a chart  $\psi$  and  $p \in U$  is a critical point. Then the linear map  $d\varphi_t|_p : T_p M \rightarrow T_{\varphi_t(p)} M$  has a local representation in the coordinates induced by  $\psi$  given by:*

$$\frac{\partial}{\partial x} \psi \circ \varphi_t(\psi^{-1}(x)) = \exp \left( - \left( \sum_k g^{ki}(0) \frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} \right)_{ij} \cdot t \right).$$

*Proof.* For this we consider the function  $(t, x) \mapsto \psi \circ \varphi_t(\psi^{-1}(x)) : \mathbb{R} \times U \rightarrow U$ . For fixed  $x$  we can calculate

$$q_\psi \frac{d}{dt} \varphi_t(\psi^{-1}(x)) = -q_\psi \circ \text{grad}(f)(\varphi_t(\psi^{-1}(x)))$$

and by changing the order of integration we have:

$$\begin{aligned} \frac{\partial}{\partial t} \frac{\partial}{\partial x} \psi \circ \varphi_t(\psi^{-1}(x)) &= \frac{\partial}{\partial x} \frac{\partial}{\partial t} \psi \circ \varphi_t(\psi^{-1}(x)) \\ &= \frac{\partial}{\partial x} q_\psi \frac{d}{dt} \varphi_t(\psi^{-1}(x)) \\ &= -\frac{\partial}{\partial x} q_\psi \circ \text{grad}(f)(\varphi_t(\psi^{-1}(x))) \\ &= -\frac{\partial}{\partial x} q_\psi \circ \text{grad}(f)(\psi^{-1} \circ \psi \circ \varphi_t(\psi^{-1}(x))) \\ &= -\frac{\partial}{\partial x} (q_\psi \circ \text{grad}(f) \circ \psi^{-1}) \frac{\partial}{\partial x} (\psi \circ \varphi_t(\psi^{-1}(x))) \end{aligned}$$

Hence by the theory of linear differential equation with  $\frac{\partial}{\partial x} \psi \circ \varphi_0(\psi^{-1}(x)) = \mathbb{1}_m$  we have:

$$\frac{\partial}{\partial x} \psi \circ \varphi_t(\psi^{-1}(x)) = \exp \left( -\frac{\partial}{\partial x} (q_\psi \circ \text{grad}(f) \circ \psi^{-1}) \cdot t \right).$$

The preceding Lemma 5.0.15 now finishes the proof. □

Hence in a refined Morse chart we have:

rem:morse-jacobianOfTheFlowInRefinedMorsechart

**Remark 5.0.17.** Let  $\psi$  be a refined Morse chart, and hence:

- (a)  $g^{ik}(0) = \delta_{ik} \mu_k$  with  $\mu_k > 0$  for any  $k$ .
- (b)  $\frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} = s_k 2\delta_{jk}$  where  $s_k = -1$  for  $k \leq \lambda_p$  and  $s_k = 1$  else.

Then the Jacobian of the flow in the critical point  $p$  reads:

$$\begin{aligned} \frac{\partial}{\partial x} \psi \circ \varphi_t(\psi^{-1}(x)) &= \exp(-J(\text{grad}(f))(p) \cdot t) \\ &= \exp \left( - \sum_k \underbrace{g^{ki}}_{=\delta_{ik}\mu_k} (0) \underbrace{\frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k}}_{=s_k 2\delta_{jk}} \cdot t \right) \\ &= \text{diag} \left( e^{2\mu_1 t}, \dots, e^{2\mu_{\lambda_p} t}, -e^{2\mu_{\lambda_p+1} t}, \dots, -e^{2\mu_m t} \right) \end{aligned}$$

In a refined Morse chart we have that the (un-)stable manifold embedding:

**Definition 5.0.18** (Stable and Unstable Manifold). Let  $p \in M$  be a critical point of  $f : M \rightarrow \mathbb{R}$ . Then we define:

1. The **stable set** of  $p$  as

$$W(\rightarrow p) := \{x \in M \mid \lim_{t \rightarrow \infty} \varphi_t(x) = p\}.$$

That is the points that flow towards  $p$ .

2. The **unstable set** of  $p$  as

$$W(p \rightarrow) := \{x \in M \mid \lim_{t \rightarrow -\infty} \varphi_t(x) = p\}.$$

That is the points that came from  $p$ .

Those sets will turn out to be manifolds, which explains the names **(un-)stable manifolds**.

thm:morse-stableAndUnstableManifoldTheorem

**Theorem 5.0.19** (Stable and unstable manifold theorem). *Let  $f : M \rightarrow \mathbb{R}$  be a smooth function on a smooth compact Riemannian manifold  $(M, g)$ . Let  $p$  be a critical point of  $f$ . Then the tangent space splits as:*

$$T_p M = T_p^s(M) \oplus T_p^u M$$

*such that the Hessian is positive definite on  $T_p^s M$  and negative definite on  $T_p^u M$ . Moreover, the stable and unstable manifolds are submanifolds and images of smooth embeddings:*

$$\begin{aligned} E_p^s : T_p^s M &\rightarrow W(\rightarrow p) \subset M, \\ E_p^u : T_p^u M &\rightarrow W(p \rightarrow) \subset M. \end{aligned}$$

*Proof.* I will summarise the proof given in Chapter 4.3 [\[banyaga2004lectures\]](#) [\[BH04\]](#). We work in the discrete setting meaning that we take the negative gradient flow  $\varphi_t$  for a fixed positive  $t$  and the limit-behaviour of the flow is captured by  $\lim_{n \rightarrow \infty} \varphi_t^n$  instead of the  $t$ -limit. First, one shows the splitting of spaces locally in a small neighbourhood  $U$  around a critical point  $p$  and then extends the splitting globally. Around  $U$  we already know by remark 5.0.17 that the tangent space  $T_p M$  splits into  $T_p^s M \oplus T_p^u M$  such that the differential of the flow is a contraction around  $T_p^s M$  and an expansion in  $T_p^u M$ . We can now map a small neighbourhood  $U$  around  $p$  into  $T_p M$  using the map  $\Psi : U \rightarrow T_p M$  given by a refined Morse chart and the identification of  $\mathbb{R}^n$  with  $T_p M$  via  $e_i \mapsto \partial_i(p)$ , where the latter uses the same refined Morse chart. Now inspect the image of the stable manifold  $W(\rightarrow p)$  under said map. This looks to be a small perturbation of the subspace  $T_p^s M$  and the local stable manifold theorem supports that intuition:

lem:morse-localUnStableManifoldTheorem

**Lemma 5.0.20** (Local (Un-)stable Manifold Theorem). *We define:*

$$T := d\varphi_t|_p, \quad T_s := T|_{T_p^s M}, \quad T_u := T|_{T_p^u M}.$$

*Let  $\lambda < 1$  such that  $\|T_s\| < \lambda$  and  $\|T_u^{-1}\| < \lambda$ , where  $\|T\| = \sup\{T(x) \mid \|x\| \leq 1\}$  denotes the operator norm. Now for this setting there exists an  $\varepsilon_\lambda$  depending only on  $\lambda$ , and there exists a  $\delta > 0$  for every  $r > 0$  which satisfy: For all  $\theta : T_p^s M \times T_p^u M \rightarrow T_p M$  that satisfies  $\theta - T$  is Lipschitz with Lipschitz constant*

$$\text{Lip}(\theta - T) < \varepsilon, \text{ and } \|\theta(0)\| < \delta.$$

*In this setting,*

$$W_r^s(\theta) := \{x \in T_p M \mid \forall n \geq 0, \theta^n(x) \text{ is defined and } \|\theta^n(x)\| \leq r\}$$

*is the graph of a Lipschitz function  $g : T_p^s M \rightarrow T_p^u M$  with  $\text{Lip}(g) \leq 1$  and:*

1. *If  $\theta$  is  $C^k$  ( $k$ -times continuously differentiable), then so is  $g$ .*
2. *If  $\theta$  is smoothly differentiable,  $\theta(0) = 0$  and  $d\theta|_0 = T$ , then  $g(0) = 0$  and  $dg|_0 = 0$ . Hence,  $T_p W_r^s(\theta) = T_p^s M$ .*

*Proof.* For a proof we refer to Chapter 4.2 [\[banyaga2004lectures\]](#) [\[BH04\]](#). □

We can now define  $\theta := \Psi \circ \varphi \circ \Psi^{-1}$ . The local (un-)stable manifold theorem now tells us that  $W_r^s(\theta)$  is the graph of a smooth function  $g : T_p^s M \rightarrow T_p^u M$ . Furthermore, it is easy to see that in a small neighbourhood of the origin we have the equality of  $W_r^s(\theta)$  and  $\text{Psi}(W(\rightarrow p) \cap U)$ . Hence the projection  $\pi$  of said graph to its domain gives us a chart of the stable manifold around  $p$ . A whole atlas of  $W(\rightarrow p)$  is obtained by  $U^n := \varphi^{-n}(U)$  which covers the whole  $W(\rightarrow p)$  and with this the maps  $\chi^n : \pi \circ \Psi \circ \varphi^n$  give us an atlas. The smooth embeddings are obtained as follows: We define  $h := \chi \circ \varphi \circ \chi^{-1}$  and  $B := W(\rightarrow p) \cap U$ . Now  $d(\varphi_p|_{T_p^s M})$  having an operator norm less than one translates

to  $dh_p$  as they are similar as matrices. Let  $0 < \alpha < 1$  and  $B_0 \subseteq B$  be a ball centred at  $p$  such that  $\|dh_x\| \leq \alpha$  for all  $x \in B_0$ . Then  $h|_{B_0}$  is a contraction, which can be extended to a global contraction  $\tilde{h}: T_p^s M \rightarrow T_p^s M$ . Now we define the embedding:

$$E^s: T_p^s M \rightarrow W(\rightarrow p)$$

$$x \mapsto \varphi^{-n} \circ \chi^{-1} \circ \tilde{h}^n(x) \quad \text{such that } \tilde{h}^n(x) \in B_0.$$

All required properties and the well-definedness of  $E^s$  are now standard calculations. The stable manifold theorem follows from the unstable one by replacing  $f$  with  $-f$ .  $\square$

rem:morse-tan-  
gendspaceOfSt-  
ableManifold

**Remark 5.0.21.** The proof also gives us the equality  $T_p W(\rightarrow p) = T_p^s M$ : near  $p$ , the stable manifold agrees with  $\Psi^{-1}(W_r^s(\theta))$ , and by part 2 of Lemma 5.0.20 (applicable since  $\theta(0) = 0$  and  $d\theta|_0 = T$ ) the set  $W_r^s(\theta)$  is the graph of a function  $g$  with  $g(0) = 0$  and  $dg|_0 = 0$ , hence has tangent space  $T_p^s M$  at the origin.

We want to refine the unstable embedding  $E_q^u: T_q^u M \rightarrow W(q \rightarrow)$ . For this we will make use of the Hartman–Grobman conjugation theorem:

**Theorem 5.0.22** (Hartman-Grobman Theorem). *Let  $E$  be an open subset of  $\mathbb{R}^n$  containing the origin, let  $f \in \mathcal{C}$ , and let  $\varphi_t$  be a flow given by  $f$ . (i.e.  $\frac{\partial}{\partial t}(\varphi_t(x)) = f(x)$ ). Assume that  $f(0) = 0$ , i.e. that 0 is a fixed point of  $\varphi_t$  and furthermore assume that fixed point to be hyperbolic, i.e.  $Df$  has no eigenvalue with zero real part. Then there exists a homeomorphism  $H$  of an open set  $U$  containing the origin onto an open set  $V$  containing the origin such that for each  $x_0 \in U$ , there is an open interval  $I_0 \subseteq \mathbb{R}$  containing zero such that for all  $x_0 \in U$  and  $t \in I_0$  the diagram*

$$\begin{array}{ccc} U & \xrightarrow{H} & V \\ \downarrow \varphi_t & & \downarrow e^{At} \\ U & \xrightarrow{H} & V \end{array}$$

*commutes, meaning that locally the flow is topologically conjugate to its linearisation.*

*Proof.* A proof is given in chapter 2.8 in [\[Perko2001\]](#) [\[Per01\]](#).  $\square$

thm:morse-ref-  
inedUnstableE-  
mbedding

**Theorem 5.0.23** (Refined Stable Embedding). *In the setting of the stable manifold theorem, we can refine the embedding*

$$E_q^u: T_q^u M \rightarrow W(q \rightarrow),$$

such that it resembles the flow, meaning that for  $x = E_p^s(v)$  the flow is given by

$$\varphi_t(x) = E_p^s(A_t v),$$

where  $A_t$  is a linear map defined as:  $A_t = B_{\mathfrak{b}} \circ \text{diag}(e^{-2\mu_1 t}, \dots, e^{-2\mu_{\lambda_p} t}) \circ B_{\mathfrak{b}}^{-1}$ , where  $\mathfrak{b}$  is the basis of  $T_q^u M$  that comes from the basis of  $T_q M$  which is induced by a refined Morse chart. The diagonal matrix is the stable part of the Jacobian of the flow in said refined Morse chart  $\psi$ .

*Proof.* In the proof of the unstable manifold theorem from [\[banyaga2004lectures\]](#) [\[BH04\]](#) which was summarised above, we started with a local map  $\rho' : U \rightarrow V$ , where  $U \subseteq W(q \rightarrow)$  is a neighbourhood of  $q$  and  $V \supseteq T_q^u$ . (This was a refined Morse chart together with a projection once realised that the stable part gets mapped to the graph of a smooth function). Now we define

$$\begin{aligned} \theta_t : V &\rightarrow V \\ v &\mapsto \rho' \circ \varphi_t \circ \rho'^{-1}(v) \end{aligned}$$

for all  $t$  such that the map is well defined. Now with a basis  $\mathfrak{b}_p^s$  of  $T_p^s M$  we can push these considerations into  $\mathbb{R}^n$  and by abuse of notation assume that  $\rho$  maps to  $\mathbb{R}^n$  and hence  $\theta$  is a flow around the origin of  $\mathbb{R}^n$ . Now by the Hartman–Grobman Theorem we find a homeomorphism  $H$  around the origin, such that  $H \circ \theta_t = e^{At} \circ H$ , where  $A = d\theta_t =$ . To calculate said differential we look into the map  $\rho'$  again. For this we give names to the functions:

- $\psi : U \rightarrow V$  is the refined Morse chart around  $q$ .
- $g : T_p^s M \rightarrow T_p^u M$  is the map that defines the image of the stable part under  $\psi$  to be a graph. I.e.  $\psi(W(\rightarrow p)) = \Gamma_g$ . We know that  $g(0) = 0$  and  $dg|_0 = 0$ .
- $\pi : \Gamma_g \rightarrow T_p^s M$  is the projection. Its inverse is obviously  $\text{id} \times g$ .

We know from the local stable manifold theorem 5.0.20, that  $dg|_0 = 0$ . With this we can calculate:

$$\begin{aligned} d\theta_t|_0 &= d(\pi \circ \psi \circ \varphi_t \circ \psi^{-1} \circ \pi^{-1})|_0 \\ &= d\pi|_{(0,0)} \circ d\psi \circ \varphi_t \circ \psi^{-1}|_{0,0} \circ d\text{id} \times g|_0 \\ &= (\text{id}_{m-\lambda_p}, 0_{\lambda_p}) \circ \text{diag}(e^{-2\mu_1 t}, \dots, e^{-2\mu_{\lambda_p} t}, e^{2\mu_{\lambda_q+1} t}, \dots, e^{2\mu_m t}) \circ (\text{id}_{m-\lambda_p}, 0_{\lambda_p})^T \\ &= \text{diag}(e^{-2\mu_1 t}, \dots, e^{-2\mu_{\lambda_p} t}) \end{aligned}$$

Now define for a small neighbourhood  $V \subseteq \mathbb{R}^{m-\lambda_p}$  the map

$$\rho := \rho' \circ H : V \rightarrow U.$$

$U$  is chosen such that the map is onto. Then we have that  $\varphi_t \circ \rho^{-1} = \rho^{-1} \circ e^{At}$ , in that small neighbourhood of  $p$ . Now finally we can extend this property to be true for the whole stable set: For this we define the embedding

$$E_p^s : T_p^s M \rightarrow W(\rightarrow p) \\ x \mapsto \varphi_{-n} \circ \rho^{-1} \circ A_n(x) \quad \text{where } n \in \mathbb{N} \text{ is big enough such that } A_n(x) \in V.$$

This map is in fact well defined since

$$\begin{aligned} \varphi_n \circ \rho^{-1} \circ A_n(x) &= \varphi_{-n} \circ \varphi_{-1} \circ \varphi_1 \circ \rho^{-1} \circ A_n(x) \\ &= \varphi_{-n} \circ \varphi_{-1} \circ \rho^{-1} \circ A_{n+1}(x) \\ &= \varphi_{-(n+1)} \circ \rho^{-1} \circ A_{n+1}(x) \end{aligned}$$

And obviously the flow compatibility that is given locally by  $\rho$  is now true globally: Let  $x \in W(\rightarrow p)$  and assume  $x = E_p^s(y)$ . Then:

$$\begin{aligned} \varphi_t(x) &= \varphi_t(E_p^s(y)) = \varphi_t \circ \varphi_{-n} \circ \rho^{-1} \circ A_n(y) \\ &= \varphi_{-n} \circ \varphi_t \circ \rho^{-1} \circ A_n(y) \\ &= \varphi_{-n} \circ \rho^{-1} \circ A_n(\varphi_t(y)) \end{aligned}$$

The details of the proof are exactly the same as in the standard (un-)stable manifold theorem.  $\square$

**Definition 5.0.24.** We say that a flow line  $\gamma$  with respect to the negative gradient flow of a Morse function  $f$  starts in  $q$ , if  $\lim_{t \rightarrow -\infty} \gamma(t) = q$ , and ends in  $p$ , if  $\lim_{t \rightarrow \infty} \gamma(t) = p$ . We denote the space of flow lines starting in  $q$  by  $\mathcal{M}(q \rightarrow)$ , the ones ending in  $p$  by  $\mathcal{M}(\rightarrow p)$  and the flow lines from  $q$  to  $p$   $\mathcal{M}(q \rightarrow p)$ .

## 5.1 Signed Flow Lines

**Definition 5.1.1** (Morse-Smale). A Morse function  $f : M \rightarrow \mathbb{R}$  such that all stable and unstable manifolds intersect transversally is called a **Morse-Smale** function.

**Definition 5.1.2.** Let  $f : M \rightarrow \mathbb{R}$  be a Morse-Smale function on a Riemannian manifold  $(M, g)$ . Furthermore let  $\mathfrak{o}$  be a choice of orientations for all unstable tangent spaces  $(T_q^u M \text{ for all } q \in \text{Crit}(f))$ . The tuple  $\mathfrak{A} := (g, f, \mathfrak{o})$  is a **Morse datum**.

**Proposition 5.1.3** (Oriented Unstable Manifolds). All unstable manifolds of a manifold with Morse function are orientable which can be deduced from the (un-)stable embedding theorem. Hence, the orientations from a Morse datum  $\mathfrak{A} := (g, f, \mathfrak{o})$  induce orientations in all unstable manifolds by enforcing Lemma 4.1.10.

prop:morse-OrientedUnstableSpaces

lem:morse-tan-  
gendBundleSMo-  
othEmbeddingI-  
sSmooth

**Lemma 5.1.4.** *Let  $M, N$  be smooth manifolds and  $f : N \rightarrow M$  be a smooth embedding. Then the tangent bundle  $T(f(N))$  of the smooth submanifold  $f(N)$  is a smooth subbundle of  $TM|_{f(N)}$ .*

*Proof.* Write  $S := f(N)$ , and let  $n = \dim N$  and  $m = \dim M$ . Since  $f$  is an embedding,  $S$  satisfies the local slice condition: around every  $p \in S$  there is a chart  $\varphi: U \rightarrow \mathbb{R}^m$  of  $M$  with  $\varphi(S \cap U) = \varphi(U) \cap (\mathbb{R}^n \times \{0\})$ , and the restriction  $\varphi|_{S \cap U}: S \cap U \rightarrow \mathbb{R}^n \times \{0\} \cong \mathbb{R}^n$  is a chart for  $S$ . With respect to these two charts, the inclusion  $S \cap U \hookrightarrow U$  has the coordinate representation  $v \mapsto (v, 0)$ , hence for every  $q \in S \cap U$  we have

$$T_q S = \text{span}(\partial_1(q), \dots, \partial_n(q)) \subset T_q M,$$

where  $\partial_1, \dots, \partial_m$  denotes the coordinate frame of  $\varphi$ . Hence the trivialisation of  $TM|_U$  induced by  $\varphi$  maps  $TS|_{S \cap U}$  onto  $(S \cap U) \times (\mathbb{R}^n \times \{0\})$ , i.e.  $TS$  is locally the standard subbundle. Moreover, any transition map between two slice charts sends slices to slices, so its Jacobian along  $S$  is block upper triangular; the transition functions of  $TS$  are the smooth upper-left  $n \times n$  blocks. Thus  $TS$  is a smooth subbundle of  $TM|_S$ .  $\square$

prop:morse-or-  
ientTheQuotie-  
nt

**Proposition 5.1.5.** Let  $\mathfrak{A} := (g, f, \mathfrak{o})$  be a Morse Datum. Then for all critical points  $p$  the quotient bundles  $TM|_{W(\rightarrow p)} / TW(\rightarrow p)$  inherit canonical orientations from .

*Proof.* By lemma 5.1.4 we know that the bundle  $TW(\rightarrow p)$  is a smooth subbundle of  $TM|_{W(\rightarrow p)}$  and by Lemma 2.0.40 we know that the quotient bundle  $TM|_{W(\rightarrow p)} / TW(\rightarrow p)$  is smooth. Finally, we have a natural vectorspace isomorphism

$$\left( TM|_{W(\rightarrow p)} / TW(\rightarrow p) \right)_p = T_p M / T_p W(\rightarrow p) \cong T_p^u M,$$

because  $T_p M \cong T_p^u M \oplus T_p^s M$  and by Remark 5.0.21  $T_p^s M = T_p W(\rightarrow p)$ . This lets us induce an orientation in the quotient bundle over  $p$  and by Lemma 4.1.10 this orientation orients the whole bundle.  $\square$

def:morse-mod-  
uliSpaceOfFlo-  
wLines

**Definition 5.1.6.** For a Morse datum  $(g, f, \mathfrak{o})$  and two critical points  $p, q$  we define the spaces

$$W(p \rightarrow q) := W(p \rightarrow) \cap W(\rightarrow q) \quad \text{and} \quad \mathcal{M}(p \rightarrow q) := W(p \rightarrow q) / \mathbb{R},$$

where the quotient is defined to be the set of orbits of the  $\mathbb{R}$ -action  $t \mapsto \varphi_t(x)$  for all  $x \in W(p \rightarrow q)$ . Hence,  $\mathcal{M}(p \rightarrow q)$  is the set of flow lines and is called the **moduli space of flow lines**.

def:morse-sig-  
nedFlowLines

**Definition 5.1.7** (Signed Flow Lines). Let  $\mathfrak{A} := (g, f, \mathfrak{o})$  be a Morse Datum. Let  $q, p$  be two critical points with  $\text{ind}(q) - 1 = \lambda(p)$ . Then we have for all  $\gamma \in \mathcal{M}(q \rightarrow p)$  the exact sequence of bundles

$$0 \longrightarrow T\gamma \xrightarrow{i} TW(q \rightarrow)|_\gamma \xrightarrow{p} TM/TW(\rightarrow p)|_\gamma \longrightarrow 0$$

where the surjectivity of  $p$  follows from the smale condition. Let  $T\gamma$  be oriented by the section  $x \mapsto -\text{grad}_x(f)$ , and the quotient be oriented as in Propostion 5.1.5. Then the short exact sequence orients  $TW(q \rightarrow)$  from the orientation in  $T_p^u M$  and on the other hand we can orient  $TW(q \rightarrow)$  from  $T_q^u M$  by Proposition 5.1.3. We define the Mapping

$$\varepsilon : \bigcup_{\substack{p, q \in \text{Crit}(f) \\ \text{ind}(q) - 1 = \text{ind}(p)}} \mathcal{M}(q \rightarrow p) \rightarrow \{-1, 1\}$$

$$\gamma \mapsto \begin{cases} +1 & \text{If the two induced orientations agree.} \\ -1 & \text{If the two orientations disagree.} \end{cases}$$

Furthermore, we define the counting of signed flow lines:

$$n : \{(q, p) \in \text{Crit}(f)^2 \mid \text{ind}(q) - 1 = \text{ind}(p)\} \rightarrow \mathbb{Z}$$

$$(q, p) \mapsto \sum_{\gamma \in \mathcal{M}(q \rightarrow p)} \varepsilon(\gamma)$$

def:Morsehomology

**Definition 5.1.8** (Morse Homology). Let  $\mathfrak{A} := (g, f, \mathfrak{o})$  be a Morse datum. With this we define now the **Morse chain groups** for a given commutative ring with one ( $\Lambda$ ):

$$C_k(M, \mathfrak{A}, \Lambda) := \bigoplus_{\text{Crit}(f)} \Lambda$$

The boundary operator is given by counting the flow lines:

$$\partial : C_k(M, \mathfrak{A}, \Lambda) \rightarrow C_{k-1}(M, \mathfrak{A}, \Lambda)$$

$$q \mapsto \sum_{p \in \text{Crit}_{k-1}(f)} n(q, p) \cdot p$$

**Theorem 5.1.9.** *The Morse chain groups form a chain complex, meaning that  $\partial^2 = 0$ .*

*Proof.* This will be a corollary of theorem . □

reference  
zu com-  
muting  
boundary  
homomor-  
phisms  
einfügen

## 5.2 A Morse-Theoretik Künneth Formula

def:DefinitionproduceMorsedatum

**Definition 5.2.1.** Let  $(M, \mathfrak{A}, \Lambda)$  and  $(M', \mathfrak{A}', \Lambda)$  be two Morse data with  $\mathfrak{A} = (g, f, \mathfrak{o})$  and  $\mathfrak{A}' = (g', f', \mathfrak{o}')$ . On  $M \times M'$  we define the Morse function  $f + f'$  and the metric  $g \oplus g'$ . Then,  $d_{(x, x')}(f + f') = d_x(f) + d_{x'}(f')$  and hence the critical points of  $f + f'$  are pairs  $(p, p')$  such that  $p$  is critical for  $f$  and  $p'$  is critical for  $f'$ . By corollary 5.0.5 we conclude that the Hessian splits into block diagonal form and hence the property of non degeneracy follows. Since we use the product metric  $g \oplus g'$  we also get a splitting of the gradient of  $f + f'$ :

$$\begin{aligned} d_{(x, x')}(f + f')(V, V') &= d_x(f)(V) + d_{x'}(f')(V') \\ &= g(\text{grad}_x(f), V) + g'(\text{grad}_{x'}(f'), V') \\ &= (g \oplus g')(\underbrace{(\text{grad}_x(f), \text{grad}_{x'}(f'))}_{\text{grad}_{(x, x')}(f + f')}, (V, V')). \end{aligned}$$

Hence, the flow also splits in the following sense: Assume  $\tilde{\varphi}$  is the flow of  $f + f'$  on  $M \times M'$ ,  $\varphi$  is the flow of  $f$  on  $M$  and  $\varphi'$  is the flow of  $f'$  on  $M'$ . Then  $\varphi_t(x, x') = (\varphi_t(x), \varphi_t(x'))$ , as the right sight satisfies the differential equation. Thereby, the (un-)stable manifolds split, letting us orient all unstable manifolds via that splitting which gives us  $\mathfrak{o} \oplus \mathfrak{o}'$ . Finally, transversal intersection is a property that carries over to products and hence the Morse datum  $(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda)$  with  $\mathfrak{A} \times \mathfrak{A}' := (g \oplus g', f + f', \mathfrak{o} \oplus \mathfrak{o}')$  indeed constitutes a valid Morse datum. The splitting of the flow gives a splitting of the connecting flow lines and hence of the moduli spaces for two critical points  $(q, q')$  and  $(p, p')$ :

$$\mathcal{M}((q, q') \rightarrow (p, p')) = \mathcal{M}(q \rightarrow p) \times \mathcal{M}(q' \rightarrow p')$$

If  $p \neq q$  and  $p' \neq q'$ , then for the moduli space to be non-empty we need

$$\text{ind}(q) > \text{ind}(p) \quad \text{and} \quad \text{ind}(q') > \text{ind}(p').$$

This results in the index of  $(q, q')$  being at least two bigger than the index of  $(p, p')$ . Hence, if we want to count flow lines between two consecutive indices, either  $q$  equals  $p$  or  $q'$  equals  $p'$ . In this case we have the moduli space

$$\mathcal{M}((q, q') \rightarrow (p, p')) = \begin{cases} p \times \mathcal{M}(q' \rightarrow p') & \text{if } p = q \\ \mathcal{M}(q \rightarrow p) \times p' & \text{if } p' = q'. \end{cases}$$

Now, we want to analyse how the splitting affects the counting of flow lines. For this we denote the flow lines  $\tilde{\gamma} \in \mathcal{M}((q, q') \rightarrow (p, p'))$ ,  $\gamma \in \mathcal{M}(q \rightarrow p)$  and  $\gamma' \in \mathcal{M}(q' \rightarrow p')$  such that  $\tilde{\gamma} = \gamma \times \gamma'$ . Then we get a short exact sequence of bundles:

$$0 \longrightarrow T\tilde{\gamma} \longrightarrow TW((q, q') \rightarrow)|_{\tilde{\gamma}} \longrightarrow \left( T(M \times M') / TW(\rightarrow (p, p')) \right)|_{\tilde{\gamma}} \longrightarrow 0$$

This sequence, however, is graded in the sense that all spaces (with the exception of the 0) are direct sums and all maps respect them, meaning it is the direct sum of the two short exact sequences:

$$0 \longrightarrow T\gamma \text{ or } 0 \longrightarrow TW(q \rightarrow)|_{\gamma} \longrightarrow \left( TM / TW(\rightarrow p) \right)|_{\gamma} \longrightarrow 0 \quad (31)$$

{eq:morse-fir  
stExactSequen  
ce}

and

$$0 \longrightarrow T\gamma' \text{ or } 0 \longrightarrow TW(q' \rightarrow)|_{\gamma'} \longrightarrow \left( TM' / TW(\rightarrow p') \right)|_{\gamma'} \longrightarrow 0 \quad (32)$$

{eq:morse-sec  
ondExactSeque  
nce}

Now, we have the two orientations of  $TW((q, q') \rightarrow)|_{\tilde{\gamma}}$ : The first one comes from the orientation of  $T_{(q, q')}^u M \times M'$  and is given as the wedge product of two non-vanishing sections into the determinant bundle from the orientations of  $T_q^u M$  and  $T_{q'}^u M'$  and thereby represented by  $s_q \otimes s_{q'}$  (Remember that the determinant of a direct sum is the tensor product of the determinants; this is also true for bundles. Furthermore, we omit

the needed inclusions or lifts to pair them with a  $\wedge$ .) The other way of orienting comes from the exact sequences. Let  $-\text{grad} \wedge s_p$  denote a representative of the orientation induced by the short exact sequence (31) and similarly denote the orientation induced in the middle of the second short exact sequence (32) by  $-\text{grad}' \wedge s_{p'}$ . By definition, the comparison of  $s_q$  with  $-\text{grad} \wedge s_p$  gives the sign  $\varepsilon(\gamma)$  and similarly the comparison of  $s_{q'}$  with  $-\text{grad}' \wedge s_{p'}$  gives the sign  $\varepsilon(\gamma')$ . For ease of notation we assume that the sections live in the right space.

Now we assume that  $q' = p'$  resulting in the first element of the short exact sequence (32) being trivial and hence giving us the need to compare those two sections:

$$(-\text{grad}) \wedge (s_p \wedge s_{p'}).$$

Hence we need to compare the two orientations:

$$s_q \wedge s_{q'} \quad \text{with} \quad (-\text{grad}) \wedge (s_p \wedge s_{q'}) = (-\text{grad} \wedge s_p) \wedge s_{q'}.$$

It is now clear that the sign we get from comparing can be expressed as  $\varepsilon(\gamma)$ .

If  $q = p$ , the first short exact sequence (31) has a trivial first bundle and hence we need to compare the two orientations:

$$s_q \wedge s_{q'} \quad \text{with} \quad (-\text{grad}') \wedge (s_q \wedge s_{q'}) = (-1)^{\text{ind}_q(f)} (s_q \wedge -\text{grad} \wedge s_{p'}),$$

where the sign comes from moving  $s_q$  to the front. It is now clear that the sign we get from comparing can be expressed as  $(-1)^{\text{ind}_q(f)} \varepsilon(\gamma')$ .

This results for critical points  $(q, q')$  and  $(p, p')$  with consecutive indices in the formula:

$$n((q, q'), (p, p')) = \begin{cases} (-1)^{\text{ind}_q(f)} n(q', p') & \text{if } q = p \\ n(q, p) & \text{if } q' = p' \end{cases}$$

**Remark 5.2.2.** Let  $(M, \mathfrak{A}, \Lambda)$  and  $(M', \mathfrak{A}', \Lambda)$  be two Morse data with  $\mathfrak{A} = (g, f, \mathfrak{o})$  and  $\mathfrak{A}' = (g', f', \mathfrak{o}')$ . Then the map (given on the generators of the pure tensors)

$$\Phi : \bigoplus_{i+j=k} C_i(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C_j(M', \mathfrak{A}', \Lambda) \rightarrow C_k(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda), \quad (p \otimes p') \mapsto (p, p')$$

is an isomorphism of groups. Furthermore, if we denote the boundary morphism of the chain complexes corresponding to  $M, M'$  and  $M \times M'$  by  $\partial_f, \partial_{f'}$  and  $\partial_{f+f'}$  respectively, we obtain the following isomorphism of chain complexes induced by  $\Phi$ :

$$\left( \bigoplus_{i+j=*} C_i(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C_j(M', \mathfrak{A}', \Lambda), \partial_f \otimes 1 + (-1)^i 1 \otimes \partial_{f'} \right) \rightarrow (C_*(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda), \partial_{f+f'})$$

Here, the left side is, by definition, the tensor product of chain complexes. Hence, we get a natural isomorphism of chain complexes:

$$C_*(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C_*(M', \mathfrak{A}', \Lambda) \rightarrow C_*(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda).$$

*Proof.* The isomorphism of groups is easy to see, since we have a bijection of the generators

$$(p \otimes q) \leftrightarrow (p, q).$$

Since both sides are free  $\Lambda$ -modules we can apply the universal property of free modules, giving us a canonical isomorphism of  $\Lambda$ -modules and hence a group isomorphism. Now we want to show that we get an isomorphism of chain complexes and hence need to check if the boundary map commutes with  $\Phi$ . Let  $q$  be a critical point of index  $i$  of  $f$  and let  $q'$  be a critical point of index  $j$  of  $f'$ . On the one hand, we can calculate

$$\begin{aligned} \Phi(\partial_f \otimes 1 + (-1)^i 1 \otimes \partial_{f'})(q \otimes q') &= \Phi(\partial_f(q) \otimes q' + (-1)^i q \otimes \partial_{f'}(q')) \\ &= \Phi\left(\sum_{p \in \text{Crit}_{i-1}(f)} n(q, p) p \otimes q' + (-1)^i q \otimes \sum_{p' \in \text{Crit}_{j-1}(f')} n(q', p') p'\right) \\ &= \sum_{p \in \text{Crit}_{i-1}(f)} n(q, p)(p, q') + \sum_{p' \in \text{Crit}_{j-1}(f')} (-1)^i n(q', p')(q, p'), \end{aligned}$$

and on the other hand,

$$\begin{aligned} \partial_{f+f'}(\Phi(q \otimes q')) &= \partial_{f+f'}(q, q') \\ &= \sum_{(p, p') \in \text{Crit}_{i+j-1}(f+f')} n((q, q'), (p, p'))(p, p') \\ &= \underbrace{\sum_{p \in \text{Crit}_{i-1}(f)} n(q, p)(p, q')}_{q'=p'} + \underbrace{\sum_{p' \in \text{Crit}_{j-1}(f')} (-1)^i n(q', p')(q, p')}_{q=p}. \end{aligned}$$

□

lem:morse-Alg  
KnForm

**Lemma 5.2.3** (Algebraic Künneth Formula). *If  $\Lambda$  is a principal ideal domain, and  $C$  and  $C'$  are chain complexes over  $\Lambda$  such that all  $C_n$  are free  $\Lambda$ -modules, then for each  $n$  there is a natural short exact sequence*

$$0 \longrightarrow \oplus_i (H_i(C) \otimes_{\Lambda} H_{n-i}(C')) \longrightarrow H_n(C \otimes_{\Lambda} C') \longrightarrow \oplus_i (\text{Tor}_{\Lambda}(H_i(C), H_{n-i-1}(C'))) \longrightarrow 0$$

*that splits.*

*Proof.* This is a standard result of homological algebra, found in most algebraic topology textbooks. Compare Theorem 3B.5. in [Hat02] for example. □

hatcher2002algebraic

**Theorem 5.2.4** (Morse-theoretic Künneth Formula). *Let  $\Lambda$  be a principal ideal domain, and let  $(M, \mathfrak{A}, \Lambda)$  and  $(M', \mathfrak{A}', \Lambda)$  be two Morse data. Then for each  $n$  there is a splitting short exact sequence*

$$\begin{aligned} 0 \rightarrow \bigoplus_i (H_i(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} H_{n-i}(M', \mathfrak{A}', \Lambda)) &\rightarrow H_n(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda) \\ &\rightarrow \bigoplus_i (\text{Tor}_{\Lambda}(H_i(M, \mathfrak{A}, \Lambda), H_{n-i-1}(M', \mathfrak{A}', \Lambda))) \rightarrow 0 \end{aligned}$$

*Proof.* The theorem follows directly from Lemma 5.2.3 above together with the isomorphism of chain complexes

$$C_*(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C_*(M', \mathfrak{A}', \Lambda) \rightarrow C_*(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda).$$

□

**Remark 5.2.5.** One might wonder, why there is no *natural* short exact sequence in the Künneth Formula for Morse homology. This is due to the missing functoriality of Morse homology at the moment.

**Definition 5.2.6** (Morse Cohomology). Let  $G$  be a commutative ring with one. To get the cohomology we have to dualize the chain complex. Let  $\text{Crit}_k(f) = \{p_1, \dots, p_l\}$  be a basis of the  $k$ -th chain group  $C_k(M, \mathfrak{A}, G)$ . Then, the functionals

$$\eta_i : C_k(M, \mathfrak{A}, G) \rightarrow G; p_j \mapsto \begin{cases} 1 & \text{if } j=i \\ 0 & \text{else.} \end{cases}$$

are a basis of the dual group  $C^k(M, \mathfrak{A}, G)$ . The coboundary map is easy to obtain too: Let  $q_j$  for  $j = 1, \dots, r$  be the critical points with index  $k+1$  and let  $\xi_i$  be the corresponding forms. Let  $n(q_j, p_i)$  be the number of flow lines from  $q_j$  to  $p_i$  counted with the sign. Then the coboundary map is given as:

$$\begin{aligned} \partial^k : C^k(M, \mathfrak{A}, G) &\rightarrow C^{k+1}(M, \mathfrak{A}, G) \\ \eta_i &\mapsto \partial^k \eta_i = \eta_i \circ \partial_{k+1} = \sum_j n(q_j, p_i) \xi_j. \end{aligned}$$

**Corollary 5.2.7** (Algebraic Künneth Formula for Cohomology). If  $\Lambda$  is a principal ideal domain, and  $C$  and  $C'$  are cochain complexes over  $\Lambda$  such that all  $C^n$  are free  $\Lambda$ -modules, then for each  $n$  there is a natural short exact sequence

$$0 \longrightarrow \bigoplus_i (H^i(C) \otimes_{\Lambda} H^{n-i}(C')) \longrightarrow H^n(C \otimes_{\Lambda} C') \longrightarrow \bigoplus_i (\text{Tor}_{\Lambda}(H^i(C), H^{n-i+1}(C'))) \longrightarrow 0$$

that splits.

*Proof.* This is a corollary of the Algebraic Künneth Formula. For this we define the chain complexes

$$D_{-i} := C^i \quad \text{and} \quad D'_{-i} := C'^i$$

with the same differentials

$$d_i : D_i \rightarrow D_{i-1}, c \mapsto d^{-i}(c) \quad \text{and} \quad d'_i : D'_i \rightarrow D'_{i-1}, c' \mapsto d'^{-i}(c')$$

Then we have the equalities

$$H_i(D) = H^{-i}(C), \quad \text{and} \quad H_i(D') = H^{-i}(C').$$

Furthermore, the redefinition is well behaved with tensor products in the sense that

$$D \otimes_{\Lambda} D' = \bigoplus_{i+j=m} D_i \otimes D_j = \bigoplus_{i+j=m} C^{-i} \otimes C^{-j} = (C \otimes_{\Lambda} C')^{-m}$$

and the boundary map of the tensor product chain complex is the same as the coboundary map of the tensor product of the cochain complexes (with the inverted degree). This leads to the equality

$$H_m(D \otimes_{\Lambda} D') = H^{-m}(C \otimes_{\Lambda} C').$$

Now we can apply the Algebraic Künneth Formula to  $D, D'$  to get the short splitting exact sequence

$$0 \longrightarrow \bigoplus_i (H_i(D) \otimes_{\Lambda} H_{m-i}(D')) \longrightarrow H_m(D \otimes_{\Lambda} D') \longrightarrow \bigoplus_i (\text{Tor}_{\Lambda}(H_i(D), H_{m-i-1}(D'))) \longrightarrow 0$$

We can now define  $n := -m$  and reindex the summations from  $i$  to  $-i$ , giving us the desired sequence.  $\square$

**Theorem 5.2.8** (Morse-theoretic Künneth Formula for Cohomology). *Let  $\Lambda$  be a principal ideal domain, and let  $(M, \mathfrak{A}, \Lambda)$  and  $(M', \mathfrak{A}', \Lambda)$  be two Morse data. Then for each  $n$  there is a splitting short exact sequence*

$$\begin{aligned} 0 \rightarrow \bigoplus_i (H^i(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} H^{n-i}(M', \mathfrak{A}', \Lambda)) &\rightarrow H^n(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda) \\ &\rightarrow \bigoplus_i (\text{Tor}_{\Lambda}(H^i(M, \mathfrak{A}, \Lambda), H^{n-i+1}(M', \mathfrak{A}', \Lambda))) \rightarrow 0 \end{aligned}$$

*Proof.* We show this by giving an isomorphism of the cochain complexes

$$C^*(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C^*(M', \mathfrak{A}', \Lambda) \rightarrow C^*(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda).$$

First we need to give a group isomorphism

$$\Phi : \bigoplus_{i+j=k} C^i(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C^j(M', \mathfrak{A}', \Lambda) \rightarrow C^k(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda)$$

This isomorphism, however, is clear and given on the generators: Let  $p$  and  $p'$  be critical points of index  $i$  and  $j$  with respect to  $f$  and  $f'$ . Let  $\eta$  and  $\eta'$  be the corresponding generators of  $C^i(M, \mathfrak{A}, \Lambda)$  and  $C^j(M', \mathfrak{A}', \Lambda)$ . Then we have the bijection

$$\eta \otimes \eta' \leftrightarrow (\eta, \eta').$$

To get a chain isomorphism we need to check the compatibility with the coboundary maps. For each  $q_l \in \text{Crit}_{i+1}(f)$  let  $\xi_l$  be the corresponding form and similarly for each  $q'_k \in \text{Crit}_{j+1}(f')$  let  $\xi'_k$  be the corresponding form. Then we calculate

$$\begin{aligned} \Phi \left( \partial^f \otimes 1 + (-1)^i 1 \otimes \partial^{f'} \right) (\eta \otimes \eta') &= \Phi \left( \partial^f(\eta) \otimes \eta' + (-1)^i \eta \otimes \partial^{f'}(\eta') \right) \\ &= \Phi \left( \sum_l n(q_l, p) \xi_l \otimes \eta' + \sum_k (-1)^i n(q'_k, p') \eta \otimes \xi'_k \right) \\ &= \sum_l n(q_l, p) (\xi_l, \eta') + \sum_k (-1)^i n(q'_k, p') (\eta, \xi'_k) \end{aligned}$$

and the other way around, if we define for  $(q, q')$  with  $\text{ind}_{f+f'}(q, q') = \text{ind}_{f+f'}(p, p') + 1$  the corresponding form to be  $\nu$  we have:

$$\begin{aligned} \partial^{f+f'} \Phi(\eta \otimes \eta') &= \partial^{f+f'}(\eta, \eta') \\ &= \sum_{(q, q') \in \text{Crit}_{i+j+1}(f+f')} n((q, q'), (p, p')) (\nu) \\ &= \underbrace{\sum_l n(q_l, p) (\xi_l, \eta')}_{q'=p'} + \underbrace{\sum_k (-1)^i n(q'_k, p') (\eta, \xi'_k)}_{q=p} \end{aligned}$$

Now finally we can apply the Algebraic Künneth Formula for Cohomology and we are done. The latter formula is almost the same as the homological one but with a different index shift in the Tor functor. Again this is standard homological algebra.  $\square$

## 6 A Morse-Theoretic Filtration

### 6.1 Conley index

```
def:morseFilt
r-compactInva
riantIsolated
Subset
```

**Definition 6.1.1** (Compact Invariant Isolated Subset). For a flow  $\varphi_t : M \rightarrow M$  on a locally compact metric space, we call a subset  $S \subseteq M$  an **invariant subset** if and only if  $\varphi_t(S) = S$  for all  $t \in \mathbb{R}$ . For any subset  $N \subseteq M$  we define the **maximal invariant subset**

$$\begin{aligned} I(N) &= \{x \in N \mid \varphi_t(x) \in N \forall t \in \mathbb{R}\} \\ &= \bigcap_{t \in \mathbb{R}} \varphi_t(N). \end{aligned}$$

A **compact invariant subset**  $S$  is called **isolated** if a compact neighbourhood  $N$  exists such that  $I(N) = S$ .

```
def:morseFilt
r-indexPairs
```

**Definition 6.1.2** (Index Pairs). Let  $S$  be an isolated compact invariant subset. A topological pair  $(N, L)$  of compact subsets of  $M$  where  $L \subseteq N$  is called an **index pair of  $S$** , if it satisfies the following:

1.  $S = I(\overline{N \setminus L}) \subseteq \text{int}(N \setminus L)$ .
2.  $x \in L$  and  $\varphi_{[0,t]}(x) \subseteq N$  implies that  $\varphi_{[0,t]}(x) \subseteq L$ . We call  $L$  **positively invariant in  $N$** . Here  $\varphi_{[0,t]}(x) := \{\varphi_{\tilde{t}}(x) \mid \tilde{t} \in [0, t]\}$ .
3. For all  $x \in N$  such that a  $t$  exists, with  $\varphi_t(x) \notin N$ , there exists a  $t'$  with  $\varphi_{[0,t']}(x) \subseteq N$  and  $\varphi_{t'}(x) \in L$ .

This definition captures how the flow lines leave the invariant space  $S$ . Due to the third property we call  $L$  the **exit set**.

```
def:morseFilt
r-regularInde
xPairs
```

**Definition 6.1.3** (Regular Index Pairs). We call an index pair  $(N, L)$  **regular** if and only if the inclusion  $i : L \hookrightarrow N$  is a cofibration.

```
thm:morseFilt
r-homotopyEqu
ivalenceOfInd
exPairs
```

**Theorem 6.1.4** (Homotopy Equivalence of Index Pairs). *If  $S$  is an isolated compact invariant subset and  $(N, L)$  and  $(\tilde{N}, \tilde{L})$  are two index pairs of  $S$ , then  $N/L$  and  $\tilde{N}/\tilde{L}$  are homotopy equivalent as pointed spaces. This homotopy equivalence is canonical, and transitive up to homotopy.*

*Proof.* For a proof of the homotopy equivalence we refer to [banyaga2004lectures](#) [BH04], Theorem 7.15. The constructed homotopy equivalence is given as follows: Choose  $T \geq 0$  such that the following implications hold for  $t \geq T$ :

$$\begin{aligned} \varphi_{[-t,t]}(x) \subseteq N \setminus L &\Rightarrow x \in \tilde{N} \setminus \tilde{L}, \\ \varphi_{[-t,t]}(x) \subseteq \tilde{N} \setminus \tilde{L} &\Rightarrow x \in N \setminus L, \end{aligned}$$

where  $\varphi_{[a,b]}(x) := \{\varphi_t(x) \mid t \in [a, b]\}$ . Then the map:

$$h : N/L \times [T, \infty) \rightarrow \tilde{N}/\tilde{L}$$

$$([x], t) \mapsto \begin{cases} [\varphi_{3t}(x)] & \text{if } \varphi_{[0,2t]}(x) \subseteq N \setminus L \text{ and } \varphi_{[t,3t]}(x) \subseteq \tilde{N} \setminus \tilde{L} \\ [\tilde{L}] & \text{otherwise,} \end{cases}$$

is continuous and for any fixed  $t$  this gives our homotopy equivalence. Obviously, such a map is canonical and transitive up to homotopy.  $\square$

```
def:morseFilt
r-connectedSi
mpleSystem
```

**Definition 6.1.5** (Connected Simple System). A **connected simple system** consists of a set  $I_0$  of pointed spaces along with a set  $I_m$  of homotopy classes of maps between these spaces such that

- (i)  $\text{hom}(X, Y) := \{[f] \in [X, Y] \mid [f] \in I_m\}$  consists of exactly one element for each pair of spaces  $(X, Y) \in I_0^2$ , where  $[X, Y]$  denotes the set of homotopy classes of maps between  $X$  and  $Y$ .
- (ii) If  $X, Y, Z \in I_0$ ,  $[f] \in \text{hom}(X, Y)$ , and  $[g] \in \text{hom}(Y, Z)$ , then  $[g \circ f] \in \text{hom}(X, Z)$ .
- (iii)  $\text{hom}(X, X) = \{[\text{id}_X]\}$  for all  $X \in I_0$ .

```
rem:morseFilt
r-connectedSi
mpleSystemSal
monmorseFilt
r-indexPairCo
nectedSimple
System
```

**Remark 6.1.6.** This definition comes from Salamon in [\[Sal85\]](#) <sup>[Salamon1985](#)</sup>.

**Remark 6.1.7.** Let  $S$  be an isolated compact invariant subset of  $M$  with respect to the Morse flow  $\varphi$ . Then the collections

$$I_0 := \{N/L \mid (N, L) \text{ is an index pair of } S\}$$

$$I_m := \{[f] \mid f \text{ is a homotopy equivalence from Theorem 6.1.4}\},$$

make up a connected simple system. We denote this by  $\mathcal{IP}_S$ . If  $S = \text{Crit}_k(f)$  for some  $k \in \mathbb{N}$ , we denote  $\mathcal{IP}_{\text{Crit}_k(f)} = \mathcal{IP}_k$ .

```
def:morseFilt
r-morphismInd
ConnectedSimp
leSystems
```

**Definition 6.1.8** (Morphisms of Connected Simple Systems). A **morphism**  $\Phi : I \rightarrow J$  between two connected simple systems  $I = (I_0, I_m)$  and  $J = (J_0, J_m)$  is a set of homotopy classes of maps between spaces in  $I_0$  and  $J_0$  such that

- (i) For every  $X \in I_0$  and  $Y \in J_0$  the set  $\Phi(X, Y) := \{[\varphi] \in [X, Y] \mid [\varphi] \in \Phi\}$  consists of exactly one element.
- (ii) If  $X, \bar{X} \in I_0$  and  $Y, \bar{Y} \in J_0$  and if  $[\varphi] \in \Phi(X, Y)$ ,  $[f] \in \text{hom}(\bar{X}, X)$ ,  $[g] \in \text{hom}(Y, \bar{Y})$ , then  $[g \circ \varphi \circ f] \in \Phi(\bar{X}, \bar{Y})$ .

```
rem:morseFilt
r-morphismInd
ucedBySingleM
ap
```

**Remark 6.1.9.** By the second property of a morphism between connected simple systems  $I$  and  $J$ , we can induce such a morphism by a single map  $\varphi : X \rightarrow Y$  where  $X \in I_0$  and  $Y \in J_0$ .

```
def:morseFilt
r-canonically
omorphismClas
s
```

**Definition 6.1.10** (Canonical Isomorphism Class). A **canonical isomorphism class** of Abelian groups (or other algebraic structures) is a set of groups  $R_0$  together with a set of group homomorphisms  $R_m$  such that

- (i) For  $R, L \in R_0$  the set  $\text{hom}(R, L) := \{f : R \rightarrow L : f \in R_m\}$  consists of exactly one element.
- (ii) If  $R, L, S \in R_0$  and  $f \in \text{hom}(R, L)$  and  $g \in \text{hom}(L, S)$ , then  $g \circ f \in \text{hom}(R, S)$ .
- (iii)  $\text{hom}(R, R) = \{\text{id}_R\}$

A **morphism** between two canonical isomorphism classes of Abelian groups  $\Psi : (R_0, R_m) \rightarrow (S_0, S_m)$  is a collection of homomorphisms such that

- (i) For any  $R \in R_0$  and  $S \in S_0$  the set  $\Psi(R, S) := \{f : R \rightarrow S : f \in \Psi\}$  contains exactly one element.
- (ii) If  $R, \bar{R} \in R_0$  and  $S, \bar{S} \in S_0$ ,  $f \in \text{hom}(\bar{R}, R)$ ,  $g \in \text{hom}(S, \bar{S})$  and  $\psi \in \Psi(R, S)$ , then  $g \circ \psi \circ f \in \Psi(\bar{R}, \bar{S})$ .

**Remark 6.1.11.** Again the second property lets us induce a morphism between two canonical isomorphism classes of Abelian Groups  $\Psi : (R_0, R_m) \rightarrow (S_0, S_m)$  by a single isomorphism

$$\psi : R \rightarrow S$$

for  $R \in R_0$  and  $S \in S_0$ . We denote the induced morphism by  $\psi_{\#} := \Psi$ .

**Example 6.1.12.** Any pointed topological space  $X$  defines a simple connected system  $(\{X\}, [\text{id}_x])$ . Similar, any Abelian group  $G$  defines a canonical isomorphism class of Abelian groups  $(\{G\}, \{\text{id}_R\})$ . Let  $\varphi : S \rightarrow L$  be a second homomorphism that induces a morphism of canonical isomorphism classes. It is then obvious that

$$(\varphi \circ \psi)_{\#} = \varphi_{\#} \circ \psi_{\#}.$$

```
def:morseFilt
r-systemCate
ories
```

**Definition 6.1.13.** We denote the category of connected simple systems with their morphisms by **CSS** and similar we call the category of canonical isomorphism classes of Abelian groups by **CIAB**, respectively, the category for rings by **CIRing**.

```
def:morseFilt
r-postcompose
dFunctor
```

**Definition 6.1.14.** Let  $G$  be an Abelian group and  $X$  be any set. We can **tag**  $G$  with  $X$  by defining the set  $G \times \{X\}$  and endowing it with the algebraic information from  $G$ . (for example we define  $(g, X) \cdot (g', X) = (g \cdot_G g', X)$ ). Let  $f : G \rightarrow L$  be a homomorphism. Then  $f$  induces a homomorphism between the tagged  $G \times \{X\}$  and  $L \times \{Y\}$  by sending  $(g, X) \mapsto (f(g), Y)$ . Let  $F : \mathbf{CGWH}^* \rightarrow \mathbf{Ab}$  be a (contravariant) functor from pointed topological spaces to Abelian groups that factors over the homotopy category, i.e. is homotopy invariant. We first define the tagged functor  $F^{tag}$  to map spaces to groups that are tagged with the spaces they come from:

$$\begin{aligned}
F^{tag} : \mathbf{CGWH}^* &\rightarrow \mathbf{Ab} \\
X &\mapsto F(X) \times \{X\} \\
(f : X \rightarrow Y) &\mapsto (F^{tag}(f) : F(Y) \times \{Y\} \rightarrow F(X) \times \{X\})
\end{aligned}$$

For covariant functors the procedure is analogous. Now we can define the **postcomposed functor**  $F_{\#}$  from the category of connected simple systems to the canonical isomorphism classes as follows:

$$\begin{aligned} F_{\#} : \mathbf{CSS} &\rightarrow \mathbf{CIAB} \\ I_0 &\mapsto \{F^{tag}(X) \mid X \in I_0\} \\ I_m &\mapsto \{F^{tag}(g) \mid [g] \in I_m\} \end{aligned}$$

Let  $\Phi : (I_0, I_m) \rightarrow (J_0, J_m)$  be a morphism in  $\mathbf{CSS}$ . Then we define

$$F_{\#}(\Phi) : F_{\#}(J_0, J_m) \rightarrow F_{\#}(I_0, I_m), \quad F_{\#}(\Phi) := \{F^{tag}(g) \mid [g] \in \Phi\}.$$

This obviously becomes a new functor, where the tagging removed uniqueness problems for non-injective (in the literal sense not up to isomorphism) functors  $F$ . If  $F(X) = F(Y)$  for different  $X, Y \in I_0$  with  $g : X \rightarrow Y \in I_m$ , then without tagging there could be two different morphisms  $F(X) \rightarrow F(X)$ : the identity and  $F(g)$ . Since our functors are almost always injective, we will ignore the tagging from now on.

**Remark 6.1.15.** Obviously, the procedure of postcomposing a functor works for any homotopy invariant functor from pointed topological spaces into any algebraic category and not just Abelian groups.

**Remark 6.1.16** (Critical points as isolated compact invariant subsets). Let  $f : M \rightarrow \mathbb{R}$  be a Morse function on a smooth Riemannian manifold  $(M, g)$ . Let  $q \in \text{Crit}_k(f)$ . Around  $q$  we have refined Morse charts  $\varphi : U \rightarrow \mathbb{R}^m \cong \mathbb{R}^{m-\lambda_q} \oplus \mathbb{R}^{\lambda_q}$  where the splitting of  $\mathbb{R}^m$  is taken such that  $d\varphi_q$  is expanding in  $\mathbb{R}^{m-\lambda_q}$  and contracting in  $\mathbb{R}^{\lambda_q}$ . With this we can define the balls:

$$\begin{aligned} D_{\varepsilon}^s &:= \{v \in \mathbb{R}^{m-\lambda_q} \mid \|v\| \leq \varepsilon\}, \\ D_{\varepsilon}^u &:= \{v \in \mathbb{R}^{\lambda_q} \mid \|v\| \leq \varepsilon\}. \end{aligned}$$

For epsilon small enough, they give rise to the index pair  $N_q := \varphi^{-1}(D_{\varepsilon}^s \times D_{\varepsilon}^u)$  and  $L_q := \varphi^{-1}(D_{\varepsilon}^s \times \partial D_{\varepsilon}^u)$ . To see this, notice that in such a refined Morse chart  $\varphi$ , the differential of the flow in the critical point splits the tangent space into  $T_q^u M \times T_q^s M$ . Finally, since the property of  $d\varphi_x$  being a contraction on  $T_q^s M$  is an open condition and thereby there is an  $\varepsilon > 0$  such that  $d\varphi_x$  is a contraction on  $T_q^s M$  for all  $x$  with  $\|\varphi^{-1}(x)\| < \varepsilon$ . Thereby, the norm of  $\varphi_t(x)$  projected to  $T_q^s M$  is strictly decreasing in  $t$ . This proves that  $L_q$  is an exit set of  $N_q$ . Similarly, one sees that  $L_q$  is positively invariant in  $N_q$ .

This index pair is in fact regular, since  $(N_q, L_q) \simeq (D_{\varepsilon}^s \times D_{\varepsilon}^u, D_{\varepsilon}^s \times \partial D_{\varepsilon}^u) \simeq (D_{\varepsilon}^u, \partial D_{\varepsilon}^u)$  and the latter's regularity is easy to see by showing that it is an NDR pair. We will call such an index pair a **standart index pair of  $q$** .

Furthermore, the index pair  $(N_q, L_q)$  is homotopically equivalent to  $\overline{T_q^u M}$ , which is clear, as  $(D_{\varepsilon}^u, \partial D_{\varepsilon}^u) \simeq \overline{T_q^u M}$  by mapping  $\mathbb{R}^{\lambda_q} \rightarrow T_q^u M$  via  $d\varphi^{-1}|_0 \circ R_0$ , where  $R_0 : \mathbb{R}^m \rightarrow$

rem:morseFilt  
r-criticalPoi  
ntsAsIsolated  
CompactInvari  
antSubsets

$T_0\mathbb{R}^m$  is the linear identification from Definition 2.0.45. Finally, we just need to stretch the ball to infinity.

We know the K-group of such a pair to be:

$$K^i(N_q, L_q) \cong K^i(\overline{T_q^u M}) \cong \begin{cases} \mathbb{Z} & \text{if } i - k \in 2\mathbb{Z}, \\ 0 & \text{else.} \end{cases} \quad (33)$$

{eq:homologyof  
findexpairsof  
criticalpoint}

Notice that for  $k = 0$  we need to consider  $\partial D^k = \emptyset$  to get an index pair. That's why we let  $\partial$  denote the manifold boundary instead of the topological boundary. However, now we can identify for all  $k$ :

$$C^k(M, \mathfrak{A}, \tilde{K}^l(pt)) = \bigoplus_{\text{Crit}_k(f)} \mathbb{Z} \cong \bigoplus_{q \in \text{Crit}_k(f)} K^{-k}(N_q, L_q). \quad (34)$$

{criticalpoint  
tsasdirectsum  
ofhomologygroups}

Or using the language of connected simple systems, we have the identification

$$C^k(M, \mathfrak{A}, \tilde{K}^l(pt)) \cong \bigoplus_{q \in \text{Crit}_k(f)} K_{\#}^{-k}(\mathcal{IP}_q), \quad (35)$$

where the isomorphism can be read as an isomorphism of canonical isomorphism classes with the left-hand side viewed as an object of **CIAB** consists of only one group.

**Remark 6.1.17.** For any chart  $\chi$  that is centered at a critical point  $q$  and  $\varepsilon > 0$  we define the topological pair

$$(N_\chi, L_\chi) := (\chi^{-1}(D_\varepsilon^s \times D_\varepsilon^u), \chi^{-1}(D_\varepsilon^s \times \partial D_\varepsilon^u)).$$

The argument that shows that  $(N_\psi, L_\psi)$  is a regular index pair of  $q$  only needed that  $d\psi|_q$  splits. Hence if  $d\chi|_q = d\psi|_q$ , we can conclude, that there is an  $\varepsilon > 0$  such that  $(N_\chi, L_\chi)$  is also a regular index pair.

**Remark 6.1.18.** Let  $\underline{\mathbb{Z}}$  be the canonical isomorphism class of Abelian groups consisting only of the group  $\mathbb{Z}$ , and let  $q$  be a critical point of index  $\lambda_q$  in  $M$ . Then, Remark 6.1.9, together with Remark 6.1.16 gives us the existence of a morphism for all  $i \in \mathbb{N}$ :

$$\tilde{K}^{-\lambda_q-2i}(\mathcal{IP}_{\{q\}}) \rightarrow \underline{\mathbb{Z}}.$$

Furthermore, this map is given by mapping a generator of  $\tilde{K}^{-\lambda_q-2i}(N_q/L_q)$  to one, and hence uniquely determined, provided we have a canonical way to choose a generator of  $\tilde{K}^{-\lambda_q-2i}(N_q/L_q) \cong \tilde{K}^{-\lambda_q-2i}(\overline{T_q^u M})$ , where the isomorphism is natural and described in Remark 6.1.16.

Hence, if  $T_q^u M$  is oriented, Corollary 4.2.5 gives us just what we need, making the map canonical. The generator is then given by pulling back the generator of  $K^{-\text{ind}_p}(\overline{T_q^u M})$  along the map of pairs:

$$\kappa_q := d\varphi^{-1}|_0 \circ R_0 \circ s \circ \text{pr}_u \circ \varphi : (\varphi^{-1}(D_\varepsilon^s \times D_\varepsilon^u), \varphi^{-1}(D_\varepsilon^s \times \partial D_\varepsilon^u)) \rightarrow (\overline{T_q^u M}, \{\infty\})$$

where  $\text{pr}_u$  is the projection onto the unstable coordinates and  $s$  is the stretching  $x \mapsto x \cdot (\frac{1}{\varepsilon} - \|x\| - \frac{1}{\varepsilon})$  with  $ds|_0 = \frac{1}{\varepsilon} \mathbb{1}$ . We call such a generator the **standart generator at  $q$** .

rem:morseFiltr  
r-indexPairFor  
DifferentCharts

rem:morseFiltr  
r-morphismToCanonicalIntegers

**Example 6.1.19** (A Map of K-Groups). Let  $f : M \rightarrow \mathbb{R}$  be a Morse-Smale function on a smooth compact Riemannian manifold  $(M, g)$ . Let  $q \in \text{Crit}_k(f)$  and  $p \in \text{Crit}_{k-1}(f)$ . Assume for a moment we already have a regular index pair  $(N_2, N_0)$  of the isolated compact subset  $S = W(p \rightarrow q) \cup \{p, q\}$ . For a  $c \in (f(p), f(q))$ , we set  $N_1 = N_0 \cup (N_2 \cap M^c)$ , where  $M^c$  is the sublevel set. Clearly now  $(N_2, N_1)$  is a regular index pair of  $q$  and  $(N_1, N_0)$  is a regular index pair of  $p$ .

Furthermore, we have the triple connecting homomorphism  $\delta_{triple}^{p,q} : K^{-\text{ind}_p}(N_1, N_0) \rightarrow K^{-\text{ind}_p+1}(N_2, N_1)$ , which we can compose with the Bott periodicity map  $b := \beta \boxtimes (-)$ . This lets us induce the map:

$$(b \circ \delta_{triple}^{p,q})_{\#} : K_{\#}^{-\text{ind}_p}(\mathcal{IP}_{\{p\}}) \rightarrow K_{\#}^{-\text{ind}_q}(\mathcal{IP}_{\{q\}})$$

Now the disjoint union of the index pairs  $(N_q, L_q)$  for all  $q \in \text{Crit}_k(f)$  is an index pair of  $\text{Crit}_k(f)$  itself. The collapsing maps  $c_p : \bigvee_{p \in \text{Crit}_{k-1}} (N_p/L_p) \rightarrow (N_p/L_p)$  together with the inclusions  $i_q : (N_q/L_q) \rightarrow \bigvee_{q \in \text{Crit}_k} (N_q/L_q)$  let us define the map

$$\Delta^{k-1} := \left( \sum_{q \in \text{Crit}_k} \sum_{p \in \text{Crit}_{k-1}} c_p \circ (b \circ \delta_{triple}^{p,q} \circ i_q) \right)_{\#} : K_{\#}^{-(k-1)}(\mathcal{IP}_{k-1}) \rightarrow K_{\#}^{-k}(\mathcal{IP}_k).$$

Due to the naturality of the triple connecting homomorphism, the inclusion and the collapsing, any list of topological triples  $(X, Y, Z)$  such that  $(X, Y)$  is a regular index pair of  $\{q\}$ ,  $(Y, Z)$  of  $\{p\}$  and  $(X, Z)$  of  $W(q \rightarrow p) \cup \{q\} \cup \{p\}$  for all  $q \in \text{Crit}_k(f), p \in \text{Crit}_{k-1}(f)$  gives rise to the same morphism of canonical isomorphism classes.

## 6.2 Commuting Boundary Maps

thm:morseFilt  
-commutingMor  
secoboundary

**Theorem 6.2.1.** *Let  $M$  be a smooth manifold with the Morse datum  $\mathfrak{A}$ , denote the one pointed topological space by  $\{pt\}$ . Then, for any  $l \in \mathbb{Z}$  the following diagram comutes:*

$$\begin{array}{ccc} C^{k-1}(M, \mathfrak{A}, K^l(pt)) & \xrightarrow{\partial^{k-1}} & C^k(M, \mathfrak{A}, K^l(pt)) \\ \downarrow & & \downarrow \\ K_{\#}^{-(k-1)-l}(\mathcal{IP}_{k-1}) & \xrightarrow{\Delta^{k-1+l}} & K_{\#}^{-(k)-l}(\mathcal{IP}_k) \end{array} \quad (36)$$

{eq:morseFilt  
-commutingDia  
gramTheorem}

Before we can prove this theorem we need the following Lemma:

def:morseFilt  
-setupCommuti  
ngDiff

**Definition 6.2.2** (Definitions and Setup). We work in a compact, Riemannian, smooth manifold  $(M, g)$  of dimension  $m$  together with a Morse-Smale function

$$f : M \rightarrow \mathbb{R} \quad (37)$$

and orientations  $\mathfrak{o}$  of all unstable tangent spaces. We denote the flow corresponding to the negative gradient by  $\varphi_t(x)$ . We assume that  $q$  is a critical point of index  $k$  and  $p$  is

a critical point of index  $k - 1$ . We define  $f(q) = b$  and  $f(p) = a$  and assume that there is no critical point in  $f^{-1}(a, b) \subseteq M$ . We define the sublevel and superlevel sets

$$M^t := \{x \in M \mid f(x) \leq t\} \quad \text{and} \quad M_t := \{x \in M \mid f(x) \geq t\}, \quad (38)$$

and the constants

$$c \in (a, b) \quad , \quad \varepsilon > 0 \text{ small} \quad , \quad T > 0 \text{ big} . \quad (39)$$

With this we define the sets:

$$N_q := \{x \in M_c \mid f(\varphi_{-T}(x)) \leq b + \varepsilon\}, \quad (40)$$

$$L_q := \{x \in N_q \mid f(x) = c\}, \quad (41)$$

$$N_p := \{x \in M^c \mid f(\varphi_T(x)) \geq a - \varepsilon\}, \quad (42)$$

$$L_p := \{x \in N_p \mid f(\varphi_T(x)) = a - \varepsilon\}, \quad (43)$$

and finally:

$$C := N_p \cup N_q, \quad (44)$$

$$B := N_p \cup L_q, \quad (45)$$


$$A := L_p \cup \text{int}(L_q \setminus N_p) \quad (46)$$

$$S(q \rightarrow) := W(q \rightarrow) \cap f^{-1}(c) \quad (47)$$

$$D_q := W(q \rightarrow) \cap M_c. \quad (48)$$

Furthermore, we assume  $W(q \rightarrow)$  to be oriented with respect to the orientation of  $T_q^u M$  as in Proposition 5.1.3. Let  $Q^p := TM|_{W(\rightarrow p)} / TW(\rightarrow p)$  be oriented with respect to  $T_p^u M$  as in Proposition 5.1.5. For an oriented real vector space  $V$  of dimension  $n$ , let  $\beta_V \in K^{-n}(\bar{V})$  denote its K-theoretic orientation from Corollary 4.2.5. For an index pair  $(N, L)$  of  $\{p\}$  let  $\alpha_p^{(N, L)} \in K^{-\text{ind}_p}(N, L)$  denote the canonical generator from Remark 6.1.18, i.e. the pullback of  $\kappa_p^*(\beta_{T_p^u M})$  along the homotopy equivalence of Theorem 6.1.4 from  $(N, L)$  to the standart index pair and  $\kappa_v = \kappa_q := R_0 \circ d\varphi^{-1}|_0 \circ s \circ \text{pr}_u \circ \varphi$ . später

Let  $(\alpha_p)_\#$  be a generator of  $K_\#^{\lambda_p}(\mathcal{IP}_p)$  meaning that  $(\alpha_p)_\#$  is a family of generators for all groups in  $K_\#^{\lambda_p}(\mathcal{IP}_p)$  that transform into each other under the canonical group isomorphisms. Notice that  $(\alpha_p)_\#$  is induced by any of its elements.



lem:morseFilt  
-npIsTubular

lem:enum:1

lem:enum:2

lem:enum:3

**Lemma 6.2.3.** *In the setting of Definition 6.2.2, we claim that*

1.  $(N_q, L_q)$  is a regular index pair for  $q$  .
2.  $(C, B)$  is an index pair for  $q$  .
3.  $(N_p, L_p)$  is a regular index pair for  $p$  .




Figure 5: The map  $\tau$  and  $\rho$  from Lemma 6.2.4 (a) is shown on the left and the map  $G$  from (c) is shown on the right.

lem:enum:4

4.  $(B, A)$  is an index pair for  $p$ .

lem:enum:5

5.  $N_p$  is a tubular neighbourhood of  $W(\rightarrow p) \cap M^c$ .

lem:morseFilt  
-flowTimeAndP  
rojectionToLe  
velSets

**Lemma 6.2.4.** Let  $\text{ind}_q(f) \geq 1$ .

- (a) For  $x \in D_q \setminus \{q\}$  there is a unique  $\tau(x) \geq 0$  with  $f(\varphi_{\tau(x)}(x)) = c$ . The resulting map  $\tau : D_q \setminus \{q\} \rightarrow [0, \infty)$  is smooth,  $\tau^{-1}(0) = S(q \rightarrow)$ ,  $\tau(\varphi_s(x)) = \tau(x) - s$ ,  $d\tau|_x(-\text{grad}_x(f)) = -1$  and  $\tau(x) \rightarrow \infty$  for  $x \rightarrow q$ . We set  $\tau(q) = \infty$ .
- (b) The map  $\rho : D_q \setminus \{q\} \rightarrow S(q \rightarrow)$ ,  $\rho(x) := \varphi_{\tau(x)}(x)$  is a smooth retraction with  $\rho \circ \varphi_s = \rho$ . In particular,  $d\rho|_x(\text{grad}_x(f)) = 0$  and for  $y \in S(q \rightarrow)$ ,  $w \in T_y(S(q \rightarrow))$  and  $s \geq 0$ :

$$d\rho(d\varphi_{-s}|_y w) = w, \quad d\tau(d\varphi_{-s}|_y w) = 0.$$

- (c)  $S(q \rightarrow)$  is a closed submanifold of  $W(q \rightarrow)$  of dimension  $\text{ind}_q - 1$ . There are homeomorphisms  $g_q : S^{\text{ind}_q} \rightarrow D_q/S(q \rightarrow)$  with  $g_q(S) = [S(q \rightarrow)]$  and a constant  $T_q \geq 0$  such that, with the open neighbourhood

$$W_q := \{x \in D_q \mid \tau(x) > T_q\} \ni q$$

of  $q$ , the map  $g_q$  restricts to an orientation preserving diffeomorphism from  $U_q := g_q^{-1}(W_q)$  onto  $W_q$ .

Bild

*Proof.* (a) The function  $f$  strictly decreases along flow lines, which approach critical points. As there is no critical point in  $f^{-1}[c, b]$  our flow line needs to pass through  $f^{-1}(c)$  giving us a  $\tau(x)$ , where uniqueness follows again from strict monotonicity. This uniqueness lets us deduce that  $\tau(\varphi_s(x)) = \tau(x) - s$ . The smoothness follows from the implicit function theorem applied to  $G(x, t) := f(\varphi_t(x)) - c$ , since  $\partial_t G(x, t) = -\|\text{grad}_{\varphi_t(x)}(f)\|^2 \neq 0$ . (compare Proposition 2.0.60 for the calculation) Differentiating  $\tau(\varphi_s(x)) = \tau(x) - s$  in  $s = 0$  gives  $d\tau(-\text{grad}(f)) = -1$ . Finally assume that  $x_n \rightarrow q$  whilst  $\tau(x_n) \leq T$  for some  $T > 0$ . Then we can pass to a subsequence such that  $\tau(x_n) \rightarrow t$  for  $t \leq T$ . In this subsequence we then have that  $\lim_{n \rightarrow \infty} f(\varphi_{\tau(x_n)}(x_n)) = f(\varphi_t(q)) = b$  contradicting  $f(\varphi_{\tau(x_n)}(x_n)) = c < b$  for all  $n \in \mathbb{N}$ .

(b) By (a) we have that  $\rho \circ \varphi_s = \rho$ , and differentiating it in  $s$  gives us  $d\rho(\text{grad}(f)) = 0$ . The last two identities are the differentials of  $\rho \circ \varphi_{-s} = \text{id}$  and  $\tau \circ \varphi_{-s} = \text{const}_s$  on  $S(q \rightarrow)$ .

(c)  $\text{grad}(f)$  is tangent to  $W(q \rightarrow)$  (meaning that it is contained in the tangent bundle of  $W(q \rightarrow)$  as a subbundle of  $TM$ ) and non-zero away from  $q$ . Hence,  $c$  is a regular value of  $f|_{W(q \rightarrow)}$ , (d  $f|_{W(q \rightarrow)}(\text{grad}(f))$  is nonzero outside of critical points) hence  $S(q \rightarrow)$  is a submanifold of  $W(q \rightarrow)$ .

Let  $\psi = (\psi_s, \psi_u)$  be a refined Morse chart around  $q$  such that the unstable coordinates correspond to a positively oriented basis of  $T_q^u M = T_q W(q \rightarrow)$ . Here,  $\psi_u$  denotes  $\pi_u \circ \psi$  where  $\pi_u$  projects to the unstable coordinates (and  $\pi_s$  is defined analogously). By Lemma 5.0.20 the unstable manifold is, in these coordinates the graph of a smooth function  $g$  with  $g(0) = 0$ , that is Lipschitz with Lipschitz constant less than one. (This  $g$  does not denote the Riemannian metric, which doesn't appear by name in this proof.) Hence,  $\psi_u$  restricts to a diffeomorphism from a neighbourhood  $O$  of  $q$  in  $W(q \rightarrow)$  onto an open ball around 0 and  $\|\psi(x)\| \leq \sqrt{2} \|\psi_u(x)\|$  on  $O$  (The latter is deduced from the Lipschitz constant of  $g$  being less than or equal to one). Write  $u(t) := \psi_u(\varphi_t(x))$  for  $x \in O$  (and  $t$  small enough such that  $\varphi_t(x) \in O$ ) and denote pullback of the gradient via the chart  $\psi$  by  $X := -q_\psi \circ \text{grad}(f) \circ \psi^{-1}$ , where  $q_\psi$  denotes the coordinate map on the tangent bundle induced by the frame  $\partial_i$  (which comes from the Morse chart. Compare Definition 5.0.14). By Lemma 5.0.15 together with Remark 5.0.17 we know that  $dX|_0 = J(\text{grad}(f))_\psi(0)$  is diagonal. Now we define

$$Y : \mathbb{R}^{\text{ind}_q} \supseteq B \rightarrow \mathbb{R}^{\text{ind}_q}, \quad v \mapsto Y(v) := \pi_u \circ (X_{g(v), v})$$

and Taylor-expand said function in the origin. By the chain rule and splitting of the tangent space at the origin we get that  $Y(v) = \Lambda v + R(v)$  with  $\Lambda = \text{diag}(2\mu_1, \dots, 2\mu_{\text{ind}_q})$  and  $\|R(v)\| = o(\|v\|)$ . Now

$$\frac{d}{dt}(u(t)) = \pi_u \circ q_\psi \circ \frac{d}{dt}\varphi_t(x) = \pi_u \circ q_\psi \circ -\text{grad}_{\varphi_t(x)}(f) = Y(u(t)).$$

Thus, with Cauchy-Schwarz we can calculate that for  $v$  close enough to zero:

$$\frac{d}{dt} \frac{1}{2} \|u(t)\|^2 = \underbrace{\langle u(t), \Lambda u(t) \rangle + \langle u(t), R(u(t)) \rangle}_{\geq -\|u(t)\| \cdot \underbrace{\|R(u(t))\|}_{\leq \mu_{\min} \|u(t)\|}} \geq 2\mu_{\min} \|u(t)\|^2 - \mu_{\min} \|u(t)\|^2 > 0.$$

Now choose  $r$  small enough such that the estimate holds and the ball  $B_r(0) \subseteq \psi_u(O)$ , and  $f > c$  on  $E_r := (\psi_u|_O)^{-1}(B_r(0))$ . Then:

- (i) For  $y \in \partial E_r$  and  $t > 0$  we have  $\varphi_{-t}(y) \in \text{int}(E_r)$ , since  $\psi_u(\varphi_{-t}(y))$  strictly decreases in  $t$ , by the estimate above.
- (ii) Every flow line in  $D_q \setminus \{q\}$  meets  $\partial E_r$  exactly once: It meets the boundary at least once, as it starts in  $q \in \text{int}(E_r)$  and ends in  $S(q \rightarrow)$  which is disjoint from  $E_r$ . If a flow line met the boundary at multiple times, there would be a time  $t_0$  where the flow line reenters  $E_r$ , which forces the  $t$  differential of said flow line in the chart  $\psi_u$  to be less than or equal to 0 (the norm needs to be non-increasing), which contradicts the estimate above.

Now we come to the homeomorphism: By (ii), the map

$$\Theta : \{(y, s) \mid y \in \partial E_r, 0 \leq s \leq \tau(y)\} \rightarrow D_q \setminus \text{int}(E_r), \quad \Theta(y, s) := \varphi_s(y)$$

is a continuous bijection from a compact set into a Hausdorff space and hence a homeomorphism. Now we define

$$G : D_q \rightarrow D_1 \subseteq \mathbb{R}^{\text{ind}_q}, \quad \begin{cases} G(x) := \frac{1}{2r} \psi_u(x) & \text{for } x \in E_r, \\ G(\varphi_s(y)) := \left(\frac{1}{2} + \frac{s}{2\tau(y)}\right) \frac{\psi_u(y)}{r} & \text{for } y \in \partial E_r, 0 \leq s \leq \tau(y). \end{cases}$$

Both formulas define something continuous (even smooth on their domain) and agree on  $\partial E_r$ . Hence  $G$  is continuous. Furthermore,  $G$  is bijective and the domain is compact, whilst the image is Hausdorff, giving us a homeomorphism that maps  $\partial D_q$  onto  $\partial D_1$ . Now let  $\overline{\pi_S} : S^{\text{ind}_q} \rightarrow \overline{\mathbb{R}^{\text{ind}_q}}$  denote the function from Definition 3.3.21 and with this define

$$g_q^{-1} := \overline{\pi_S^{-1}} \left( \frac{G}{1 - \|G\|} \right) : D_q / S(q \rightarrow) \rightarrow S^{\text{ind}_q} \quad \text{where } \frac{c}{0} := \infty. \text{ for } c \in \mathbb{R}$$

Now only the statement about the orientation preservation of  $g_q$  at a neighbourhood of  $q$  is missing: Define  $T_q := \max_{\partial E_r} \tau$ . If  $\tau(x) > T_q$  and  $x \neq q$ , let  $y$  be the point, where the flow line through  $x$  passes through the boundary  $\partial E_r$ . Then  $x = \varphi_{-t}(y)$  with  $t = \tau(x) - \tau(y) > 0$  and thereby  $x \in \text{int}(E_r)$  by (i). Hence,  $W_q \subseteq \text{int}(E_r)$  and in said interior of  $E_r$   $G$  is a diffeomorphism that is orientation preserving at  $q$  by the definition of  $\psi_u$  and by Lemma 4.1.17 we know that the sign of the determinant of a diffeomorphism of oriented manifolds is computed by a smooth procedure, and thereby it is locally constant, making  $G$  orientation preserving on the whole  $\text{int}(E_r)$ . The map




Figure 6: The setup of Lemma 6.2.5.

$\pi_S^{-1}$  is orientation preserving: This follows from the orientation of  $S^{\text{ind}_q}$  by Remark 4.4.3. Furthermore, the stretching  $v \mapsto \frac{v}{1-\|v\|}$  is orientation preserving where  $D_1$  is oriented as a subspace of  $\mathbb{R}^{\text{ind}_q}$  (this can be shown by looking at the sign of the determinant of its Jacobian). Hence the whole composition  $g_q^{-1}$  is orientation preserving and thereby also  $g_q$  is.  $\square$

lem:morseFilt  
r-orientation  
ClassOfADisk

**Lemma 6.2.5.** *Let  $n \geq 1$  and let  $(\Sigma, \partial\Sigma)$  be a compact pair ( $\partial\Sigma$  denotes a closed subspace of  $\Sigma$ , and not some kind of boundary). In applications of the Lemma, it will be a boundary) such that  $(\Sigma \setminus \partial\Sigma)$  is an oriented  $C^1$ -manifold of dimension  $n$ . Let  $g : S^n \rightarrow \Sigma / \partial\Sigma$  be a homeomorphism with  $g(S) = [\partial\Sigma]$  (where  $S$  denotes the south pole), and let  $U \subseteq S^n \setminus S$  be open such that  $g|_U$  is an orientation preserving  $C^1$ -diffeomorphism onto an open subset of  $\Sigma \setminus \partial\Sigma$ . Define*

$$o_\Sigma := (g^{-1})^*(\beta_n) \in \tilde{K}^{-n}(\Sigma / \partial\Sigma) = K^{-n}(\Sigma, \partial\Sigma),$$

*with  $\beta_n$  being the canonical generator defined in Definition 3.5.28. Let  $V$  be an oriented  $n$ -dimensional vector space with  $K$ -theoretic orientation  $\beta_V$  and  $h : \Sigma / \partial\Sigma \rightarrow \bar{V}$  be a pointed map such that for some  $z \in V$  we have that  $h^{-1}(z) = \{y\}$  with  $y \in g(U)$ ,  $h$  is  $C^1$  near  $y$  and  $dh|_y$  is invertible. Then*

$$h^*(\beta_V) = \text{sign det}(dh|_y) \cdot o_\Sigma$$

*Proof.* Let  $\kappa_V : V \rightarrow \mathbb{R}^n$  be the coordinate map of a positively oriented basis, meaning that  $(\overline{\pi_S^{-1}} \circ \overline{\kappa_V})^* (\beta_n) = \beta_V$  (compare Corollary 4.2.5). Then  $\xi := \overline{\pi_S^{-1}} \circ \overline{\kappa_V} \circ h \circ g : S^n \rightarrow S^n$  satisfies:

- (i)  $\xi^{-1}(\pi_S^{-1} \circ \kappa_V(z)) = \{g^{-1}(y)\}$ ,
- (ii)  $\xi$  is  $C^1$  near  $g^{-1}(y) \in U$ ,
- (iii) by the chain rule, Lemma 4.1.17 and since  $\pi_S^{-1}$ ,  $\kappa_V$  and  $g$  are orientation preserving we have:

$$\text{sign det}(\text{d}\xi|_{g^{-1}(y)}) = \text{sign det}(\text{d}h|_y).$$

Hence by the Single Regular Preimage Lemma 4.4.5 applied to  $\xi^*$  we can calculate

$$g^* h^* (\beta_V) = \xi^* (\beta_n) = \text{sign det}(\text{d}h|_y) \cdot \beta_n.$$

Finally we can apply  $(g^{-1})^*$  from the left to finish the proof.  $\square$

**Remark 6.2.6.** We can apply the Lemma above to the pair  $(D_q, S(q \rightarrow))$  together with  $g = g_q$ , and  $U = U_q$  from Lemma 6.2.4 (c).

lem:filt-orientatedTubularEmbedding


**Lemma 6.2.7** (Oriented Tubular Embedding). *Let  $B_p := W(\rightarrow p) \cap M^c$  and let  $J : D_\varepsilon(\mathcal{N}) \rightarrow N_p$  be the tubular embedding from Lemma 6.2.3, where  $\mathcal{N} \rightarrow B_p$  denotes the normal bundle. Now we can orient the normal bundle via the bundle isomorphism  $l : \mathcal{N} \rightarrow Q^p$  where  $Q^p$  denotes the quotient bundle from Definition 6.2.2 (the map is given by including the normal bundle into the tangent bundle and projecting then onto the quotient bundle). Then  $J$  can be chosen to be fibrewise orientation preserving, meaning that with the linear identification  $R_0 : \mathcal{N}_x \rightarrow T_0 \mathcal{N}_x$  and the projection bundle map  $pr : TM|_{W(\rightarrow p) \cap M^c} \rightarrow Q^p$  we have that for all  $x$  the map*

$$pr \circ \text{d}J_x|_0 \circ R_0 : \mathcal{N}_x \rightarrow Q_x^p$$

*is orientation preserving.*

*Proof.* This follows directly from  $pr \circ \text{d}J_x|_0 \circ R_0 : \mathcal{N}_x \rightarrow Q_x^p$  being a bundle isomorphism. Hence it either is orientation preserving in all fibers, or reversing in all fibers (due to the connected base space). If the latter happens, we redefine  $J$  by composing it with a map  $\rho$  that reflects, say the first coordinate: Here we choose a global orthonormal trivialisation of  $\mathcal{N}$  (which exists since the base space  $B_p$  is contractible) such that  $\rho$  maps the domain of the tubular embedding isomorphically onto itself.  $\square$

**Remark 6.2.8.** For a fiber  $V_j$  of the tubular neighbourhood from Lemma 6.2.3 with center  $x_j$ , we can apply the Lemma 6.2.5 with  $U = S^n \setminus \{S\}$  and  $g_j$  given by the oriented tubular embedding composed with the inverse of  $v \mapsto \pi_S^{-1}\left(\frac{v}{1-\|v\|}\right)$ . Then  $g_j(N) = x_j$  for

Figure 7: The definition of the transverse  $\Sigma$  from Lemma 6.2.9.

the north pole  $N$  and  $h_j := \overline{\kappa_V^{-1}} \circ \overline{\pi_S} \circ g_j^{-1}$  is a map where we can apply the Lemma 6.2.5 above with  $h^{-1}(0) = \{x_j\}$  to conclude that  $h_j^*(\beta_V) = o_{V_j}$ . (It is easy to see that  $h_j$  is locally a diffeomorphism and  $h$  as invertible differential in the origin.)

lem:morseFilt  
-canonicalGen  
eratorsOnTran  
sversalDisks

**Lemma 6.2.9.** *Let  $q \in \text{Crit}(f)$  with  $n := \text{ind}(q) \geq 1$ , let  $(N, L)$  be an index pair of  $\{q\}$  and let  $\Sigma \subseteq N$  with  $\partial\Sigma \subseteq L$  be as in Lemma 6.2.5. Assume*

- (i)  $\Sigma \cap W(\rightarrow p) = \{x\}$  with  $x \in g(U)$  and  $T_x\Sigma \oplus T_xW(\rightarrow p) = T_xM$ , ( $\Sigma$  is transversal)
- (ii) for every  $s \geq 0$  there is a neighbourhood  $U_s$  of  $x$  in  $\Sigma$  with  $\varphi_{[0,s]}(U_s) \subseteq N \setminus L$ .

*Orient  $\Sigma \setminus \partial\Sigma$  such that the isomorphism  $T_x\Sigma \rightarrow Q_x^p$  is orientation preserving. Then for the inclusion  $i : (\Sigma, \partial\Sigma) \rightarrow (N, L)$*

$$i^* \left( \alpha_q^{(N,L)} \right) = o_\Sigma.$$

**Remark 6.2.10.** The two cases where we apply said lemma are:

1.  $\Sigma = D_q$ ,  $x = q$ ,  $(N, L) = (C, B)$  (from Lemma 6.2.4) where  $Q_q^q = T_qM/T_q^sM \cong T_q^uM$ , so  $o_{D_q}$  is taken with respect to the orientation of  $T_q^uM$ .
2.  $\Sigma = V_j$   $x = x_j$  and  $(N, L) = (B, A)$ .

*Proof of Lemma 6.2.9.* We prove the Lemma in three steps. First we restricted the statement to the easier case, where  $\Sigma \subseteq W(q \rightarrow)$ . Then we define a new chart such that the stable manifold is not just a graph of a function, but a linear subspace. This makes sure

that the projection onto the unstable coordinates collapses the stable part to a point which finally in step 3 generalizes the statement for arbitrary  $\Sigma$ :

*Step 1.* Let  $\chi = (\chi_s, \chi_u)$  be a chart around  $q$  such that

1.  $W(q \rightarrow)$  is the graph of function over the unstable coordinates,
2. the box  $(N_\chi, L_\chi)$  defined as

$$(\chi^{-1}(D_\varepsilon^s \times D_\varepsilon^u), \chi^{-1}(D_\varepsilon^s \times \partial D_\varepsilon^u))$$

is a regular index pair of  $\{q\}$ .

3.  $\varepsilon$  is small enough such that  $W(q \rightarrow) \cap N_\chi$  is the graph of the function from above.
4.  $d\chi_q = d\psi_q$  where  $\psi$  is a refined Morse chart.

The existence of such a chart follows from the local unstable manifold theorem 5.0.20. Let  $\kappa_\chi$  be the map defined as

$$\kappa_\chi := d\chi^{-1}|_0 \circ R_0 \circ s \circ \text{pr}_u \circ \chi$$

similar to the construction of  $\kappa_q$  in Remark 6.1.18. Let  $\Sigma$  be a disk contained in  $W(q \rightarrow)$  and hence  $\Sigma \cap W(\rightarrow q) = q$ . Near  $q$  and for large  $t > 0$  the homotopy equivalence  $c : (N, L) \rightarrow (N_\chi, L_\chi)$  is given by  $y \mapsto [\varphi_{3t}(y)]$  (Compare Theorem 6.1.4 and notice that  $q$  is in the interior of both  $(N \setminus L)$  and  $(N_\chi \setminus L_\chi)$ ). Define  $i : (\Sigma, \partial\Sigma) \rightarrow (N, L)$  to be the inclusion. Then consider the function

$$h := \underbrace{d\chi^{-1}|_0 \circ R_0 \circ s \circ \text{pr}_u \circ \chi \circ c \circ i}_{\kappa_\chi} : (\Sigma/\partial\Sigma) \rightarrow (\overline{T_q^u M})$$

If  $h(y) = 0$ , then  $y$  cannot be  $[\partial\Sigma]$  as that point end at infinity. By definition of  $\kappa_\chi$  we know that  $\chi \circ c \circ i(y) \in D_\varepsilon^s \times \{0\}$ . And since  $y \in \Sigma \subseteq W(q \rightarrow)$ , and  $c$  restricts to a map from  $W(q \rightarrow)$  to  $W(q \rightarrow)$  (restricted to the right domain, and image), we know that  $\chi \circ c \circ i(y) \in \chi(W(q \rightarrow)) \cap D_\varepsilon^s \times \{0\}$ . Finally, we assumed that  $W(q \rightarrow)$  was the graph of the unstable coordinates in  $\chi$  and hence, there is only one  $x \in D_\varepsilon^s \times \{0\} \cap \chi(W(q \rightarrow))$ , which is  $(0, 0)$ . Hence  $\varphi_{3t}(y) = q$  which is a fixpoint of the flow and thereby,  $y = q$  and 0 has a unique preimage under  $h$ . Furthermore,  $dh|_q$  is an endomorphism on  $T_q^u M$ . We can calculate its determinant by calculating the jacobian determinants of  $c \circ i$  and  $\kappa_\chi$  in  $q$  (resp. in the origin) in a refined Morse chart seperately:

1. The function  $d(c \circ i)|_q$  (restricted in its image to  $T_q^u M$ ) has positive determinant which follows from the flow in a refined morse chart in a critical point restricted to the unstable space has a differential that is diagonal with positive entries. This was calculated in Remark 5.0.17.

2. The function  $\kappa_\chi$  can be differentiated in  $q$ :

$$d\kappa_\chi = d\left(d\chi^{-1}|_0 \circ R_0 \circ s \circ \text{pr}_u \circ \chi\right)$$

where  $\iota := d\chi^{-1}|_0 \circ R_0 : \{0\} \times \mathbb{R}^n \rightarrow T_q^u M$  is a linear isomorphism (since  $d\chi|_q$  maps  $T_q^u M$  onto the unstable coordinate plane) and  $s$  is the radial stretching with  $ds|_0 = \lambda \text{id}$ , where  $\lambda > 0$ . On  $T_q^u M$  the map  $\text{pr}_u \circ d\chi|_q$  is the inverse of  $\iota$ , so, identifying  $T_0 T_q^u M$  with  $T_q^u M$ ,

$$dh|_q = \lambda d\varphi_{3t}|_q \Big|_{T_q^u M} \in \text{End}(T_q^u M).$$

Its determinant is again positive, as the flow restricted to the unstable tangent space of a critical point is diagonal with positive entries.

Hence,  $dh|_q$  has positive determinant and Lemma 6.2.5 gives us the equality

$$i^* \circ c^* \circ \kappa_\chi^*(\beta_{T_q^u M}) = h^*(\beta_{T_q^u M}) = o_\Sigma.$$

This concludes the proof for the special case, where  $\Sigma$  is contained in  $(q \rightarrow)$ .

So far we have seen that for any chart  $\chi$  that satisfies the properties from Step 1 there is the equality above. If we choose  $\chi$  to be a refined morse chart, then  $\kappa_\chi^*(\beta_{T_q^u M}) = \alpha_q^{(N_\chi, L_{chi})}$  is a canonical generator and a member of  $(\alpha_q)_\#$ .

*Step 2* We know by the local unstable manifold theorem, that for our refined morse chart  $\psi$  there exists a smooth  $g_s$  such that  $\psi(W(\rightarrow q))$  is the graph of  $g_s$  with  $g_s(0) = 0$  and  $dg_s|_0 = 0$ . The chart  $\tilde{\psi} := (\psi_s, \psi_u - g_s \circ \psi_s)$  satisfies  $d\tilde{\psi}|_q = d\psi|_q$ ,  $W(\rightarrow q) = \tilde{\psi}^{-1}(\mathbb{R}^{m-n} \times \{0\})$  (locally),  $W(q \rightarrow)$  is still the graph of a function over the unstable coordinates and by Remark 6.1.17, for small  $\varepsilon$ , its box  $(N_{\tilde{\psi}}, L_{\tilde{\psi}})$  is an index pair. We claim now, that  $\kappa_{\tilde{\psi}}^*(\beta_{T_q^u M})$  is a member of the canonical generator  $(\alpha_q)_\#$ . Let  $E := W(q \rightarrow) \cap N_\psi$ , be the unstable disk in the box of  $\psi$ . In the chart  $\psi$ ,  $E$  is the graph of a map

$$e : D_\varepsilon^u \rightarrow D_\varepsilon^s$$

Now we define the deformation homotopy for  $t \in [0, 1]$

$$H_t(l, x) := ((1-t)l + te(x), x)$$

that shows that  $(E, \partial E)$  is a strong deformation retract of  $(N_\psi, L_\psi)$  onto and hence induces an isomorphism  $K^{-n}(N_\psi, L_\psi) \rightarrow K^{-n}(E, \partial E)$ . Now we apply Step 1 to the two charts  $\chi = \psi$  and  $\chi = \tilde{\psi}$  with  $\Sigma = E$ , and  $(N, L) = (N_\psi, L_\psi)$ . This tells us that pulling back the canonical generator along both paths in the diagram

$$\begin{array}{ccccc} & & (N_{\tilde{\psi}}, L_{\tilde{\psi}}) & & \\ & \nearrow c_1 & & \searrow \kappa_{\tilde{\psi}} & \\ (E, \partial E) & \xleftarrow{i} & (N_\psi, L_\psi) & & (\overline{T_q^u M}, \{\infty\}) \\ & \searrow c_2 \simeq \text{id} & & \nearrow \kappa_\psi = \kappa_q & \\ & & (N_\psi, L_\psi) & & \end{array}$$

Bild  
einfügen

induce the same  $o_E$ . Now  $c_2 \simeq \text{id}$  and the inclusion  $i$  induces an isomorphism by the deformation above. Hence,

$$c_1^* \circ \kappa_{\tilde{\psi}}^*(\beta_{T_q^u M}) = \kappa_q^*(\beta_{T_q^u M}) = \alpha_q^{(N_{\psi}, L_{\psi})}.$$

But the canonical generators are compatible with the maps  $c$  and  $c_1^*$  is an isomorphism, we know that  $\alpha_q^{(N_{\tilde{\psi}}, L_{\tilde{\psi}})}$  is the unique element with  $c_1^*(\alpha_q^{(N_{\tilde{\psi}}, L_{\tilde{\psi}})}) = \alpha_q^{(N_{\psi}, L_{\psi})}$ . Thereby

$$\kappa_{\tilde{\psi}}^*(\beta_{T_q^u M}) = \alpha_q^{(N_{\tilde{\psi}}, L_{\tilde{\psi}})}.$$

*Step 3* Now we assume  $\Sigma$  to just satisfy the Lemmas conditions, use again  $\tilde{\psi}$  and define the map on the quotients

$$h := \underbrace{d\tilde{\psi}^{-1}|_0 \circ R_0 \circ s \circ \text{pr}_u \circ \tilde{\psi} \circ c \circ i}_{\kappa_{\tilde{\psi}}}$$

with  $c : (N, L) \rightarrow (N_{\tilde{\psi}}, L_{\tilde{\psi}})$  being the canonical homotopy equivalence as in Step 1. If  $h(y) = 0$ , then similar to before, we argue, that  $h(y) = 0$  means that  $\tilde{\psi} \circ c \circ i(y) \in D_{\varepsilon}^s \times \{0\}$  which in turn means that  $c(y) = \varphi_{3t}(y) \in \tilde{\psi}^{-1}(D_{\varepsilon}^s \times \{0\})$ . By definition of  $\tilde{\psi}$  we made sure that  $W(\rightarrow q) \supseteq \tilde{\psi}^{-1}(D_{\varepsilon}^s \times \{0\})$ . Thereby, our  $y \in \Sigma \cap W(\rightarrow q) = \{x\}$  by requirements on  $\Sigma$ . By (ii) and since  $\varphi_{[t, 3t]}(x)$  are close to  $q$  for large  $t$  our map  $h$  equals  $h = \kappa_{\tilde{\psi}} \circ \varphi_{3t}$  near  $x$ . Due to the projection  $\text{pr}_u$ ,  $\kappa_{\tilde{\psi}}$  is constant on  $W(\rightarrow q)$  (for a small enough box), and the flow  $\varphi_{3t}$  preserves  $W(\rightarrow q)$ . Thereby, the differential factors through the kernel of  $d(\text{pr}_u \circ \tilde{\psi})_{\varphi_{3t}(x)} = T_{\varphi_{3t}(x)} W(\rightarrow q)$ . And since  $\Sigma$  intersects  $W(\rightarrow q)$  transversally we factor through the composition

$$\begin{array}{ccc} T_x \Sigma & \xrightarrow{dh|_x} & T_q^u M \\ \cong \downarrow & & \uparrow [d\kappa_{\tilde{\psi}}] \\ Q_x^q & \xrightarrow{[d\varphi_{3t}]} & Q_{\varphi_{3t}(x)}^q \end{array}$$

Now the first isomorphism is orientation preserving, by the choice of orientation on  $\Sigma$ .

Since  $\varphi_s$  preserves  $W(\rightarrow q)$ , its differential induces a bundle morphism  $[d\varphi_s] : Q^q \rightarrow Q^q$  over  $\varphi_s|_{W(\rightarrow q)}$  which is an isomorphism on every fibre. As  $W(\rightarrow q)$  is connected, by Proposition 4.1.12 it suffices to check that  $[d\varphi_s]$  is orientation preserving at a single point. We take  $q$ : since  $\varphi_s(q) = q$ , the map  $[d\varphi_s]_q$  is an endomorphism of  $Q_q^q$ , and with respect to the basis  $([\partial_i(q)])_i$  given by the unstable coordinate vectors of the refined Morse chart it is diagonal with entries  $e^{2\mu_i s} > 0$  by Remark 5.0.17. Hence it has positive determinant, and  $[d\varphi_{3t}]$  is orientation preserving (note  $3t > 0$ ).

Similarly,  $\kappa_{\tilde{\psi}}$  is constant 0 on the local stable disk  $W(\rightarrow q) \cap N_{\tilde{\psi}}$ , and its differential induces a bundle morphism  $[d\kappa_{\tilde{\psi}}]$  from  $Q^q|_{W(\rightarrow q) \cap N_{\tilde{\psi}}}$  to the oriented vector space  $T_q^u M$  (a bundle over the point 0). It is an isomorphism on every fibre, since  $\text{pr}_u \circ d\tilde{\psi}|_y$  is

surjective with kernel  $T_y W(\rightarrow q)$  and  $\tilde{\psi}_u(y) = 0$ . The local stable disk is connected, and at  $y = q$ , under the identification  $Q_q^q = T_q M / T_q^s M \cong T_q^u M$  by which  $Q^q$  is oriented (Proposition 5.1.5),  $[d\kappa_{\tilde{\psi}}]_q$  is the endomorphism  $\lambda \text{id}$  of  $T_q^u M$  with  $\lambda > 0$ . Hence  $[d\kappa_{\tilde{\psi}}]$  is orientation preserving, and so is the composition  $dh|_x$ .

Lemma 6.2.5 gives us

$$o_\Sigma = h^*(\beta_{T_q^u M}) = i^* \circ c^* \circ \kappa_{\tilde{\psi}}^*(\beta_{T_q^u M})$$

Then, Step 2 gives us that  $\kappa_{\tilde{\psi}}^*(\beta_{T_q^u M}) = \alpha_q^{(N_{\tilde{\psi}}, L_{\tilde{\psi}})} \in (\alpha_q)_\#$ . Thereby  $c^* \circ \kappa_{\tilde{\psi}}^*(\beta_{T_q^u M}) = \alpha_q^{(N, L)}$  by definition. This finally concludes the proof.  $\square$

lem:morseFilt  
-connectingHo  
morphismAlongCollar

**Lemma 6.2.11** (Connecting Homomorphism along a Collar). *Let  $(X, Y)$  be a compact pair (such that the inclusion  $Y \rightarrow X$  is a cofibration) and  $X$  well pointed by  $y_0 \in Y$ . Let  $s : X \rightarrow [0, 1]$  be continuous with  $s^{-1}(0) = Y$  and let  $r : s^{-1}([0, 1]) \rightarrow Y$  be a continuous retraction. Define, with  $S^1 = I/\partial I$ ,*

$$\bar{k} : X/Y \rightarrow \Sigma Y = S^1 \wedge Y, \quad \bar{k}([z]) := \begin{cases} [1 - s(z), r(z)] & \text{if } s(z) < 1, \\ * & \text{if } s(z) = 1. \end{cases}$$

*Then for all  $n \geq 1$  the connecting homomorphism  $\delta : \tilde{K}^{-n}(Y) \rightarrow K^{-n+1}(X, Y)$  of Definition 3.5.9 satisfies*

$$\delta = (-1)^n \cdot \bar{k}^* \circ \sigma_n,$$

*where  $\sigma_n := (\varphi_{n-1,1} \wedge \text{id}_Y)^* : \tilde{K}^{-n}(Y) = \tilde{K}(S^n \wedge Y) \rightarrow \tilde{K}(S^{n-1} \wedge S^1 \wedge Y) \cong \tilde{K}^{-(n-1)}(\Sigma Y)$  is the identification via the map  $\varphi_{l,r} : S^l \wedge S^r \rightarrow S^{l+r}$  from Definition 3.3.27 together with the associativity of the wedge product and  $\bar{k}^*$  denotes  $(\text{id}_{S^{n-1}} \wedge \bar{k})^*$ .*

*Proof.* Let  $n = 1$ . With the notation of Definition 3.5.9 let

(i)  $\bar{m} : X \cup CY \rightarrow (X \cup CY)/X \cong CY/Y$

(ii)  $l : CY/Y \rightarrow \Sigma Y$ , given by  $[y, t] \mapsto [t, y]$  as in Proposition 3.5.7.

(iii)  $\delta = (\pi^*)^{-1} \circ \bar{m}^* \circ l^* \circ \sigma_1$ . This is the same map as in Definition 3.5.9. We split  $\theta$  into two parts and moved one part to our definition of  $\bar{m}$  (The part that used to be the map  $T : (X \cup CY)/(X) \cong CY/Y$ ) and the second part is our map  $l$ .

Now for  $\mu \in [0, 1]$  we define the map  $H_\mu : X \cup CY \rightarrow \Sigma Y$  by

$$H_\mu([y, t]) := [(1 - \mu)t, y] \text{ on } CY, \quad H_\mu(z) := [(1 - \mu) + \mu s(z), r(z)] \text{ on } X$$

(and  $*$  if  $s(z) = 1$ ). The map  $H_\mu$  is well-defined as both formulas agree on  $Y$ , continuous (for all, and in  $\mu$ ) as  $s(z) \rightarrow 1, H_\mu(z) \rightarrow [\partial I]$  and pointed. Furthermore,  $H_0 = l \circ \bar{m}$ :

- On  $X$ :  $l \circ \overline{m}(x) = *$  and  $H_0(x) = [1, r(x)] = *$
- On  $CY$ :  $l \circ \overline{m}([y, t]) = [t, y]$  and  $H_0([y, t]) = [t, y]$ ,

and  $H_1 = (R \wedge 1) \circ \overline{k} \circ \pi$  with the Reflection  $R(t) = (1 - t)$  and the quotient map  $\pi : X \cup CY \rightarrow X/Y$ :

- On  $CY$ :  $(R \wedge 1) \circ \overline{k} \circ \pi([y, t]) = (R \wedge 1) \circ \overline{k}(\underbrace{[y, t]}_{=s^{-1}(0)}) = (R \wedge 1)([1, \underbrace{y'}_{y' \in Y}]) = [0, y'] = [0, y] = H_1([y, t])$
- On  $X$ :  $(R \wedge 1) \circ \overline{k} \circ \pi(x) = (R \wedge 1)[1 - s(x), r(x)] = [s(x), r(x)] = H_1(x)$ .

Hence, by the homotopy invariance and Corollary 3.4.16:

$$\overline{m}^* \circ l^* = \pi^* \circ \overline{k}^* \circ (R \wedge 1)^* = -\pi^* \circ \overline{k}^* \quad \Rightarrow \quad \delta = -\overline{k}^* \circ \sigma_1.$$

For  $n \geq 2$ ,  $\delta$  is by definition the connecting homomorphism of  $(S^{n-1} \wedge X, S^{n-1} \wedge Y)$ . Define the map  $H'_\mu : (S^{n-1} \wedge X) \cup C(S^{n-1} \wedge Y) \rightarrow S^1 \wedge S^{n-1} \wedge Y$ :

$$H'_\mu([v, y], t) := [(1 - \mu)t, v, y], \quad H'_\mu([v, x]) := \begin{cases} [(1 - \mu) + \mu s(x), v, r(x)] & \text{if } s(x) \neq 1 \\ * & \text{if } s(x) = 1 \end{cases}$$

Similar to  $H_\mu$ , the map  $H'$  is a pointed homotopy from  $l \circ \overline{m}$  to  $(R \wedge 1) \circ \overline{k}_n \circ \pi$  where

$$\overline{k}_n : S^{n-1} \wedge (X/Y) \rightarrow S^1 \wedge S^{n-1} \wedge Y, [v, x] \mapsto [1 - s(x), v, r(x)],$$

so  $\delta = -\overline{k}_n^* \circ (\varphi_{1,n-1} \wedge \text{id}_Y)^*$ . Now  $\overline{k}_n = (P \wedge \text{id}_Y) \circ (\text{id}_{S^{n-1}} \wedge \overline{k})$  where  $P : S^{n-1} \wedge S^1 \rightarrow S^1 \wedge S^{n-1}$  is given by swapping the factors. Now consider the following commuting diagram

$$\begin{array}{ccc} S^{n-1} \wedge S^1 \wedge Y & \xrightarrow{P \wedge \text{id}_Y} & S^1 \wedge S^{n-1} \wedge Y \\ \varphi_{n-1,1} \wedge \text{id}_Y \downarrow & & \downarrow \varphi_{1,n-1} \wedge \text{id}_Y \\ S^n \wedge Y & \xrightarrow{B} & S^n \wedge Y \\ \overline{\pi}_S \wedge \text{id}_Y \downarrow & & \downarrow \overline{\pi}_S \wedge \text{id}_Y \\ \overline{\mathbb{R}}^n \wedge Y & \xrightarrow{\overline{A} \wedge \text{id}_Y} & \overline{\mathbb{R}}^n \wedge Y \end{array}$$

The middle arrow  $B := (\varphi_{1,n-1} \circ P \circ (\varphi_{n-1,1})^{-1}) \wedge \text{id}_Y$  is given by going around the top and commutes by definition. The bottom square commutes by definition of  $A$  together with the naturality of the map  $\Xi$  that builds up the  $\varphi_{l,r} = (\overline{\pi}_S^{l+r})^{-1} \circ \Xi^{-1} \circ (\overline{\pi}_S^l \wedge \overline{\pi}_S^r) = (\overline{\pi}_S^{l+r})^{-1} \circ (\pi_S^l \times \pi_S^r) \circ \Xi^{-1}$ . (The notation and definition is from Definition 3.3.27) (We omit the identification of  $\mathbb{R}^l \times \mathbb{R}^r = \mathbb{R}^{l+r}$ ). Now  $B = B' \wedge \text{id}_Y$  with

$$B' = (\overline{\pi}_S^n)^{-1} \circ \underbrace{(\pi_S^{n-1} \times \pi_S^1) \circ \Xi^{-1} \circ P \circ \Xi \circ (\pi_S^1 \times \pi_S^{n-1})^{-1}}_{A: \mathbb{R}^n \rightarrow \mathbb{R}^n} \circ (\overline{\pi}_S^n)$$

Finally the linear map  $A$  is given by moving the last coordinate to the front and hence has determinant  $(-1)^{n-1}$ . Hence, by Lemma 4.2.3 we conclude that  $(\bar{A} \wedge \text{id}_Y)^* = (-1)^{n-1} \cdot \text{id}$ . Hence  $B^* = (\pi_{\bar{S}} \wedge \text{id}_Y)^* \circ (-1)^{n-1} \cdot \text{id} \circ (\pi_{\bar{S}} \wedge \text{id}_Y)^{* -1} = (-1)^{n-1} \cdot \text{id}$ , as it is the conjugation of  $(\bar{A} \wedge \text{id}_Y)^*$  via an isomorphism (Alternatively one could say that  $\pm \text{id}$  commutes with group homomorphisms). As the top square commutes we have

$$(P \wedge \text{id}_Y)^* \circ (\varphi_{1,n-1} \wedge \text{id}_Y)^* = (-1)^{n-1} (\varphi_{n-1,1} \wedge \text{id}_Y)^*.$$

Putting everything together we conclude that

$$\begin{aligned} \delta &= -\overline{k_n}^* \circ (\varphi_{1,n-1} \wedge \text{id}_Y)^* \\ &= -(\text{id}_{S^{n-1}} \wedge \bar{k})^* \circ \underbrace{(P \wedge \text{id}_Y)^* \circ (\varphi_{1,n-1} \wedge \text{id}_Y)^*}_{=(-1)^{n-1} (\varphi_{n-1,1} \wedge \text{id}_Y)^*} \\ &= (-1)^n (\text{id}_{S^{n-1}} \wedge \bar{k})^* \circ \underbrace{(\varphi_{n-1,1} \wedge \text{id}_Y)^*}_{\sigma_n} \end{aligned}$$

□

**Lemma 6.2.12.** *Let  $V$  be an oriented real vector space of dimension  $n \geq 1$ , let  $g : (0, 1) \rightarrow \mathbb{R}$  be an orientation preserving diffeomorphism where  $(0, 1)$  is oriented as a subbase of  $\mathbb{R}$  (e.g.  $g(t) := \tan(\pi(t - \frac{1}{2}))$ ) and*

$$G_V : \Sigma \bar{V} = S^1 \wedge \bar{V} \rightarrow \mathbb{R} \oplus \bar{V}, \quad [t, v] \mapsto (g(t), v).$$

*Then  $G_V$  is a homeomorphism by Lemma 3.3.24. We can orient  $\mathbb{R} \oplus V$  by putting  $\mathbb{R}$  in front. Let  $b = \beta \boxtimes (-)$  be the Bott map. Then there is an  $e_n \in \{\pm 1\}$ , depending only on  $n$ , with*

$$b(\sigma_n(\beta_V)) = e_n \cdot G_V^*(\beta_{\mathbb{R} \oplus V}) \in \tilde{K}^{-(n+1)}(\Sigma \bar{V}),$$

*where  $\sigma_n := (\varphi_{n-1,1} \wedge \text{id}_Y)^* : \tilde{K}^{-n}(Y) = \tilde{K}(S^n \wedge Y) \rightarrow \tilde{K}(S^{n-1} \wedge S^1 \wedge Y) \cong \tilde{K}^{-(n-1)}(\Sigma Y)$  similar to Lemma 6.2.11.*

*Proof.* Both sides of the equation are generators of  $\tilde{K}^{-(n+1)}(\Sigma \bar{V}) \cong \mathbb{Z}$ , as isomorphisms map generators to generator and the isomorphism to  $\mathbb{Z}$  follows from  $\Sigma \bar{V} \simeq S^{n+1}$ . Hence both sides agree up to a sign  $e(V)$ :

$$b(\sigma_n(\beta_V)) = e(V) G_V^*(\beta_{\mathbb{R} \oplus V}) \in \tilde{K}^{-(n+1)}(\Sigma \bar{V})$$

In the notation of Lemma 3.3.24, the map  $G_V$  is given by  $\Xi^{-1} \circ (\bar{g} \wedge \text{id}_{\bar{V}})$ , where  $\bar{g}$  is the extension  $I/\partial I \rightarrow \bar{R}$  of  $g$ . Let  $A$  be an isomorphism  $V \rightarrow W$ . Then using the naturality

of  $\Xi$  (we omit the indices of  $\Xi$ ) and that the identity commutes with all maps we can conclude that:

$$\begin{aligned}
 G_W \circ (\text{id} \wedge \overline{A}) &= \Xi^{-1} \circ (\overline{g} \wedge \text{id}_{\overline{W}}) \circ (\text{id} \wedge \overline{A}) \\
 &= (\overline{g} \times \text{id}_W) \circ \Xi^{-1} \circ (\text{id} \wedge \overline{A}) \\
 &= (\overline{g} \times \text{id}_W) \circ (\text{id} \times A) \circ \Xi^{-1} \\
 &= (\text{id} \oplus A) \circ (\overline{g} \times \text{id}_V) \circ \Xi^{-1} \\
 &= (\text{id} \oplus A) \circ \underbrace{\Xi^{-1} \circ (\overline{g} \wedge \text{id}_V)}_{G_V}
 \end{aligned}$$

Now by Proposition 4.2.6 we know that  $\overline{A}^* \beta_W = \text{sign}(\det(A)) \beta_V$  and  $\overline{\text{id} \oplus A}^* (\beta_{\mathbb{R} \oplus W}) = \text{sign} \det(A) \beta_{\mathbb{R} \oplus V}$ . Now  $b$  and  $\sigma_n$  are natural and thereby commute with topological induced maps. Now we pull back the statement via  $(\text{id} \wedge \overline{A})^*$ :

$$\begin{aligned}
 (\text{id} \wedge \overline{A})^* \circ b \circ \sigma_n (\beta_W) &= b \circ \sigma_n \circ (\overline{A}^* \beta_W) = \text{sign} \det(A) \cdot b \circ \sigma_n (\beta_V) \\
 (\text{id} \wedge \overline{A})^* \circ G_W^* (\beta_{\mathbb{R} \oplus W}) &= (G_W \circ (\text{id} \wedge \overline{A}))^* \beta_{\mathbb{R} \oplus W} = G_V^* \circ (\overline{\text{id} \oplus A})^* \beta_{\mathbb{R} \oplus W} \\
 &= \text{sign}(\det(A)) \cdot G_V^* \beta_{\mathbb{R} \oplus V}
 \end{aligned}$$

I now claim that for fixed  $n$  the defined  $e(V)$  agrees with  $e(W)$  (and is thereby independent of the choosen vector space). For this we consider the diagram below, that commutes by the considerations above (naturality of  $b$  and  $\sigma$  and  $\Xi$ ):

$$\begin{array}{ccccc}
 \tilde{K}^{-n}(\overline{W}) & \xrightarrow{b \circ \sigma_n} & \tilde{K}^{-(n+1)}(\Sigma \overline{W}) & \xleftarrow{G_W^*} & \tilde{K}^{-(n+1)}(\overline{\mathbb{R} \oplus W}) \\
 \overline{A}^* \downarrow & & \downarrow (\text{id} \wedge \overline{A})^* & & \downarrow \overline{\text{id} \oplus A}^* \\
 \tilde{K}^{-n}(\overline{V}) & \xrightarrow{b \circ \sigma_n} & \tilde{K}^{-(n+1)}(\Sigma \overline{V}) & \xleftarrow{G_V^*} & \tilde{K}^{-(n+1)}(\overline{\mathbb{R} \oplus V})
 \end{array}$$

This lets us compute:

$$\begin{aligned}
 (\text{id} \wedge \overline{A})^* \circ (b \circ \sigma_n) (\beta_W) &= \text{sign} \det(A) \cdot \underbrace{b \circ \sigma_n (\beta_V)}_{e(V) G_V^* (\beta_{\mathbb{R} \oplus V})} \\
 &= e(V) \cdot \underbrace{\text{sign} \det(A) \cdot G_V^* (\beta_{\mathbb{R} \oplus V})}_{(\text{id} \wedge \overline{A})^* \circ G_W^* (\beta_{\mathbb{R} \oplus W})} \\
 &= e(V) \cdot (\text{id} \wedge \overline{A})^* \circ G_W^* (\beta_{\mathbb{R} \oplus W})
 \end{aligned}$$

Now applying  $((\text{id} \wedge \overline{A})^*)^{-1}$  from the left (which exists as it is induced by a homeomorphism) gives us that the sign  $e(V)$  satisfies the definition of  $e(W)$  making them equal. Thereby  $e(V)$  is invariant of  $V$  and only depends on  $n$ .  $\square$

*Proof of Theorem 6.2.1.* For all  $l \in \mathbb{Z}$  we get the commuting square

$$\begin{array}{ccc}
K_{\#}^{-(k-1)-l}(\mathcal{IP}_{k-1}) & \xrightarrow{\Delta^k} & K_{\#}^{-(k)-l}(\mathcal{IP}_k) \\
\downarrow b & & \downarrow b \\
K_{\#}^{-(k-1)-l-2}(\mathcal{IP}_{k-1}) & \xrightarrow{\Delta^k} & K_{\#}^{-(k)-l-2}(\mathcal{IP}_k)
\end{array}$$

Hence, we only need to proof the original statement for  $l = 0, 1$ . For  $l = 1$  the bottom row of the diagram 36 is trivial, as the connected simple systems  $\mathcal{IP}_k$  consists of spaces that are homotopically equivalent to  $k$ -spheres and we know by 3.5.27 that  $K^{-k-1}(S^k) \cong \tilde{K}(S^{2k+1}) = 0$ . The same argument works for the image of  $\Delta^k$  for  $l = 1$ . So for the rest of the proof we assume  $l = 0$ . Now since  $N_p$  is a tubular neighbourhood of  $W(\rightarrow p) \cap M^c$  we know that  $N_p \cap S(q \rightarrow)$  consists of finitely many components  $V_j$  and each component has one  $\{x_j\} = W(q \rightarrow p) \cap V_j$ . In fact, the proof of Lemma ?? shows that the  $V_j$  are fibers of the normal neighbourhood of  $W(\rightarrow p)$  inside  $W(q \rightarrow)$  and hence homeomorphic to  $\mathbb{R}$ . Now we take the tuple  $(C, B, A)$  from Lemma ??. We have a map of triples:

referenz!

$$\left( W(q \rightarrow) \cap N_q, S(q \rightarrow), S(q \rightarrow) \setminus \bigcup_j \text{int}(V_j) \right) \rightarrow (C, B, A)$$

giving us a commuting diagramm (the square brackets for the input of the K functors  $K[-]$  are choosen for readability and have no deeper meaning)

$$\begin{array}{ccc}
K^{-\lambda_p}[B, A] & \xrightarrow{\delta_{triple}^1} & K^{-\lambda_q}[C, B] \\
\downarrow i_{B,A}^* & & \downarrow i_{C,B}^* \\
K^{-\lambda_p}\left[S(q \rightarrow), S(q \rightarrow) \setminus \bigcup_j \text{int}(V_j)\right] & \xrightarrow{\delta_{triple}^2} & K^{-\lambda_q}[W(q \rightarrow) \cap N_q, S(q \rightarrow)] \\
\downarrow i_j^* \uparrow c_j^* & \nearrow f^j & \\
K^{-\lambda_p}[V_j, \partial V_j] & & 
\end{array}$$

Diagram 49

Now  $i_{C,B}^*$  is an isomorphism, because the collapsing of the tubular neighbourhood  $N_q$  induces a homotopy equivalence


$$\frac{N_q \cup N_p}{N_p \cap L_q} \cong \frac{N_q}{L_q} \rightarrow \frac{W(q \rightarrow) \cap M_c}{L_q \cap W(q \rightarrow)} \cong \frac{W(q \rightarrow) \cap N_q}{S(q \rightarrow)}$$

that is inverse to the inclusion (once the pairs are viewed as pointed quotients). For the rest of the proof we will often argue in similar ways by identifying pairs and there pointed spaces given by quotienting the subspace. This is justified, as the K-theory of a pair is defined to be the K theory of said quotient, and this helps us in the proof as we can write down maps of pointed spaces, that would not be continuous as maps of pairs.

Due to the quotient

$$S(q \rightarrow) / S(q \rightarrow) \setminus \bigcup_j \text{int}(V_j) \cong \bigvee_j (V_j / \partial V_j)$$

being a pointed sum, the map  $\delta_{triple}^2$  is a sum of the functions  $f^j \circ i_j^*$  where the  $f^j : K^{-\lambda_p} [V_j, \partial V_j] \rightarrow K^{-\lambda_q} [W(q \rightarrow) \cap N_q, S(q \rightarrow)]$  are defined to be  $\delta_{triple}^2 \circ c_j^*$ . Now we will inspect said composition more closely, for which we need to dive into the definition of the triple connecting homomorphism, which was given by the composition of the connecting homomorphism of the pair after the pullback along the inclusion:



If we already start by identifying  $K^{-\lambda_p} [V_j, \partial V_j]$  with  $K^{-\lambda_{p+1}} [S \wedge (V_j, \partial V_j)]$  we have that the composition


def wedge of a pair!

$$(\text{id} \wedge l_1)^* := i_1^* \circ c_j^* : K^{-\lambda_{p+1}} [S \wedge (V_j, \partial V_j)] \rightarrow K^{-\lambda_{p+1}} [S \wedge (S(q \rightarrow))]$$

is induced by the collapsing map


$$l_1 : (S(q \rightarrow)) \rightarrow (V_j / \partial V_j).$$

This brings us into the following setup:



The dashed arrows denote functions that we still need to find, to make the diagram commute. The functions  $f_p$  and  $f_q$  will be given below. We will later see, that the notation of  $s^*$  and  $r^*$  is justified in the sense that those are topologically induced funktions.

**ALTERNATIVE**



Notice how by definition of the reduced suspension, there is a natural homomorphism:  $S \wedge (V_j / \partial V_j) \cong ([-1, 1] \times V_j / \partial([-1, 1] \times V_j))$ . Now  $V_j$  is the image of  $\mathcal{N}_{x_j} \gamma^{W(q \rightarrow)}$  under the tubular neighbourhood embedding  $J$ . And  $\text{int}([-1, 1])$  is homeomorphic to  $\mathbb{R}$  via  $x \mapsto \frac{-x}{x^2-1}$ . This gives us a mapping

notation  
einführen

$$f_p : S \wedge (V_j / \partial V_j) \rightarrow \overline{\mathbb{R} \oplus \mathcal{N}_{x_j}^{W(q \rightarrow)} \gamma}$$

With this we can define the homomorphism  $\tilde{f}_p : (I \times V_j / \partial(I \times V_j)) \rightarrow \overline{T_{x_j} W(q \rightarrow)}$  as the closure of the map  $(x, v) \mapsto -\frac{-x}{x^2-1} \cdot \text{grad}_{x_j}(f) + J(v)$ . Then we define  $f_p := (\overline{g_{b_q}} \circ \tilde{f}_p)^*$ , where the map  $g_{b_q} : \mathbb{R}^{\lambda_p+1} \rightarrow T_{x_j} W(q \rightarrow)$  is the inverse of the koordinate map  $\kappa_{b_q}$  where  $b_q$  is the basis of  $T_{x_j} W(q \rightarrow)$  induced by Proposition 5.1.7 that comes from the short exact sequence.

Now lets fix  $(\alpha_p)_\#$  to be the generator induced in Remark 6.1.18 and hence given by the generator  $\alpha_p = \kappa_p^*(b_q)$  of  $K^{\lambda_p}(\varphi^{-1}(D_\varepsilon^s \times D_\varepsilon^u), \varphi^{-1}(D_\varepsilon^s \times \partial D_\varepsilon^u))$ , where  $b_q$  is the generator of  $\tilde{K}^{\lambda_p}(\overline{T_q^u M})$  induced by the orientation of  $T_q^u M$ . Furthermore, let  $\alpha_p^{(B,A)} \in (\alpha_p)_\#$  be the generator of  $K^{-\lambda_p}[B, A]$ . Let  $\mathfrak{e}$  denote the generator of  $K^{-\lambda_p+1}(\mathbb{R}^{\lambda_p+1})$  induced by the standart bases (and composed with a bott periodicity map). Then  $f^p(\mathfrak{e}) = i_j^* \circ i_{B,A}^*(\alpha_p^{(B,A)})$  □

## References

- [Ati89] M. F. Atiyah. *K-theory*. Second. Advanced Book Classics. Notes by D. W. Anderson. Redwood City, CA: Addison-Wesley Publishing Company Advanced Book Program, 1989, pp. xx+216. ISBN: 0-201-09394-4.
- [BH04] A. Banyaga and D. Hurtubise. *Lectures on Morse Homology*. Texts in the Mathematical Sciences. Springer Netherlands, 2004. ISBN: 9781402026959. URL: [https://books.google.de/books?id=AX-\\_sbMj0K4C](https://books.google.de/books?id=AX-_sbMj0K4C).
- [Bro96] Jacek Brodzki. *An Introduction to K-theory and Cyclic Cohomology*. 1996. arXiv: [funct-an/9606001](https://arxiv.org/abs/funct-an/9606001). URL: <https://arxiv.org/abs/funct-an/9606001>.
- [Cou04] Dennis Courtney. *A Brief Glance at K-theory*. 2004. URL: <https://math.berkeley.edu/~hutching/teach/215b-2004/courtney.pdf>.
- [CP61] A. H. Clifford and G. B. Preston. *The Algebraic Theory of Semigroups*. Vol. I. Mathematical Surveys 7. Providence, R.I.: American Mathematical Society, 1961.
- [Gre65] J. A. Green. *Sets and Groups*. London: Routledge & Kegan Paul, 1965.
- [Hat02] Allen Hatcher. *Algebraic Topology*. Cambridge: Cambridge University Press, 2002. ISBN: 978-0-521-79540-1.
- [Lan05] Gregory D. Landweber. *K-Theory and Elliptic Operators*. 2005. arXiv: [math/0504555](https://arxiv.org/abs/math/0504555). URL: <https://arxiv.org/abs/math/0504555>.
- [Per01] Lawrence Perko. *Differential Equations and Dynamical Systems*. 3rd ed. Vol. 7. Texts in Applied Mathematics. Softcover / Paperback edition. New York / Berlin: Springer, 2001, p. 557. ISBN: 978-1461265269.
- [Sal85] Dietmar Salamon. “Connected simple systems and the Conley index of isolated invariant sets”. In: *Transactions of the American Mathematical Society* 291.1 (1985), pp. 1–41. DOI: [10.1090/S0002-9947-1985-0797044-3](https://doi.org/10.1090/S0002-9947-1985-0797044-3).
- [Ste67] N. E. Steenrod. “A convenient category of topological spaces”. In: *Michigan Mathematical Journal* 14.2 (1967), pp. 133–152. DOI: [10.1307/mmj/1028999711](https://doi.org/10.1307/mmj/1028999711).
- [Str] Neil P. Strickland. *The Category of CGWH Spaces*. Unpublished note, <https://ncatlab.org/nlab/files/StricklandCGWHSpaces.pdf>. abgerufen am 25. Juni 2026.
- [tom08] Tammo tom Dieck. *Algebraic Topology*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008. ISBN: 978-3-03719-048-7. DOI: [10.4171/048](https://doi.org/10.4171/048).