
Contents

1	Differentiable structures on topological manifolds	2
2	Morse Theory	9
3	Morse-Theoretic Atiyah-Hirzebruch Spectralsequence	29
3.1	Conley index	29
4	The Spectral Sequence using Exact Couples	35
5	Higher Differentials	43
6	The prove that N_p is a tubular neighbourhood	44
7	A Künneth Formular	67
8	Examples	68

1 Differentiable structures on topological manifolds

This section aims to set the playing field for the thesis. Due to the theory being of interest for mathematicians and physicists alike there are many (equivalent) definitions used to introduce the objects. The main goal of this section is to avoid confusion, by giving the definitions instead relying on the reader to guess what a certain symbol stands for.

Definition 1.0.1 (Topological Manifold). A second countable Hausdorff space M is called **topological manifold** of dimension $m \in \mathbb{N}$, if it is locally homeomorphic to $\mathbb{R}^m$. To be precise, if for all $p \in M$ there exists an open neighborhood $U \subseteq M$ of p , an open set $V \subseteq \mathbb{R}^m$ and a map $\varphi : U \rightarrow V$ that is a homeomorphism. We call the map $\varphi : U \rightarrow V$ a **chart around p on M** and φ^{-1} a **local coordinate system around p on M** .

Definition 1.0.2 (Differentiable Manifold). Let M be a topological manifold of dimension m .

1. A **differentiable atlas of class $r \in \mathbb{N} \cup \{\infty\}$** is a family of charts $\mathfrak{A} = (\varphi_i : U_i \rightarrow V_i)_{i \in I}$ such that
 - a) $\bigcup_{i \in I} U_i = M$, meaning that (U_i) is an open covering of M .
 - b) For every pair $(i, j) \in I^2$ the **transition function**:

$$\begin{aligned} \varphi_{ij} : \varphi_j(U_i \cap U_j) &\rightarrow \varphi_i(U_i \cap U_j) \\ x &\mapsto (\varphi_i \circ \varphi_j^{-1})(x) \end{aligned}$$

is differentiable of class r .

We call such an atlas a C^r -atlas.

2. Two C^r -atlases $\mathfrak{A}$ and $\mathfrak{B}$ are called **equivalent** if the family $\mathfrak{A} + \mathfrak{B} = (\varphi_i, \varphi_j)_{ij}$ is a C^r -atlas.

A **differentiable structure of class r on M** is an equivalence class c of C^r -atlases. For $r = \infty$ we call the pair (M, c) a smooth manifold.

Corollary 1.0.3. Every transition functions φ_{ij} where $(i, j) \in I^2$ of a differentiable atlas $\mathfrak{A} = (\varphi_i)_{i \in I}$ is not just a homeomorphism but also a diffeomorphism due to $\varphi_{ji} = \varphi_{ij}^{-1}$.

Definition 1.0.4. Let (M, c) be a differentiable manifold of class r and $U \subseteq M$ open. We call a continuous function

$$f : U \rightarrow \mathbb{R}$$

differentiable of class r , if for any one (and hence for all) $(\varphi_i)_{i \in I} \in c$ the compositions $f \circ \varphi_i^{-1}$ are differentiable of class r . For $r = \infty$ we define:

$$\mathcal{E}(U) = \{f \in C(U, \mathbb{R}) \mid f \text{ is differentiable of class } \infty\}.$$

Corollary 1.0.5. Let (M, c) be a smooth manifold of dimension m and $U \subseteq M$ be an open subset. With pointwise defined operations, the set $(\mathcal{E}(U), +, \cdot, \circ)$ becomes an $\mathbb{R}$ -algebra. Furthermore, $\mathcal{E}$ becomes a sheaf of $\mathbb{R}$ -algebras.

Proof. There is not really a need for a proof. However, it might help to work through the definition of a sheaf as a reminder. First, $\mathcal{E}$ is a presheaf, where the restriction in the domain of a function gives the needed restriction homomorphism:

$$\begin{aligned} \text{res}_V^U : \mathcal{E}(U) &\rightarrow \mathcal{E}(V) \\ f &\mapsto f|_V . \end{aligned}$$

The required properties of a presheaf are trivial. Furthermore, this gives a sheaf as the requirement of locality is trivial for functions and the property of gluing is also trivial for functions, since differentiability is a local property. $\square$

Definition 1.0.6. If $p \in M$ is fix, $f \in \mathcal{E}(U)$ and $g \in \mathcal{E}(U')$ such that $p \in U \cap U'$ we say that f and g have the same **germ in** p , if there is another open neighborhood $W \subseteq U \cap U'$ of p such that $f|_W = g|_W$. This defines an equivalence relation $\sim_p$. An equivalence class s of local functions around p is called a **germ in** p . We write $s = f_p$, if $s = [f]$ with $f \in \mathcal{E}(U)$. We write

$$\mathcal{E}_p(M) = \left(\bigsqcup_{U \text{ open}, p \in U} \mathcal{E}(U) \right) / \sim_p .$$

For the set of germs and call it the **stalk in** p . Here $\sum$ denotes the co-product (also called sum) in **Top** and hence the disjoint union.

Corollary 1.0.7. For a smooth manifold (M, c) the set $\mathcal{E}_p(M)$ inherits an $\mathbb{R}$ -algebra structure from the $\mathcal{E}(U)$. Furthermore, it carries a natural (evaluation-)homomorphism:

$$\begin{aligned} \text{eval}_p : \mathcal{E}_p(M) &\rightarrow \mathbb{R} \\ f_p &\mapsto f(p) =: f_p(p) \end{aligned}$$

The stalks are also local rings with maximal ideal $\mathfrak{m}_p = \ker(\text{eval}_p)$. Hence, the pair $(M, \mathcal{E})$ gives us a locally ringed space.

Proof. Here, we only need to prove the statement about the locality of the stalks. This follows from $f_p \in \mathcal{E}_p(M)$ being invertible if and only if $f(p) \neq 0$ which is the same as $f_p \notin \ker(\text{eval}_p)$. $\square$

Definition 1.0.8. Let (M, c) be a smooth manifold of dimension m and $p \in M$. We call an $\mathbb{R}$ -linear map $\delta : \mathcal{E}_p(M) \rightarrow \mathbb{R}$ a **derivation**, if it satisfies the Leibnitz-rule:

$$\delta(f_p \cdot g_p) = \delta(f_p)g_p(p) + f_p(p)\delta(g_p) \quad \text{for all } f, g \in \mathcal{E}_p(M) .$$

We call $\text{Der}_{\mathbb{R}}(\mathcal{E}_p(M), \mathbb{R})$ the set of derivations and give it a $\mathbb{R}$ vector space structure by pointwise operations. We define the **tangent space of** M **at** p to be the vector space

$$TM_p := \text{Der}_{\mathbb{R}}(\mathcal{E}_p(M), \mathbb{R}) .$$

Corollary 1.0.9. Let (M, c) be a smooth manifold and $\varphi : U \rightarrow V$ be a chart around p with $x_0 = \varphi(p)$ ($\varphi \in \mathfrak{A} \in c$). Then

$$\xi = \frac{\partial}{\partial x_j} \Big|_p : \mathcal{E}_p(M) \rightarrow \mathbb{R}$$

$$f_p \mapsto \xi(f_p) = \frac{\partial}{\partial x_j} \Big|_{x_0} (f \circ \varphi^{-1})$$

is well-defined and a tangent vector. In fact, the family

$$\left(\frac{\partial}{\partial x_1} \Big|_p, \dots, \frac{\partial}{\partial x_m} \Big|_p \right)$$

defines a basis of TM_p . Hence, the dimension of TM_p is m .

Definition 1.0.10. For a smooth manifold (M, c) the sum $\bigsqcup_p TM_p$ comes with a natural projection

$$\pi : TM \rightarrow M$$

$$\xi \mapsto p \text{ where } \xi \in TM_p$$

Furthermore, the **local vector fields** with respect to a chart $\varphi : U \rightarrow V$

$$\frac{\partial}{\partial x_j} : U \rightarrow \pi^{-1}(U)$$

$$p \mapsto \frac{\partial}{\partial x_j} \Big|_p$$

induce a local trivialization:

$$\pi^{-1}(U) \cong U \times \mathbb{R}^m.$$

We can induce a topology on TM such that all those trivializations are continuous. This then gives an atlas for TM such that we have a $2m$ -dimensional manifold. To be precise, the atlas is given by the maps $\pi^{-1}(U_i) \rightarrow \mathbb{R}^m \times \mathbb{R}^m; x \mapsto (\pi(x), q_{\varphi \circ \pi}(x))$ where q_p denotes the coordinate map corresponding to the basis $(\frac{\partial}{\partial x_1} \Big|_p, \dots, \frac{\partial}{\partial x_m} \Big|_p)$ that depends on the chart φ_i . In fact, this yields a smooth manifold and a (smooth) vector bundle of dimension m . We call TM the **tangent bundle**.

Definition 1.0.11. For a m dimensional vector space V with basis $\mathfrak{b}$ we define the two maps:

$$B_{\mathfrak{b}} : \mathbb{R}^m \rightarrow V : e_i \mapsto \mathfrak{b}_i$$

$$q_{\mathfrak{b}} : B_{\mathfrak{b}}^{-1} : V \rightarrow \mathbb{R}^m.$$

Definition 1.0.12 (The Derivative). Let (M, c) and (M', c') be smooth manifolds (from now on we suppress the differentiable structure in our notation). We call a continuous function $f : M \rightarrow M'$ **smooth**, if for all $\varphi \in \mathfrak{A} \in c$ and $\varphi' \in \mathfrak{A}' \in c'$ the maps

$$\varphi' \circ f \circ \varphi^{-1} : V \rightarrow V'$$

are smooth. A given smooth function induces a smooth function between the tangent bundles as follows:

$$Df : TM \rightarrow TM', Df_p(\xi)(g_p) = \xi((g \circ f)_p)$$

Here, $\xi \in T_p M$, $g_p \in E_p(M')$.

Corollary 1.0.13. Let $f : M \rightarrow \mathbb{R}$ be smooth and $\varphi : U \rightarrow V$ be a chart. Then we can interpret df as a one-form. To be precise assume q to be the coordinate function $T\mathbb{R} \rightarrow \mathbb{R}$ from the basis induced by the identity as a chart. $v \in \Gamma TM$ we have

$$q \circ df(v) = v \cdot f$$

Here on the right side, f denotes a map $p \mapsto f_p$ such that $v(f)(p) := v_p(f_p)$ is well defined. We keep this notations so vector fields can take functions as an input.

Proof. Let $\varphi : U \rightarrow V$ be a chart of M . We proof this by showing it for the local sections $\frac{\partial}{\partial x_i}$ and thereby for any, since those sections form a basis of the space of sections as a $C^\infty(M, \mathbb{R})$ vector space. Now let $g_p \in \mathcal{E}_p(\mathbb{R})$ and $\varphi(p) = x_0$. Then

$$df_p(\partial_i(p))(g_p) = \partial_i((g \circ f)_p) = \left. \frac{\partial}{\partial x_i} \right|_{x_0} (g \circ f \circ \varphi^{-1}) = \left. \frac{\partial}{\partial x} \right|_p (g_p) \cdot \left. \frac{\partial}{\partial x_i} \right|_p (f_p)$$

Hence, $df(v) = v(f) \frac{\partial}{\partial x}$ letting us conclude the statement. $\square$

Definition 1.0.14. Let V be a linear manifold (i.e. a finite dimensional vector space). Then for any $v \in V$ we have the canonical maps

$$Q_v : V \rightarrow T_v V$$

$$x \mapsto \left(f_v \mapsto \left. \frac{\partial}{\partial t} \right|_{t=0} f(v + tx) \right)$$

Let $L : V \rightarrow V$ be a linear map viewed as a smooth map between manifolds. Then we have the identity:

$$dL|_v \circ Q_v = Q_{L(v)} \circ L.$$

This is proven by a short calculation:

$$\left(dL|_v (Q_v(x)) \right) (f_{L(v)}) = Q_v(x) (f \circ L) = \left. \frac{\partial}{\partial t} \right|_{t=0} f \circ L(v + tx) = \underbrace{\left. \frac{\partial}{\partial t} \right|_{t=0} f(L(v) + tL(x))}_{=Q_{L(v)} \circ L(x)}$$

Now let $v, v' \in V$ be two points in V . Then we have the translation

$$tr_{v,v'} : V \rightarrow V; x \mapsto (x + (v' - v))$$

With this we can now find a connection between Q_v and $Q_{v'}$ as follows:

$$\left(dtr_{v,v'}|_v (Q_v(x)) \right) (f_{v'}) = \frac{\partial}{\partial t} \Big|_{t=0} (f \circ tr_{v,v'}(v + tx)) = \frac{\partial}{\partial t} \Big|_{t=0} (f(v' + tx)) = Q_{v'}(x)(f_{v'}) .$$

Hence:

$$Q_{v'} = dtr_{v,v'} \circ Q_v .$$

The map Q then denotes the map on the whole tangent bundle:

$$Q : TV \rightarrow V; \xi_p \mapsto Q_p(\xi_p)$$

Definition 1.0.15 (A Metric). Let M be a smooth manifold. A section $g \in \Gamma(TM^* \otimes TM^*)$ into the tensor product of the dual tangent bundle with itself is a **Riemannian metric** if the following are satisfied for all $p \in M$:

1. g is **non degenerate**, meaning for a $v_p \in T_p M$ $g_p(v_p, w_p) = 0 \ \forall w_p \in T_p M$ then $v_p = 0$.
2. g is **symmetric**, meaning $g_p(v_p, w_p) = g_p(w_p, v_p) \ \forall v_p, w_p \in T_p M$.
3. g is **positive definite**, meaning that $g_p(v_p, v_p) \geq 0 \ \forall v_p$ and vanishes only for $v_p = 0$.

We call a tuple (M, g) a **Riemannian manifold**.

Definition 1.0.16 (The Gradient). Assume that (M, g) is a smooth manifold together with a Riemannian metric. Let $f : M \rightarrow \mathbb{R}$ be a smooth function. We define its **gradient** to be a vector field $\text{grad}(f) \in \Gamma(TM)$ such that for any vector field V :

$$Q \circ df(V) = g(\text{grad}(f), V) .$$

Definition 1.0.17. Let (M, g) be a Riemannian manifold and $\varphi : U \rightarrow V$ be a chart. We define the smooth functions $g_{ij} : U \rightarrow \mathbb{R}$ to be :

$$g_{ij} = g\left(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j}\right) .$$

Then if two vector fields v, w over U are given with $v = \sum v^i \frac{\partial}{\partial x_i}$ and $w = \sum w^i \frac{\partial}{\partial x_i}$ we can calculate the metric locally:

$$g(v, w) = \sum_{ij} g_{ij} v_i w_j : U \rightarrow \mathbb{R}$$

Furthermore, we can invert the matrices $(g_{ij}(p))_{ij}$ for all p (which is a smooth procedure which can be seen by the construction via Cramers rule) to get smooth functions

$$g^{ij} : U \rightarrow \mathbb{R}$$

$$p \mapsto g^{ij}(p) = \left((g_{kl}(p))_{kl}^{-1} \right)_{ij}$$

Remark 1.0.18. Due to the rare appearance of big summations with vectors and co-vectors in this work, we omit the Einstein summation convention and give all coordinate entries lower indices.

Lemma 1.0.19. *Let $\varphi : U \rightarrow V$ be a chart. Then the gradient has the local form:*

$$\text{grad}(f) = \sum_{i,j} g^{ij} \left(\frac{\partial}{\partial x_i}(f) \right) \frac{\partial}{\partial x_j} =: \sum_{i,j} g^{ij} \frac{\partial f}{\partial x_i} \frac{\partial}{\partial x_j}.$$

Proof. Assume that $v, w \in \Gamma(TM)$ such that $v = \sum_i v^i \frac{\partial}{\partial x_i}$ and $w = \sum_i w^i \frac{\partial}{\partial x_i}$. Then $g(v, w) = \sum_{i,j} g_{ij} v^i w^j$. Suppose that $\text{grad}(f) = \sum_j G^j \frac{\partial}{\partial x_j}$ and q is the coordinate map $T\mathbb{R} \rightarrow \mathbb{R}$ in each tangend space. Then by definition of the gradient we have:

$$\frac{\partial}{\partial x_j}(f) = q \circ df \left(\frac{\partial}{\partial x_j} \right) = q \circ \frac{\partial f}{\partial x_j} = g(\text{grad}(f), \frac{\partial}{\partial x_j}) = \sum_{ij} g_{ij} G^i$$

Hence,

$$\begin{aligned} (G^1, \dots, G^m)(g_{ij}) &= \left(\frac{\partial}{\partial x_1}(f), \dots, \frac{\partial}{\partial x_m}(f) \right) \\ &\Leftrightarrow (G^1, \dots, G^m) = \left(\frac{\partial}{\partial x_1}(f), \dots, \frac{\partial}{\partial x_m}(f) \right) (g^{ij}). \end{aligned}$$

But then we can conclude that $G^j = \sum_i g^{ij} \frac{\partial}{\partial x_i}(f)$. □

Definition 1.0.20 (A Dynamical System). Let M be a smooth manifold. A **dynamical system on M** is given by a smooth map $\varphi : \Omega \rightarrow M$ such that:

1. $\{0\} \times M \subseteq \Omega \subseteq \mathbb{R} \times M$ open, and for any $p \in M$ the set $I(p) := \pi_1(\Omega \cup \mathbb{R} \times \{p\})$ is connected. (π_1 is the projection onto the first factor)
2. For all $p \in M$ and $t \in I(p)$ we have that $s \in I(\varphi(t, p))$ if and only if $s + t \in I(p)$. Then we have:

$$\varphi(s, (\varphi(t, p))) = \varphi(s + t, p)$$

3. For all p we require $\varphi(0, p) = p$.

We will write $\varphi_t(p) := \varphi(t, p)$ to keep the notation bearable. If for all p $I(p) = \mathbb{R}$ we call the system **global**.

Corollary 1.0.21. For a global dynamical system φ on a smooth manifold M the map

$$\rho : \mathbb{R} \rightarrow \text{Diff}, \quad t \mapsto \varphi_t$$

is a homomorphism. To see the well definition notice how for all $t \in \mathbb{R}$ φ_{-t} is an inverse of φ_t .

Definition 1.0.22 (Vector Field of a Dynamical System). Let φ be a dynamical system on a smooth manifold M . Then we call the mapping

$$p \mapsto X(p) := \left. \frac{d}{dt} \right|_{t=0} \varphi_t(p) \in T_p M$$

the **associated vector field**. This is in fact a smooth vector field.

Definition 1.0.23 (Dynamical System of a Vector Field). Let $X \in \Gamma(M)$ be a global vector field on the smooth manifold M . Then we call a dynamical system φ on M the **associated dynamical system** to X , if for all $t \in \mathbb{R}$ and $x \in M$

$$d_t \varphi_t(x)|_{t=t_0} = X(\varphi_{t_0}(x)).$$

d_t denotes the differential in t , meaning for fixed x we differentiate

$$\varphi(x) : \mathbb{R} \rightarrow M, \quad t \mapsto \varphi_t(x)$$

and evaluate that in the constant trivial vector field $E : \mathbb{R} \rightarrow T\mathbb{R}$ with $E(t) = 1_t$.

Those two concepts are inverse to one another or in other words:

Theorem 1.0.24. *Let M be a smooth manifold. Then there is a one to one correspondence between global maximal flows and vector fields.*

Remark 1.0.25. Let $X \in \Gamma(TM)$ be a vector field on the smooth manifold M , and φ be the associated dynamical system. For any fixed $t_0 \in \mathbb{R}$ and any $x_0 \in M$ we have the identity

$$d\varphi_{t_0}|_{x_0} \circ (X(x_0)) = X(\varphi_{t_0}(x_0)).$$

Proof. This follows directly from the second property of flows as follows: We first make use of the identity $\varphi_{t_0}(x_0) = \varphi_{t_0} \circ \varphi_0(x_0)$ and $X(x_0) = d_t \varphi_t(x_0)|_{t=0}$ to calculate using the chain rule:

$$d_t \varphi_{t_0+t}(x_0)|_{t=0} = d_t \varphi_{t_0} \circ \varphi_t(x_0)|_{t=0} = d_x \varphi_{t_0}|_{x=\varphi_0(x_0)} \circ d\varphi_t(x_0)|_{t=0} = d_x \varphi_{t_0}|_{x=x_0} \circ X(x_0),$$

and alternatively

$$d_t \varphi_{t_0+t}(x_0)|_{t=0} = d_t \varphi_t(\varphi_{t_0}(x_0))|_{t=0} = X(\varphi_{t_0}(x_0)).$$

□

2 Morse Theory

For everything concerning Morse theory our main reference will be the wonderful book [BH04]. In Morse theory, we will always deal with a smooth function $f : M \rightarrow \mathbb{R}$ from a smooth Riemannian manifold (M, g) that is also compact. We will denote its dimension by $m = \dim(M)$.

Definition 2.0.1 (Critical Points). We call a point $p \in M$ **critical**, if $df|_p$ vanishes meaning that

$$df|_p : T_p M \rightarrow T_{f(p)} \mathbb{R}$$

is trivial. By $\text{Crit}(f)$ we denote the **set of critical points**.

Definition 2.0.2 (The Hessian). Let $f : M \rightarrow \mathbb{R}$ be a smooth function. The **Hessian** of f at $p \in \text{Crit}(f)$ is a symmetric bilinear function

$$\text{Hess}_p(f) : T_p M \times T_p M \rightarrow \mathbb{R}$$

that is defined as follows. Let $\xi_p, \zeta_p \in T_p M$ be two tangent vectors. Now choose two vector fields Ξ and Z in a neighbourhood of p such that $\Xi(p) = \xi_p$ and $Z(p) = \zeta_p$. With this define the Hessian to be:

$$\text{Hess}_p(f)(\xi_p, \zeta_p) = (\Xi \cdot (Z \cdot f))(p)$$

Compare 1.0.13 for the notation. In corollary 2.0.5 we will give a local description telling us that this definition is well defined, meaning independent of the Extensions Ξ and Z .

Definition 2.0.3. A smooth function $f : M \rightarrow \mathbb{R}$ is called **Morse**, if for any critical point.

The Hessian is a symmetric bilinear and non-degenerate Map. For such a map there exists the following theorem:

Theorem 2.0.4 (Sylvester's Law of Inertia). *Let $A \in \text{Mat}(n, \mathbb{R})$ be symmetric and let $T, T' \in \text{GL}(n, \mathbb{R})$ and $k, k', l, l' \in \mathbb{N}$ such that*

$$T^t \circ A \circ T = \begin{pmatrix} \mathbb{1}_k & 0 & 0 \\ 0 & -\mathbb{1}_l & 0 \\ 0 & 0 & 0_{n-k-l} \end{pmatrix} \text{ and } T'^t \circ A \circ T' = \begin{pmatrix} \mathbb{1}_{k'} & 0 & 0 \\ 0 & -\mathbb{1}_{l'} & 0 \\ 0 & 0 & 0_{n-k'-l'} \end{pmatrix}$$

then $k = k'$, $l = l'$ and $\text{rank}(A) = k + l$.

Proof. Since T, T' are invertible we have that

$$k + l = \text{rank}(T^t \circ A \circ T) = \text{rank}(A) = \text{rank}(T'^t \circ A \circ T') = k' + l'. \quad (1)$$

Hence it suffices to show that $k = k'$. Which we will do by proofing the claim:

$$k = \max \{ \dim(U) \mid U \subseteq \mathbb{R}^n \text{ subspace such that } x^t A x > 0 \ \forall x \in U \setminus \{0\} \}$$

So we start by showing „ $\leq$ “: Denote the first k colons of T with $x_1, \dots, x_k$. They form a basis of $\mathbb{R}^n$ and with $0 \neq x = \sum_{i=1}^k \lambda_i x_i$ we have that by bilinearity

$$x^t A x = \sum_{i=1}^k \lambda_i x_i^t A x = \sum_{i,j=1}^k \lambda_i \lambda_j x_i^t A x_j = \sum_{i=1}^k (\lambda_i)^2 \geq 0. \quad (2)$$

This concludes the first inequality.

Now let U be any k -dimensional subspace such that for all non zero $x \in U$ $x^t A x > 0$. By a calculation analog to the one above we have that for any $x \in W := \text{span}(x_{k+1}, \dots, x_n)$ the number $x^t A x$ is less or equal to zero. Hence, $W \cap U = \{0\}$ and with this we conclude:

$$\dim(U) = \dim(U + W) - \dim(W) + \dim(U \cap W) \leq (k + (n - k)) - (n - k) + 0 = k. \quad (3)$$

This is the last inequality proving the statement. $\square$

Corollary 2.0.5. Let $\varphi : U \rightarrow V$ be a chart around a critical point $p \in M$. Then in said chart the matrix corresponding to the Hessian has the form:

$$M_p(f) = \left(\frac{\partial^2 (f \circ \varphi^{-1})}{\partial x_i \partial x_j} \right)_{i,j}$$

Proof. This is just an easy calculation, since $\text{Hess}_p(f) \left(\frac{\partial}{\partial x_i} \Big|_p, \frac{\partial}{\partial x_j} \Big|_p \right)$ is given by

$$\left(\frac{\partial}{\partial x_i} \Big|_p \left(\frac{\partial}{\partial x_j} (f) \right) \Big|_p \right) = \left(\frac{\partial^2 (f \circ \varphi^{-1})}{\partial x_i \partial x_j} \right)$$

$\square$

The Hessian matrix $M_p(f)^i$ in two different charts $\varphi_i : U_i \rightarrow V_i$ for $i = 1, 2$ transforms via a congruence transformation by the Jacobian of the transition function. If we have two different charts. Then the Matrix $M_p(f)$ of the Hessian transforms from one base to the other via conjugation of the differential of the transition function $\varphi_{12} := \varphi_1 \circ \varphi_2^{-1}$:

$$M_p^1(f) = d\varphi_{12}^t \cdot M_p^2(f) \cdot d\varphi_{12}$$

Hence we can define the following:

Definition 2.0.6. By Sylvesters Law of inertia, the number l that denotes the dimension of the maximal subspace on which a matrix is negative definite, is a invariant under conjugation. We call that number the index of said matrix. Now in the setting of Morse theory we assign to each kritical point p the index of $M_p(f)$ and call that the **(Morse-)index** of p , denoted with λ_p . If we cannot suppress the function without loss of clarity we will write λ_p^f . We denote the set of all critical points of index k by $\text{Crit}_k(f)$.

Theorem 2.0.7 (Morse Lemma). *Let p be a critical point of index k of the Morse function $f : M \rightarrow \mathbb{R}$. There exists a centered chart $\varphi : U \rightarrow \mathbb{R}^m$ around p such that*

$$(f \circ \varphi^{-1})(x_1, \dots, x_m) = f(p) + \sum_{i=1}^k -x_i^2 + \sum_{i=k+1}^m x_i^2$$

Proof. Compare for example Lemma 3.11 in [BH04]. $\square$

hier noch
mehr
Quellen!

Definition 2.0.8. Let $\varphi_t : M \rightarrow M$ be the local one-parameter group of diffeomorphisms generated by $-\text{grad}(f)$, i.e. if $e : \mathbb{R} \rightarrow T\mathbb{R}$ denotes the canonical unit vector field that comes from the identity as the global chart on $\mathbb{R}$, we have that

$$\frac{d}{dt} \varphi_t(x) \circ e = -\text{grad}(f)(\varphi_t(x)), \quad \text{and} \\ \varphi_0(x) = x.$$

We call that group the **negative gradient flow** of f with respect to the metric g . The integral curve $\gamma_x : (a, b) \rightarrow M$ given by

$$\gamma_x(t) = \varphi_t(x)$$

is called a **gradient flow line**.

Next we will deepen our **local** understanding of said vector field. Notice how the Morse lemma is a statement about smooth manifolds no metric needed nor mentioned. In fact there are many Morse charts and we can find one, where the Riemannian metric in p is diagonal with non-zero. To proof such a statement we first need a somewhat obvious lemma:

Lemma 2.0.9. *Assume that $\psi : U \rightarrow V$ is a chart and $L : \mathbb{R}^m \rightarrow \mathbb{R}^m$ a linear function. Denote the local coordinate maps of the tangent bundle in a point p that come from a basis in said point induced from a chart χ by q_χ . Then the diagram*

$$\begin{array}{ccc} T_p M & \xrightarrow{q_\psi|_p} & \mathbb{R}^m \\ & \searrow q_{L \circ \psi}|_p & \downarrow L \\ & & \mathbb{R}^m \end{array}$$

commutes for all $p \in U$.

Proof. We show this by showing the inverse: For any $v \in \mathbb{R}^m$ we have the two $(q_\psi|_p)^{-1} \circ L^{-1}(v)$ and $(q_{\psi \circ L}|_p)^{-1}$ and claim that they are the same. Assume that $L^{-1} = (\lambda_{ij})_{ij}$, $\psi(p) = x_0$, $L(x_0) = \tilde{x}_0$, $(v_1, \dots, v_m) = v \in \mathbb{R}^m$ and denote the basis of $T_p M$ coming from

the chart ψ by $\left(\frac{\partial}{\partial x_i}\right)_p$ and the one coming from $L^{-1} \circ \psi$ by $\left(\frac{\partial}{\partial \tilde{x}_i}\right)_p$. We now calculate for $f_p \in \mathcal{E}_p M$:

$$\begin{aligned}
 \left[(q_{\psi \circ L}|_p)^{-1}(v)\right](f_p) &= \sum_i v_i \frac{\partial}{\partial \tilde{x}_i} \Big|_p (f_p) \\
 &= \sum_i v_i \frac{\partial}{\partial x_i} \Big|_{\tilde{x}_0} (f \circ \psi^{-1} \circ L^{-1}) \\
 &= \sum_i v_i \sum_j \frac{\partial f \circ \psi^{-1}}{\partial x_j} \Big|_{x_0} \cdot \frac{\partial L_j^{-1}}{\partial x_i} \Big|_{\tilde{x}_0} \\
 &= \sum_{i,j} v_i \frac{\partial}{\partial x_j} \Big|_p (f_p) \cdot \lambda_{ji} \\
 &= \sum_{i,j} v_i q_{\psi}|_p^{-1}(e_j)(f_p) \cdot \lambda_{ji} \\
 &= \left[q_{\psi}|_p^{-1} \left(\sum_{i,j} v_i \lambda_{ji} e^j \right) \right] (f_p) \\
 &= \left[q_{\psi}|_p^{-1} (L^{-1}(v)) \right] (f_p)
 \end{aligned}$$

This concludes the proof. $\square$

Now we can proof that a Morse chart exists where the gradient is of particular easy form:

Theorem 2.0.10 (Simultaneous Diagonalisation of Quadratic Forms). *Let p be a critical point of a Morse function f on a manifold M and let $g(p)$ be the Riemannian metric on $T_p M$. Then there exists a Morse chart ψ around p such that the representation of $g(p)$ in the basis induced by the Morse chart is given by a diagonal matrix $\text{diag}(\mu_1, \dots, \mu_m)$ with $\mu_i > 0$ for all i .*

Proof. The quadratic form of $f \circ \psi^{-1} - f(p)$ in the coordinates of any Morse chart ψ is $q_H(v) = v^T H v$. The quadratic form induced by the Riemannian metric $g(p)$ is $q_G(v) = v^T G v$, where $v \in \mathbb{R}^m$ are the coordinate vectors with respect to the Morse chart. To be precise this means for $v, w \in \mathbb{R}^m$:

$$g \circ (q_{\psi}|_p \oplus q_{\psi}|_p)(v, w) = v^T G w \text{ and } f \circ \psi^{-1}(v) = f(p) + v^T H v.$$

We now aim to manipulate ψ such that G becomes diagonal: Since $g(p)$ is positive definite, G is also positive definite, and H is symmetric.

Consider the generalized eigenvalue problem $Hv = \lambda Gv$. Since H and G are real symmetric matrices and G is positive definite, this problem has m real eigenvalues $\lambda_1, \dots, \lambda_m$ and corresponding eigenvectors $v_1, \dots, v_m$, which can be chosen to be orthogonal with

respect to the bilinear form defined by G , such that $v_i^T G v_j = \delta_{ij}$, since if v_i and v_j are different eigenvectors with respect to different eigenvalues, we can calculate:

$$v_j^T H v_i = (H v_j)^T v_i = \lambda_j (G v_j)^T v_i = \lambda_j v_j^T G(v_i) \quad \text{and} \quad v_j^T H v_i = v_j^T \lambda_i G(v_i) = \lambda_i v_j^T G(v_i).$$

Hence we have the equality

$$\lambda_j (G v_j)^T v_i = \lambda_i v_j^T G(v_i) \Leftrightarrow \underbrace{(\lambda_j - \lambda_i)}_{\neq 0} v_j^T G(v_i) = 0$$

Hence all eigenspaces are orthogonal and we can use Gram Schmitt to make the bases of the eigenspaces orthogonal and by rescaling orthonormal.

Let $L = [v_1 | \dots | v_m]$ be the matrix whose columns are these G -orthonormal eigenvectors. The linear change of coordinates $y = Lz$ leads to new coordinates z . In these new coordinates, the quadratic forms transform as follows:

$$\begin{aligned} q_G(y) &= y^T G y = (Lz)^T G(Lz) = z^T L^T G L z = z^T I z = \sum_{i=1}^m z_i^2 \\ q_H(y) &= y^T H y = (Lz)^T H(Lz) = z^T L^T H L z \end{aligned}$$

Since $H v_i = \lambda_i G v_i$, we have $L^T H L = \text{diag}(\lambda_1, \dots, \lambda_m)$. The eigenvalues λ_i are real and have the same signature as the eigenvalues of H (l negative, $m-l$ positive). This is true to Sylvester's law of inertia. By a further scaling of the v_i and a reordering the matrix $L^T H L$ can be brought into the form $\text{diag}(-1, \dots, -1, 1, \dots, 1)$. (by doing this however, the matrix $L^T G L$ becomes $L^T G L = \text{diag}(\mu_1, \dots, \mu_m)$ with $\mu_j > 0 \ \forall j$). If ψ is the Morse chart we started with, then $L^{-1} \circ \psi$ is the new Morse chart:

$$f \circ (\psi^{-1} \circ L)(y) = f \circ \psi^{-1}(L(y)) = q_H(L(y)) = \sum_{i=1}^l -y_i^2 + \sum_{i=l+1}^m y_i^2$$

Furthermore, we want to inspect the metric $g|_p : T_p M^2 \rightarrow \mathbb{R}$ with respect to the chart $L^{-1} \circ \psi$ -induced basis $\left(\frac{\partial}{\partial x_1} \Big|_p, \dots, \frac{\partial}{\partial x_m} \Big|_p \right)$. By the lemma 2.0.9

$$g(q_{L^{-1} \circ \psi} \Big|_p^{-1}, q_{L^{-1} \circ \psi} \Big|_p^{-1})(v, w) = g(q_\psi \Big|_p^{-1}, q_\psi \Big|_p^{-1})(Lv, Lw) = (Lv)^T G L w = v^T L^T G L w$$

□

Definition 2.0.11. We call a Morse chart where the Matrix corresponding to the gradient in the critical point is of diagonal form a **refined Morse chart**.

Using such a chart, we can start looking at the flow itself and try to find easy local formulas around critical points. For this we start by looking at the Jacobian of the gradient in a critical point:

Definition 2.0.12 (Jacobian of the Gradient). The gradient is a section into the tangent bundle, $\text{grad}(f) : M \rightarrow TM$. Let $\psi : M \supseteq U \rightarrow V \subseteq \mathbb{R}^m$ be a chart. The coordinate map q_ψ on TU assigns to a vector $v \in T_p M$ (where $p \in U$ with $\psi(p) = x$) its components with respect to the basis $\left(\frac{\partial}{\partial x_i}\big|_p\right)$, i.e., $q_\psi(v) = (v_1, \dots, v_m)$ if $v = \sum_{i=1}^m v_i \frac{\partial}{\partial x_i}\big|_p$. We define the Jacobian of the gradient with respect to the chart ψ as the Jacobian matrix of the coordinate representation of the gradient in this chart:

$$J(\text{grad}(f))_\psi(x) := J(q_\psi \circ \text{grad}(f) \circ \psi^{-1})(x) = \left(\frac{\partial(q_\psi \circ \text{grad}(f) \circ \psi^{-1})_i}{\partial x_j}(x) \right)_{ij}$$

Lemma 2.0.13. Let ψ be a coordinate system around a critical point p . I.e. $\psi : p \in U \rightarrow V$ with $\psi(p) = 0$ and let $\left(\frac{\partial}{\partial x_1}\big|_x, \dots, \frac{\partial}{\partial x_m}\big|_x\right)$ be the induced basis of $T_x M$. Then the Jacobian of the gradient reads

$$J(\text{grad}(f))_\psi(0) = \left(\sum_k g^{ki}(0) \frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} \right)_{ij}, \quad (4)$$

where g^{ij} is defines as in definition 1.0.17.

Proof. Let q_ψ denote the coordinate funktion $TU \rightarrow \mathbb{R}^m$ induced from the basis $\left(\frac{\partial}{\partial x_1}\big|_x, \dots, \frac{\partial}{\partial x_m}\big|_x\right)$. then the gradient in ψ reads:

$$q_\psi \circ \text{grad}(f) = \left(\sum_k g^{k1} \frac{\partial f}{\partial x_k}, \dots, \sum_k g^{km} \frac{\partial f}{\partial x_k} \right)$$

Hence, we can differentiate:

$$\begin{aligned} \frac{\partial}{\partial x} q_\psi \circ \text{grad}(f) \circ \psi^{-1} &= \left(\frac{\partial}{\partial x_j} \sum_k g^{ki} \frac{\partial f \circ \psi^{-1}}{\partial x_k} \right)_{ij} \\ &= \left(\sum_k \left[\frac{\partial g^{ki}}{\partial x_j} \frac{\partial f \circ \psi^{-1}}{\partial x_k} + g^{ki} \frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} \right] \right)_{ij}. \end{aligned}$$

and now in $p = \psi^{-1}(0)$ we have

$$\frac{\partial}{\partial x} q_\psi \circ \text{grad}(f) \circ \psi^{-1}\big|_0 = \left(\sum_k \left[g^{ki}(0) \frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} \right] \right)_{ij}.$$

□

Now that we now how the gradient locally looks like its reasonable to assume, that the flow locally looks like its exponential:

Lemma 2.0.14. *Let $t \in \mathbb{R}$ be small enough and $\varphi_t : M \rightarrow M$ be the flow map corresponding to the negative gradient. Assume that U is the domain of a chart ψ and $p \in U$ is a critical point. Then the linear map $d\varphi_t|_p : T_p M \rightarrow T_{\varphi_t(p)} M$ has a local representation in the coordinates induced by ψ given by:*

$$\frac{\partial}{\partial x} \psi \circ \varphi_t(\psi^{-1}(x)) = \exp \left(- \left(\sum_k g^{ki}(0) \frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} \right)_{ij} \cdot t \right).$$

Proof. For this we consider the function $(t, x) \mapsto \psi \circ \varphi_t(\psi^{-1}(x)) : \mathbb{R} \times U \rightarrow U$. For fixed x we can calculate

$$q_\psi \frac{d}{dt} \varphi_t(\psi^{-1}(x)) = -q_\psi \circ \text{grad}(f)(\varphi_t(\psi^{-1}(x)))$$

and by changing the order of integration we have:

$$\begin{aligned} \frac{\partial}{\partial t} \frac{\partial}{\partial x} \psi \circ \varphi_t(\psi^{-1}(x)) &= \frac{\partial}{\partial x} \frac{\partial}{\partial t} \psi \circ \varphi_t(\psi^{-1}(x)) \\ &= \frac{\partial}{\partial x} q_\psi \frac{d}{dt} \varphi_t(\psi^{-1}(x)) \\ &= -\frac{\partial}{\partial x} q_\psi \circ \text{grad}(f)(\varphi_t(\psi^{-1}(x))) \\ &= -\frac{\partial}{\partial x} q_\psi \circ \text{grad}(f)(\psi^{-1} \circ \psi \circ \varphi_t(\psi^{-1}(x))) \\ &= -\frac{\partial}{\partial x} (q_\psi \circ \text{grad}(f) \circ \psi^{-1}) \frac{\partial}{\partial x} (\psi \circ \varphi_t(\psi^{-1}(x))) \end{aligned}$$

Hence by the theory of linear differential equation with $\frac{\partial}{\partial x} \psi \circ \varphi_0(\psi^{-1}(x)) = \mathbb{1}_m$ we have:

$$\frac{\partial}{\partial x} \psi \circ \varphi_t(\psi^{-1}(x)) = \exp \left(-\frac{\partial}{\partial x} (q_\psi \circ \text{grad}(f) \circ \psi^{-1}) \cdot t \right).$$

The preceding lemma now finishes the proof. □

Hence in a refined Morse chart we have:

Remark 2.0.15. Let ψ be a refined Morse chart, and hence:

- (a) $g^{ik}(0) = \delta_{ik} \mu_k$ with $\mu_k > 0$ for any k .
- (b) $\frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k} = s_k 2\delta_{jk}$ where $s_k = -1$ for $k \leq \lambda_p$ and $s_k = 1$ else.

Then the Jacobian of the flow in the critical point p reads:

$$\begin{aligned} \frac{\partial}{\partial x} \psi \circ \varphi_t(\psi^{-1}(x)) &= \exp(-J(\text{grad}(f))(p) \cdot t) \\ &= \exp \left(- \sum_k \underbrace{g^{ki}}_{=\delta_{ik}\mu_k} (0) \underbrace{\frac{\partial^2 f \circ \psi^{-1}}{\partial x_j \partial x_k}}_{=s_k 2\delta_{jk}} \cdot t \right) \\ &= \text{diag} \left(e^{2\mu_1 t}, \dots, e^{2\mu_{\lambda_p} t}, -e^{2\mu_{\lambda_p+1} t}, \dots, -e^{2\mu_m t} \right) \end{aligned}$$

In a refined Morse chart we have that the (un-)stable Manifold embedding:

Definition 2.0.16 (Stable and Unstable Manifold). Let $p \in M$ be a critical point of $f : M \rightarrow \mathbb{R}$. Then we define:

1. The **stable set** of p as

$$W(\rightarrow p) := \{x \in M \mid \lim_{t \rightarrow \infty} \varphi_t(x) = p\}.$$

That is the points that flow towards p .

2. The **unstable set** of p as

$$W(p \rightarrow) := \{x \in M \mid \lim_{t \rightarrow -\infty} \varphi_t(x) = p\}.$$

That is the points that came from p .

Those sets will turn out to be manifolds, which explains the names **(un-)stable manifolds**.

The main theorem of this chapter will be the:

Theorem 2.0.17 (Stable and unstable manifold theorem). *Let $f : M \rightarrow \mathbb{R}$ be a smooth function on a smooth compact Riemannian manifold (M, g) . Let p be a critical point of f . Then the tangent space splits as:*

$$T_p M = T_p^s(M) \oplus T_p^u M$$

such that the Hessian is positive definite on $T_p^s M$ and negative definite on $T_p^u M$. Moreover, the stable and unstable manifolds are submanifolds and images of smooth embeddings:

$$\begin{aligned} E_p^s : T_p^s M &\rightarrow W(\rightarrow p) \subset M, \\ E_p^u : T_p^u M &\rightarrow W(p \rightarrow) \subset M. \end{aligned}$$

Proof. For the proof compare [BH04]. $\square$

We want to refine the unstable embedding $E_q^u : T_q^u M \rightarrow W(q \rightarrow)$. For this we will make use of the Grobman-Hartman conjugation theorem:

Theorem 2.0.18 (Hartman-Grobman Theorem). *Let E be an open subset of $\mathbb{R}^n$ containing the origin, let $f \in \mathcal{C}$, and let φ_t be a flow given by f . (i.e. $\frac{\partial}{\partial t}(\varphi_t(x)) = f(x)$). Assume that $f(0) = 0$, i.e. that 0 is a fixpoint of φ_t and furthermore assume that fixpoint to be hyperbolic, i.e. Df has no eigenvalue with zero real part. Then there exists a homeomorphism H of an open set U containing the origin onto an open set V containing the origin such that for each $x_0 \in U$, there is an open Interval $I_0 \subseteq \mathbb{R}$ containing zero such that for all $x_0 \in U$ and $t \in I_0$ the diagram*

$$\begin{array}{ccc} U & \xrightarrow{H} & V \\ \downarrow \varphi_t & & \downarrow e^{At} \\ U & \xrightarrow{H} & V \end{array}$$

commutes, meaning that locally the flow is a topological conjugate to its linearisation.

Proof. Compare chapter 2.8 in [Per01]. $\square$

Theorem 2.0.19 (Refined Stable Embedding). *In the setting of the stable manifold theorem, we can refine the embedding*

$$E_q^u : T_q^u M \rightarrow W(q \rightarrow),$$

such that it resembles the flow, meaning that for $x = E_p^s(v)$ the flow is given by

$$\varphi_t(x) = E_p^s(A_t v),$$

where A_t is a linear map defined as: $A_t = B_{\mathfrak{b}} \circ \text{diag}(e^{-2\mu_1 t}, \dots, e^{-2\mu_{\lambda_p} t}) \circ B_{\mathfrak{b}}^{-1}$, where $\mathfrak{b}$ is the bases of $T_q^u M$ that comes from the basis of $T_q M$ which is induced by a refined Morse chart. The diagonal matrix is the stable part of the jacobian of the flow in said refined Morse chart ψ .

Proof. In the proof of the unstable manifold theorem we started with a local map $\rho' : U \rightarrow V$, where $U \subseteq W(q \rightarrow)$ is an neighbourhood of q and $V \supseteq T_q^u$. (This was a refined Morse chart together with a projection once realized that the stable part gets mapped

to the graph of a smooth funktion). Now we define

$$\begin{aligned}\theta_t : V &\rightarrow V \\ v &\mapsto \rho' \circ \varphi_t \circ \rho'^{-1}(v)\end{aligned}$$

for all t such that the map is well defined. Now with a basis $\mathfrak{b}_p^s$ of $T_p^s M$ we can push these considerations into $\mathbb{R}^n$ and by abuse of notation assume that ρ maps to $\mathbb{R}^n$ and hence θ is a flow around the origin of $\mathbb{R}^n$. Now by the Hartman-Grobman Theorem we find a homeomorphism H around the origin, such that $H \circ \theta_t = e^{At} \circ H$, where $A = d\theta_t =$. To calculate said differential we look into the map ρ' again. For this we give names to the funktions:

- $\psi : U \rightarrow V$ is the refined morse chart around q .
- $g : T_p^s M \rightarrow T_p^u M$ is the map that defines the image of the stable part under ψ to be a graph. I.e. $\psi(W(\rightarrow p)) = \Gamma_g$. We know that $g(0) = 0$ and $dg|_0 = 0$.
- $\pi : \Gamma_g \rightarrow T_p^s M$ is the projection. Its inverse is obviously $\text{id} \times g$.

We know from the local stable manifold theorem, that $dg|_0 = 0$. With this we can calculate:

$$\begin{aligned}d\theta_t|_0 &= d(\pi \circ \psi \circ \varphi_t \circ \psi^{-1} \circ \pi^{-1})|_0 \\ &= d\pi|_{(0,0)} \circ d\psi \circ \varphi_t \circ \psi^{-1}|_{0,0} \circ d\text{id} \times g|_0 \\ &= (\text{id}_{m-\lambda_p}, 0_{\lambda_p}) \circ \text{diag}(e^{-2\mu_1 t}, \dots, e^{-2\mu_{\lambda_p} t}, e^{2\mu_{\lambda_q+1} t}, \dots, e^{2\mu_m t}) \circ (\text{id}_{m-\lambda_p}, 0_{\lambda_p})^T \\ &= \text{diag}(e^{-2\mu_1 t}, \dots, e^{-2\mu_{\lambda_p} t})\end{aligned}$$

Now define for a small neighbourhood $V \subseteq \mathbb{R}^{m-\lambda_p}$ the map

$$\rho := \rho' \circ H : V \rightarrow U.$$

U is choosen such that the map is onto. Then we have that $\varphi_t \circ \rho^{-1} = \rho^{-1} \circ e^{At}$, in a that small neighbourhood of p . Now finally we can extend this property to be true for the whole stable set: For this we define the embedding

$$\begin{aligned}E_p^s : T_p^s M &\rightarrow W(\rightarrow p) \\ x &\mapsto \varphi_{-n} \circ \rho^{-1} \circ A_n(x) \quad \text{where } n \in \mathbb{N} \text{ is big enough such that } A_n(x) \in V.\end{aligned}$$

This map is in fact well defined since

$$\begin{aligned}\varphi_n \circ \rho^{-1} \circ A_n(x) &= \varphi_{-n} \circ \varphi_{-1} \circ \varphi_1 \circ \rho^{-1} \circ A_n(x) \\ &= \varphi_{-n} \circ \varphi_{-1} \circ \rho^{-1} \circ A_{n+1}(x) \\ &= \varphi_{-(n+1)} \circ \rho^{-1} \circ A_{n+1}(x)\end{aligned}$$

And obviously the flow compatibility that is given locally by ρ is now true globally: Let $x \in W(\rightarrow p)$ and assume $x = E_p^s(y)$. Then:

$$\begin{aligned}\varphi_t(x) &= \varphi_t(E_p^s(y)) = \varphi_t \circ \varphi_{-n} \circ \rho^{-1} \circ A_n(y) \\ &= \varphi_{-n} \circ \varphi_t \circ \rho^{-1} \circ A_n(y) \\ &= \varphi_{-n} \circ A_n(A_t(y))\end{aligned}$$

□

Now we want to understand the flow in an unstable set and analyze two maps:

Remark 2.0.20. Let q be a critical point and $c < f(q)$ such that no other critical point lies in $f^{-1}[c, f(q)]$ then, the map

$$\begin{aligned}\tau : W(q \rightarrow) \setminus \{q\} &\rightarrow \mathbb{R} \\ x &\mapsto \tau(x) \text{ such that } f \circ \varphi_{\tau(x)}(x) = c\end{aligned}$$

is smooth.

Proof. We know that such a map exists and is well defined since every flow line passes through $f^{-1}(c)$. Now let $x \in W(q \rightarrow) \setminus \{q\}$, V be a neighbourhood of x and U be a neighbourhood of $\varphi_{\tau(x)}(x)$. Define the map

$$\begin{aligned}G : V \times I &\rightarrow \mathbb{R} \\ (x, t) &\mapsto f \circ (\varphi_t(x)) .\end{aligned}$$

Now the partial derivative reads

$$\frac{\partial}{\partial t} G|_{x, t} = df|_{\varphi_t(x)} \circ \frac{\partial}{\partial t} \varphi_t(x)|_t = df|_{\varphi_t(x)} \circ (-\text{grad}_{\varphi_t(x)}(f)) .$$

For $\varphi_{\tau(x)}$ we have that $T_{\varphi_{\tau(x)}(x)} f^{-1}(c) = \ker df|_{\varphi_{\tau(x)}(x)}$. Since the flow line is transversal to the level set or said differently, the gradient is orthogonal to level sets, we have that the latter does not live in the kernel and hence

$$\frac{\partial}{\partial t} G|_{x, \tau(x)} = df|_{\varphi_{\tau(x)}(x)} \left(-\text{grad}_{\varphi_{\tau(x)}(x)}(f) \right) \neq 0 .$$

This lets us use the implicit function theorem to deduce that the level set $G = c$ is locally the graph of a function $\tau : U_x \rightarrow \mathbb{R}$ where U_x is a neighbourhood of x . Hence the function τ is globally smooth since this argument can be done everywhere. □

Remark 2.0.21. Now we care for the differential of τ in a point $x_j \in W(q \rightarrow) \cap f^{-1}(c)$. For this we choose a chart of $W(q \rightarrow)$ around x_j such that the induced bases of $T_{x_j} W(q \rightarrow) = \langle -\text{grad}_{x_j}(f) \rangle \oplus T_{x_j} V_j$ that respects the splitting. for example $\frac{\partial}{\partial x^1}|_{x_j} =$

$-\text{grad}_{x_j}(f)$ and $\frac{\partial}{\partial x^i}|_{x_j} \in T_{x_j}V_j$ for i bigger than one. With this we can start our calculations: Notice how $x \mapsto f \circ \varphi_{\tau(x)}(x)$ is a constant function and hence its differential is 0.

$$\begin{aligned} 0 = d_x(f \circ \varphi_\tau) &= \overbrace{df}^{\text{1-form}}(d\varphi_\tau) = df\left(d_t\varphi_t \circ d_x\tau(x) + d_x\varphi_{\tilde{t}=\tau(x)}\right) \\ &= df\left(d_t\varphi_t \circ d_x\tau(x)\right) + df\left(d_x\varphi_{\tilde{t}=\tau(x)}\right) \\ &= d_t(f \circ \varphi_t) \circ d_x\tau(x) + df\left(d_x\varphi_{\tilde{t}=\tau(x)}\right) \end{aligned}$$

Hence we know that

$$d_x\tau = -\frac{df\left(d_x\varphi_{\tilde{t}=\tau(x)}\right)}{d_t(f \circ \varphi_t)} = -\frac{df\left(d_x\varphi_{\tilde{t}=\tau(x)}\right)}{-\|\text{grad}(f)^2\|} \quad (5)$$

Now we can calculate in x_j where $\tau(x_j) = 0$

$$d_x\tau|_{x_j} \frac{\partial}{\partial x^i}|_{x_j} = \frac{df|_{x_j} \frac{\partial}{\partial x^i}|_{x_j}}{\|\text{grad}(f)^2\|} = \frac{g\left(\text{grad}_{x_i}(f), \frac{\partial}{\partial x^i}|_{x_j}\right)}{\|\text{grad}(f)^2\|} = \begin{cases} -1 & \text{if } i = 1 \\ 0 & \text{else.} \end{cases}$$

This result can be extended to also work outside of x_j as follows: Notice the identity for all $s \in \mathbb{R}$ and $x \in W(q \rightarrow) \setminus \{q\}$:

$$\tau(\varphi_s(x)) = \tau(x) - s.$$

Differentiating this in the variable s and in for s' yealds

$$d\tau|_{\varphi_{s'}(x)} \circ \underbrace{d\varphi_s(x)|_{s=s'}}_{=-\text{grad}_{\varphi_s(x)}(f)} = d\tau(\varphi_{s'}(x))|_{s'} = d\tau(x) - s|_{s'} = -1.$$

And hence in $s' = 0$ we have for all $x \in W(q \rightarrow) \setminus \{q\}$:

$$d\tau|_x(-\text{grad}(x)(f)) = -1.$$

We can furthermore define a basis of $T_yW(q \rightarrow)$ with $\mathbf{b}_1 := -\text{grad}_y(f)$ and $\mathbf{b}_i = d\varphi_{\tau(y)}|_y^{-1}(w_i)$ where (w_i) denotes a basis of $T_{x_j}V_j$. Hence w_i is orthogonal on $-\text{grad}_{x_j}(f)$ for all $i > 1$. Hence by the equation (5) we can calucate:

$$d_x\tau(\mathbf{b}_i) = -\frac{df\left(d_x\varphi_{\tilde{t}=\tau(x)}(\mathbf{b}_i)\right)}{-\|\text{grad}(f)^2\|} = -\frac{df(w_i)}{-\|\text{grad}(f)^2\|} = 0 \quad \text{for all } i > 1.$$

Definition 2.0.22. Let q be a critical point and c such that $c < f(q)$ and $f^{-1}[c, f(q)] = q$. Then we define the map

$$\begin{aligned} \rho : W(q \rightarrow) \setminus \{q\} &\rightarrow W(q \rightarrow) \cap f^{-1}(c) \\ x &\mapsto \varphi_{\tau(x)}(x). \end{aligned}$$

Lemma 2.0.23. Assume that $\mathfrak{d} = (\mathfrak{d}_i)$ is a basis of $T_q^u M$. Let $y := E_q^{u-1}(x_j)$ and $y' \in (\rho \circ E_q^u)^{-1}(x_j)$. Then the map

$$dE_q^u|_y^{-1} \circ d\rho \circ dE_q^u|_{y'} \circ Q_{y'} \circ B_{\mathfrak{d}} : \mathbb{R}^{\lambda_q} \rightarrow T_y T_q^u M$$

can be written as

$$\begin{aligned} \pi_{-\text{grad}_{x_j}(f)}^\top \circ (B_{\mathfrak{b}_{x_j}^q}) : \mathbb{R}^{\lambda_q} &\rightarrow T_{x_j} V_j \\ e_i &\mapsto \begin{cases} 0 & \text{for } i = 1 \\ (\mathfrak{b}_{x_j}^q)_i & \text{for } i \neq 1. \end{cases} \end{aligned}$$

where $(\mathfrak{b}_{x_j}^q)_j$ is basis of $T_{x_j} W(q \rightarrow)$ that is induced from $\mathfrak{d}$ and $(\mathfrak{b}_{x_j}^q)_1 = -\text{grad}_{x_j}(f)$. In other words, this is a basis compatible with the one used in the counting of flow lines in the Morse boundary operator.

Proof Strategy:

To prove this, we first remind ourself how the basis $(\mathfrak{b}_{x_j}^q)_{j>1}$ is constructed. It is a basis that when extended with $-\text{grad}_{x_j}(f)$ as the first vector is compatible with $\mathfrak{d}$. Hence, we can construct it from $\mathfrak{d}$, by assuming that $dE_q^u|_y \circ Q_y \circ \mathfrak{d}_1 = -\text{grad}_y(f)$. Then, if we push $\mathfrak{d}$ to $T_{x_j} W(q \rightarrow)$ via $dE_q^u|_y \circ Q_y$ and delete the first vector, we get the Basis we wanted. So what we need to check in order to proof this lemma is, if the maps

$$\begin{aligned} dE_q^u|_y \circ Q_y \circ B_{\mathfrak{d}} : \mathbb{R}^{\lambda_q-1} &\rightarrow \left(-\text{grad}_{x_j}(f)\right)^\top \\ dE_q^u|_y^{-1} \circ d\rho \circ dE_q^u|_{y'} \circ Q_{y'} \circ B_{\mathfrak{d}} : \mathbb{R}^{\lambda_q-1} &\rightarrow \left(-\text{grad}_{x_j}(f)\right)^\top \end{aligned}$$

induce the same orientation when restricted to the same domain such that they become vector space isomorphisms. To proof this we will work from the bottom up: We adapt the basis $\mathfrak{d}$ to work with the lower map and show, later, that the adapted basis fits the requirements above:

Proof. We start by defining the space $(\tau \circ E_q^u)^{-1}(\tilde{t})$, where $\tilde{t} := \tau \circ E_q^u(y')$. As a level set, this is a submanifold of $T_q^u M$ and its tangend space in y' is given by the basis $E_q^u|_{y'}^{-1}(\mathfrak{b}_i)$ for $i > 1$, where $\mathfrak{b}_i = d\varphi_{\tilde{t}}|_{y'}^{-1}(w_i)$ where (w_i) denotes a basis of $T_{x_j} V_j$. Now we have the equality of maps:

$$A_{\tilde{t}}|_{(\tau \circ E_q^u)^{-1}(\tilde{t})} = (E_q^{u-1} \circ \rho \circ E_q^u)|_{(\tau \circ E_q^u)^{-1}(\tilde{t})}$$

Hence for a basis $\mathfrak{d}_i$ of $T_q^u M$ such that $dE_q^u|_{y'} \circ Q_{y'}(\mathfrak{d}_i)$ is $-\text{grad}_{y'}(f)$ for $i = 1$ and $\mathfrak{b}_i$ else,

we have:

$$d(E_q^{u-1} \circ \rho \circ E_q^u)|_{y'} \circ Q_{y'}(\mathfrak{d}_i) = Q_y \circ A_{\tilde{t}}(\mathfrak{d}_i).$$

For $i = 1$, the left side maps to 0, and the right side:

$$\begin{aligned} Q_y \circ A_{\tilde{t}}(\mathfrak{d}_1) &= dA_{\tilde{t}}|_{y'} \circ Q_{y'}(\mathfrak{d}_1) \\ &= dA_{\tilde{t}}|_{y'} \circ dE_q^u|_{y'}^{-1}(-\text{grad}_{y'}(f)) \\ &= dA_{\tilde{t}} \circ dE_q^u|_{y'}^{-1}(-\text{grad}_{y'}(f)) \\ &= d(E_q^u)^{-1} \circ \varphi_{\tilde{t}}|_{y'}^{-1}(-\text{grad}_{y'}(f)) \\ &= dE_q^u|_{x_j}^{-1} \circ d\varphi_{\tilde{t}}|_{y'}^{-1}(-\text{grad}_{y'}(f)) \\ &= dE_q^u|_{x_j}^{-1}(-\text{grad}_{x_j}(f)) \end{aligned}$$

The last equation is the general fact proven in remark 2.0.15, that for fixed t the identity holds:

$$d_x \varphi_t|_y(-\text{grad}_y(f)) = -\text{grad}_{\varphi_t(y)}(f)$$

Hence if π denotes the projection onto the complement of $dE_q^u|_{x_j}^{-1}(-\text{grad}_{x_j}(f))$ we have an equality of maps

$$d(E_q^{u-1} \circ \rho \circ E_q^u)|_{y'} \circ Q_{y'} = \pi \circ Q_y \circ A_{\tilde{t}}.$$

Or by the above calculation, we can throw out π , if we exclude $\mathfrak{d}_1$. Furthermore, the equation shows that our requirement $dE_q^u|_{y'} \circ Q_{y'}(\mathfrak{d}_1) = -\text{grad}_{y'}(f)$ implies that $\mathfrak{d}$ is compatible with a basis where $dE_q^u|_y \circ Q_y \circ \mathfrak{d}_1 = -\text{grad}_y(f)$, since the sign of the determinant of $A_{\tilde{t}}$ is always positive. This concludes the prove! $\square$

Definition 2.0.24 (Morse Homology). Let $f : M \rightarrow \mathbb{R}$ be a Morse-Smale function on a Riemannian manifold (M, g) . Furthermore let $\mathfrak{o}$ be an choice of orientations for all unstable Tangendspaces ($T_q^u M$ for all $q \in \text{Crit}(f)$). The tuple $\mathfrak{A} := (g, f, \mathfrak{o})$ is a **Morse datum**. With this we define now the **Morse chain groups** for a given commutative ring with one (Λ) :

$$C_k(M, \mathfrak{A}, \Lambda) := \bigoplus_{\text{Crit}(f)} \Lambda$$

To define a boundary operator between the chain groups on the generators we count flow lines between them. Lets say that p is a critical point of index k and q one of index $k + 1$. Then we have the set of flow lines from q to p denoted by $\mathcal{M}(q \rightarrow p)$. For each $\gamma \in \mathcal{M}(q \rightarrow p)$ we assign a sign $n_\gamma \in \{\pm 1\}$. Let $x \in \gamma$. Then the tangent space in x splits $T_x W(q \rightarrow) = N_x(\gamma) \oplus \langle -\text{grad}(f)(x) \rangle$. Furthermore, we can parallel transport $N_x(\gamma)$ to $T_p^u M$. Hence, the orientation in $T_q^u M$ can be compared to the one in $T_p^u M$ and the sign n_γ can be chosen accordingly. The **boundary operator** on a generator is defined as:

$$\partial_{k+1}(q) := \sum_{p \in \text{Crit}_k(f)} \sum_{\gamma \in \mathcal{M}(q \rightarrow p)} n_\gamma$$

This gives rise to a chain complex and hence gives rise to a homology theory, and a cohomology theory by applying the hom functor to the chain complex.

Definition 2.0.25 (Orientation and Exact Sequences). As stated before, an orientation of a Vector space V of dimension m is given by an ordered basis $\mathbf{b} = (b_1, \dots, b_m)$ of V and hence the produkt $b_1 \wedge \dots \wedge b_m$ is an element of the maximal exterior product $\det(V) = \Lambda^m V$. Two ordered bases define different elements in $\det(V)$ where their difference is given by the determinant of the base change matrix. (This is just a calculation using multilinearity and the alternating property). Hence an orientation is a choice of one connected component of $\det(V) \setminus \{0\}$. Now a short exact sequence of vector spaces

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

induces a canonical isomorphism

$$\det(A) \otimes \det(C) \rightarrow \det(B) \\ (a_1 \wedge \dots \wedge a_l) \otimes (c_1 \wedge \dots \wedge c_k) \mapsto (f(a_1) \wedge \dots \wedge f(a_l) \wedge \tilde{c}_1 \wedge \dots \wedge \tilde{c}_k),$$

where $\tilde{c}_j \in g^{-1}(c_j)$. The well definition of this map (invariance under chosen lift of c) can be calculated, since two different choices differ in an element in the kernel of g which equals the image of f and hence the different choices vanish in the wedge produkt. By this isomorphism we get an orientation induced in the middle from the outside.

Remark 2.0.26 (The Counting of Flow Lines). There are several ways to formulate the counting of flow lines that happens in the boundary operator. Let $\gamma \in \mathcal{M}(q \rightarrow p)$ and $x \in \gamma$. Instead of using the parallel transport, we can introduce two different ordered bases in $T_x W(q \rightarrow)$. We use the following notation: Let $\gamma \in \mathcal{M}(q \rightarrow p)$ and $x \in \gamma$. For this we define $N_x^{W(q \rightarrow)} \gamma$ be the normal bundle of γ as a submanifold of $W(q \rightarrow)$. Since γ is contractible we can trivialize said normal bundle to get a bundle isomorphism:

$$\nu : N^{W(q \rightarrow)} \gamma \rightarrow \gamma \times T_p^u M.$$

Let $\nu|_x$ denote the isomorphism $N_x^{W(q \rightarrow)} \gamma \rightarrow T_p^u M$. An orientation of $T_p^u M$ induces then one on $N_x^{W(q \rightarrow)} \gamma$ and if we define the bases $(-\text{grad}_x(f))$ of its own span to be positive, we get the exact sequence

$$0 \longrightarrow \langle -\text{grad}_x(f) \rangle \longrightarrow T_x W(q \rightarrow) \longrightarrow N_x^{W(q \rightarrow)} \gamma \longrightarrow 0 \\ \uparrow (\nu_x)^{-1} \\ T_p^u M$$

That introduces an orientation in the middle that can be represented by the basis $(-\text{grad}_x(f), i \circ \nu|_x)(\mathbf{b}^p)$, where i is the inclusion $N_x^{W(q \rightarrow)} \gamma \rightarrow T_x W(q \rightarrow)$ and $(\mathbf{b}^p)$ is an ordered basis of $T_p^u M$. Another way to get an orientation there is to trivialize the bundle $TW(q \rightarrow)$ via the isomorphism:

$$\mu : TW(q \rightarrow) \rightarrow W(q \rightarrow) \times T_q^u M; \xi_y \mapsto \left(y, Q_{E_q^{u-1}(y)}^{-1} \circ dE_q^{u-1}|_y(\xi_y) \right).$$

This trivialization together with an orientation on $T_q^u M$ orients the middle in the above exact sequence. Now the deciding factor that gives γ its sign is whether the two orientations agree. This can be checked by calculating the determinant of a certain map. To formulate the map somewhat readably, we assume that $\mathfrak{b}^q$ is a basis of $T_q^u M$ such that for $y = E_q^{u-1}(x)$ we have $dE_q^u|_y \circ Q_y(\mathfrak{b}_i^q) = -\text{grad}_x(f)$ for $i = 1$, and for all $i > 1$ the image lives in the orthogonal complement of $-\text{grad}_x(f)$ and hence in the domain of ν_x . Then we can define the map

$$B_{\mathfrak{b}^p}^{-1} \circ \nu_x \circ \mu_x^{-1} \circ B_q : \mathbb{R}^{\lambda_q} \supseteq \mathbb{R}^{\lambda_q-1} = \langle e_2, \dots, e_{\lambda_q} \rangle \rightarrow \mathbb{R}^{\lambda_q-1}$$

$$e_i \mapsto B_{\mathfrak{b}^p}^{-1} \circ \nu_x \circ dE_q^u|_y \circ Q_y(\mathfrak{b}_i^q),$$

and the determinant of said map gives us n_γ .

Remark 2.0.27. Notice how we already saw a different way to calculate the sign by the map from lemma 2.0.23.

Definition 2.0.28. Let $(M, \mathfrak{A}, \Lambda)$ and $(M', \mathfrak{A}', \Lambda)$ be two Morse data with $\mathfrak{A} = (g, f, \mathfrak{o})$ and $\mathfrak{A}' = (g', f', \mathfrak{o}')$. Then on $M \times M'$ we define the Morse function $f + f'$ and the metric $g \oplus g'$. Then $d_{(x, x')}(f + f') = d_x(f) + d_{x'}(f')$ and hence the critical points of $f + f'$ are pairs (p, p') such that p is critical for f and p' is critical for f' . By corollary 2.0.5 we conclude that the hessian splits into block diagonal form and hence the property of non degeneracy follows. Since we use the product metric $g \oplus g'$ we also get a splitting gradient of $f + f'$:

$$\begin{aligned} d_{(x, x')}(f + f')(V, V') &= d_x(f)(V) + d_{x'}(f')(V') \\ &= g(\text{grad}_x(f), V) + g'(\text{grad}_{x'}(f'), V') \\ &= (g \oplus g')(\underbrace{(\text{grad}_x(f), \text{grad}_{x'}(f'))}_{\text{grad}_{(x, x')}(f + f')}, (V, V')). \end{aligned}$$

Hence the flow also splits and thereby the (un-)stable manifolds split. Hence we can orient all unstable manifolds via that splitting which gives us $\mathfrak{o} \oplus \mathfrak{o}'$. Finally transversal intersection is a property that carries over to products and hence the Morse datum $(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda)$ with $\mathfrak{A} \times \mathfrak{A}' := (g \oplus g', f + f', \mathfrak{o} \oplus \mathfrak{o}')$ really gives a valid Morse datum. The splitting of the flow gives a splitting of the connecting flow lines and hence of the modular space:

$$\mathcal{M}((q, q') \rightarrow (p, p')) = \mathcal{M}(q \rightarrow p) \times \mathcal{M}(p' \rightarrow p')$$

Assume that $p \neq p'$ and $q \neq q'$. Then for the modular space to be non empty we need

$$\text{ind}(q) > \text{ind}(p) \quad \text{and} \quad \text{ind}(q') > \text{ind}(p').$$

This results in the index of (q, q') being at least two bigger than the index of (p, p') . Hence, if we want to count flow lines between two consecutive indices, either $p = p'$ or $q = q'$. Then we have the modular space

$$\mathcal{M}((q, q') \rightarrow (p, p')) = \begin{cases} p \times \mathcal{M}(q', p') & \text{if } p = q \\ \mathcal{M}(q, p) \times p' & \text{if } p' = q'. \end{cases}$$

Now we want to analyse how the splitting affects the counting of flow lines. For this we go through our classical procedure given in the definition 2.0.24. Assume that $q' = p'$, and let γ be an element of $\mathcal{M}(q, p)$ with $x \in \gamma$. Then $(\gamma, q') \in \mathcal{M}((q, q'), (p, q'))$. Now let (b, b') be a basis of

$$T_{(x, q')}W((q, q') \rightarrow) = T_xW(q \rightarrow) \times T_{q'}W(q' \rightarrow).$$

that is oriented with respect to $T_{q, q'}^u M \times M'$ and assume that $-\text{grad}_x(f)$ is the last vector in b . Now manipulate this basis such that it fits into the splitting

$$T_{(x, q')}W((q, q') \rightarrow) = N_{(x, q')}(\gamma, q') \oplus -\text{grad}_{x, q'}(f + f').$$

by moving $-\text{grad}_x(f)$ to the back. This requires $\text{ind}_{q'}(f')$ many transpositions. Hence if $\tilde{b}$ denotes b without its last vector we have the equality :

$$(b, b') \text{ carries the same orientation as } (-1)^{\text{ind}_{q'}(f')}(\tilde{b}, b', -\text{grad}_{(x, x')}(f + f')).$$

Now moving the basis $(\tilde{b}, b, -\text{grad}_{(x, x')}(f + f'))$ down along the flow and comparing it to the orientation of

$$T_{p, q'}W((p', q') \rightarrow) = T_pW(p \rightarrow) \times T_{q'}W(q' \rightarrow)$$

is the same just comparing $\tilde{b}$ with the orientation of $T_pW(p \rightarrow)$ which again is the sign n_γ . Hence: $n_{\gamma, q'} = (-1)^{\text{ind}_{q'}(f')} n_\gamma$. Going through the same argument for the case that $q = p$ is easier, since we can assume $-\text{grad}_x(f)$ to already be in the back and hence we get for critical points (q, q') and (p, p') with consecutive indices:

$$n((q, q'), (p, p')) = \begin{cases} n(q', p') & \text{if } q = p \\ (-1)^{\text{ind}_{q'}(f')} n(q, p) & \text{if } q' = p' \\ 0 & \text{else.} \end{cases}$$

The consideration of products stems from [AD14]. However she is missing the sign in her consideration of the boundary map of the product.

Remark 2.0.29. Let $(M, \mathfrak{A}, \Lambda)$ and $(M', \mathfrak{A}', \Lambda)$ be two Morse data with $\mathfrak{A} = (g, f, \mathfrak{o})$ and $\mathfrak{A}' = (g', f', \mathfrak{o}')$. Then the map

$$\Phi : \bigoplus_{i+j=k} C_i(M, \mathfrak{A}, \Lambda) \otimes_\Lambda C_j(M', \mathfrak{A}', \Lambda) \rightarrow C_k(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda)$$

Is an isomorphism of groups and furthermore if we denote the boundary morphism in the chaincomplex corresponding to M by ∂_f , the one corresponding to M' by $\partial_{f'}$ and the one corresponding to the product $M \times M'$ by $\partial_{f+f'}$ we get an isomorphism of chain complexes induced by Φ :

$$\left(\bigoplus_{i+j=*} C_i(M, \mathfrak{A}, \Lambda) \otimes_\Lambda C_j(M', \mathfrak{A}', \Lambda), (-1)^j \partial_f \otimes 1 + 1 \otimes \partial_{f'} \right) \rightarrow (C_*(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda), \partial_{f+f'})$$

Here the left side defines the tensor product of chain complexes. Hence we get an natural isomorphism of chain complexes:

$$C_*(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C_*(M', \mathfrak{A}', \Lambda) \rightarrow C_*(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda).$$

Proof. The isomorphism of groups is easy to see, since we have an bijection of the generators

$$(p \otimes q) \leftrightarrow (p, q).$$

Since both sides are free Λ -modules we can apply the universal property of free modules giving us an canonical isomorphism of Λ modules and hence an group isomorphism. Now we want to show that we get a isomorphism of chain complexes and hence need to check if the boundary map commutes with Φ . Let a be a critical point of index i of f and let a' be a critical point of index j of f' . On the one hand we can calculate

$$\begin{aligned} \Phi((-1)^j \partial_f \otimes 1 + 1 \otimes \partial_{f'})(a \otimes a') &= \Phi((-1)^j \partial_f(a) \otimes a' + a \otimes \partial_{f'}(a')) \\ &= \Phi\left(\sum_{b \in \text{Crit}_{i-1}(f)} (-1)^j n(a, b) b \otimes a' + a \otimes \sum_{b' \in \text{Crit}_{j-1}(f')} n(a', b') b'\right) \\ &= \sum_{b \in \text{Crit}_{i-1}(f)} (-1)^j n(a, b) (b, a') + \sum_{b' \in \text{Crit}_{j-1}(f')} n(a', b') (a, b'), \end{aligned}$$

and on the other hand

$$\begin{aligned} \partial_{f+f'}(\Phi(a \otimes a')) &= \partial_{f+f'}(a, a') \\ &= \sum_{(b, b') \in \text{Crit}_{i+j-1}(f+f')} n((a, a'), (b, b')) (b, b') \\ &= \underbrace{\sum_{b \in \text{Crit}_{i-1}(f)} (-1)^j n(a, b) (b, a')}_{a'=b'} + \underbrace{\sum_{b' \in \text{Crit}_{j-1}(f')} n(a', b') (a, b')}_{a=b}. \end{aligned}$$

□

Lemma 2.0.30 (Algebraic Künneth Formula). *If Λ is a principle ideal domain, and C and C' are chain complexes over Λ such that all C_n are free Λ modules, then for each n there is a natural short exact sequence*

$$0 \longrightarrow \oplus_i (H_i(C) \otimes_{\Lambda} H_{n-i}(C')) \longrightarrow H_n(C \otimes_{\Lambda} C') \longrightarrow \oplus_i (\text{Tor}_{\Lambda}(H_i(C), H_{n-i-1}(C'))) \longrightarrow 0$$

that splits.

Proof. The proof is purely homological algebra and standard in algebraic topology textbooks. Compare Theorem 3B.5. in [Hat02] for example. □

Theorem 2.0.31 (Morse-theoretic Künneth Formula). *If Λ is a principle ideal domain, and let $(M, \mathfrak{A}, \Lambda)$ and $(M', \mathfrak{A}', \Lambda)$ be two Morse data. Then for each n there is a splitting short exact sequence*

$$\begin{aligned} 0 \rightarrow \bigoplus_i (H_i(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} H_{n-i}(M', \mathfrak{A}', \Lambda)) &\rightarrow H_n(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda) \\ &\rightarrow \bigoplus_i (\text{Tor}_{\Lambda}(H_i(M, \mathfrak{A}, \Lambda), H_{n-i-1}(M', \mathfrak{A}', \Lambda))) \rightarrow 0 \end{aligned}$$

Proof. The theorem follows directly from the Lemma above together with the natural isomorphism of chain complexes

$$C_*(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C_*(M', \mathfrak{A}', \Lambda) \rightarrow C_*(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda).$$

□

Definition 2.0.32 (Morse Cohomology). Let G be a commutative ring with one. To get the cohomology we have to dualize the chain complex. Let $\text{Crit}_k(f) = \{p_1, \dots, p_l\}$ be a basis of the k -th chain group $C_k(M, \mathfrak{A}, G)$. Then the functionals

$$\eta_i : C_k(M, \mathfrak{A}, G) \rightarrow G; p_j \mapsto \begin{cases} 1 & \text{if } j=i \\ 0 & \text{else.} \end{cases}$$

are a basis of the dual group $C^k(M, \mathfrak{A}, G)$. The coboundary map is easy to obtain too. Let q_j for $j = 1, \dots, r$ be the critical points with index $k+1$ and let ξ_i be the corresponding forms. Then let $n_f(q_j, p_i)$ be the number of flow lines from q_j to p_i counted with the sign. Then the coboundary map is given as:

$$\begin{aligned} \partial^k : C^k(M, \mathfrak{A}, G) &\rightarrow C^{k+1}(M, \mathfrak{A}, G) \\ \eta_i &\mapsto \partial^k \eta_i = \eta_i \circ \partial_{k+1} = \sum_j n_j(q_j, p_i) \xi_j. \end{aligned}$$

Theorem 2.0.33 (Morse-theoretic Cohomological Künneth Formular). *If Λ is a principle ideal domain, and let $(M, \mathfrak{A}, \Lambda)$ and $(M', \mathfrak{A}', \Lambda)$ be two Morse data. Then for each n there is a splitting short exact sequence*

$$\begin{aligned} 0 \rightarrow \bigoplus_i (H^i(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} H^{n-i}(M', \mathfrak{A}', \Lambda)) &\rightarrow H^n(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda) \\ &\rightarrow \bigoplus_i (\text{Tor}_{\Lambda}(H^i(M, \mathfrak{A}, \Lambda), H^{n-i+1}(M', \mathfrak{A}', \Lambda))) \rightarrow 0 \end{aligned}$$

Proof. We show this by giving a natural isomorphism of the chain complexes

$$C^*(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C^*(M', \mathfrak{A}', \Lambda) \rightarrow C^*(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda).$$

First we need to give an group isomorphism

$$\Phi : \bigoplus_{i+j=k} C^i(M, \mathfrak{A}, \Lambda) \otimes_{\Lambda} C^j(M, \mathfrak{A}', \Lambda) \rightarrow C^k(M \times M', \mathfrak{A} \times \mathfrak{A}', \Lambda)$$

This isomorphism however is clear and given on the generators : Let a and a' be critical points of index i and j with respect to f and f' . Let η and η' be the corresponding generators of $C^i(M, \mathfrak{A}, \Lambda)$ and $C^j(M, \mathfrak{A}', \Lambda)$. Then we have the bijection

$$\eta \otimes \eta' \leftrightarrow (\eta, \eta').$$

To get an chain isomorphism we need to check the compatibility with the coboundary maps. For each $b_l \in \text{Crit}_{i+1}(f)$ let ξ_l be the corresponding form and similar for each $z_k \in \text{Crit}_{j+1}(f')$ let ζ_k be the corresponding form. Then we calculate

$$\begin{aligned} \Phi \left((-1)^j \partial^f \otimes 1 + 1 \otimes \partial^{f'} \right) (\eta \otimes \eta') &= \Phi \left((-1)^j \partial^f (\eta) \otimes \eta' + \eta \otimes \partial^{f'} (\eta') \right) \\ &= \Phi \left(\sum_l (-1)^j n(b_l, a) \xi_l \otimes \eta' + \sum_k n(z_k, a') \eta \otimes \zeta_k \right) \\ &= \sum_l (-1)^j n(b_l, a) (\xi_l, \eta') + \sum_k n(z_k, a') (\eta, \zeta_k) \end{aligned}$$

and the other way around, if we define for (x, x') with $\text{ind}_{f+f'}(x, x') = \text{ind}_{f+f'}(a, a') + 1$ the corresponding form to be ν we have:

$$\begin{aligned} \partial^{f+f'} \Phi(\eta \otimes \eta') &= \partial^{f+f'} (\eta, \eta') \\ &= \sum_{(x, x') \in \text{Crit}_{i+j+1}(f+f')} n((a, a'), (x, x')) (\nu) \\ &= \underbrace{\sum_l (-1)^j n(b_l, a) (\xi_l, \eta')}_{a'=x'} + \underbrace{\sum_k n(z_k, a') (\eta, \zeta_k)}_{a=x} \end{aligned}$$

Now finally we can apply the Algebraic Künneth Formular for Cohomology and we are done. The latter formular is almost the same is the homological one but with a different index shift in the Tor functor. Again this is standart homological algebra. $\square$

Corollary 2.0.34. If M is an oriented manifold, orientations from the unstable tangend space in critical points $T_q^u M$ induce Orientations in the stable tangend space since

$$T_q M = T_q^u M \otimes T_q^s M.$$

3 Morse-Theoretic Atiyah-Hirzebruch Spectralsequence

3.1 Conley index

This is just a short summary. Prooves are omitted and the goal is to remaind ourself of the definitinos.

Definition 3.1.1 (Compact invariant isolated subset). For a flow $\varphi_t : M \rightarrow M$ on a locally compact metric space we call a subset $S \subseteq M$ **invariant subspace** if and only if $\varphi_t(S) = S$ for all $t \in \mathbb{R}$. For any subset $N \subseteq M$ we define the **maximal invariant subset**

$$\begin{aligned} I(N) &= \{x \in N \mid \varphi_t(x) \in N \forall t \in \mathbb{R}\} \\ &= \bigcap_{t \in \mathbb{R}} \varphi_t(N). \end{aligned}$$

A **compact invariant subset** S is called **isolated**, if a compact neighborhood N exists, such that $I(N) = S$.

Definition 3.1.2 (Index pairs). Let S be an isolated compact invariant subset. A topological pair (N, L) of compact subsets of M where $L \subseteq N$ is called an **index pair of S**, if it satisfies the following:

1. $S = I(\overline{N \setminus L}) \subseteq (N \setminus L)$.
2. $x \in L$ and $\varphi_{[0,t]}(x) \subseteq N$ implies that $\varphi_{[0,t]}(x) \subseteq L$. We call L **positively invariant in N** . Here $\varphi_{[0,t]}(x) := \{\varphi_{\tilde{t}}(x) \mid \tilde{t} \in [0, t]\}$.
3. For all $x \in N$ such that a t exists, with $\varphi_t(x) \notin N$, there exists a t' with $\varphi_{[0,t']}(x) \subseteq N$ and $\varphi_{t'}(x) \in L$.

This definition captures how the flow lines leave the invariant space S . Due to the third property we call L the **exit set**.

Definition 3.1.3 (Regular index pairs). We call an index pair (N, L) **regular** if and only if the inclusion $I : L \hookrightarrow N$ is a cofibration.

Theorem 3.1.4 (Homotopy equivalence of index pairs). *If S is an isolated compact invariant set and (N, L) and $(\tilde{N}, \tilde{L})$ are two index pairs of S . Then N/L and $\tilde{N}/\tilde{L}$ are homotopy equivalent as pointed spaces. This homotopie is canonical in transitive up to homotopy.*

Definition 3.1.5 (Connected Simple System). A **connected simple system** consists of a clection I_0 of pointed spaces along with a collection I_m of homotopy classes of maps between these spaces such that

- (i) $\text{hom}(X, Y) := \{[f] \in [X : Y] : [f] \in I_m\}$ consists of exactly one element for each pair of spaces $(X, Y) \in I_0^2$, where $[X : Y]$ denotes the set of homotopy classes of maps between X and Y .
- (ii) If $X, Y, Z \in I_0$, $[f] \in \text{hom}(X, Y)$, and $g \in \text{hom}(Y, Z)$, then $[g \circ f] \in \text{hom}(X, Z)$.
- (iii) $\text{hom}(X, Y) = \{[\text{id}_X]\}$ for all $X \in I_0$.

Remark 3.1.6. This definitions comes from Salamon in [Sal85].

Remark 3.1.7. Let S be a isolated, compact, invariant subset of M with respect to the Morse flow φ . Then the collections

$$I_0 := \{N/L : (N, L) \text{ is index pair for } S\}$$

$$I_m := \{[f] : f \text{ is the homotopy equivalence of index pairs from the proof of theorem 3.1.4}\},$$

make up a connected simple system. We call this $\mathcal{IP}_S$.

Definition 3.1.8 (Morphisms of Connected Simple Systems). A **morphism** $\Phi : I \rightarrow Y$ between two connected simple systems $I = (I_0, I_m)$ and $J = (J_0, J_m)$ is a collection of homotopy classes of maps between spaces in I_0 and J_0 such that

- (i) For every $X \in I_0$ and $Y \in J_0$ the set $\Phi(X, Y) := \{[\varphi] \in [X : Y] : [\varphi] \in \Phi\}$ consists of exactly one element.
- (ii) If $X, \bar{X} \in I_0$ and $Y, \bar{Y} \in J_0$ and if $[\varphi] \in \Phi[X, Y], [f] \in \text{hom}[\bar{X}, X], g \in \text{hom}(Y, \bar{Y})$, then $[g \circ \varphi \circ f] \in \Phi(\bar{X}, \bar{Y})$.

Remark 3.1.9. By the second property of a morphism between connected simple systems I and J we can induce such a single map $\varphi : X \rightarrow Y$ where $X \in I_0$ and $Y \in J_0$.

Remark 3.1.10 (Critical points as isolated compact invariant subsets). Let $f : M \rightarrow \mathbb{R}$ be a Morse function on a smooth Riemannian manifold (M, g) . Let $q \in \text{Crit}_k(f)$. Around q we have Morse charts $\varphi : U \rightarrow T_q M$ (after identifying q_i and ∂q_i) where $\varphi(W(q \rightarrow) \cap U) \subseteq T_q^u M$ and $\varphi(W(\rightarrow q) \cap U) \subseteq T_q^s M$. With this we can define the balls:

$$D_\varepsilon^s = \{v \in T_q^s M \mid \|v\| \leq \varepsilon\},$$

$$D_\varepsilon^u = \{v \in T_q^u M \mid \|v\| \leq \varepsilon\}.$$

They give rise to the index pair $N_q := \varphi^{-1}(D^s \times D^u)$ and $L_q = \varphi^{-1}(D^s \times \partial D^u)$. This index pair is in fact regular, since $(N_q, L_q) \cong (D^s \times D^u, D^s \times \partial D^u) \cong (D_\varepsilon^u, \partial D_\varepsilon^u)$. Now we know the K-group of such a pair:

$$K^i(N_q, L_q) = K^i(D^k, \partial D^k) = \begin{cases} \mathbb{Z} & \text{if } i - k \in 2\mathbb{N}, \\ 0 & \text{else.} \end{cases} \quad (6)$$

Notice that for $k = 0$ we need to consider $\partial D^k = \emptyset$, to get an index pair. That's why we let ∂ denote the manifold boundary insted of the topological boundary. However, now we can identify for all k :

$$C^k(M, \mathfrak{A}, \tilde{K}^l(pt)) = \bigoplus_{\text{Crit}_k(f)} \mathbb{Z} \cong \bigoplus_{q \in \text{Crit}_k(f)} K^k(N_q, L_q; \mathbb{Z}). \quad (7)$$

Or using the language of connected simple systems, we have the identification

$$C^k(M, \mathfrak{A}, \tilde{K}^l(pt)) = \bigoplus_{q \in \text{Crit}_k(f)} K^k(\mathcal{IP}_q). \quad (8)$$

define
what
K(IP)
should
mean

Definition 3.1.11 (Canonical Isomorphism Class). A **canonical isomorphism class** of rings (or other algebraic structures) is a set of rings R_0 together with a set of ring homomorphisms R_m such that

- (i) For $R, L \in R_0$ the set $\text{hom}(X, Y) := \{f : X \rightarrow Y : f \in R_m\}$ consists of exactly one element.
- (ii) If $R, L, S \in R_0$ and $f \in \text{hom}(R, L)$ and $g \in \text{hom}(L, S)$, then $g \circ f \in \text{hom}(X, Z)$.
- (iii) $\text{hom}(X, X) = \{\text{id}_x\}$

A **morphism** between two canonical isommorphism classes of rings $\Psi : (R_0, R_m) \rightarrow (S_0, S_m)$ is a collection of homomorphisms such that

1. For any $R \in R_0$ and $S \in S_0$ the set $\Psi(R, S) := \{f : R \rightarrow S : f \in \Psi\}$ contains exactly one element.
2. If $R, \bar{R} \in R_0$ and $S, \bar{S} \in S_0$, $f \in \text{hom}(\bar{R}, R)$, $g \in \text{hom}(S, \bar{S})$ and $\psi \in \Psi(R, S)$, then $g \circ \psi \circ f \in \Psi(\bar{R}, \bar{S})$.

Remark 3.1.12. Again the second property lets us induce a morphism between two canonical isommorphism classes of rings $\Psi : (R_0, R_m) \rightarrow (S_0, S_m)$ by a single isomorphism

$$\psi : R \rightarrow S$$

for $R \in R_0$ and $S \in S_0$.

Remark 3.1.13. We call the category of simple connected systems **SCS** and the categorie of canonical isomorphism classes of rings **CICR**

Definition 3.1.14. The classical complex topological K-functors $\tilde{K}^i$ define a family of kontravariant functors

$$\text{SCS} \rightarrow \text{CICR}$$

that is fiven on the objects as follows: For (I_0, I_m) being a family of pointed topological spaces, we get a new family of rings as:

$$R_0 := \{\tilde{K}^i(X) : X \in I_0\},$$

and a family of ringhomomorphisms

$$R_m := \{\tilde{K}^i(f) : [f] \in I_m\}.$$

Obviously, the tuple (R_0, R_m) is an object in **CICR**. Now a morphism in $\Phi : (I_0, I_m) \rightarrow (J_0, J_m)$ in **SCS** gives rise to a collecting of ringisomorphisms

$$\Psi := \{\tilde{K}^i(\varphi) : [\varphi] \in \Phi\}.$$

By the homotopy property of reduced K-theory, this becomes a morphism in **CICR**.

Example 3.1.15. Any pointed topological space X defines a simple connected system $(\{X\}, [\text{id}_x])$. Similar, any ring R defines a canonical isomorphism class of rings $(\{R\}, \{\text{id}_R\})$.

Definition 3.1.16. Let (M, f, g) be a smooth manifold M together with a Morse Smale function f . Let $S \subseteq M$ be a compact invariant subset. Then the index pairs of S together with the canonical isomorphisms between them define a simple connected system which we will denote with $\mathcal{IP}_S$.

Remark 3.1.17. Let $\underline{\mathbb{Z}}$ be the canonical equivalence class of rings consisting only of the ring $\mathbb{Z}$, and let p be a critical point of index λ_p in M . Then remark 3.1.10 gives us the existence of a morphism for all $i \in \mathbb{N}$:

$$\tilde{K}^{\lambda_p+2i}(\mathcal{IP}_p) \rightarrow \underline{\mathbb{Z}}.$$

The next remark will help us to get a canonical isomorphism.

Remark 3.1.18. Furthermore, this isomorphism

$$C^k(M, \mathfrak{A}, \tilde{K}^l(pt)) = \bigoplus_{\text{Crit}_k(f)} \mathbb{Z} \cong \bigoplus_{q \in \text{Crit}_k(f)} K^k(N_q, L_q; \mathbb{Z}). \quad (9)$$

can be made canonically, using orientations: Let q be a critical point in index λ_q . Let $\mathfrak{b}$ be a ordered basis of $T_q^u M$. Let ψ be a refined Morse chart around q , such that the induced basis in $T_q M = T_q^u M \oplus T_q^s M$ projects onto a basis of the first factor $T_q^u M$ that is compatible with $\mathfrak{b}$. Using this Morse chart do define the generator of the regular index pair of q gives a unique generator.

Proof. Let ψ_1 and ψ_2 be two such Morse charts. Notice that they only differ by order of the directions and their signs, meaning that for all i there is a j such that $\psi_1^{-1}(e_i) = \pm \psi_2^{-1}(e_j)$. Hence, viewed as maps into the tangent space we have that $\psi_1(N_p, L_q) = \psi_2(N_p, L_q) \subseteq T_q M$. We define $B_\varepsilon^{\lambda_q}(0) \subseteq \mathbb{R}^{\lambda_q}$ to be the sphere. The generator of $\psi_1(N_p, L_q)$ comes as the pullback of the canonical Bott generator of $K^i(B_\varepsilon^{\lambda_q}(0), \partial B_\varepsilon^{\lambda_q}(0))$ via the basis map after the collapse map $c : (D^s \times D^u, D^s \times \partial D^u) \cong (D_\varepsilon^u, \partial D_\varepsilon^u)$. Let α be a generator of $\psi_1(N_p, L_q)$ induced from the basis $\mathfrak{b}_1$ that comes from ψ_1 . I.e.

$$\alpha = (B_{\mathfrak{b}_1} \circ c)^*(\beta)$$

By assumption, the map $B_{\mathbf{b}_1} \circ B_{\mathbf{b}}^{-1}$ has determinant one. Hence it is homotopic to the identity, and this homotopy can be reduced to a well defined homotopy on the quotient space $(D_\varepsilon^u / \partial D_\varepsilon^u)$. Hence

$$\alpha = (B_{\mathbf{b}} \circ c)^* (\beta)$$

And thereby, the generator only depends on $\mathbf{b}$ concluding the proof. $\square$

Example 3.1.19 (A Map of K-Groups). Let $f : M \rightarrow \mathbb{R}$ be a Morse Smale function on a smooth compact Riemannian manifold (M, g) . Let $q \in \text{Crit}_k(f)$ and $p \in \text{Crit}_{k-1}(f)$. Assume for a moment we already have an isolated regular compact index pair (N_2, N_0) of $S = W(p \rightarrow q) \cup \{p, q\}$. For a $c \in (f(p), f(q))$ we set $N_1 = N_0 \cup (N_2 \cap M^c)$, where M^c is the sublevel set. Clearly now (N_2, N_1) is an index pair for q and (N_1, N_0) is a pair for p . Assuming we have regular index pairs (N_q, L_q) and (N_p, L_p) for p, q we can now define a map:

$$\Delta^k(q \rightarrow p) : K^{k-1}(N_p, L_p) \rightarrow K^k(N_q, L_q)$$

as the composition:

$$K^{k-1}(N_p, L_p; \mathbb{Z}) \xrightarrow{\cong} K^{k-1}(N_1, N_0; \mathbb{Z}) \xrightarrow{\delta_{\text{triple}}} K^k(N_2, N_1; \mathbb{Z}) \xrightarrow{\cong} K^k(N_q, L_q; \mathbb{Z})$$

where δ_{triple} is the connecting homomorphism, and the other two maps are induced by the homotopic equivalence of two index pairs corresponding to the same isolated compact invariant subset. Putting all together we can define a morphism:

$$\Delta^k : \bigoplus_{p \in \text{Crit}_{k-1}(f)} K^{k-1}(N_p, L_p) \rightarrow \bigoplus_{q \in \text{Crit}_k(f)} K^k(N_q, L_q) \quad (10)$$

We can define this also with canonical isomorphism classes:

$$\underline{\Delta}^k : \bigoplus_{p \in \text{Crit}_{k-1}(f)} K^{k-1}(\mathcal{IP}_p) \rightarrow \bigoplus_{q \in \text{Crit}_k(f)} K^k(\mathcal{IP}_q)$$

Definition 3.1.20 (Regular Index Pairs and a Filtration on M). As always let $f : M \rightarrow \mathbb{R}$ be a Morse-Smale function on a compact smooth Riemannian manifold (M, g) of dimension m . Let $\varphi_t : M \rightarrow M$ be the flow given by $-\text{grad} f$. For $0 \leq j \leq k \leq m$ define:

$$W(k, j) = \bigcup_{j \leq \lambda_p \leq \lambda_q \leq k} W(q \rightarrow p).$$

There exists a filtration

$$\emptyset =: N_{-1} \subset N_0 \subset N_1 \subset \dots \subset N_{m-1} \subset N_m = M, \quad (11)$$

such that (N_k, N_{j-1}) is a regular index pair for $W(k, j)$ for all $0 \leq j \leq k \leq m$. If we define $\mathcal{IP}_i := \mathcal{IP}_{\text{Crit}_i(f)}$ we can induce a filtration of conected simple systems:

$$\emptyset \rightarrow \mathcal{IP}_0 \rightarrow \mathcal{IP}_1 \rightarrow \dots \rightarrow \mathcal{IP}_{m-1} \rightarrow \mathcal{IP}_m$$

define direkt sum of canonical isomorphism classes

Proof. The proof is done by Dietmar Salamon and can be checked in [Sal85] corollary 4.4. $\square$

Remark 3.1.21. Let M and M' be smooth manifolds with $f : M \rightarrow \mathbb{R}$ and $f' : M' \rightarrow \mathbb{R}$ Morse smale functions for the metrics g on M and g' on M' . Let $\mathfrak{o}$ be a choice of orientations on all unstable manifolds of M and $\mathfrak{o}'$ on M' . Let $(M \times M')$ be equipped with the Morse function $f + f'$ and the metric $g \oplus g'$ together with the orientations $\mathfrak{o} \oplus \mathfrak{o}'$. In other words, we work in a setting that we established in definition 2.0.28. Let $(N_i)_i$ be a filtration on M that satisfies that $(N_k, N_j - 1)$ is a regular index pair for $W(k, j)$ for all $0 \leq j \leq k \leq m = \dim(M)$. Let $(N'_j)_j$ be one on M' that also suffices those conditions. Then the Filtration $N_* \otimes N'_*$ defined by

$$(N_* \otimes N'_*)_k := \bigcup_{i+j \leq k} N_i \times N'_j$$

gives a filtration on $M \times M'$ satisfying the same condition.

Proof. We need to show that $((N_* \otimes N'_*)_k, (N_* \otimes N'_*)_{j-1})$ gives a regular index pair for $W(k, j)$ for all $0 \leq j \leq k \leq m$. Hence we need to go through the three properties. The first property states that the maximal invariant subset of the closure of $((N_* \otimes N'_*)_k \setminus (N_* \otimes N'_*)_{j-1})$ is $W(k, j)$ and exists in the interior of the set-difference. This is clear, since the space $W(k, j)$ splits nicely:

$$\begin{aligned} W(k, j) &= \bigcup_{(q, q') \in \text{Crit}_k(f+f')} W((q, q') \rightarrow) \cap \bigcup_{(p, p') \in \text{Crit}_j(f+f')} W(\rightarrow (p, p')) \\ &= \bigcup_{\substack{q \in \text{Crit}_i(f), \\ q' \in \text{Crit}_{i'}(f'), \\ i+i'=k}} W(q \rightarrow) \times W(q' \rightarrow) \cap \bigcup_{\substack{p \in \text{Crit}_l(f), \\ p' \in \text{Crit}_{l'}(f'), \\ l+l'=j}} W(\rightarrow p) \times W(\rightarrow p') \\ &= \bigcup_{\substack{i+i'=k \\ l+l'=j}} \left(\bigcup_{\substack{q \in \text{Crit}_i(f) \\ p \in \text{Crit}_l(f)}} W(q \rightarrow p) \times \bigcup_{\substack{q' \in \text{Crit}_{i'}(f') \\ p' \in \text{Crit}_{l'}(f')}} W(q' \rightarrow p') \right) \\ &= \bigcup_{\substack{i+i'=k \\ l+l'=j}} W(i, l) \times W(i', l'). \end{aligned}$$

With this it is easy to see the first property: Since the flow splits into the two factors, we have that the maximal invariant subset I also splits meaning

$$I((N_* \otimes N'_*)_k) = I\left(\bigcup_{i+j \leq k} N_i \times N'_j\right) = \bigcup_{i+j \leq k} I(N_i \times N'_j) = \bigcup_{i+j \leq k} (I(N_i) \times I(N'_j)).$$

Furthermore the same argument works on the difference $((N_* \otimes N'_*)_k, (N_* \otimes N'_*)_{j-1})$ letting us deduce the first property for the tensorproduct filtration from said property for all factors. The same splitting idea lets us deduce the positive invariance of $(N_* \otimes N'_*)_k$ in $(N_* \otimes N'_*)_{j-1}$ and shows the exit set property of $(N_* \otimes N'_*)_{j-1}$. Now finally we need to proof the regularity, meaning that the inclusion $(N_* \otimes N'_*)_{j-1} \rightarrow (N_* \otimes N'_*)_k$

is a cofibration. For this we notice that the cofibration property is stable under finite products, since we can just build the product homotopy of each factor. Now a standard result is that cofibrations are stable under pushouts and hence especially under unions concluding the proof. Compare [May99] Chapter 6 on the claim about cofibrations. $\square$

Remark 3.1.22. Now $\mathcal{IP}_k$ has a pointed spaces as a representative, that if of the form $\bigvee_{p \in \text{Crit}_k} (N_p / L_p)$. Hence, $\tilde{K}^k(\mathcal{IP}_k)$ has a representative that omits an isomorphism to $\bigoplus_{p \in \text{Crit}_k(f)} \tilde{K}^k(N_p / L_p)$. Hence as canonical isomorphism classes of rings we get the isomorphism:

$$\tilde{K}^k(\mathcal{IP}_k) \rightarrow \bigoplus_{q \in \text{Crit}_k(f)} \tilde{K}^k(\mathcal{IP}_p)$$

and with this we can define the morphism $\underline{\Delta}^k : \tilde{K}^{k-1}(\mathcal{IP}_{k-1}) \rightarrow \tilde{K}^k(\mathcal{IP}_k)$. I think with this, i can give the definition of the following spectral sequence in the language of canonical isomorphism classes!

4 The Spectral Sequence using Exact Couples

Definition 4.0.1. Let M be a smooth manifold with Morse data and

$$\emptyset =: N_{-1} \subset N_0 \subset N_1 \subset \dots \subset N_{m-1} \subset N_m = M,$$

be a filtrations constructed in 3.1.20. Then we define the Groups

$$\begin{aligned} D^{p,q} &:= K^{p+q}(N_p) & \text{and} & \quad D := \bigoplus_{p,q} D^{p,q} \\ E^{p,q} &:= K^{p+q}(N_p, N_{p-1}) & \text{and} & \quad E := \bigoplus_{p,q} E^{p,q} \end{aligned}$$

Now we have the exact sequence from the pair (N_p, N_{p-1}) :

$$\dots \longrightarrow K^{p+q-1}(N_{p-1}) \xrightarrow{\delta_{p,q}^*} K^{p+q}(N_p, N_{p-1}) \xrightarrow{j_{p,q}^*} K^{p+q}(N_p) \xrightarrow{i_{p,q}^*} K^{p+q}(N_{p-1}) \dots$$

With this we define the maps

$$\begin{array}{lcl} \alpha & := \bigoplus_{p,q} i_{p,q}^* : D \rightarrow D & (-1, 1) \\ \beta & := \bigoplus_{p,q} \delta_{p,q}^* : D \rightarrow E & (1, 0) \\ \gamma & := \bigoplus_{p,q} j_{p,q}^* : E \rightarrow D & (0, 0) \end{array} \quad \left| \begin{array}{l} \\ \\ \text{bidegree} \end{array} \right.$$

Obviously by definition the compositions in the triangle are exact:

$$\begin{array}{ccc} D & \xrightarrow{\alpha} & D \\ & \nwarrow \gamma & \nearrow \beta \\ & E & \end{array}$$

Finally we define $d := \beta \circ \gamma$.

Definition 4.0.2 (Exact Couples). Two groups D, E and maps

$$\begin{array}{lcl} \alpha: D \rightarrow D & D & \xrightarrow{\alpha} D \\ \beta: D \rightarrow E & \swarrow \gamma & \searrow \beta \\ \gamma: E \rightarrow D & E & \end{array}$$

such that the compositions in the triangle are exact are called an **exact couple**. we define $d := \beta \circ \gamma$. In this case it is obvious that $d^2 = 0$.

Definition 4.0.3 (Categorie of Exact Couples). We define the categorie **ExCoup** as follows: The objects are exact couples, i.e. quintuples $(D, E, \alpha, \beta, \gamma)$ where D, E are groups (or objects in an abelian categorie), such that the triangle is exact:

$$\begin{array}{ccc} D & \xrightarrow{\alpha} & D \\ & \swarrow \gamma & \searrow \beta \\ & E & \end{array}$$

.Given another exact couple $(D', E', \alpha', \beta', \gamma')$ we define a morphism from one to another to be a tuple of morphisms $(f: D \rightarrow D', g: E \rightarrow E')$ such that the following diagram commutes:

$$\begin{array}{ccccccc} D & \xrightarrow{\alpha} & D & \xrightarrow{\beta} & E & \xrightarrow{\gamma} & D \\ \downarrow f & & \downarrow f & & \downarrow g & & \downarrow f \\ D' & \xrightarrow{\alpha'} & D' & \xrightarrow{\beta'} & E' & \xrightarrow{\gamma'} & D' \end{array}$$

Corollary 4.0.4 (An Endofunktor in the Categorie of Exact Couples). We can derive a new exact couple with the following definitions:

$$\begin{array}{lll} E' & := \ker d / \text{imd} & \alpha' : D' \rightarrow D' \quad a \mapsto \alpha(a) \\ D' & := \alpha(A) & \beta' : D' \rightarrow E' \quad a = \alpha(\tilde{a}) \mapsto [\beta(\tilde{a})] \\ & & \gamma' : E' \rightarrow D' \quad [b] \mapsto \gamma(b) \end{array} \quad \begin{array}{ccc} D' & \xrightarrow{\alpha'} & D' \\ & \swarrow \gamma' & \searrow \beta' \\ & E' & \end{array}$$

We again define $d' := \beta' \circ \gamma'$. The deriving of a new exact couple as defined above defines an endofunktor which we denote by Der .

Proof. Let $(D, E, \alpha, \beta, \gamma)$ be the exact couple we started with and call the derived couple $(D', E', \alpha', \beta', \gamma')$. First we show the well definition of β . Let $a \in D' = \alpha(a_1) = \alpha(a_2)$. Then $a_1 - a_2 \in \ker \alpha = \text{imd}$. Hence $\beta(a_1) - \beta(a_2) = \beta(a_1 - a_2) \in \text{imd}(\beta \circ \gamma) = \text{imd}$. Furthermore, since $\beta(a_1) \in \text{imd} = \ker \gamma \subseteq \ker d$ we have that the map β' is well defined. For the well definition of γ' we assume that $[b_1] = [b_2] \in E'$. But then the difference of b_1 and b_2 is zero in E' meaning in the image of d , which is again contained in the image of β and thereby in the kernel of α . This means that under α the image of b_1 and b_2 agree. Now to show that the derived couple is also exact one continues the diagram yoga. We do exemplify the equality of the kernel of β' with the image of α' : Assume

that $\beta'(x) = 0$. That means that $\beta(\tilde{x}) \in \text{imd}$ with $\alpha(\tilde{x}) = x$. We now want to show that $x \in \alpha^2 D$, and do this by showing that $\tilde{x} = \alpha(z)$ for some z . Now since $\beta(\tilde{x}) \in \text{imd}$ we have that $\beta(\tilde{x}) = d(y) = \beta \circ \gamma(x)$ for some y . In other words, $\tilde{x} - \gamma(y) \in \ker \beta = \text{im} \alpha$. Hence, $\tilde{x} - \gamma(y) = \alpha(z)$ for some z . But now applying α to both side and using that $\alpha \circ \gamma = 0$ we can deduce that $\alpha^2(z) = \alpha(\tilde{x})$. This shows that the kernal of β' is contained in the image of α' . The other direction of the inclusion is easier: Let $a = \alpha(\tilde{a}) \in D'$. Then $\beta' \circ \alpha'(z) = [\beta(z)] = \beta \circ \alpha(\tilde{a}) = 0$. Similary one checks for exactnes in the other corners. Now to show that we have an endofunktor we need to define what to do on morphisms. So assume that we have the two exact couples $(D, E, \alpha, \beta, \gamma)$ and $(\tilde{D}, \tilde{E}, \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma})$ with the morphisms $(f : D \rightarrow \tilde{D}, g : E \rightarrow \tilde{E})$. We now need to define a morphism of exact couples $((f' : D' \rightarrow \tilde{D}', g' : E' \rightarrow \tilde{E}'))$ between the derived couples. For this we define:

$$f' : D' \rightarrow \tilde{D}'; x \mapsto f(x) \quad \text{and} \quad g' : E' \rightarrow \tilde{E}'; [x] \mapsto [g(x)]$$

and check for well definition as maps and that the pair is a morphism of exact couples. The well definition of f' follows directly from $f \circ \alpha = \tilde{\alpha} \circ f$. For the well definition of g' asume that $[x] = [y] \in E'$, meaning that $x - y \in \text{imd} = \text{im} \beta \circ \gamma$ hence for some z we have that $x - y = \beta \circ \gamma(z)$ and hence $g(x - z) = g \circ \beta \circ \gamma(z) = \tilde{\beta} \circ f \circ \gamma(z) = \tilde{\beta} \circ \tilde{\gamma} \circ g(z) \in \text{imd}$. This proofes the well definition. $\square$

Definition 4.0.5 (Spectral Sequence From an Exact Couple). We can now iterate this procedure to get a list of spaces with maps $(E, d), (E', d'), (E'', d'') \dots$ or often written as $(E_1, d_1), (E_2, d_2), \dots$ which is called a **spectral sequence**. Notice how E_{r+1} is given by (E_r, d_r) whilst d_{r+1} needs further information. Whilst the spectral sequence is defined inductively, the needed information is not captured in $(E_r, d_r)_r$ but rather in the exact couple. Or said differently, we get a sequence of exact couples $(E_r, D_r, \alpha_r, \beta_r, \gamma_r)$ from which we can get the spectral sequence. We can reformulate the maps as follows:

$$D_r = \alpha^{r-1}(D), \quad E_r = \ker d_{r-1} / \text{imd}_{r-1}, \quad \alpha_r = \alpha|_{\alpha^{r-1}(D)}, \quad \beta_r = [\beta \circ \alpha^{1-r}], \quad \gamma_r = \gamma, \quad d_r = \beta_r \circ \gamma_r$$

α^{1-r} means taking the preimage by alpha $1-r$ times. The well definition of this formulation was proven earlier. The bidegree of the maps is easy to deduct from the formulations above:

$$\begin{array}{ll} \alpha_r & \text{has bidegree} \quad (-1, 1) \\ \beta_r & \text{has bidegree} \quad (-r + 2, r - 1) \\ \gamma_r & \text{has bidegree} \quad (0, 0) \\ d_r & \text{has bidegree} \quad (r, +r + 1) \end{array}$$

Corollary 4.0.6. We can define the categorie of spectral sequences **SpS** where the objects consist of families of pairs $(E_r, d_r)_{r \in \mathbb{Z}}$ where E_r are groups (or objects in an abelian categorie) and d_r are endomorphisms in E_r , such that $E_{r+1} \cong \frac{\ker d_r}{\text{imd}_r}$. Morphisms between two spectral sequences $(E_r, d_r)_{r \in \mathbb{N}} \rightarrow (\tilde{E}_r, \tilde{d}_r)_{r \in \mathbb{N}}$ are families of morphisms $(f_r : E_r \rightarrow \tilde{E}_r)$ such that for all r we have that $f_r \circ d_r = \tilde{d}_r \circ f_r$.

Corollary 4.0.7. We have a functor from **ExCoup** to **SpS** given as follows: Let $(E, D, \alpha, \beta, \gamma)$ be an exact couple. Let $(E_r, d_r, \alpha_r, \beta_r, \gamma_r) = \text{Der}^r((E, D, \alpha, \beta, \gamma))$ where

Der^r denotes the r -th derivation of the exact couple. Then $(E_r, d_r := \beta_r \circ \gamma_r)_r$ is an exact sequence as described above. Assume that we have the two exact couples $(D, E, \alpha, \beta, \gamma)$ and $(\tilde{D}, \tilde{E}, \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma})$ with the morphisms $(f : D \rightarrow \tilde{D}, g : E \rightarrow \tilde{E})$. Then the family $(Der^r(f))_r$ is a morphism of spectral sequences.

Corollary 4.0.8 (Explicit Pages). For a spectral sequence arising from an exact couple defined as above we have a natural isomorphism

$$E_{r+1} = \frac{\gamma^{-1}(\alpha^r D)}{\beta(\ker \alpha^r)}$$

Proof. We proceed by induction starting with $n = 1$. Here we have by exactness:

$$\ker(\beta \circ \gamma) = \gamma^{-1}(\ker(\beta)) = \gamma^{-1}(\operatorname{im}(\alpha)), \quad \text{and} \quad \operatorname{im}(\beta \circ \gamma) = \beta(\ker(\alpha)).$$

Hence:

$$E_2 = \frac{\ker(d_1)}{\operatorname{im}(d_1)} = \frac{\ker(\beta \circ \gamma)}{\operatorname{im}(\beta \circ \gamma)} = \frac{\gamma^{-1}(\alpha D)}{\beta(\ker(\alpha))}.$$

Now Assume that

$$E_r = \frac{\gamma^{-1}(\alpha^{r-1} D)}{\beta(\ker \alpha^{r-1})}$$

Then we have

$$\ker d_r = \ker \beta_r \gamma_r = \gamma_r^{-1}(\operatorname{im}(\alpha_r)) = \gamma_r^{-1}(\alpha^r(D)).$$

Now $\gamma_r : E_r \rightarrow D_r$ maps $[x] \mapsto \gamma(x)$, hence by induction assumption

$$\ker d_r = \frac{\gamma^{-1}(\alpha^r D)}{\beta(\ker \alpha^r)}.$$

Notice that $\alpha^{r-1}(\ker(\alpha_r))$ contains those elements in D , that are killed by the α^n , in other words this equals $\ker \alpha^r$. Now $\beta_r = [\beta \circ \alpha^{r-1}]$ and hence using the induction assumption again we can conclude:

$$\operatorname{im}(\beta_r \gamma_r) = [\beta(\alpha^{r-1} \ker(\alpha_r))] = \frac{\beta(\ker \alpha^r D)}{\beta(\ker \alpha^{r-1})}.$$

Now using the isomorphism theorem we get the natural isomorphism from the statement. $\square$

Definition 4.0.9. In our case all pages of the spectral sequence inherit a double grading and the boundary maps to. Hence we will denote

$$E_r = \bigoplus_{p,q} E_r^{p,q} \quad \text{and} \quad d_r = \oplus_{p,q} d_r^{p,q}$$

Corollary 4.0.10. The grading carries through all pages and we can reformulate

$$E_{r+1}^{p,q} = \frac{\gamma^{-1} \alpha^r D}{\beta(\ker \alpha^r)} \cap E_{r+1}^{p,q} = \frac{\gamma^{-1}((\alpha^r D) \cap D^{p,q})}{\beta((\ker \alpha^r) \cap D^{p-1,q})} = \frac{\gamma^{-1} \alpha^r D^{p+r, q-r}}{\beta(\ker \alpha^r|_{D^{p-1,q}})}$$

Corollary 4.0.11 (The Exact Sequence Stabilizes). For all $r \geq m + 1$ it holds that

$$E_r = E_{r+1} = \dots$$

For this great enough r we define $E_\infty := E_r$.

Proof. Notice how our original $E^{p,q}$ is supportet in a strip regarding p which is fancy language for saying that $E^{p,q} = 0$ for $p > 0$ and for $p \geq m$. These conditinos cary through the whole spectral sequence, as succeeding pages are quotients of preceeding ones respecting the grading. Now the map d_r has bidegree $(r, -r + 1)$, and hence $d_r = 0$ since the domain or codomain is zero, if the p grading is greater then the p -length of the supporting strip meaning $p > m$ which gives us the claimes $r \geq m + 1$. $\square$

Now we refomulate the corollary 4.0.8 for the infinity page. We have the constant sequence of groups $E_r = E_{r+1} = \dots$ and by the corollary we have

$$\frac{\gamma^{-1}(\alpha^r D)}{\beta(\ker \alpha^r)} = \frac{\gamma^{-1}(\alpha^{r+1} D)}{\beta(\ker \alpha^{r+1})} = \frac{\gamma^{-1}(\alpha^{r+2} D)}{\beta(\ker \alpha^{r+2})} = \dots$$

Hence in our later inspections of E_∞ we can assume r to be arbitrary big.

Lemma 4.0.12 (The last page). *In our spectral sequence, we have that*

$$E_\infty^{p,q} = \frac{\ker[K^{p+q}(M) \rightarrow K^{p+q}(N_{p-1})]}{\ker[K^{p+q}(M) \rightarrow K^{p+q}(N_{p1})]}$$

Proof. For each fixed (p, q) there is an r big enough such that $\alpha^r D^{p,q} = 0$, since α^r has degree $-r$ in p , hence for $r > p$, α^r maps $D^{p,q}$ to something negative in p and hence to the trivial group. Thus $D = \cup_{r \geq 0} \ker \alpha^r$. Now the denominator of $E_r^{p,q}$ equals $\beta(\ker(\alpha^r|_{D^{p+q-1, q-r}}))$ and hence for r big enough this equals the image of $\beta|_{D^{p-1, q}}$ which in turn is the kernal of $\gamma|_{E^{p, q}}$. For the numerator, we start by looking at $(\alpha^r D)^{p, q} = \alpha^r D^{p+r, q-r}$. Now $D^{p+r, q-r} = K^{p+q}(N_{p+n})$. Hence for n big enough (such that $N_{p+n} = M$) we can calculate $\alpha^n(K^{p+q}(N_{p+n})) = \text{im}(K^{p+q}(M) \rightarrow K^{p+q}(N_p))$. This results in the description

$$E_\infty^{p,q} = \frac{\gamma^{-1}(\text{im}(K^{p+q}(M) \rightarrow K^{p+q}(N_p)))}{\ker \gamma|_{E^{p, q}}}$$

Next we define the groups

$$L^{p,q} := \text{im}(K^{p+q}(M) \rightarrow K^{p+q}(N_p)) \cap \text{im} \gamma = \text{im}(K^{p+q}(M) \rightarrow K^{p+q}(N_p)) \cap \ker \alpha$$

And since we have for any f and any subset S of the codomain of f the equality $f^{-1}(S) = f^{-1}(S \cap \text{im} f)$ holds, we have in our setting $E_\infty^{p,q} = \frac{L^{p,q}}{\ker(\gamma|_{E^{p, q}})}$. Now the kernal of the funktion

$$\tilde{\gamma} : \gamma^{-1}(L^{p,q}) \rightarrow L^{p,q}; x \mapsto \gamma(x)$$

is das wahr mit gamma eingeschränkt auf $E^{p,q}$?

equals $\ker \gamma|_{E^{p,q}}$: Since $\gamma^{-1}(L^{p,q}) \subseteq E^{p,q}$ the kernel of $\tilde{\gamma}$ is clearly contained in $\ker \gamma|_{E^{p,q}}$. If on the other hand $\gamma(x) = 0$ for $x \in E^{p,q}$ then $x \in \gamma^{-1}(0)$ and since zero is contained in $L^{p,q}$ we conclude that the two kernels are equal and hence we can apply the isomorphism theorem to get the isomorphism that is induced from γ :

$$E_{\infty}^{p,q} = \frac{L^{p,q}}{\ker(\gamma|_{E^{p,q}})} \rightarrow L^{p,q}.$$

Now finally we define the mapping

$$\ker(K^{p+q}(M) \rightarrow K^{p+q}(N_{p-1})) \rightarrow L^{p,q} = \text{im}(K^{p+q}(M) \rightarrow K^{p+q}(N_p)) \cap \ker \alpha$$

That maps an element $x \in K^{p,q}(M)$ to the pullback of the inclusion $N_p \rightarrow M$. This map is well defined, since applying α to the image of an element from the domain is the same as the pullback of x via the inclusion $N_{p-1} \rightarrow M$ and equals zero by definition of the domain. Hence the original map maps into the kernel of alpha which proofs the well definition. This line of reasoning also shows the surjectivity since every element in $K^{p,q}(N_p)$ that comes from the whole M and maps to zero if restricted to N_{p-1} can be expressed as the pullback from the whole M restricted to those, that vanish over N_{p-1} . Its easy to see that the kernel of this surjection equals the kernel of the pullback

$$K^{p,q}(M) \rightarrow K^{p,q}(N_p),$$

because ever element that pulls back to zero over N_p also pulls back to zero over N_{p-1} , and the other containemend is trivially true. To the surjection we can again apply the isomorphism theorem to get the isomorphism

$$\frac{\ker(K^{p+q}(M) \rightarrow K^{p+q}(N_{p-1}))}{\ker(K^{p,q}(M) \rightarrow K^{p,q}(N_p))} \cong L^{p,q},$$

that is induced by the pullback of the inclusion. Together with the γ induced isomorphism from $E_{\infty}^{p,q} \rightarrow L^{p,q}$ we conclude the proof. $\square$

Definition 4.0.13 (Associated Graded Group). Let G be a group together with a finite filtration of subgroups $\{F^i\}_{i=-1, \dots, -m}$ for $i = -1, \dots, m$ such that

$$G = F^{-1} \subseteq F^0 \subseteq F^1 \subseteq \dots \subseteq F^{m-1} \subseteq F^m = 0$$

Then we define the **associated graded group of G with respect to the Filtration $\{F_i\}$** to be

$$\text{gr}(G) := \bigoplus_{i=0}^m F^{i-1} / F^i.$$

From $\text{gr}(G)$ we can rebuild G by rebuilding $\{F^i\}$ up to extension problems in the following way:

$$\begin{array}{ll}
F^{m-1} & = F^{m-1}/F^m = \text{gr}(G)_m \\
F^{m-2} & \text{ is an extension } 0 \longrightarrow F^{m-1} \longrightarrow F^{m-2} \longrightarrow F^{m-2}/F^{m-1} = \text{gr}(G)_{m-1} \longrightarrow 0 \\
F^{m-1} & \text{ is an extension } 0 \longrightarrow F^{m-2} \longrightarrow F^{m-3} \longrightarrow F^{m-3}/F^{m-2} = \text{gr}(G)_{m-2} \longrightarrow 0 \\
\vdots & \dots \\
F^{-1} & \text{ is an extension } 0 \longrightarrow F^0 \longrightarrow F^{-1} \longrightarrow F^{-1}/F^0 = \text{gr}(G)_0 \longrightarrow 0
\end{array}$$

Corollary 4.0.14. Define the groups

$$F^l K^n(M) := \ker(K^n(M) \rightarrow K^n(N_l)) .$$

Those give an decending filtration of the $n - th$ K-group of M :

$$K^n(M) = F^{-1} K^n(M) \supseteq F^0 K^n(M) \supseteq \dots \supseteq F^{m-1} K^n(M) \supseteq F^m K^n(M) = 0 .$$

Then by the corollary above we can conculde that

$$E_r^{p,q} = \frac{F^{p-1} K^{p+q}(M)}{F^p K^{p+q}(M)} .$$

Hence, the diagonals on the infinity page are the associated graded groups to $K^{p+q}(M)$ with respect to the filtration above.

Corollary 4.0.15 (The first page). In our setting we get the first page to be defined as the quotient

$$E^{p,q} := K^{p+q}(N_p, N_{p-1})$$

Now since (N_p, N_{p-1}) is a regular index pair for $\text{Crit}_p(f)$ we have a flow induced homotopy equivalentce:

$$N_p/N_{p-1} \simeq \left(\bigcup_{w \in \text{Crit}_p(f)} D_\varepsilon^u(w) \right) / \left(\bigcup_{w \in \text{Crit}_p(f)} \partial D_\varepsilon^u(w) \right) \cong \bigvee_{w \in \text{Crit}_p(f)} S^p$$

Here, we use the definitions from above

$$D_\varepsilon^u(w) := \varphi_w^{-1}(\{v \in T_w^u M \mid \|v\| \leq \varepsilon\}),$$

and the last isomorphism is induced by orientations in the unstable manifolds. We can continue our inspection:

$$E_1^{p,q} \cong \tilde{K}^{p+q}(N_p/N_{p-1}) \tag{12}$$

$$\cong \bigoplus_{w \in \text{Crit}_p(f)} \tilde{K}^q(pt) = C^k(M, f, g, \mathfrak{o}; \tilde{K}^q(pt)) \tag{13}$$

$$\tag{14}$$

Notice that this is trivial for all odd q and naturally isomorphic for all even q . In sum we have the chain of horizontal isomorphisms:

$$\begin{array}{ccccccc}
 E_1^{p,q} & \xrightarrow{\alpha} & \tilde{K}^{p+q}(N_p/N_{p-1}) & \xrightarrow{\beta} & \oplus_{w \in \text{Crit}_p(f)} \tilde{K}^q(pt) & \longleftarrow & C^p(M, \mathfrak{A}, \tilde{K}^q(pt)) \\
 \downarrow d_1^{p,q} & & \downarrow & & \downarrow & & \downarrow \\
 E_1^{p+1,q} & \xrightarrow{\alpha'} & \tilde{K}^{p+1+q}(N_{p+1}/N_p) & \xrightarrow{\beta'} & \oplus_{w \in \text{Crit}_{p+1}(f)} \tilde{K}^q(pt) & \longleftarrow & C^{p+1}(M, \mathfrak{A}, \tilde{K}^q(pt))
 \end{array}$$

Diagram 15

By naturality of the triple boundary operator, that the second vertical map is the boundary operator. Now on the next rug we can induce two maps. From the left to make the diagram commute and from the right. Do they agree? Or asked differently, is the map induced from the d_1 map the boundary operator of morse homology? We already answered this in theorem 6.0.5. This theorem lets us define a new spectral sequence as follows:

Definition 4.0.16. Let $(D, E, \alpha, \beta, \gamma)$ be the exact couple from definition 4.0.1. Furthermore, let $\Psi := \oplus_{p,q} \Psi^{p,q}$ be the isomorphism from $E = \oplus_{p,q} E^{p,q} \rightarrow \oplus_{p,q} C^p(M, \mathfrak{A}, \tilde{K}^q(pt))$ that is defined in equation 12. Furthermore, let $\beta : K^{2s}(pt) \rightarrow \mathbb{Z}$ be the isomorphisms induced from the bott generator. Then we define

$$\Lambda : E \rightarrow \tilde{E} := \bigoplus_{p,q} C(M, \mathfrak{A}, \mathbb{Z}) \otimes_{\mathbb{Z}} \begin{cases} \mathbb{Z} & \text{if } q \text{ is even,} \\ 0 & \text{if } q \text{ is odd.} \end{cases}$$

Then define $\tilde{D} := D$ and the new exact couple $(\tilde{D}, \tilde{E}, \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma})$ as follows:

$$\begin{array}{lll}
 \tilde{\alpha} & := \alpha & : \tilde{D} \rightarrow \tilde{E} \\
 \tilde{\beta} & := \Lambda \circ \beta & : \tilde{D} \rightarrow \tilde{E} \\
 \tilde{\gamma} & := \gamma \circ \Lambda^{-1} & : \tilde{E} \rightarrow \tilde{D}
 \end{array}$$

We get a morphism of exact couples from $(D, E, \alpha, \beta, \gamma)$ to $(\tilde{D}, \tilde{E}, \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma})$ that is given as (Λ, id_D) . This new exact couple gives rise to a spectral sequence $(\tilde{E}_r, \tilde{d}_r)_r$, where the first page is the Morse-chain group in all even q . Hence, the second page is the Morse homology.

Corollary 4.0.17 (Periodicity in q). The exact couple $(\tilde{D}, \tilde{E}, \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma})$ has periodicity in q meaning that we have the isomorphism of exact couples given by the identity functions

$$f^{p,q} : \tilde{E}^{p,q} \rightarrow \tilde{E}^{p,q+2}$$

Here the notation might seem a bit strange. However, the summands $\tilde{E}^{p,q}$ and $\tilde{E}^{p,q+2}$ are actually the same. They only differ by there incslion map into $\tilde{E}$. Now the pair $(f : \oplus_{p,q} \tilde{E} \rightarrow \tilde{E}, g := \text{id } \tilde{D} \rightarrow \tilde{D})$ defines an isomorphism of exact couples. The compatibility with morphisms is clear. Now since building up the spectral sequence is functorial we

know that for all p, q we have canonical (in the sence that they traslate $\widetilde{d}_r^{p,q}$ to $\widetilde{d}_r^{p,q+2}$) isomorphisms

$$\widetilde{E}_r^{p,q} \cong \widetilde{E}_r^{p,q+2}.$$

Furthermore, $\widetilde{E}_r^{p,q} = 0$ for all odd q .

Definition 4.0.18. For $r > 0$ we define $EM_r^p := \widetilde{E}_r^{p,0}$ and

$$dM_r : EM_r^p \rightarrow EM_r^{p+r}; x \mapsto \begin{cases} 0 & \text{if } r \text{ is even} \\ \widetilde{d}_r^{p,0}(x) \in \widetilde{E}_r^{p,-r+1} = EM_r^p & \text{else.} \end{cases}$$

The last equality comes from the corollary above. Furthermore, we have a morphism of spectral sequences $(\widetilde{E}_r, \widetilde{d}_r) \rightarrow (EM_r, dM_r)$ by the maps

$$f_r^{p,q} : \widetilde{E}_r^{p,q} \rightarrow EM_r^p; x \mapsto \begin{cases} 0 & \text{if } q \text{ is odd} \\ x \in EM_r = \widetilde{E}_r^{p,0} \cong \widetilde{E}_r^{p,q} & \text{if } q \text{ is even.} \end{cases}$$

Theorem 4.0.19 (Summary of the Spectral Sequence). *For each Morse datum $(M, f, g, \mathfrak{A})$ together with a filtration $N_{-1} \subseteq N_0 \subseteq \dots \subseteq N_{m-1} \subseteq N_m = M$ from (11) there exists a Spectralsequence (EM_r^p, dM_r^p) where each page is graded in p such that*

- *The first page is the morse complex, hence the second page is the Morse homology*
- *The spectral sequence becomes stationary and the infinity page is the associated graded group of the K -groups meaning:*

$$\bigoplus_{p \in 2\mathbb{N}} E_\infty^p \text{ is a associated graded group to } K^0(M)$$

$$\bigoplus_{p \in 2\mathbb{N}+1} E_\infty^p \text{ is a associated graded group to } K^1(M).$$

Remark 4.0.20 (Recover from Associated Graded Groups). Let G be a group and

5 Higher Differentials

6 The prove that N_p is a tubular neighbourhood

So to proof this, we first formulate two lemmas. First we restate/ refine the λ -lemma:

parameter
depen-
dence
einfügen

Lemma 6.0.1 (The Technical λ -Lemma). *Let M be a smooth manifold and let p be a hyperbolic fixpoint of a diffeomorphism φ on M with $\dim W(p \rightarrow) = r$ for $0 < r < m = \dim M$. Let N be an injective immersed submanifold of M that is invariant under φ and has a point of transverse intersection q with $W(\rightarrow p)$. Then for any $\varepsilon > 0$, any r -cell neighbourhood D of q in N that also intersects $W(\rightarrow p)$ transversal and any r -cell neighbourhood B of p in $W(p \rightarrow)$ there is a smaller cell $\tilde{D} \subset D$, an $n \in \mathbb{N}$ such that the following is true:*

We can assume that B and $\varphi^n(\tilde{D})$ lives in a chart centered at p with a product structure $U \cong W(\rightarrow p) \cap B_\mu(0) \times W(p \rightarrow) \cap B_\mu(0)$. Here, the invars of the projection into $B \subseteq W(\rightarrow p)$:

$$h : B \rightarrow \varphi^n(\tilde{D})$$

satisfies:

1. $\sup_{x \in B} \|x - h(x)\| < \varepsilon$ and
2. For the inclusions $i_1 : B \rightarrow M$ and $i_2 : \varphi^n(\tilde{D}) \rightarrow M$ we have that

$$\sup_{x \in B} \|d(i_1)_x - d(i_2 \circ h)_x\| < \varepsilon.$$

Proof. Let q be a point of transversal intersection of $W(\rightarrow p)$ with N . Since both N and $W(\rightarrow p)$ are invariant under φ , it follows that $\varphi^n(q) \in N \cap W(\rightarrow p)$ for all $n \in \mathbb{N}$. Moreover, the intersection remains transversal: the direct sum decomposition $T_q M = T_q W(\rightarrow p) \oplus T_q N$ is preserved under $d\varphi_q$, which implies transversality at $\varphi^n(q)$ as well.

Therefore, we may assume that q lies in a coordinate chart centered at the hyperbolic fixed point p , and identify a neighborhood of p with an open subset of $\mathbb{R}^m$, where $m = \dim M$. By hyperbolicity of p , the tangent space at p splits as

$$T_p M = T_p^s M \oplus T_p^u M,$$

and this splitting is preserved by $d\varphi_p$. As discussed earlier, there exists a chart in which $d\varphi_p$ is represented by a real diagonal matrix (in suitable coordinates). Hence, we may choose an orthonormal basis of coordinates $(x_s, x_u) \in \mathbb{R}^s \times \mathbb{R}^u$, adapted to the stable and unstable subspaces, such that

$$d\varphi_0 = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix},$$

where the eigenvalues of A have modulus less than 1, and those of B greater than 1.

reference
einfügen

in fact we
can choose
a Morse
chart that
satisfies
this!

Since we are working in $\mathbb{R}^m$, we identify the manifold and its tangent space locally around 0. We then expand φ in a Taylor series at the origin:

$$\varphi(x_s, x_u) = \varphi(0) + d\varphi_0(x_s, x_u) + f(x_s, x_u),$$

where $f(0) = 0$ and $df(0) = 0$. Writing this in coordinates gives:

$$\varphi(x_s, x_u) = (Ax_s + f_s(x_s, x_u), Bx_u + f_u(x_s, x_u)). \quad (16)$$

Furthermore, since $f = \varphi - d\varphi|_0$ we know that:

$$\frac{\partial f_s}{\partial x_u}|_{(0, x_u)} = \frac{\partial \varphi_s}{\partial x_u}|_{(0, x_u)} = 0 = \frac{\partial f_u}{\partial x_s}|_{(x_s, 0)} = \frac{\partial \varphi_u}{\partial x_s}|_{(x_s, 0)}.$$

Similar to Corollary ?? we can define an inner product such that for the operator norm we have

$$\|A\| \leq \alpha < 1 \quad \text{and} \quad \|B^{-1}\| \leq \frac{1}{\beta} < 1.$$

Since $1 = \|\text{id}\| = \|BB^{-1}\| \leq \|B\| \cdot \|B^{-1}\|$ we know that $\|B\| \geq \beta$. For a neighbourhood $U_1 \subseteq U$ of the origin we define

$$k := \max_{i,j=s,u} \left\{ \sup_{U_1} \left\| \frac{\partial f_i}{\partial x_j} \right\| \right\},$$

and choose U_1 small enough such that k satisfies:

$$\begin{aligned} 0 &< k < 1 \\ \beta_1 &:= \beta - k > 1 \\ \alpha_1 &:= \alpha + k < 1 \\ k &< \frac{1}{4}(b_1 - 1)^2. \end{aligned}$$

This can be achieved since $df|_0 = 0$. Let B be a cell neighbourhood of the origin in $W(p \rightarrow)$. For some $n \in \mathbb{N}$ we have that $\varphi^{-n}(B) \subseteq U_1$. If the λ -lemma holds for $\varphi^{-n}(B)$ for all ε it also holds for B , since the metric and φ are continuous and the r -cell is compact. Therefore, the change of distance after applying φ^{-n} is finite and we can choose the distance from $\varphi^{-n}(B)$ to D small enough such that B is ε C^1 -close to D . Hence, without restrictions $B \subseteq U_1$ and $q \in U_1$. Next we define the inclination of N with respect to the (un-)stable manifolds as follows:

Let $x \in N \cap U_1$ and $v^0 \in T_x N \subseteq T_0 M = T_0^s M \oplus T_0^u M$ and let $v^0 = (v_s^0, v_u^0)$ be the coordinate entries. Furthermore let v be a unit vector. Now we define the **inclination** of v^0 :

$$\lambda(v^0) = \frac{\|v_s^0\|}{\|v_u^0\|}.$$

In other words: The smaller the inclination, the more parallel we are to $W(p \rightarrow)$. Notice how we can only define the inclination if $v_u^0 \neq 0$. Furthermore define:

$$x^n := \varphi^n(x), \quad v^n := d\varphi_{x^{n-1}}(v^{n-1}) \text{ and } \varphi(x) = (\varphi_s(x), \varphi_u(x)) \in W(\rightarrow 0) \times W(0 \rightarrow).$$

Now we want to calculate the derivative $d\varphi|_{q^n}$. But first inspect:

$$d\varphi|_x = \begin{pmatrix} \frac{\partial \varphi_s}{\partial x_s}|_x & \frac{\partial \varphi_s}{\partial x_u}|_x \\ \frac{\partial \varphi_u}{\partial x_s}|_x & \frac{\partial \varphi_u}{\partial x_u}|_x \end{pmatrix} \stackrel{(16)}{=} \begin{pmatrix} A + \frac{\partial f_s}{\partial x_s}|_x & \frac{\partial f_s}{\partial x_u}|_x \\ \frac{\partial f_u}{\partial x_s}|_x & B + \frac{\partial f_u}{\partial x_u}|_x \end{pmatrix}$$

Since $q^n \in W(\rightarrow 0)$ and $\frac{\partial f_u}{\partial x_s}(x_s, u) = 0$ this gives us:

$$d\varphi|_{q^n} = \begin{pmatrix} A + \frac{\partial f_s}{\partial x_s}|_{q^n} & \frac{\partial f_s}{\partial x_u}|_{q^n} \\ 0 & B + \frac{\partial f_u}{\partial x_u}|_{q^n} \end{pmatrix}$$

Let $B_s(\delta)$ denote the ball of radius δ around the origin that lives in $W(\rightarrow p)$ and $U_2 := B_s(\delta) \times B$. Since $\frac{\partial f_s}{\partial x_u}|_{(0, x_u)} = 0$ and by the smoothness of φ , for any $\varepsilon > 0$ there exists a δ such that

$$\sup_{U_2} \underbrace{\left\| \frac{\partial f_s}{\partial x_u} \right\|}_{:=k_1} < \min\{\varepsilon, k\}.$$

Again, we assume $q \in D \subseteq U_2$. Now we want to get a hold of the inclination: Let $x \in N$ such that $x^j \in U_2$ for all $j \leq n$ and let $v^0 \in T_x N$. Denote $v^n = (v_s^n, v_u^n) \in T_0 W(\rightarrow 0) \times T_0 W(0 \rightarrow)$.

$$\begin{aligned} \lambda(v^n) &= \frac{\|d\varphi_s|_{x^{n-1}}(v^{n-1})\|}{\|d\varphi_u|_{x^{n-1}}(v^{n-1})\|} \\ &= \frac{\left\| A v_s^{n-1} + \frac{\partial f_s}{\partial x_s}|_{x^{n-1}} v_s^{n-1} + \frac{\partial f_s}{\partial x_u}|_{x^{n-1}}(v_u^{n-1}) \right\|}{\left\| \frac{\partial f_u}{\partial x_s}|_{x^{n-1}}(v_s^{n-1}) + B v_u^{n-1} + \frac{\partial f_u}{\partial x_u}|_{x^{n-1}}(v_u^{n-1}) \right\|}} \\ &\leq \frac{(\alpha + k) \|v_s^{n-1}\| + k_1 \|(v_u^{n-1})\|}{k \|(v_s^{n-1})\| + (\beta + k) \|v_u^{n-1}\|} \\ &\leq \frac{(\alpha + k) \|v_s^{n-1}\| + k_1 \|(v_u^{n-1})\|}{(\beta - k) \|v_u^{n-1}\| - k_1 \|(v_s^{n-1})\|} && \text{since } \beta - k > 1 \text{ and } k_1 < k < 1 \\ &= \frac{(\alpha + k)\lambda(v^{n-1}) + k_1}{\beta_1 - k_1 \lambda(v^{n-1})} && \text{we divided by } \|v_u^{n-1}\| \\ &\leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1 - k_1 \lambda(v^{n-1})} \end{aligned} \tag{17}$$

If $x \in W(\rightarrow 0)$ the blue part above disappears and we get:

$$\lambda(v^n) \leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1}.$$

By induction we get:

$$\begin{aligned} \lambda(v^n) &\leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1} = \frac{\frac{\lambda(v^{n-2}) + k_1}{\beta_1} + k_1}{\beta_1} = \frac{\frac{\lambda(v^{n-2}) + \frac{k_1\beta_1}{\beta_1}}{\beta_1} + k_1}{\beta_1} = \frac{\lambda(v^{n-2}) + k_1 + k_1\beta_1}{\beta_1^2} \leq \dots \\ &\leq \frac{\lambda(v^0) + \sum_{l=0}^{n-1} k_1\beta_1^l}{\beta_1^n}. \end{aligned} \quad (18)$$

By using the geometric series we see:

$$\begin{aligned} \lambda(v^n) &\leq \frac{\lambda(v^0)}{\beta_1^n} + k_1 \sum_{l=1}^n \frac{1}{\beta_1^l} \leq \frac{\lambda(v^0)}{\beta_1^n} + k_1 \left(\frac{1}{1 - \beta_1} - 1 \right) = \frac{\lambda(v^0)}{\beta_1^n} + k_1 \left(\frac{b}{\beta_1 - 1} - \frac{\beta_1 - 1}{\beta_1 - 1} \right) \\ &= \frac{\lambda(v^0)}{\beta_1^n} + k_1 \left(\frac{1}{\beta_1 - 1} \right). \end{aligned}$$

Since β_1 was greater then 1 we can conclude that $\lim_{n \rightarrow \infty} \frac{\lambda(v^0)}{\beta_1^n} = 0$ and since $k < \frac{1}{4}(\beta_1 - 1)^2$ we know that $\frac{k}{\beta_1 - 1} < \frac{b-1}{4}$. Because of that, a n_0 exists such that for all $n > n_0$:

$$\lambda(v^n) \leq \frac{\lambda(v^0)}{\beta_1^n} + \frac{k}{b-1} < \frac{\beta_1 - 1}{3}.$$

Lets collect what we have done so far: We take a point x from a transverse intersection of N and $W(\rightarrow p)$. Then, we inspect the inclination of a tangent vector v in $T_x N \subseteq T_0^s M \times T_0^u M$ and move the tangent vector via $d\varphi$ alongside of the stable set and realize that the inclination gets smaller, the more we move it.

For any $v \in T_{q^n} N$ there exists a $\tilde{v} \in T_q N$ such that $v = \tilde{v}^n$, since $d\varphi|_x : T_x M \rightarrow T_{\varphi(x)} M$ is bijective. If we take an arbitrary local section $f : M \rightarrow TM$ that denotes the choice of $v^0 \in T_x N$ we know that $\lambda \circ f$ is continuous, as it is the composition of continuous functions. In other words, the inclination smoothly depends on v^0 and x . This tells us that for a small enough cellular neighbourhood $D_0 \subseteq D \subseteq N \cap U_2$ such that for any $x \in D_0$ and $v_0 \in T_x N$ of length 1 we have

$$\lambda(v^0) \leq \frac{1}{2}(\beta_1 - 1). \quad (19)$$

(This is true as long as the invlination is well defined) Now we claim that if $\varphi^j(v^0) \in U_2$ for all $j = 0, \dots, n$ then $\lambda(v^n) \leq \frac{1}{2}(\beta_1 - 1)$. We prove this by induction: The case $n = 0$ is already done. For the induction step we use the estimate (17), $k_1 < k < 1$ and $k < \frac{1}{4}(\beta_1 - 1)^2$:

$$\lambda(v^{n+1}) \leq \frac{\lambda(v^n) + k_1}{\beta_1 - k_1\lambda(v^n)} \leq \frac{\frac{1}{2}(\beta_1 - 1) + \frac{1}{4}(\beta_1 - 1)^2}{\beta_1 - \frac{1}{2}(\beta_1 - 1)} = \frac{1}{2}(\beta_1 - 1).$$

Using the above estimate and (17) again we conclude that for all $x \in D_0$ such that $\varphi^j(x) \in U_2$ for all $j = 0, \dots, n$ and for any unit vector $v^0 \in T_x N$ we have:

$$\begin{aligned} \lambda(v^n) &\leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1 - k_1 \lambda(v^{n-1})} \\ &\leq \frac{\lambda(v^{n-1}) + k_1}{\beta_1 - \frac{1}{2}(\beta_1 - 1)} \\ &\leq \frac{\lambda(v^{n-1}) + k_1}{(\frac{1}{2}(1 + \beta_1))}. \end{aligned}$$

Notice that for $\beta_2 := \frac{1}{2}(1 + \beta_1)$ we have $1 < \beta_2 < \beta_1$. Again like in the equation (18) we can use induction to conclude:

$$\lambda(v^n) \leq \frac{\lambda(v^0)}{\beta_2^n} + \frac{k_1}{\beta_2 - 1},$$

and since $k_1 \leq \varepsilon$ we have for any n large enough, any $x \in D_0$ such that $\varphi^j(x) \in U_2$ for $j = 0, \dots, n$ and any unit vector $v^0 \in T_x N$:

$$\lambda(v^b) \leq \varepsilon(1 + \frac{1}{\beta_2 - 1}).$$

Let $\pi_u : U \rightarrow W(0 \rightarrow)$ be the projection. Since φ is expanding on $W(0 \rightarrow)$ we know that a $n_2 \geq n_1$ exists such that $B \subseteq \pi_u(D_{n_2})$, where D_{n_2} denotes the connected component of $\varphi^{n_2}(D_0) \cap U$ that contains q^{n_2} . If $\dim N > \dim W(0 \rightarrow)$ we can choose a subset of $\dim W(0 \rightarrow)$ and then the projection restricted to D_{n_2} : $\pi_u|_{D_{n_2}}$ can be modified to be one to one. For this we use the fact that we are an embedded manifold. Hence, we are given by an implicit function where the derivatives in $W(\rightarrow p)$ -direction are invertible as a matrix. By the implicit function theorem we can take any point x from D_{n_2} and conclude that there is a neighbourhood $U(x) \subset W(p \rightarrow)$ around $\pi_u(x)$ in D_{n_2} and a smooth function $G : W(p \rightarrow) \rightarrow W(\rightarrow p)$, such that around x D_{n_2} looks like $(y, G(y))$ telling us that the projection onto the domain of G is one to one after restricting D_{n_2} to $U(x) \times \text{im}(G)$. Now we can define h to be the inverse of that projection (that is $G \times \text{id}$) and get the closeness of B to D_{n_2} . To see that we first check the distances and the difference after differentiating. For this we need the maps: $i_1 : B \rightarrow M, x \mapsto (0, x)$, and $i_2 : D_{n_2} \rightarrow M, x \mapsto x$ and $h : B \rightarrow D_{n_2}, x \mapsto \pi_u^{-1}(x) = (h_s(x), x) = (h_s(x), h_u(x))$:

$$\begin{aligned}
\sup_{x \in B} \|d(i_1)_x - d(i_2 \circ h)_x\| &= \sup_{x \in B} \left\| \begin{pmatrix} 0 \\ 1 \end{pmatrix} - \begin{pmatrix} d(h_s)_x \\ 1 \end{pmatrix} \right\| \\
&= \sup_{x \in B} \|d(h_s)_x\| \\
&= \sup_{x \in B} \sup_{\|v\| \leq 1} \left\| \frac{d(h_s)_x}{d(h_u)_x}(v) d(h_u)_x \right\| (v) \quad \text{here: } \begin{cases} h_u \equiv \text{id}_{W(\rightarrow p)} \times h_u \\ h_s \equiv h_s \times \text{id}_{W(p \rightarrow)} \end{cases} \\
&= \sup_{x \in B} \sup_{\|v\| \leq 1} \frac{\|(d(h)_x)_s(v)\|}{\|(d(h)_x)_u(v)\|} \|v\| \\
&= \sup_{x \in B} \sup_{\|v\|=1} \lambda(dh_x(v)) \quad \text{since } dh_x \text{ is linear} \\
&\leq \varepsilon \left(1 + \frac{1}{\beta_2 - 1}\right).
\end{aligned}$$

The last inequality follows from $dh_x(v)$ being an element in $T_x N$ and the existence of a $\tilde{v} \in D_0 : \tilde{v}^n = v$. Finally since β_2 is independent of ε we can make this construction for any ε . Furthermore since we can increase n (and decrease the size of $\tilde{D}$) such that D_{n^2} is arbitrary close to the origin, the statement that

$$\sup_{x \in B} \|x - h(x)\| < \varepsilon$$

is clear.

Therefore, by definition we have that for any $\mathcal{N}^r(f, (\varphi, U), (\psi, V), \varepsilon)$ that contains i_2 there is a r -cell D_{n^2} and a h such that $i_2 \circ h \in \mathcal{N}^r(f, (\varphi, U), (\psi, V), \varepsilon)$. This tells us that every neighbourhood of i_1 contains such an h and therefore, in the C^r -generating metric d_1 , an r -cell and a h exists such that $d_1(i_1, i_2 \circ h) < \varepsilon$. This concludes the proof of the lambda lemma.

we proofed the same with the only difference, that we defined the r -cell D arbitrary. We now want to show that this r -cell can be defined to be the image of a moved smaller cell of a given r -cell. So lets inspect the already given proof:

We started by moving the point of transverse intersection q closer to the fixpoint by applying the diffeomorphism to it. Then we defined an neighbourhood D_0 of that moved point in the stable manifold (compare ??). Here we were able to get a hold of the inclination. Then we moved D_0 via applying the isomorphism closer to the fixed point and restricted it to the dimension of the unstable manifold. This restriction we can refine as follows: First we define D_{n^2} to be the moved D_0 . Lets say we moved q n -times. In other words $q \in \varphi^{-n}(D_{n^2})$. Then there exist a small r -cell neighbourhood of q that lives in $D \cap \varphi^{-n}(D_{n^2})$. That is because $\varphi^{-n}(D_{n^2})$ is an in N open neighbourhood of y , hence $\varphi^{-n}(D_{n^2}) \cap D$ is open in D . Hence we can find a ball around y in D that is contained in the intersection. We call that r -cell $\tilde{D}$. Then $\varphi^n(\tilde{D})$ is an r -dimensional subset of D_{n^2} and the rest of the proof can be done with that subset. All arguments work, as that subset is transverse to $W(\rightarrow p)$.

□

Next we prove that fiberwise the flow is monotonly decreasing in any normal-direction with respect to $W(\rightarrow p)$:

Lemma 6.0.2. *Let U be a tubular neighbourhood of $W(\rightarrow p) \cap M^c$, with $J : NW(\rightarrow p) \cap M^c \rightarrow M$ beeing the corresponding embedding. For $x \in W(\rightarrow p) \cap M^c$ we denote U_x the embedded fiber over x . Then there exists a $T > 0$ big enough and a global trivialization $\Phi : NW(\rightarrow p) \cap M^c \rightarrow (W(\rightarrow p) \cap M^c) \times \mathbb{R}^{\lambda_p}$ such that the the composition $f \circ \varphi_T \circ J \circ \Phi^{-1}$ is fibrewise monotonically decreasing meaning that for any $0 \neq v \in U_x$ and $\nu > 0$ small the function*

$$l_\nu : (1 - \nu, 1 + \nu) \rightarrow \mathbb{R}$$

$$\lambda \mapsto f \circ \varphi_T \circ J \circ \Phi^{-1}(x, \lambda \cdot v)$$

has negative derivative.

Proof. We will apply the technical λ -lemma. For this we reuse the notation as follows. The lemma tells us, that for any $\mu_T > 0$ we can find a T big enough, such that $\varphi_T(J(U_x))$ is close in the following sense: There is a neighbourhood L of p and a Morse chart

$$\Psi : L \rightarrow \mathbb{R}^{\lambda_p} \times \mathbb{R}^{m-\lambda_p}$$

such that $\Psi|_{\mathbb{R}^{\lambda_p}} \supseteq W(p \rightarrow)$. In this chart the set $\varphi_T(J(U_x))$ is the graph of a funktion $h_x^s : \mathbb{R}^{\lambda_p} \rightarrow \mathbb{R}^{m-\lambda_p}$ that satisfies:

$$\|dh_v^s\| < \mu_T$$

$$\|h_v^s\| < \mu_T$$

We define the map $h_x = h_x^s \oplus \text{id}_{\lambda_p} : \mathbb{R}^{\lambda_p} \rightarrow \mathbb{R}^m$.

Now $\varphi_T(J(U_x)) = \Gamma(h_x^s) = \text{im}(h_x)$. Using this we define our global trivialisation:

$$\Phi : NW(\rightarrow p) \cap M^c \rightarrow (W(\rightarrow p) \cap M^c) \times \mathbb{R}^{\lambda_p}$$

$$\xi \mapsto \left(\pi(\xi), h_{\pi(\xi)}^{-1} \circ \Psi \circ \varphi_T \circ J(\xi) \right)$$

In this trivialization the funktion l_ν looks like:

$$l_\nu(\lambda) = f \circ \varphi_T \circ J \circ \Phi^{-1}(x, \lambda v)$$

$$= f \circ \varphi_T \circ J \circ J^{-1} \circ \varphi_{-T} \circ \Psi^{-1} \circ h_x$$

$$= f \circ \Psi^{-1} \circ h_x.$$

In coordinates this reads:

Now we can calculate its derivative to conclude the statement:

$$\begin{aligned}
d(l_v)(\lambda) &= d(f \circ \psi^{-1})|_{h_x(\lambda v)} \circ dh_x|_{\lambda v}(v) \\
&= d(f \circ \psi^{-1})|_{h_x(\lambda v)}(v, dh_x^s(v)) \\
&= -2\lambda \sum_{i_1}^{\lambda_p} (v^i)^2 + (h_x^s(\lambda v))^T \cdot (dh_x^s)^i(\lambda v) \\
&\leq -2\lambda \|v\|^2 + \lambda^2 (\mu^T)^2 \|v\|^2 \\
&= (-2 + (\mu^T)^2 \lambda) \lambda \|v\|^2 < 0 \text{ for all } v \neq 0, \lambda \in I \text{ and for all } T \text{ big enough.}
\end{aligned}$$

□

Hence, the sets $N_p \cap U_x$ are starshaped with respect to the embedding of the normal space.

Now we want to **smoothly** rescale to the starshaped neighbourhood. This is in general not possible and homeomorphisms from starshaped neighbourhoods to R^n are hard to describe, even if they are not required to be a rescaling. The main problem in the general case is, that the following: for each $v \in \varphi_T(J(U_x))$ by the star shaped property there is a unique λ_v , such that $f \circ (\lambda_v v) = a - \varepsilon$. But the map $v \mapsto \lambda_v$ needs not to be continuous in the general starshaped setting however here it is and it even continuously depends on x :

Definition 6.0.3 (Hyperspherical coordinates). For our next proof we need to introduce spherical coordinates on R^{λ_p} . Define $s_k := \sin(\theta_k)$ and $c_k := \cos(\theta_k)$:

$$\Lambda : \mathbb{R}_{\geq 0} \times [0, \pi]^{\lambda_p-2} \times [0, 2\pi) \rightarrow \mathbb{R}^{\lambda_p}$$

$$(r, \theta_1, \dots, \theta_{\lambda_p-1}) \mapsto \begin{pmatrix} rc_1 \\ rs_1c_2 \\ \vdots \\ rs_1 \cdots s_{\lambda_p-2} c_{\lambda_p-1} \\ rs_1 \cdots s_{\lambda_p-2} s_{\lambda_p-1} \end{pmatrix}$$

With differential

$$\begin{pmatrix} c_1 & -rs_1 & 0 & 0 & \cdots & 0 \\ s_1c_2 & rc_1c_2 & -rs_1s_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \\ s_1 \cdots s_{n-2} c_{n-1} & \cdots & \cdots & \cdots & -rs_1 \cdots s_{n-2} s_{n-1} \\ s_1 \cdots s_{n-2} s_{n-1} & rc_1 \cdots s_{n-1} & \cdots & \cdots & rs_1 \cdots s_{n-2} c_{n-1} \end{pmatrix}$$

We have shown in lemma 6.0.2 that the maps l_v are monotonically decreasing. Hence the condition $f(x) \geq a - \varepsilon$ gives rise to a starshaped fiber. But we want the fibers to be homeomorphic to R^{λ_p} and the dream would be to just rescale it as follows: Assume that $\rho > 0$ is small enough, such that for all $x \in (W(\rightarrow p) \cap M^c)$ we have that $\sup_{x \in h_x(B_\rho(0))} (f(x)) \geq a - \varepsilon$. Now we have the maps:

$$\begin{aligned} r_x : h_x(\partial(B_\rho(0))) &\rightarrow \mathbb{R}_{\geq 1} \\ v &\mapsto \lambda_v \text{ such that } f(\lambda_v v) = a - \varepsilon \end{aligned}$$

Notice that for small enough ε such a λ_v exists and the lemma 6.0.2 gives us uniqueness. Now we want to have continuooity and furthermore continouos dependence on x for which we will use the parameter-dependent implicit funktion theorem:

Lemma 6.0.4. *The parameterdependent funktions r_x are continouos and depend continouos on x*

Proof. We redefine the funktion r_x by using spherical coordinates as follows: Implizitly for each x we have the Mapping:

$$\begin{aligned} F_x : \mathbb{R}_{\geq 0} \times [0, \pi]^{\lambda_p-2} \times [0, 2\pi) &\rightarrow \mathbb{R} \\ (r, \theta_1, \dots, \theta_{\lambda_p-1}) &\mapsto f \circ h_x \circ \Lambda(r, \theta_1, \dots, \theta_{\lambda_p-1}) \end{aligned}$$

Here the funktion reads:

$$f \circ h_x \circ \underbrace{\Lambda(r, \theta_1, \dots, \theta_{\lambda_p-1})}_{:=v} = f \left(\begin{pmatrix} h_x^s(\Lambda(v)) \\ \Lambda(v) \end{pmatrix} \right) = -2r^2 + 2 \sum_{i=\lambda_p+1}^m (h_x(\Lambda(v)))_i^2$$

To make use of the implizit funktion theorem, we inspect the differential in r -direktion:

$$\frac{\partial}{\partial r} (-2r^2 + 2 \sum_{i=\lambda_p+1}^m (h_x(\Lambda(v)))_i^2) = -4r + 2 \sum_{i=\lambda_p+1}^m \left(\sum_{j=1}^{\lambda_p} \left(\frac{\partial (h_x)_i}{\partial x_j} \frac{\partial \Lambda_j}{\partial r} \right) \right)$$

This does not vanish for big enough r , since the partial differentials of λ are of norm one, and those of h_x are small. Hence for big enough r , lets say $r \geq \mu > 0$ this is negative and thereby we get a funktion

$$\tilde{r}_x : [0, \pi]^{\lambda_p-2} \times [0, 2\pi) \rightarrow \mathbb{R}_{\geq \mu}$$

such that $(f \circ h_x \circ \Lambda)^{-1}(a - \varepsilon)$ is the graph of $\tilde{r}_x$. Hence, if we define $\pi_r : \mathbb{R}_{\geq 0} \times [0, \pi]^{\lambda_p-2} \times [0, 2\pi) \rightarrow [0, \pi]^{\lambda_p-2} \times [0, 2\pi)$ to be the projection we have that

$$\begin{aligned} r_x : h_x(\partial B_\rho(0)) &\rightarrow \mathbb{R} \\ v &\mapsto \tilde{r}_x \circ \pi_r(\Lambda^{-1}(h_x^{-1}v)) \end{aligned}$$

f mit $f \circ \Psi^{-1}$ ersetzen, da hier eigentlich die Morse karte genutzt wird!!

is continuous and depends continuously on x . (In reality we still need to check for smoothness in $\Lambda(\{\theta_{\lambda_p-1} = 0\})$. This can be done by a different parametrisation of the sphere meaning by different spherical coordinates.) The resulting function agrees with the earlier definition of r_x concluding the proof. $\square$

Proof That N_p is Tubular. To finish our proof we rescale the Fibers smoothly. For this we first restrict the domain of J : Let $\nu > 0$ be small enough, such that for

$$N_\nu W(\rightarrow p) \cap M^c := \{\xi \mid \text{if } \Phi(\xi) = (x, v), \|v\| < \nu\}$$

the supremum

$$\sup_{x \in \varphi_T \circ J(N_\nu W(\rightarrow p) \cap M^c)} \geq a - \varepsilon$$

Hence, $J(N_\nu W(\rightarrow p) \cap M^c)$ is included in N_p and now we rescale the fibres $U_x^\nu := U_x \cap \text{im} J(N_\nu W(\rightarrow p) \cap M^c) \supseteq U_x$ to get a surjective inclusion into N_p . For all $x \in W(\rightarrow p) \cap M^c$ we define the smooth rescaling as follows:

$$RS : U_x^\nu \rightarrow U_x$$

$$v \mapsto \varphi_{-T} \circ \Psi^{-1} \circ r_x \left(\rho \cdot \frac{h_x^{-1} \circ \Psi \circ \varphi_T(v)}{\|h_x^{-1} \circ \Psi \circ \varphi_T(v)\|} \right)$$

This rescaling is smooth, smoothly depends on x , and it maps surjectively onto $N_p \cap U_x$. $\square$

Theorem 6.0.5. *Given the filtration $N_{-1} \subseteq N_0 \subseteq \dots \subseteq N_{m-1} \subseteq N_m = M$ from (11). The following diagram commutes:*

$$\begin{array}{ccc}
 C^k(M, \mathfrak{A}, \tilde{K}^l(pt)) & \xrightarrow{\partial^p} & C^{k+1}(M, \mathfrak{A}, \tilde{K}^l(pt)) \\
 \downarrow & & \downarrow \\
 \bigoplus_{p \in \text{Crit}_k(f)} K^{k+l}(N_p, L_p) & \xrightarrow{\Delta_{k+l}} & \bigoplus_{q \in \text{Crit}_{k+1}(f)} K^{k+l+1}(N_q, L_q) \\
 \downarrow & & \downarrow \\
 K^{k+l}(N_k, N_{k-1}) & \xrightarrow{\delta_{triple}} & K^{p+l+1}(N_{k+1}, N_k)
 \end{array}$$

Proof Strategy:

We will first build up a diagram that encapsulates the connecting homomorphism from K-theory, which was build as follows: Assume that (X, Y, Z) is a topological triple. Then the first map is induced from the kollaps: $Y \rightarrow Y/Z$. Afterwards we have the connecting homomorphism of the pair (X, Y) :

The K-groups of the pointed spaces X/Y and $X \cup CY$, where the latter denotes a cone above Y agree. From the latter we can quotient out X , and $X \cup CY/X \cong SY$. Hence we have the triple connecting homomorphism as the composition:

$$K^{-1}(Y, Z) \rightarrow K^{-1}(Y) = K(SY) \cong K(X \cup CY/X) \rightarrow K(X \cup CY) \cong K(X/Y)$$

The triple connecting homomorphism for lower K-Groups is defined exactly the same via the identification $K^{-n}(X, Y) = K(S^n \vee (X, Y))$. For higher K-Groups we apply the natural Bott isomorphism and make use of the identity. By doing this we get the tripple homomorphism

$$\begin{array}{ccccc}
 K^n(Y, Z) & \xrightarrow{\delta_{triple}} & K^{n+1}(X, Y) \\
 \Downarrow & & \Downarrow \\
 K^{-n}(Y, Z) & \xrightarrow{i^*} K^{-n}(Y, p) \xrightarrow{\delta_{double}} K^{-n+1}(X, Y) & \xrightarrow{bott} & K^{-n-1}(X, Y)
 \end{array}$$


For the proof of the upper square, we actually proof that the following commutes:

$$\begin{array}{ccc}
 C^k(M, \mathfrak{A}, \tilde{K}^l(pt)) & \xrightarrow{\partial^p} & C^{k+1}(M, \mathfrak{A}, \tilde{K}^l(pt)) \\
 \downarrow & & \downarrow \\
 \bigoplus_{p \in \text{Crit}_k(f)} K^{-k}(N_p, L_p) & \xrightarrow{\Delta_k} & \bigoplus_{q \in \text{Crit}_{k+1}(f)} K^{-(k+1)}(N_q, L_q)
 \end{array}$$

Diagram 20

Furthermore we assumed that $l = 0$. This doesnt change the generality since if l was even, we can apply the Bott isomorphism and else the groups are 0. Now we choose a triple (A, B, C) in M such that:

- (a) (A, B) is a regular index pair for a critical point $q \in \text{Crit}_{k+1}$.
- (b) (B, C) is a regular index pair for a critical point $p \in \text{Crit}_k$.



From the perspective of K-theory a regular index pair for a critical point is the same as a sphere with the dimension beeing the index of said critical point. To see this just use a small index pair in a Morse chart (they are all homotopic anyway), collapse the stable part and quotient by the exit set. By doing this we get the communative diagraphm:

$$\begin{array}{ccc}
 K^{k-1}(B, C) & \xrightarrow{\delta_{triple}} & K^k(A, B) \\
 \downarrow & & \downarrow \\
 K^{k-1}(S^{k-1}) & \longrightarrow & K^k(S^{k-1}) \\
 \downarrow & \nearrow & \\
 K^k(S^k) & &
 \end{array}$$

And as a map between K Groups of spheres, we know that this map corresponds to the question of orientation. Now we have the maps

- (a) $S^k \rightarrow S \vee (B/C)$ induced from an orientation in the unstable manifold arround p together with $-\text{grad}(f)(x_i)$ where $x_i \in W(q \rightarrow p)$.
- (b) $S^k \rightarrow (A/B)$ induced from an orientation in the unstable manifold arround q .

And hence the diagram

$$\begin{array}{ccc} K^k(S \vee (B, C)) & \xrightarrow{\delta_{triple}} & K^k(A, B) \\ & \nwarrow \quad \nearrow & \\ & K^k(S^k) & \end{array}$$

The map now factors over The K-Group of a bouquet of k -spheres generated over the connecting flow lines. In each such sphere the comparison of orientations happens.

Bevore we start proving this we need the lemma:

Lemma 6.0.6. *We work in a compact, riemanian, orientable, smooth Manifold (M, g) of dimension m together with a Morse-Smale funktion*

$$f : M \rightarrow \mathbb{R} \quad (21)$$

We assume that q is a critical point of index $k+1$ and p is a critical point of index k . we define $f(q) = b$ and $f(p) = a$ and assume that there is no critical point in $f^{-1}(a, b) \subseteq M$. We define the sub and superlevelsets

$$M^t := \{x \in M | f(x) \leq t\} \quad \text{and} \quad M_t := \{x \in M | f(x) \geq t\}, \quad (22)$$

and the constants

$$c \in (a, b) \quad , \quad \varepsilon > 0 \text{ small} \quad , \quad T > 0 \text{ big} . \quad (23)$$

With this we define the sets:

$$N_q := \{x \in M_c | f(\varphi_{-T}(x)) \leq b + \varepsilon\}, \quad (24)$$

$$L_q := \{x \in N_q | f(x) = c\}, \quad (25)$$

$$N_p := \{x \in M^c | f(\varphi_T(x)) \geq a - \varepsilon\}, \quad (26)$$

$$L_p := \{x \in N_p | f(\varphi_T(x)) = a - \varepsilon\}, \quad (27)$$

and finally:

$$C := N_p \cup N_q, \quad (28)$$

$$B := N_p \cup L_q, \quad (29)$$


$$A := L_p \cup (L_q \overset{\circ}{-} N_p). \quad (30)$$

We claim that

1. (N_q, L_q) is a regular index pair for q .
2. (C, B) is an index pair for q .
3. (N_p, L_p) is a regular index pair for p .
4. (B, A) is an index pair for p .
5. N_p is a tubular neighbourhood of $W(\rightarrow p) \cap M^c$.

Proof Strategy:

The statements 1-4 are easy to proof by the definitions. The regularity can be derived from a general fact for pairs (X, Y) in metric spaces: If Y is closed in X and there is a neighbourhood of Y that is open in X such that Y is a strong deformation retract of U . So the only thing left to show is, that N_p is a tubular neighbourhood. In [Sal90] in the proof of lemma 3.2 Salamon claims this to be true without a proof (page 119 in the attached source). Similar, in [BH04] Banyaga claims this (also without any argument). The Set N_p is sketched in the figure below. The Idea of the proove is as follows: By increasing T (or decreasing ε) we can assume N_p to be an arbitrary small neighbourhood of $W(q \rightarrow) \cap M^c$. Now using the flow we can push N_p close to p ($\varphi_N(N_p)$ for big N), where we use a tubular nieghbourhood embedding J that overlaps N_p . We now need to proof that we can reduce the radii of the fibers in the domain of J in a smooth way such that J actually becomes a embedding onto the N_p (that we pushed close to p). For this we will make use of the lambda lemma (in an improved version), which lets us induce a metric from the unstable tangend space of P into the fiberes of J (since they are transversal to the stable set). With this metric we can finally smoothly reduce the radii of our domain of J to end up with the statement, that $\varphi_N(N_p)$ is a tubular neighbourhood of a small part of the stable set of p and hence N_p also. The prove is given as an appendix in section 6.



The same arguments work to show that N_q is a tubular neighbourhood of $W(q \rightarrow) \cap M_c$, by replacing f with $-f$. Furthermore, we can assume, that the V_j from below are fibers in the normal neighbourhood. This can be realized, since V_j intersect transversal and by trivialization we have a global frame of the bundle and hence can smoothly rescale the embedding along the frame to get that V_j is the image of the fibre above x_j .

Proof of Theorem 6.0.5. We first notice, that we have the map of triples given by the

inclusion:

$$\left(W(q \rightarrow) \cap N_q, S(q \rightarrow), S(q \rightarrow) \setminus \bigcup_j V_j \right) \rightarrow (C, B, A)$$

and hence by naturality we have the diagram, where the top square commutes:

$$\begin{array}{ccc}
 K^{-\lambda_p} [B, A] & \xrightarrow{\delta_{triple}^1} & K^{-\lambda_q} [C, B] \\
 \downarrow i_{B,A}^* & & \downarrow i_{C,B}^* \\
 K^{-\lambda_p} \left[S(q \rightarrow), S(q \rightarrow) \setminus \bigcup_j \text{int}(V_j) \right] & \xrightarrow{\delta_{triple}^2} & K^{-\lambda_q} [W(q \rightarrow) \cap N_p, S(q \rightarrow)] \\
 \uparrow i_j^* \downarrow c_j^* & \nearrow \delta_{triple}^j & \\
 K^{-\lambda_p} [V_j, \partial V_j] & &
 \end{array}$$

Diagram 31

Furthermore, in this diagram the tuple $\left(K^{-\lambda_p} \left[S(q \rightarrow), S(q \rightarrow) \setminus \bigcup_j \text{int}(V_j) \right], (c_j^*)_j \right)$ is a coproduct, and we define $\delta_{triple}^j := \delta_{triple}^2 \circ c_j^*$. Then we have that

$$\delta_{triple}^2 = \sum_j \delta_{triple}^j \circ i_j^*$$

Since $i_{C,B}^*$ is an isomorphism (since $W(q \rightarrow) \cap N_p$ is a deformation retract of C and the collapse collapses B to $S(q \rightarrow)$), we have that

$$\Delta(\alpha_q) = \delta_{triple}^1(\alpha_q) = i_{C,B}^* \circ \sum_j \delta_{triple}^j \circ i_j^* \circ i_{B,A}^*(\alpha_q) \quad (32)$$

The map δ_{triple}^2 is defined as the composition:

$$\begin{array}{ccc}
 K^{-\lambda_p} [V_j, \partial V_j] & & \\
 \downarrow c_j^* & & \\
 K^{-\lambda_p} \left[S(q \rightarrow), S(q \rightarrow) \setminus \bigcup_j \text{int}(V_j) \right] & \xrightarrow{\delta_{triple}^2} & K^{-\lambda_q} [W(q \rightarrow) \cap N_p, S(q \rightarrow)] \\
 \downarrow i_1^* & & \uparrow \beta_{\text{triv}_{C_1, S(q \rightarrow)}} \\
 K^{-\lambda_p} [S(q \rightarrow)] & \xrightarrow{h_1^1} K^{-\lambda_p+1} [S_1 \vee S(q \rightarrow)] \xrightarrow{h_2^1} K^{-\lambda_p+1} [W(q \rightarrow) \cap N_p \cup C_1(S(q \rightarrow)), W(q \rightarrow) \cap N_p] \xrightarrow{i_2^1} K^{-\lambda_p+1} [W(q \rightarrow) \cap N_p \cup C_1(S(q \rightarrow))] &
 \end{array}$$

Diagram 33

The downwards maps in the theorem are the induction of the canonical generators. This is done by the following assignment

Now the composition

$$h_1 * i_1 * \circ c_j^* : K^{-\lambda_p} [V_j, \partial V_j] \rightarrow K^{-\lambda_p+1} [S_1 \vee S(q \rightarrow)]$$


can be simplified by the map

$$l^* : K^{-\lambda_p} [V_j, \partial V_j] = K^{-\lambda_p+1} [S_1 \vee (V_j, \partial V_j)] \rightarrow K^{-\lambda_p+1} [S_1 \vee S(q \rightarrow)]$$

where l is the collapsing map.

$$\begin{aligned} l : S_1 \vee S(q \rightarrow) &\rightarrow S_1 \vee (V_j, \partial V_j) \\ (t, x) &\mapsto (t, \bar{x}) \end{aligned}$$

So now we are in the following situation:



The maps in the diagram are given as follows:

- The map f_p induces α_p . Remember, that α_p comes from the canonical generator α'_p of the index pair $[B, A]$ and α'_p is defined as

$(\varphi \circ \tilde{\psi}^{-1})^*(\beta)$ where φ is the homotopie equivalence of index pairs

$\tilde{\psi}$ is a Morse chart adapted to the basis of $T_p^u M$,

and β is the bott generator of the sphere $[D_\varepsilon^u \times D_\varepsilon^s / \partial D_\varepsilon^u \times D_\varepsilon^s]$.

Instead of using φ we can rescale $\tilde{\psi}$ to be a mapping to $[B, A]$ instead of first mapping to the prototype index pair $[D_\varepsilon^u \times D_\varepsilon^s, \partial D_\varepsilon^u \times D_\varepsilon^s]$. We call this rescaled Morse chart ψ . Then $\psi^{-1}(\beta) = (\varphi \circ \tilde{\psi}^{-1})^*(\beta)$ since the map $\tilde{\psi} \circ \varphi^{-1} \circ \psi^{-1}$ has the local and hence global degree one. (this can be checkt by differentiatiaion at the origin). Hence $\alpha'_p = (\psi^{-1})^*(\beta)$. The inclusion

$$i_p : (V_j, \partial V_j) \rightarrow (B, A)$$

induces an isomorphism in the K-groups since N_p is a tubular neighbourhood of the contractible $W(\rightarrow p) \cap M^c$ and the contraction to the fiber V_j contracts L_p to ∂V_j (the contraction is well defined on the quotients, and for K-Theory, we only care for the qutoient spaces). Now $\alpha''_p = i_p^*(\alpha'_p)$. Define the two funktions

$$\begin{aligned} J : \mathcal{N}^{W(q \rightarrow)} W(\rightarrow p) \cap M^c &\rightarrow N_p \text{ the normal neighbourhood embedding,} \\ \nu : \mathcal{N}^{W(q \rightarrow)} W(\rightarrow p) \cap M^c &\rightarrow W(\rightarrow p) \times T_p^u M \text{ The trivialization of the normal bundle.} \end{aligned}$$

Define $\mathfrak{b}^p$ to be an ordered basis of $T_p^u M$ that is compatible to the orientation of said tanged space. Then assume that J is defined, such that

$$(B_{\mathfrak{b}^p}^{-1} \circ \nu_{x_j} \circ J^{-1})^* (\beta) = \alpha_p''.$$

If this is not the case just compose J with a basechage of degree one. Now α_p'' generates $K^{-\lambda_p}(V_j, \partial V_j)$. The latter is isomoprhic to $K^{-\lambda_p+1}(I \times V_j, \partial(I \times V_j))$ by definition. Now we denote the suspension is the wedge produkt with a sphere. Hence for $K^{-\lambda_p+1}(I \times V_j, \partial(I \times V_j))$ we can induce the generator by the function

$$\begin{aligned} f_p : I \times V_j / \partial(I \times V_j) &\rightarrow S \wedge S^{\lambda_p} \simeq S^{\lambda_p+1} \\ (t, x) &\mapsto \Omega(t, \Gamma \circ (B_{\mathfrak{b}^p}^{-1} \circ \nu_{x_j} \circ J^{-1})(x)). \end{aligned}$$

Γ maps the disk modulo its boundary to a sphere and Ω maps the suspension of the sphere to the sphere one dimension higher These Maps are given in definitino ?? and ??. We abuse the notation a bit and identify the map $(B_{\mathfrak{b}^p}^{-1} \circ \nu_{x_j} \circ J^{-1})$ with the one defined by quotienting out the boundary. We will leave out the quotienting in the notatino from now on to make the prove easier to read. The invested reader should check, if the quotiend maps are always well defined! Now finally we have a little definition issue: In the diagramm we have that

$$f_p^* : K^{\lambda_p-1}[\mathbb{S}^{\lambda_p+1}] \rightarrow K^{-\lambda_p}[V_j, \partial V_j],$$

whilst at the moment, f_p maps to $K^{-\lambda_p-1}(I \times V_j, \partial(I \times V_j))$. Hence we need to compose this f_p^* with the isomorphism that comes from the suspension axiom:

$$K^{-\lambda_p}(X) \cong K^{-\lambda_p+1}(SX).$$

And then add the bott isomorphism to rais th eindex by two. By abuse of notation we dont write that idetification and rather think of the top left corner as $K^{-\lambda_p+1}(I \times V_j, \partial(I \times V_j))$.

- Again, we want f_q to be the map that induces the generator of $K^{-\lambda_q}[W(q \rightarrow) \cap N_q, S(q \rightarrow)]$ which comes from the canonical generator of $[C, B]$ together with the inclusion. Again instead of first introducing the generator on the prototype index pair, we can use E_q^{u-1} instead of the Morse chart. Again we can argue here, that the local degree of changing between the two is one, since E_q^{u-1} is a refined morse chart which can be choosen to be adapted to the ordered basis $\mathfrak{b}^q$ of $T_q^u M$. Hence we can define f_q to be the quotient map:

$$\begin{aligned} W(q \rightarrow) \cap N_q / S(q \rightarrow) &\rightarrow S^{\lambda_p+1} \\ x &\mapsto \Gamma \circ B_{\mathfrak{b}^q}^{-1} \circ E_q^{u-1}(x). \end{aligned}$$

Here Γ maps the disk modulo its boundary to the sphere. Hence $f_q = (B_{\mathfrak{b}^q}^{-1} \circ E_q^{u-1})^* : K$

- The map l_1 is the collapsing map

$$S_1 \wedge S(q \rightarrow) \rightarrow S_1 \wedge (V_j / \partial V_j)$$

- The map l_2 is not topologically induced.
- The map s is given by $f_p \circ l_1$. Again we abuse the notation and leave out the suspension isomorphism when

$$s^* : K^{-\lambda_p+1}[S_1 \wedge S(q \rightarrow)] \rightarrow K^{-\lambda_p-1}[\mathbb{S}^{\lambda_p+1}].$$

- To find the map r that fits the diagram (to make it commute) is the next goal. First we need to dig into l_2 : From diagram 33 we know that $l_2 = \Sigma \circ \text{triv}_{C_1 S(q \rightarrow)} \circ i_2^* \circ h_2^*$, Where Σ denotes the suspension isomorphism. Hence, if the function r^* makes the diagram commutative we need to ensure, that $f_q^* \circ r^* = l_2$. If we define the function

$$\text{col} : W(q \rightarrow) \cap N_p \cup C_1 S(q \rightarrow) \rightarrow S_1 \wedge S(q \rightarrow)$$

to be the collision, of $W(q \rightarrow) \cap N_p$ to a point, then $i_2^* h_2^* = \text{col}^*$. Now while the function $\text{triv}_{C_1 S(q \rightarrow)}$ is not topologically induced, its inverse is. Lets call the collaps of $C_1 S(q \rightarrow)$:

$$k_1 : (W(q \rightarrow) \cap N_p) \cup C_1 S(q \rightarrow) \rightarrow (W(q \rightarrow) \cap N_p / S(q \rightarrow))$$

π_1 . Then since Σ is a natural transformation we have

$$f_q^* \circ r^* = l_2 \Leftrightarrow k_1^* \circ f_q^* \circ r^* = \Sigma \circ \text{col}^*$$

Now all maps are topologically induced except Σ and the latter is just used to geht the index in the K group right. Hence, if r would be such that

$$r \circ f_q \circ c_1 = \text{col},$$

then the requirement on the induced maps would be satisfied. We first however need to formulate a little lemma:

Lemma 6.0.7. *Let $S^k := \{x \in \mathbb{R}^{k+1} \mid \|x\| = 1\}$, $H_+ := \{x \in S^k \mid x_0 \geq 0\}$ denote the upper hemissphere, H_- the lower hemissphere, $\pi : S^k \rightarrow D^k$ with $\pi(x_0, \dots, x_k) = (x_1, \dots, x_k)$ be the projection and Λ be the homeomorphism from $D^k / \partial D^k \rightarrow S^k$. Let furthermore $c_{Eq} : S^k \rightarrow \overline{H_+} \vee \overline{H_-}$ be the collaps of the equator, where $\overline{H_+}$ denotes the upper hemisphere modulo the equator and $\overline{H_-}$ is defined analog. Let finally $c_2 : \overline{H_+} \vee \overline{H_-} \rightarrow \overline{H_+}$ be the collaps of the second summand and analogusly $c_1 : \overline{H_+} \vee \overline{H_-} \rightarrow \overline{H_-}$. Then we have that $\Lambda \circ \pi \circ c_2 \circ c_{Eq} \simeq -\Lambda \circ \pi \circ c_1 \circ c_{Eq}$, where $-$ denotes any smooth homotopie equivalence of negative degree.*

Proof. This lemma is the concrete version of a standart lemma that is often formulated along the lines: Let $q : S^k \rightarrow S^k \vee S^k$ be the collaps of the equator and p_i be the projection to the i -th summand, then $p_1 \circ q = -p_2 \circ q$. $\square$

Now we need to transport our situation to fit the lemma. For this we will use the maps defined in the lemma above, and need a bunch of maps again:

- Firts, the map ρ is given by

$$\begin{aligned} \rho : W(q \rightarrow) \setminus \{q\} &\rightarrow W(q \rightarrow) \cap f^{-1}(c) \setminus W(q \rightarrow) \setminus \{q\} \\ x &\mapsto \varphi_{\tau(x)}(x), \end{aligned}$$

and

$$\begin{aligned} \tau : W(q \rightarrow) \setminus \{q\} &\rightarrow \mathbb{R} \\ x &\mapsto \tau(x) \text{ such that } f \circ \varphi_{\tau(x)}(x) = c. \end{aligned}$$


- $\Phi : S^{\lambda_q} \rightarrow W(q \rightarrow) \cap N_p \cup C_1 S(q \rightarrow)$ is defined with a case distinction. We define

$$\begin{aligned} \Phi|_{H_+}(x) &= E_q^u \circ B_{\mathfrak{b}^q}^{-1} \circ \pi(x) \\ \Phi|_{H_-}(x) &= (e^{\tau(y)}, \rho(y)) \text{ where } y = E_q^u \circ B_{\mathfrak{b}^q}(\pi_-(x)). \end{aligned}$$

Notice, how this function is continuous and does not depend on the choice taken in the definition of ρ' .

- $\Psi : S^{\lambda_q} \vee S^{\lambda_q} \rightarrow (W(q \rightarrow) \cap N_p / S(q \rightarrow)) \vee \mathbb{S} \wedge S(q \rightarrow)$ is given by mapping $x \mapsto \overline{\Phi(x)}$. The well definition is easy to see.
- p_i for $i = 1, 2$ denotes the projection onto the first or second summand in a pointed sum.
- $C_{S(q \rightarrow)} : W(q \rightarrow) \cap N_q \cup C_1 S(q \rightarrow) \rightarrow (W(q \rightarrow) \cap N_p / S(q \rightarrow)) \vee \mathbb{S} \wedge S(q \rightarrow)$ is the collaps of $S(q \rightarrow)$.
- $R : \mathbb{S}^{\lambda_q} \rightarrow \mathbb{S} \wedge S(q \rightarrow)$ is defined to be $col \circ \Phi$

This gives the commutative diagram below, where by the lemma above the upper path is the negative of the lower path.



Now define $r := -R$. For example apply R after the reflection $(x_0, \dots, x_{\lambda_q}) \mapsto (-x_0, \dots, x_{\lambda_q})$. Then by the lemma above and commutativity we have up to homotopy:

$$\begin{aligned} r \circ f_q \circ k_1 &= -R \circ f_q \circ k_1 \circ \Phi \circ \Phi^{-1} \\ &= -R \circ \Lambda \circ \pi \circ c_2 \circ C_{Eq} \circ \Phi^{-1} \\ &= R \circ \underbrace{\Lambda \circ \pi \circ c_1 \circ C_{Eq}}_{\simeq \text{id}} \circ \Phi^{-1} \\ &= col. \end{aligned}$$

With r we finally have all the maps. Now we want to find out what degree the composition $s \circ r$ has. For this we calculate the local degree in some point by checking the sign of the determinant of the Jacobean. This strategy works out since we are in the situation of lemma ???. Before we start differentiating, we expand the map around $x \in \mathbb{R}^{\lambda_q}$, such that $x' := E_{bq}^u \circ B_{bq}(x) \in \rho^{-1}(x_j)$, meaning that x' is on the flow line towards x_j and

assume that $\tau(x') = 1$.

$$\begin{aligned}
& \pi_+ \circ s \circ r \circ \pi_+^{-1}(x_1, \dots, x_{\lambda_q}) \\
&= \pi_+ \circ s \circ R(-x_0, x_1, \dots, x_{\lambda_q}) \quad x_0 > 0 \\
&= \pi_+ \circ s \circ \text{col} \circ \Phi|_{H_-}(-x_0, x_1, \dots, x_{\lambda_q}) \\
&= \pi_+ \circ f_p \circ l_1 \circ \text{col}(e^{\tau(x')}, \rho(x')) \quad \text{col and } l_q \text{ are the identity arround } (e^{\tau(x')}, \rho(x')) \\
&= \pi_+ \circ \Omega \left(e^{\tau(E_q^u \circ B_{\mathfrak{b}^q}(x))}, \underbrace{\Gamma \circ B_{\mathfrak{b}^p}^{-1} \circ \nu_{x_j} \circ J^{-1} \circ \rho \circ E_q^u \circ B_{\mathfrak{b}^q}(x)}_{:=Z(x)} \right) \\
&= \sin(\pi e^{\tau(E_q^u \circ B_{\mathfrak{b}^q}(x))}) \cdot \begin{pmatrix} -\cos(\pi \cdot \|Z(x)\|) \\ \sin(\pi \cdot \|Z(x)\|) \cdot \frac{Z(x)}{\|Z(x)\|} \end{pmatrix}
\end{aligned}$$

Here Z is a funktion $Z : \text{int}(I \times D^{\lambda_q}) \rightarrow \mathbb{R}^{\lambda_q-1}$ and we can calculate its jacobean Notice how $Z(y) = 0$ where y is the same as above. We only care for the sign of its jacobean, so we will manipulate the funktion in a way that might change the funktion, but not the sign of ots Jacobean determinant. Let $Q_y : V \rightarrow T_y V$ be the canonical map, that identifies a linear space with the tangendspace of a point from inside said space. Now we calculate:

$$\begin{aligned}
& Q_0^{-1} \circ d(B_{\mathfrak{b}^p}^{-1} \circ \nu_{x_j} \circ J^{-1} \circ \rho \circ E_q^u \circ B_{\mathfrak{b}^q})|_y \circ Q_y(e_i) \\
&= Q_0^{-1} \circ d(B_{\mathfrak{b}^p}^{-1} \circ \nu_{x_j} \circ J^{-1} \circ \rho \circ E_q^u)|_{B_{\mathfrak{b}^q}(y)} \circ Q_{B_{\mathfrak{b}^q}(y)} \circ B_{\mathfrak{b}^q}(e_i) \\
&= Q_0^{-1} \circ d(B_{\mathfrak{b}^p}^{-1} \circ \nu_{x_j} \circ J^{-1})|_{x_j} \circ d(\rho \circ E_q^u)|_{B_{\mathfrak{b}^q}(y)} \circ Q_{B_{\mathfrak{b}^q}(y)} \circ B_{\mathfrak{b}^q}(e_i)
\end{aligned}$$

Now the term $d(\rho \circ E_q^u)|_{B_{\mathfrak{b}^q}(y)} \circ Q_{B_{\mathfrak{b}^q}(y)} \circ B_{\mathfrak{b}^q}$ maps e_1 to zero. and $(e_2, \dots, e_{\lambda_q})$ induces a basis in $T_{x_j} V_j$ that is compatible with a basis $(\mathfrak{d}_i)_{i=2, \dots, \lambda_q}$ that satisfies the following: Adding $-\text{grad}_{x_j}(f)$ as the first basis vector to $(\mathfrak{d}_i)_{i=2, \dots, \lambda_q}$ gives a basis of $T_{x_j} W(q \rightarrow)$ that is compatible with the basis induced from $\mathfrak{b}^q$ that appears in the counting of flow lines in the Morse boundary operator. This was proven earlier in lemma 2.0.23. Now we can look at the first part:

$$\begin{aligned}
Q_0^{-1} \circ d(B_{\mathfrak{b}^p}^{-1} \circ \nu_{x_j} \circ J^{-1})|_{x_j} &= \left(dJ|_{x_j} \circ d\nu_{x_j}^{-1}|_0 \circ dB_{\mathfrak{b}^p}|_0 \circ Q_0 \right)^{-1} \\
&= \underbrace{\left(dJ|_{x_j} \circ Q_0 \circ \nu_{x_j}^{-1} \circ B_{\mathfrak{b}^p} \right)^{-1}}_{=\text{id}_{T_{x_j} V_j}} \\
&= \left(\nu_{x_j}^{-1} \circ B_{\mathfrak{b}^p} \right)^{-1}
\end{aligned}$$

Notice how $\nu_{x_j}^{-1} \circ B_{\mathfrak{b}^p}$ induces a basis on $T_{x_j} V_j$ from a basis in $T_p^u M$. Comparing this basis to the basis $(\mathfrak{d}_i)_{i=2, \dots, \lambda_q}$ is what assigned the sign to the flow line in the Morse boundary

operator. By the calculations we know that the Jacobean of the function Z

$$dQ_0^{-1} \circ d(B_{\mathfrak{b}^p}^{-1} \circ \nu_{x_j} \circ J^{-1} \circ \rho \circ E_q^u \circ B_{\mathfrak{b}^q})|_y \circ Q_y = \begin{pmatrix} 0 & \tilde{Z} \end{pmatrix},$$

where $\tilde{Z}$ is a λ_{q-1} matrix with the sign of its determinant being given by the sign in the Morse boundary operator n_γ .

Now we can calculate the differential

$$\begin{aligned} & d(\pi_+ \circ s \circ r \circ \pi_+^{-1}(x_1, \dots, x_{\lambda_q}))|_x \\ &= d\left(\sin(\pi e^{\tau(E_q^u \circ B_{\mathfrak{b}^q}(x))}) \cdot \begin{pmatrix} -\cos(\pi \cdot \|Z(x)\|) \\ \sin(\pi \cdot \|Z(x)\|) \cdot \frac{Z(x)}{\|Z(x)\|} \end{pmatrix}\right)|_x \end{aligned}$$

To keep it readable we will first calculate

$$d\left(\sin(\pi e^{\tau(E_q^u \circ B_{\mathfrak{b}^q}(x))})\right)|_x = \cos(\pi \cdot e) \cdot \pi e \cdot d\tau|_{E_q^u \circ B_{\mathfrak{b}^q}(x)} = \underbrace{(\cos(\pi \cdot e) \cdot \pi e, 0, \dots, 0)}_{<0}$$

In this calculation we used that x was choosen such that $\tau \circ E_q^u \circ B_{\mathfrak{b}^q} = 1$ and $\mathfrak{b}^q$ was such that $dE_q^u|_y \circ Q_y(\mathfrak{b}_i^q)$ equals $-\text{grad}_y(f)$ for $i = 1$ and for $i > 1$ it equals $d\varphi_{\tau(y)}|_y^{-1}(w_i)$ where (w_i) denotes a basis of $T_{x_j}V_j$. Hence applying the product rule to the differential above yealds a sum with the first summand being

$$(-1, 0, \dots, 0) \cdot \underbrace{(\cos(\pi \cdot e) \cdot \pi e, 0, \dots, 0)}_{<0} = \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} \quad \text{where } a > 0.$$

Here we just used, that $Z(x) = 0$. The second summand is:

$$\underbrace{\sin(\pi \cdot e)}_{>0} \begin{pmatrix} \sin(0) \cdot * = 0 \\ \pi \cdot 1_{\lambda_q \times \lambda_q} \circ dZ|_x \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & b \cdot Z \end{pmatrix} \quad \text{where } b > 0).$$

Here we used that

$$d \frac{\sin(\pi \|x\|)}{\|x\|} \cdot x \Big|_{x \rightarrow 0} = \pi \cdot 1_{\lambda_q \times \lambda_q}$$

But with this we have the jacobian

$$d(\pi_+ \circ s \circ r \circ \pi_+^{-1}(x_1, \dots, x_{\lambda_q}))|_x = \begin{pmatrix} a & 0 \\ 0 & bZ \end{pmatrix},$$

and know that its determinant has a positive that equals n_γ which is the assigned sign to the flow line containing x_j in the Morse Boundary operator. Or said differently: The local and hence global degree of the map $r^* \circ s^*$ equals n_γ . With this we can calculate

$$\delta_{triple}^j(\alpha_p) = \sum_j \delta_{triple}^j \circ f_p^*(\beta) = n_\gamma f_q^*(\beta) = n_\gamma(\alpha_q)$$

Furthermore: $\Delta(\alpha_p) = \delta_{triple}^2(\alpha_p) = \sum_{\gamma \in \epsilon} n_\gamma(\alpha_q)$. This concludes the prove. $\square$

7 A Künneth Formular

Lemma 7.0.1. *Let M, f be a smooth manifold with a Morse-Smale function f and an adapted filtration $\{N_i\}$. Let furthermore X be a topological space such that $K^*(X)$ is free. Then the external Product*

$$K^*(M) \otimes K^*(X) \rightarrow K^*(M \times X)$$

is an isomorphism.

Proof. We start by considering the spectral sequence for the filtration $\{N_i\}$ on M and $\{N_i \times X\}$ on $M \times X$. Notice if we denote the pointed X by $X_+ := X \sqcup \{+\}$ we have

$$\begin{aligned} (N_i \times X / N_{i-1} \times X) &\simeq (N_i / N_{i-1}) \times X \\ &\simeq \bigvee_{p \in \text{Crit}_i} (D^p \times X / \partial D^p \times X) \\ &\simeq \bigvee_{p \in \text{Crit}_i} (D^p \times X_+ / \partial D^p \times X_+ \cup D^p \times \{+\}) \\ &\simeq \bigvee_{p \in \text{Crit}_i} \left((D^p / \partial D^p) \times X_+ / D^p \vee X_+ \right) \\ &= \bigvee_{p \in \text{Crit}_i} (S^p \wedge X_+) \end{aligned}$$

Since the homotopys here are natural, we have commuting differentials □

8 Examples

We begin with a minimal example that has non trivial higher differential:

Example 8.0.1 (The Sphere). If we take the three-sphere $S^3 \subseteq \mathbb{R}^4$ and denote the north pole with N and the south pole with S we get the Morse chain complex as

$$0 \longrightarrow \langle S \rangle_{\mathbb{Z}} \longrightarrow 0 \longrightarrow 0 \longrightarrow \langle N \rangle_{\mathbb{Z}} \longrightarrow 0$$

Then for the second page we get the Morse homology together with the 0 maps and on the third page we have:

$$\begin{array}{ccc} \langle S \rangle & \begin{array}{cc} 0 & 0 \end{array} & \langle N \rangle \\ & \searrow \quad \quad \nearrow & \\ & d_3^0 & \end{array}$$

Now we want to check what $d_3^0(S)$ is. For this we want to use its definition and hence we need to give the Morse filtration. For this we can use:

$$N_{-1} := \emptyset, \quad N_{0,1,2} := H_- \text{ the southern hemisphere}, \quad N_3 = S^3.$$

First we translate E_r into the language where the definition was formulated. We know that EM_3^p is naturally isomorphic to

$$E_3^{p,0} = \frac{\gamma^{-1}(\alpha^2 D^{p,-2})}{\beta(\ker(\alpha^2|_{D^{p+3-1,-2}}))}$$

Now for $r = 3$ and $p = 0$

$$\alpha^2 D^{2,-2} = \text{im}(K^0(N_2) \rightarrow K^0(N_0)) = \text{im}(K^0(H_-) \rightarrow K^0(H_-)) \cong \mathbb{Z}$$

Hence $E_3^{0,0} = \gamma^{-1}(\alpha^2 D^{2,-2})^{0,0} \cap E_0^{0,0} = K^0(N_0, \emptyset) \cong \mathbb{Z}$. And similar we conclude that $E_3^{3,0} = K^3(N_3, N_2) \cong \mathbb{Z}$. Now we analyze the map

$$d_3^{3,0} : K^3(N_3, N_2) \rightarrow K^0(N_0, \emptyset)$$

So far we were able to ignore all orientation, (which we formally need for the morse boundary operator). So let's fix the orientations of the unstable part by saying the unstable part of the north pole (i.e. the whole S^3) is the canonical orientation of $S^3 \subset \mathbb{R}^4$, whilst the unstable part of the south pole (i.e. the south pole) has the orientation -1 .

Now $d_r = \beta \circ \alpha^2 \circ \gamma$ and $\gamma : K^3(N_3, N_2) \rightarrow K^0(N_3)$ maps our generator induced by N to the standard generator of $K^3(S^3)$ induced by the chosen orientation, due to N_2 being a contractible subset. Now $\alpha^2 : K^3(N_3) \rightarrow K^3(N_1)$ is the zero map, since its range is 0 due to N_1 being homotopically equivalent to the point and 3 being odd. Finally $\beta : K^3(N_1) \rightarrow K^4(N_0, \emptyset)$ is the connecting homomorphism that is trivial here. Hence $d_3^{3,0}$ is the zero map. And thereby EM_4 which equals E_∞ is

$$\langle S \rangle \quad 0 \quad 0 \quad \langle N \rangle$$

Now how to we get information about our beloved K groups back? For this we remind ourself about the filtration

$$K^0(S^3) = F^{-1}K^0(S^3) \supseteq F^0K^0(S^3) \supseteq F^1K^0(S^3) \supseteq F^2K^0(S^3) \supseteq F^3K^0(S^3) = 0$$

Now using this filtration we have the short exact sequence

$$0 \longrightarrow F^0K^0(S^3) \longrightarrow K^0(S^3) \longrightarrow E_4^{0,0} = \frac{F^{-1}K^0(S^3)}{F^0K^0(S^3)} \longrightarrow 0$$

Hence, $K^0(S^3)$ is an extension of $F^0K^0(S^3)$ and $E_4^{0,0} \cong \mathbb{Z}$. Furthermore, we have the exact sequence

$$0 \longrightarrow F^1K^0(S^3) \longrightarrow F^0K^0(S^3) \longrightarrow E_4^{1,-1} = \frac{F^0K^0(S^3)}{F^1K^0(S^3)} = 0 \longrightarrow 0$$

Hence, $F^0K^0(S^3) \cong F^1K^0(S^3)$ and by the same argument we can continue this chain

$$F^0K^0(S^3) \cong F^1K^0(S^3) \cong F^2K^0(S^3) \cong F^3K^0(S^3) = 0$$

Hence, the first short exact sequence gives is an isomorphism between $K^0(S^3) \cong \mathbb{Z}$.

Now we want to calculate the K group of a non trivial example:

Example 8.0.2 (The Real Projective Space). Here, we define $\mathbb{RP}^n$ to be $\mathbb{R}^{n+1} \setminus 0$ modulo scalar multiples. Hence it is the space of lines in $\mathbb{R}^{n+1}$. Alternatively we can view it as the n -sphere where the operation from $\{\text{id}, -\text{id}\}$ is quotient out. From this description we get a metric, since $S^n \supseteq \mathbb{R}^{n+1}$ inherits the subspace metric in which all operations from $\{\text{id}, -\text{id}\}$ are isometries. Hence, this metric projects down to $\mathbb{RP}^n$. Now let

$$A = \text{diag}(\lambda_0, \lambda_1, \dots, \lambda_n)$$

be a symmetric matrix with real $\lambda_i \in \mathbb{R}$. Then we define the map

$$\Lambda : \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{R}; \Lambda(x) := \frac{\langle x, Ax \rangle}{\|x\|^2}.$$

Notice that this function descends to a well defined function $\tilde{\Lambda} : \mathbb{RP}^n \rightarrow \mathbb{R}$. On the whole $\mathbb{R}^{n+1}$ we can define the function

$$f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}; f(x) := \frac{1}{2} \langle x, Ax \rangle = \frac{1}{2} \sum_{i=0}^n \lambda_i x_i^2.$$

Now assume that $\varphi_t(x)$ is the flow associated to $-\text{grad}(f)$, meaning

$$\frac{d\varphi_t(x)}{dt} = -\text{grad}_{\varphi_t(x)}(f) = -A\varphi_t(x),$$

since

$$\frac{\partial f}{\partial x_i}(x) = \lambda_i x_i.$$

Now we have the lemma:

Lemma 8.0.3. *The flow lines of the negative gradient flow with respect to $\frac{1}{2}\tilde{\Lambda}$ on $\mathbb{RP}^n$ are the images under the quotient map*

$$\pi : \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{RP}^n$$

of the flow lines of the linear system

$$\frac{d\Phi_t(x)}{dt} = -A\Phi_t(x)$$

Proof. Let x be a point on the flow line $\Phi_t(x)$. Then $-\text{grad}(f)(x) = -Ax$. Now we split this into a part that is orthogonal to y and one that is parallel:

$$-\text{grad}(f)(x) = \underbrace{-\Lambda(x)x}_{\text{parallel to } x} + \underbrace{-Ax + \Lambda(x)x}_{\text{orthogonal to } x}.$$

The claim of orthogonality is just a calculation of the inner product $\langle -Ax + \Lambda(x)x, x \rangle = 0$ □

Now we radially project this onto the sphere via

$$\pi_r : \mathbb{R}^{n+1} \setminus \{0\} \rightarrow S^n; x \mapsto \frac{x}{\|x\|}$$

the differential of this projection kills the parallel part and hence reads:

$$d\pi_r|_x(-\text{grad}_x(f)) = \frac{-Ax + \Lambda(x)x}{\|x\|}$$

If we restrict $f|_{S^n}$ we get that the gradient $-\text{grad}_y(f|_{S^n}) = -Ay + \Lambda(y)y$. To see this, let $v \in T_y S^n \subseteq T_y \mathbb{R}^{n+1}$ and remember that the inner product on S^n is induced from the euclidean one. Then

$$\begin{aligned} \langle \text{grad}_y(f|_{S^n}), y \rangle_{S^n} &= d f|_{S^n}(y) = df(y) = \langle \text{grad}_y(f), y \rangle_{\mathbb{R}^{n+1}} \\ &= \langle \text{proj}_{T_y S^n}(\text{grad}_y(f)), y \rangle_{\mathbb{R}^{n+1}} = \langle Ay - \Lambda(y)y, y \rangle_{\mathbb{R}^{n+1}} \end{aligned}$$

Here $\text{proj}_{T_y S^n}$ denotes the projection onto the subspace $T_y S^n \subseteq T_y \mathbb{R}^{n+1}$. This however tells us that the image $\widetilde{\Phi_t(x)}$ of $\Phi_t(x)$ under the radial projection solves the equation

$$\frac{d\widetilde{\Phi_t(y)}}{dt} = -\text{grad}_y(f|_{S^n})(\widetilde{\Phi_t(y)}) = -\frac{1}{2}\text{grad}_y(\Lambda|_{S^n})(\widetilde{\Phi_t(y)}) \quad \text{since } f = \frac{1}{2}\Lambda$$

Hence descending to the quotient $\mathbb{RP}^n$ yields that $\left[\widetilde{\Phi_{2t}(y)}\right]$ is a solution of the flow associated to the negative gradient of $\tilde{\Lambda}$ in $\mathbb{RP}^n$. Now the other way round should be trivial: Take a solution $\psi_t(x)$ of $\tilde{\Lambda}$ in $\mathbb{RP}^n$. Then for fixed t, x let U be a neighbourhood

of $\psi_t(x)$. This U can be lifted to S^n and the solution there solves the lifted version of the flow induced from $\tilde{\Lambda}$. And from here we can argue that the solution is part of a solution of the linear differential equation

$$\frac{d\varphi_t(x)}{dt} = -\text{grad}_{\varphi_t(x)}(f) = -Ax.$$

This concludes the proof. Hence critical points in $\mathbb{RP}^n$ are eigenvalues of A . Hence, for the function to be Morse, we need the critical points to be isolated, meaning that all eigenspaces have dimension one. So we assume that the diagonal entries in A are all different. For simplicity we assume that $\lambda_0 < \lambda_1 < \dots < \lambda_n$. Since A is diagonal we know that $[e_i]$ are the critical points.

Lemma 8.0.4. *Each critical point $[e_i]$ is non degenerate, of index i and its unstable manifold is*

$$W([e_i] \rightarrow) =: U_i = \{[x_0 : x_1 : \dots : x_i : 0 : \dots : 0] \in \mathbb{RP}^n | x_i \neq 0\},$$

whilst the stable manifold is

$$W(\rightarrow [e_i]) =: S_i = \{[0 : \dots : 0 : x_i : x_{i+1} : \dots : x_n] \in \mathbb{RP}^n | x_i \neq 0\}$$

Proof. Let B^n be the open unit ball in $\mathbb{R}^n$ and $V_i := \{[x, \dots, x_n] \in \mathbb{RP}^n | x_i \neq 0\}$. Then we define the parametrization

$$\begin{aligned} \varphi_i : B^n &\rightarrow V_i \\ (y_1, \dots, y_n) &\mapsto [y_1, \dots, y_i, \sqrt{1 - \|y\|_2^2}, y_{i+1}, \dots, y_n]. \end{aligned}$$

Now the image gives representatives of norm one and hence in this chart we calculate:

$$\begin{aligned} \tilde{\Lambda} \circ \varphi_i(y_1, \dots, y_n) &= \sum_{j=1}^i y_j^2 \lambda_{j-1} + \lambda_i (1 - \|y\|_2^2) + \sum_{j=i+1}^n y_j^2 \lambda_j \\ &= \sum_{j=1}^i y_j^2 \lambda_{j-1} + \lambda_i (1 - \sum_{j=1}^n y_j^2) + \sum_{j=i+1}^n y_j^2 \lambda_j \\ &= \sum_{j=1}^i y_j^2 (\lambda_{j-1} - \lambda_i) + \lambda_i + \sum_{j=i+1}^n y_j^2 (\lambda_j - \lambda_i) \end{aligned}$$

With this local description we can calculate the hessian in $[e_i] = \varphi_i(0)$:

$$\left. \frac{\partial^2 \tilde{\Lambda} \circ \varphi_i}{\partial y_k \partial y_l} \right|_0 = \begin{cases} 0 & \text{if } k \neq l \\ 2(\lambda_{k-1} - \lambda_i) < 0 & \text{if } k = l \leq i \\ 2(\lambda_k - \lambda_i) > 0 & \text{if } k = l > i \end{cases}$$

Hence, $[e_j]$ is a non degenerate critical point of index i . Now concerning the unstable manifolds, we pick a point $x = x_k e_k + \dots x_l e_l \in \mathbb{R}^{n+1}$ with x_k and x_l not equal to zero. This points trajectory in $\mathbb{R}^{n+1}$ under $-\text{grad}(f)$ is

$$\varphi_t(x) = x_k e^{-\lambda_k t} e_k + \dots x_l e^{-\lambda_l t} e_l.$$

Hence for t approaching negative infinity, the term $x_l e^{-\lambda_l t} e_l$ is leading, whilst for t approaching infinity, the summand $x_k e^{-\lambda_k t} e_k$ is leading. Since projectively we don't care about absolute values but rather relations having a leading coordinate meaning that relatively all other coordinates approach zero. Hence:

$$\lim_{t \rightarrow -\infty} ([\varphi_t(x)]) = [e_l] \quad \text{and,} \quad \lim_{t \rightarrow +\infty} ([\varphi_t(x)]) = [e_k].$$

□

Definition 8.0.5 (Orientations). For orienting the unstable manifolds we make use of the projection $\mathbb{R}^{n+1} \supseteq S^n \rightarrow \mathbb{RP}^n$. Hence, we can lift $T_{[e_i]}\mathbb{RP}^n$ to $T_{\pm e_i}S^n$ and include that into $T_{\pm e_i}\mathbb{R}^{n+1}$, where we have the maps $Q_{\pm e_i} : \mathbb{R}^{n+1} \rightarrow T_{\pm e_i}\mathbb{R}^{n+1}$. In this way we get two canonical lifts of the tangent space:

$$\begin{array}{ccc} T_{[e_i]}\mathbb{RP}^n & \xrightarrow{d\pi_{e_i}^{-1}} & T_{e_i}S^n \\ \downarrow d\pi_{-e_i}^{-1} & & \downarrow Q_{e_i}^{-1} \\ T_{-e_i}S^n & \xrightarrow{Q_{-e_i}} & \langle (e_j)_{j \neq i} \rangle \end{array}$$

In general, the diagram is not commutative. To see this notice how $\pi \circ -\text{id} = \pi$. Hence $d\pi|_{-e_i} \circ d - \text{id}|_{e_i} = d\pi|_{e_i}$ and with that:

$$\begin{aligned} Q_{-e_i}^{-1} \circ d\pi|_{-e_i}^{-1} \circ d\pi|_{e_i} \circ Q_{e_i} &= Q_{-e_i}^{-1} \circ d\pi|_{-e_i}^{-1} \circ d\pi|_{-e_i} \circ d(-\text{id})|_{e_i} \circ Q_{e_i} \\ &= Q_{-e_i}^{-1} \circ d\pi|_{-e_i}^{-1} \circ d\pi|_{-e_i} \circ Q_{-e_i} \circ -\text{id} \\ &= Q_{-e_i}^{-1} \circ Q_{-e_i} - \text{id} = -\text{id}. \end{aligned}$$

Hence, we can define a canonical orientation on $T_{[e_i]}\mathbb{RP}^n$ that is given by $(e_j)_{j \neq i}$ or equivalent $(-e_j)_{j \neq i}$ depending on the chosen lift. Now in the same way we can induce orientations to the unstable manifolds U_i by two equivalent ways:

$$\begin{array}{ccc} T_{[e_i]}U_i & \hookrightarrow & T_{[e_i]}\mathbb{RP}^n \\ \uparrow & & \uparrow \\ \langle (e_0, \dots, e_{i-1}) \rangle & \hookrightarrow & T_{e_i}S^n \end{array} \quad \text{or equivalently} \quad \begin{array}{ccc} T_{[e_i]}U_i & \hookrightarrow & T_{[e_i]}\mathbb{RP}^n \\ \uparrow & & \uparrow \\ \langle -e_0, \dots, -e_{i-1} \rangle & \hookrightarrow & T_{-e_i}S^n \end{array}$$

Now we know how the groups in the Morse chain complex are looking. The next step is the boundary operator:

Lemma 8.0.6. *The differential in $C^*(\mathbb{RP}^n, \Lambda^*)$ with the right orientations act as*

$$\partial_i([e_i]) = n([e_{i-1}], e_i)[e_{i+1}] = ((-1)^i - 1)[e_{i+1}]$$

Proof. First we notice that $W([e_i] \rightarrow [e_{i-1}])$ is a one dimensional projective space with two points missing. If we would look at the Morse homology with coefficients in $\mathbb{Z}_2$ we would be done here, since $\mathbb{RP}^1 \setminus \{p, q\}$ consists of two lines. Hence, over $\mathbb{Z}_2$ the boundary operator is trivial. However we want to use the orientations from above. Now let

$$x \in W([x_i] \rightarrow [e_{i-1}]) = \{[0 : \dots : 0 : x_{i-1} : x_i : 0 : \dots : 0] \mid x_{i-1} \neq 0 \text{ and } x_i \neq 0\}.$$

Then we can choose representatives, such that $x_{i-1}^2 + x_i^2 = 1$, and (x_{i-1}, x_i) is in the upper half plane meaning that $x_i > 0$. Now take a $x \in W([e_i] \rightarrow [e_{i-1}])$. Then $T_x W([e_i] \rightarrow [e_{i-1}])$ has an oriented basis given as the pushforward of $v = x_i e_{i-1} - x_{i-1} e_i$ under the projection $T_x S^n \rightarrow T_x \mathbb{RP}^n$. Now we want to analyze how this relates to the orientation induced by the flow and the two unstable orientations: if both x_i and x_{i-1} are positive, then v points in the direction of the flow or said different, $-\text{grad}_x(f) = v$. Else, if x_{i-1} is negative, $-v$ points in the direction of the flow (i.e. $-\text{grad}_x(f) = -v$). The parallel transport of $T_{[e_{i-1}]} U_{i-1}$ to the normal space of $-\text{grad}_x(\tilde{\Lambda})$ induced by the negative flow maps the basis $(e_0, \dots, e_{i-2})$ of $T_{[e_{i-1}]} U_i$ to $(e_0, \dots, e_{i-2})$ in the orthogonal complement of $-\text{grad}_x(\tilde{\Lambda})$ in $T_x \mathbb{RP}^n$. So we need to compare in $T_x \mathbb{RP}^n$ the basis $(-\text{grad}_x(f), e_0, \dots, e_{i-2})$ to the basis $(e_0, \dots, e_{i-1})$ in $T_{[e_i]} U_i$ after transporting it to $T_x \mathbb{RP}^n$ along the flow. This latter transport maps the basis vectors as follows:

$$e_j \mapsto \begin{cases} x_i e_{i-1} - x_{i-1} e_i & \text{if } j = i-1 \\ e_j & \text{else.} \end{cases}$$

In other words, we need to ask when the two basis

$$((x_i e_{i-1} - x_{i-1} e_i), e_0, \dots, e_{i-2}) \quad \text{and} \quad (e_0, \dots, e_{i-2}, x_i e_{i-1} - x_{i-1} e_i),$$

if $x_{i-1} > 0$ are comparable. Then, the basis carry the same orientation, if and only if i is odd, since then it takes evenly many transpositions of coordinates.

If $x_{i-1} < 0$ we notice, that our representation of $[e_{i-1}]$ is given by $-e_{i-1}$ and hence the unstable tangent space has the basis $-e_0, \dots, -e_{i-2}$. Hence in x we get the oriented basis

$$(-\text{grad}_x(f), -e_0, \dots, -e_{i-2}) = (-v, -e_0, \dots, -e_{i-2}).$$

We compare this basis to basis that comes from transporting the basis $(e_0, \dots, e_{i-1})$ of $T_{[e_i]} U_{[e_i]}$ to x resulting in the basis $(e_0, \dots, e_{i-2}, v)$. Hence we compare the bases

$$(-v, -e_0, \dots, -e_{i-2}) \quad \text{and} \quad (e_0, \dots, e_{i-2}, v)$$

we manipulate the first and keep note of the sign of the alterations:

$$(-v, -e_0, \dots, -e_{i-2}) = (-1)^i (v, e_0, \dots, e_{i-2}) = (-1)^{i+i-1} (e_0, \dots, e_{i-2}, v).$$

Hence, the orientations never agree. Now if i is odd the two flow lines cancel out when counted with the sign and if i is even, we count minus two flow lines. Therefore the counting of flow lines noted by $n(p, q)$ gives:

$$n(e_{i+1}, e_i) = (-1)^i - 1,$$

and the Morse cohomology boundary operator equals:

$$\partial_i([e_i]) = n([e_{i-1}], e_i)[e_{i+1}] = ((-1)^i - 1)[e_{i+1}]$$

□

Now for even n , EM_1 of the form:

$$0 \quad \langle [e_0] \rangle \xrightarrow{0} \langle [e_1] \rangle \xrightarrow{-2} \langle [e_2] \rangle \xrightarrow{0} \dots \xrightarrow{-2} \langle [e_n] \rangle \xrightarrow{0} 0$$

and for odd n :

$$0 \quad \langle [e_0] \rangle \xrightarrow{0} \langle [e_1] \rangle \xrightarrow{-2} \langle [e_2] \rangle \xrightarrow{0} \dots \xrightarrow{0} \langle [e_n] \rangle \xrightarrow{0} 0$$

Hence we have the EM_2 pages as follows:

$$0 \quad \langle [e_0] \rangle \quad 0 \quad \frac{\langle [e_2] \rangle}{\langle -2[e_2] \rangle} \quad 0 \quad \dots \quad \frac{\langle [e_n] \rangle}{\langle -2[e_n] \rangle} \quad 0$$

if n is even, and in the odd case:

$$0 \quad \langle [e_0] \rangle \quad 0 \quad \frac{\langle [e_2] \rangle}{\langle -2[e_2] \rangle} \quad 0 \quad \dots \quad \langle [e_n] \rangle \quad 0$$

When it comes to higher differentials, we notice the following: Only for odd r the differentials are possible nontrivial Hence for $\mathbb{RP}^2$ we are done and we can get the complex k-groups: For $K^0(\mathbb{RP}^2)$ we get a associated graded group that is isomorphic to $\mathbb{Z} \oplus \mathbb{Z}_2$. Hence we need to calculate the extension

$$0 \longrightarrow \mathbb{Z} \longrightarrow K^0(\mathbb{RP}^2) \longrightarrow \mathbb{Z}_2 \longrightarrow 0$$

giving us $K^0(\mathbb{RP}^2) \cong \mathbb{Z} \oplus \mathbb{Z}_2$. And similar $K^1(\mathbb{RP}^2) = 0$. To understand the higher differentials we need to get specific about the used filtration. We define the filtration inductively as follows: For any i define $c_i \in (\lambda_i, \lambda_{i+1})$ and with this we define the filtration

$$N_{-1} := \emptyset, \quad N_i = \tilde{\Lambda}(-\infty, c_i) \text{ the sublevel sets, } \quad N_n := \mathbb{RP}^n.$$

With this filtration it is somewhat easy to see that (N_i, N_{i-1}) gives a regular index pair of $[e_i]$. One just needs to go through the definitions: First it is clear, that the only critical point inside of the closure of $N_i \setminus N_{i-1}$ is $[e_i]$ and it is in the interior. Secondly, the positiv lime invariance is clear, since $\tilde{\Lambda}$ decreases along flow lines. And N_{i-1} is an exit set also by definition. Finally the cofibration property of the pair N_i, N_{i-1} is to be shown. For this we notice, that c_i is a regular value of R . Hence there exists an $\varepsilon > 0$, such that $f^{-1}([a, a + \varepsilon))$ contains no critical point and hence this defines an open color of N_i inside N_{i+1} . This open color can be shown to be an NDR pair, which in turn is a cofibration. Those are standart arguments seen in [Whi78] Chapter 5.1.

Lemma 8.0.7. *In the filtration, each N_i is homotopically equivalent to $\mathbb{RP}^i$.*

Proof. Notice how for all i the sublevelset $\tilde{\Lambda}^{-1}(-\infty, \lambda_i]$ is an embedded projective space $\mathbb{RP}^i$. To show that N_i is homotopically equivalent we start by noticing that $N_i \simeq \tilde{\Lambda}^{-1}(-\infty, \lambda_{i-1} + \varepsilon)$ for all $\varepsilon > 0$ small enough. Now the next step is to get rid of epsilon. Let L_ε denote $\tilde{\Lambda}^{-1}(-\infty, \lambda_{i-1} + \varepsilon)$ and hence we want to show that $L_\varepsilon \simeq L_0$. For this we define

$$H : L_\varepsilon \times [0, 1] \rightarrow L_\varepsilon \quad ([x], t) \text{ with } \|x\| = 1 \mapsto \left[\sum_{j=1}^i x_j e_j + (1-t) \sum_{j>i} x_j e_j \right]$$

First we need to check if this map is well defined in two senses. Is the definition invariant under the chosen representative and do we map into L_ε . For the independence of the chosen representative we notice that the image of x and $-x$ differ by sign and hence define the same equivalence class. Also applying $\tilde{\Lambda}$ to the image we notice that the image is less than $\lambda_i + \varepsilon$. this is true, since $\tilde{\Lambda}$ is a weighted sum of the eigenvalues and the addition of $(1-t)$ decreases the weights on the bigger eigenvalues reducing the overall weighted sum.

Now obviously H_1 maps to $\tilde{\Lambda}^{-1}(-\infty, \lambda_i]$. □

Now that we have our filtration we can start looking at higher differentials. If n is even, then every other group in EM_2 is zero hence all higher differentials vanish as only odd d_r are candidates to be non trivial. So let n be odd. Then only differentials landing in the last group EM_n can possibly be non trivial. However, for all $r < n$ the differential that lands in $EM_r^m \cong \mathbb{Z}$ starts in $EM_r^{m-r} \cong \mathbb{Z}_2$. Hence they are all trivial as well. Now if $r = m$ we have the differential

$$d_m^0 : EM_m^0 \rightarrow EM_m^m$$

which is possibly non-zero. However this differential is given as follows:

$$\begin{array}{ccccc} & & & & K^0(N_m) \\ & & & & \downarrow i^* \\ K^0(N_0, \emptyset) & \xrightarrow{\gamma} & K^0(N_0) & \xrightarrow{\widetilde{\alpha^{-m}}} & K^0(N_{m-1}) \\ & \searrow d_m^0 & & & \downarrow \delta \\ & & & & K^1(N_m, N_{m-1}) \end{array}$$

Here α^{-m} is any left inverse of α^m or taking the preimage (the well definedness or rather independence of the chosen inverse was shown in the definition of the derived exact couple). The vertical sequence is exact and since $N_{m-1} = N_m \setminus D$ where D is contractible, we have that i^* is surjective and hence δ is trivial and thereby $dM_m^0 = 0$. Then for this we start with $n = 3$ and then go up. In the case of $n = 3$ we are trivial, since d_3^0 is given by the composition

in formeln
nachrechnen

d_3^0 ist
sowieso
0, da das
Bild 0 ist!!

$$\begin{array}{ccccc}
& & & K^0(N_3) & \\
& & & \downarrow i^* & \\
K^0(N_0, \emptyset) & \xrightarrow{\gamma} & K^0(N_0) & \xrightarrow{\widetilde{\alpha^{-2}}} & K^0(N_2) \\
& \searrow d_3 & & \downarrow \delta & \\
& & & K^1(N_3, N_2) &
\end{array}$$

Here α^{-1} is any left inverse of α^2 or taking the preimage (the well definedness or rather independence of the choosen inverse was shown in the definition of the derived exact couple). The vertical sequence is exact and since $N_2 = N_3 \setminus D$ where D is contractible. A contraction is given by flowing along the positive gradient of $\tilde{\Lambda}$. Hence, we have that i^* is surjective and thereby δ is trivial resulting in $dM_m^0 = 0$. For all real projective spaces the spectral sequence becomes stable on the second page and hence we can calculate the complex K-groups of $\mathbb{RP}^n$: For even n we have the two graded associated groups:

$$\begin{aligned}
\text{gr}(K^0(\mathbb{RP}^n)) &= \mathbb{Z} \oplus \underbrace{\mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2}_{\frac{n}{2} \text{ summands}} \\
\text{gr}(K^1(\mathbb{RP}^n)) &= 0.
\end{aligned}$$

For odd n we get

$$\begin{aligned}
\text{gr}(K^0(\mathbb{RP}^n)) &= \mathbb{Z} \oplus \underbrace{\mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2}_{\frac{n-1}{2} \text{ summands}} \\
\text{gr}(K^1(\mathbb{RP}^n)) &= \mathbb{Z}.
\end{aligned}$$

Hence for any n we have

$$K^0(\mathbb{RP}^n) \cong \mathbb{Z} \oplus G \text{ with } |G| = 2^{\lfloor \frac{n}{2} \rfloor}.$$

Fact 8.0.8. *In fact $K^0(\mathbb{RP}^n) \cong \mathbb{Z} \oplus \mathbb{Z}_{2^{\lfloor \frac{n}{2} \rfloor}}$.*

Example 8.0.9. Make a produkt morse funktion on $\mathbb{RP}^2 \times \mathbb{RP}^4$!. For this we assume that $\tilde{\Lambda}_2$ is a Morse function on $\mathbb{RP}^2$ and defined as above and similary $\tilde{\Lambda}_4$ is one for $\mathbb{RP}^4$. From the established Künneth formular we know that the Morse cohomology of $\mathbb{RP}^2 \times \mathbb{RP}^4$ is as follows:

$$H^n(\mathbb{RP}^2 \times \mathbb{RP}^4, \tilde{\Lambda}_2 + \tilde{\Lambda}_4) \cong \bigoplus_i (H^i(\mathbb{RP}^2, \tilde{\Lambda}_2) \otimes H^{n-i}(\mathbb{RP}^4, \tilde{\Lambda}_4)) \oplus \bigoplus_i \text{Tor}(H^i(\mathbb{RP}^2, \tilde{\Lambda}_2), H^{n-i+1}(\mathbb{RP}^4, \tilde{\Lambda}_4)).$$

First we list the Homology of $\mathbb{RP}^2$ and $\mathbb{RP}^4$:

	$\mathbb{RP}^2$	$\mathbb{RP}^4$
H^0	$\mathbb{Z}$	$\mathbb{Z}$
H^1	0	0
H^2	$\mathbb{Z}_2$	$\mathbb{Z}_2$
H^3	0	0
H^4	0	$\mathbb{Z}_2$

Furthermore the only non-zero Tor is $\text{Tor}(\mathbb{Z}_2, \mathbb{Z}_2)$. Which only appears for $n = 3$ and $n = 5$. Furthermore we have for the tensor groups for any A : $\mathbb{Z} \otimes A \cong A$ and $\mathbb{Z}_2 \otimes \mathbb{Z}_2 \cong \mathbb{Z}_2$. Hence we have the resulting isomorphisms:

$$H^i(\mathbb{RP}^2 \times \mathbb{RP}^4) \cong \begin{cases} \mathbb{Z} & \text{if } i = 0, \\ 0 & \text{if } i = 1, \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & \text{if } i = 2, \\ \mathbb{Z}_2 & \text{if } i = 3, \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & \text{if } i = 4, \\ \mathbb{Z}_2 & \text{if } i = 5, \\ \mathbb{Z}_2 & \text{if } i = 6, \\ 0 & \text{if } i > 6. \end{cases}$$

Hence the E_3 page looks with the only possible nontrivial differentials looks like

$$\begin{array}{ccccccccccc} & & & & & d_3^2 & & d_3^3 & & & \\ & & & & & \curvearrowright & & \curvearrowright & & & \\ 0 & \mathbb{Z} & 0 & \mathbb{Z}_2 \oplus \mathbb{Z}_2 & \mathbb{Z}_2 & \mathbb{Z}_2 \oplus \mathbb{Z}_2 & \mathbb{Z}_2 & \mathbb{Z}_2 & 0 \end{array}$$

Now the K theory of the product space $\mathbb{RP}^2 \times \mathbb{RP}^4$ is known and can be calculated for the K groups of the factors via the Künneth formular for K-theory that gives us two short exact sequence

$$0 \longrightarrow K^0(X) \otimes K^0(Y) \longrightarrow K^0(X \times Y) \longrightarrow \text{Tor}(K^1(X), K^1(Y)) \longrightarrow 0$$

$$0 \longrightarrow K^1(X) \otimes K^1(Y) \longrightarrow K^1(X \times Y) \longrightarrow \text{Tor}(K^0(X), K^0(Y)) \longrightarrow 0$$

In our case the two sequences lool like

$$0 \longrightarrow (\mathbb{Z} \oplus \mathbb{Z}_2) \otimes (\mathbb{Z} \oplus \mathbb{Z}_4) \longrightarrow K^0(\mathbb{RP}^2 \times \mathbb{RP}^4) \longrightarrow \text{Tor}(0, 0) \longrightarrow 0$$

$$0 \longrightarrow 0 \longrightarrow K^1(\mathbb{RP}^2 \times \mathbb{RP}^4) \longrightarrow \text{Tor}(\mathbb{Z}_2, \mathbb{Z}_4) \longrightarrow 0$$

And thereby

$$K^0(\mathbb{RP}^2 \times \mathbb{RP}^4) \cong \mathbb{Z} \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_4 \text{ and } K^1(\mathbb{RP}^2 \times \mathbb{RP}^4) \cong \mathbb{Z}_2$$

Now the order of $\mathbb{P}^4) \cong \mathbb{Z} \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_4$ and $K^1(\mathbb{RP}^2 \times \mathbb{RP}^4)$ is two. Hence assuming all higher differentials being 0 and thereby having the E_3 page as the last page this results in a contradiction since the sum over all odd entries of EM_3 gives a group of order four and the order stays invariant under group extensions.

References

- [AD14] Michèle Audin and Mihai Damian. *Morse Theory and Floer Homology*. Trans. by Reinie Ern . Universitext. London: Springer, 2014. ISBN: 978-1-4471-5495-2. DOI: 10.1007/978-1-4471-5496-9.
- [BH04] A. Banyaga and D. Hurtubise. *Lectures on Morse Homology*. Texts in the Mathematical Sciences. Springer Netherlands, 2004. ISBN: 9781402026959. URL: https://books.google.de/books?id=AX-_sbMjOK4C.
- [Hat02] Allen Hatcher. *Algebraic Topology*. Cambridge: Cambridge University Press, 2002. ISBN: 978-0-521-79540-1.
- [May99] J. P. May. *A Concise Course in Algebraic Topology*. Chicago Lectures in Mathematics. University of Chicago Press, 1999. ISBN: 978-0-226-51183-2. URL: <https://math.uchicago.edu/~may/CONCISE/ConciseRevised.pdf>.
- [Per01] Lawrence Perko. *Differential Equations and Dynamical Systems*. 3rd ed. Vol. 7. Texts in Applied Mathematics. Softcover / Paperback edition. New York / Berlin: Springer, 2001, p. 557. ISBN: 978-1461265269.
- [Sal85] Dietmar Salamon. “Connected simple systems and the Conley index of isolated invariant sets”. In: *Transactions of the American Mathematical Society* 291.1 (1985), pp. 1–41. DOI: 10.1090/S0002-9947-1985-0797044-3.
- [Sal90] Dietmar A. Salamon. “Morse theory, the Conley index and Floer homology”. In: *Bull. London Math. Soc.* 22.2 (1990). Bulletin of the London Mathematical Society, 22(2):113–140 (1990), pp. 113–140. DOI: 10.1112/blms/22.2.113.
- [Whi78] George W. Whitehead. *Elements of Homotopy Theory*. Vol. 61. Graduate Texts in Mathematics. New York–Berlin: Springer-Verlag, 1978, pp. xxi+744. ISBN: 0-387-90336-4. DOI: 10.1007/978-1-4612-6318-0.