

Product Information: Similac® for Diarrhea

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® for Diarrhea

Soy formula for dietary management of diarrhea

- 1 A 20 Cal/fl oz formula for dietary management of diarrhea to help firm loose and watery stools in infants older than 6 months and toddlers.
- 1 First and only nutritionally complete infant formula to contain added dietary fiber (soy) specifically for dietary management of diarrhea (6 g/L).
- 1 Clinically shown to reduce the duration of loose, watery stools during mild to severe diarrhea in older infants.^{1,2}
- 1 Clinically shown to be effective in the management of antibiotic-induced diarrhea.³
- 1 A unique blend of two carbohydrates using two absorptive pathways to help maximize absorption and minimize malabsorption risks.
- 1 Low osmolality (240 mOsm/kg water) to reduce the risk of osmotic diarrhea.
- 1 Halal.
- 1 Kosher.
- 1 Gluten-free.
- 1 Lactose-free.



Safety Precautions

- 1 **Never use a microwave oven to warm formula.** Serious burns can result.
- 1 Similac Expert Care for Diarrhea should not be fed to infants and toddlers with constipation.

¹ Vanderhoof JA, et al. Clin Pediatr 1997;36(3):135-139.
² Brown KH, et al. Pediatrics 1993;92(2):241-247.
³ Burks AW, et al. J Pediatr 2001;139(4):578-582.

Ingredients

Liquid Unflavored:

Water, Corn Syrup, Sugar, Soy Oil, Soy Protein Isolate, Coconut Oil, Soy Fiber. Less than 0.5% of: Calcium Citrate, Calcium Phosphate, Potassium Phosphate, Taurine, Potassium Citrate, m-Inositol, Magnesium Chloride, Monoglycerides, Soy Lecithin, Salt, Ascorbic Acid, Potassium Chloride, Zinc Sulfate, Carrageenan, L-Methionine, Niacinamide, Ferrous Sulfate, Calcium Pantothenate, Potassium Hydroxide, Choline Chloride, Cupric Sulfate, L-Carnitine, d-Alpha-Tocopheryl Acetate, Thiamine Chloride Hydrochloride, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, and Citric Acid.
Allergens: Contains soy ingredients.

Availability

List Number	Item
51276	Similac for Diarrhea Ready to Feed / 8-fl-oz (237-mL) can / 24 ct
57768	Similac for Diarrhea Ready to Feed / 1-QT (946-mL) bottle / 6 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac® for Diarrhea

Soy formula for dietary management of diarrhea

Nutrition Information - Liquid Unflavored

	100 Cal (5 fl oz)		1000 mL (1.1 QT)	
	Value	%RDI / %DV	Value	%RDI / %DV
Calories	100		676	
Volume, mL	148		1000	
Protein, g	2.66		17.99	
Fat, g	5.46		36.93	
Carbohydrate, g	10.1		68.3	
Water, g	133		899	
Linoleic Acid, mg	1300		8792	
Potential Renal Solute Load, mOsm	24.0		162.6	
Vitamin A, IU	300		2029	
Vitamin D, IU	60		406	
Vitamin E, IU	1.5		10.1	
Vitamin K, mcg	11		74	
Thiamin (Vitamin B1), mcg	60		406	
Riboflavin (Vitamin B2), mcg	90		609	
Vitamin B6, mcg	60		406	
Vitamin B12, mcg	0.45		3.04	
Niacin, mcg	1350		9130	
Folic Acid, mcg	15		101	
Pantothenic Acid, mcg	750		5072	
Biotin, mcg	4.5		30.4	
Vitamin C, mg	9		61	
Choline, mg	12		81	
Inositol, mg	5		33.8	
Calcium, mg	105		710	
Calcium, mEq	5.2		35.4	
Phosphorus, mg	75		507	
Magnesium, mg	7.5		50.7	
Iron, mg	1.8		12.2	
Zinc, mg	0.75		5.07	
Manganese, mcg	25		169	
Copper, mcg	75		507	
Iodine, mcg	15		101	
Selenium, mcg	2		14	
Sodium, mg	44		298	
Sodium, mEq	1.9		12.9	
Potassium, mg	108		730	
Potassium, mEq	2.8		18.7	
Chloride, mg	62		419	
Chloride, mEq	1.8		11.8	

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Preparation

Ready-to-Feed

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

8-fl-oz can

- 1 **Do not add water.**
- 1 Rinse can lid; shake very well.
- 1 Lift and pull tab to open.
- 1 Pour formula into bottle or cup; attach nipple or lid; once feeding begins, use within 1 hour or discard.

1-QT bottle

- 1 **Do not add water.**
- 1 **Do not use if band around cap or inner foil seal is damaged.**
- 1 Shake very well before each use.
- 1 Remove protective band; twist off and clean cap.
- 1 Invert cap; press down to pierce foil, then turn cap a half turn.
- 1 Remove foil.
- 1 Pour formula into bottle or cup; attach nipple or lid.
- 1 Once feeding begins, use within 1 hour or discard.

Storage & Handling

Ready-to-Feed

8-fl-oz can

- 1 Cover opened cans and prepared bottles not used immediately.
- 1 Store in refrigerator and **use within 48 hours.**
- 1 Store unopened cans at room temperature; avoid extreme temperatures.

1-QT bottle

- 1 Once opened, store quart bottle in refrigerator.
- 1 Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- 1 Store unopened containers at room temperature; avoid extreme temperatures.
- 1 Prepare and feed all formula from this container **within 3 days after opening.**
- 1 Do not reuse container.