

For the following tests list the number of ratio scale explanatory variables, the number of nominal scale explanatory variables (factors), and the number of interaction terms.

Write a GLM with df below each term.

Ratio	Factors	Interaction	
1. Oneway ANOVA comparing hematocrit in 3 treated groups and one control group. N = 10 in each group.	___0___	___1___	___0___

$$df_{total} = (10 \times 4) - 1$$

$$39 = 1 + 38$$

2. Twoway ANOVA for BACI design (before / after at control and impacted sites, in environmental assessment). N = 4 measurements at control before impact, 4 at impacted before impact, then after impact 4 more at control and 4 more at impact site.	___0___	___2___	___1___
--	---------	---------	---------

Y = response variable

$$df_{total} = (4 \times 4) - 1$$

$$15 = 1 + 1 + 1 + 12$$

3. Paired comparison of reaction times in 30 subjects, before and after alcohol intake.	___0___	___2___	___0___
---	---------	---------	---------

$$df_{total} = (2 \times 30) - 1$$

$$59 = 1 + 29 + 29 \quad [\text{Subject} \times \text{BeforeAfter interaction term cannot be estimated}]$$

___1___	___0___	___0___
---------	---------	---------

$$df_{total} = 30 - 1$$

$$29 = 1 + 28 \quad [\text{Regress after on before: valid. Not a paired comparison test}]$$

4. Carpal tunnel symptom severity with and without surgery, in 4 different hospitals (ntotal = 32)	___0___	___2___	___1___
--	---------	---------	---------

$$df_{total} = 32 - 1$$

$$31 = 1 + 3 + 3 + 24$$

5. Regression analysis of growth rates in 25 babies as a function of birth weight	___1___	___0___	___0___
---	---------	---------	---------

$$df_{total} = 25 - 1$$

$$24 = 1 + 23$$