

The due date for this assignment is past. Your work can be viewed below, but no changes can be made.


Important! Before you view the answer key, decide whether or not you plan to request an extension. Your Instructor may *not* grant you an extension if you have viewed the answer key. Automatic extensions are not granted if you have viewed the answer key.

[Request Extension](#)

1. 2.5/2.5 points | [Previous Answers](#)SCalcET8 4.1.003.

For each of the numbers a , b , c , d , r , and s , state whether the function whose graph is shown has an absolute maximum or minimum, a local maximum or minimum, or neither a maximum nor a minimum. (Enter your answers as a comma-separated list.)

absolute maximum	\$\$\$ <input type="text" value="s"/>
absolute minimum	\$\$\$ <input type="text" value="r"/>
local maximum	\$\$\$ <input type="text" value="c"/>
local minimum	\$\$\$ <input type="text" value="b, r"/>
neither a maximum nor a minimum	\$\$\$ <input type="text" value="a, d"/>




Solution or Explanation

Absolute maximum at s , absolute minimum at r , local maximum at c , local minima at b and r , neither a maximum nor a minimum at a and d .

2. 2/2 points | [Previous Answers](#)SCalcET8 4.1.005.

Use the graph to state the absolute and local maximum and minimum values of the function. (Assume each point lies on the gridlines. Enter your answers as a comma-separated list. If an answer does not exist, enter DNE.)



absolute maximum value

absolute minimum value

local maximum value(s)

local minimum value(s)





Solution or Explanation

Absolute maximum value is $f(4) = 5$; there is no absolute minimum value; local maximum values are $f(4) = 5$ and $f(6) = 4$; local minimum values are $f(2) = 2$ and $f(1) = f(5) = 3$.

3. 1/1 points | [Previous Answers](#)SCalcET8 4.1.007.


Sketch the graph of a function f that is continuous on $[1, 5]$ and has the given properties.

Absolute maximum at 5, absolute minimum at 2, local maximum at 3, local minima at 2 and 4







Solution or Explanation

Absolute maximum at 5, absolute minimum at 2, local maximum at 3, local minima at 2 and 4







4. 1.5/1.5 points | [Previous Answers](#)SCalcET8 4.1.011.



(a) Sketch the graph of a function that has a local maximum at 3 and is differentiable at 3.





(b) Sketch the graph of a function that has a local maximum at 3 and is continuous but not differentiable at 3.




(c) Sketch the graph of a function that has a local maximum at 3 and is not continuous at 3.







Solution or Explanation



(b)



(c)



5. 2/2 points | [Previous Answers](#)SCalcET8 4.1.017.


Sketch the graph of f by hand and use your sketch to find the absolute and local maximum and minimum values of f . (Enter your answers as a comma-separated list. If an answer does not exist, enter DNE.)

$$f(x) = \frac{2}{x}, \quad x \geq 2$$

absolute maximum value	<div> <div>1</div> </div>
absolute minimum value	<div> <div>DNE</div> </div>
local maximum value(s)	<div> <div>DNE</div> </div>
local minimum value(s)	<div> <div>DNE</div> </div>

Solution or Explanation

$f(x) = \frac{2}{x}$, $x \geq 2$. Absolute maximum $f(2) = 1$; no local maximum. No absolute or local minimum.



6. 2/2 points | [Previous Answers](#)SCalcET8 4.1.022.

Sketch the graph of f by hand and use your sketch to find the absolute and local maximum and minimum values of f . (Enter your answers as a comma-separated list. If an answer does not exist, enter DNE.)

$$f(t) = 9 \cos(t), \quad -3\pi/2 \leq t \leq 3\pi/2$$

absolute maximum value

9

✓ 9

absolute minimum value

-9

✓ -9

local maximum value(s)

9

✓ 9

local minimum value(s)

-9

✓ -9

Solution or Explanation

[Click to View Solution](#)

7. 1/1 points | [Previous Answers](#)SCalcET8 4.1.030.

Find the critical numbers of the function. (Enter your answers as a comma-separated list. If an answer does not exist, enter DNE.)

$$f(x) = x^3 + 12x^2 - 27x$$

$x =$

-9, 1

1, -9

✓ 1, -9

Solution or Explanation

$f(x) = x^3 + 12x^2 - 27x \Rightarrow f'(x) = 3x^2 + 24x - 27 = 3(x^2 + 8x - 9) = 3(x + 9)(x - 1)$. $f'(x) = 0 \Leftrightarrow x = -9, 1$. These are the only critical numbers.

8. 2/0 points | [Previous Answers](#)SCalcET8 4.1.027.

Sketch the graph of f by hand and use your sketch to find the absolute and local maximum and minimum values of f . (Enter your answers as a comma-separated list. If an answer does not exist, enter DNE.)

$$f(x) = \begin{cases} x^2 & \text{if } -1 \leq x \leq 0 \\ 8 - 9x & \text{if } 0 < x \leq 1 \end{cases}$$

\$\$\$DNE

absolute maximum value



DNE

\$\$\$-1

absolute minimum value



-1

\$\$\$DNE

local maximum value(s)



DNE

\$\$\$0

local minimum value(s)




0

Solution or Explanation

$$f(x) = \begin{cases} x^2 & \text{if } -1 \leq x \leq 0 \\ 8 - 9x & \text{if } 0 < x \leq 1 \end{cases}$$

No absolute or local maximum.

Local minimum $f(0) = 0$.Absolute minimum $f(1) = -1$.

9. 1/1 points | [Previous Answers](#)SCalcET8 4.1.037.MI.

Find the critical numbers of the function. (Enter your answers as a comma-separated list. If an answer does not exist, enter DNE.)

$$h(t) = t^{3/4} - 6t^{1/4}$$

\$\$\$0,4

$t =$

✓

Solution or Explanation

[Click to View Solution](#)

10. 1/0 points | [Previous Answers](#)SCalcET8 4.1.041.

Find the critical numbers of the function. (Enter your answers as a comma-separated list. Use n to denote any arbitrary integer values. If an answer does not exist, enter DNE.)

$$f(\theta) = 10 \cos(\theta) + 5 \sin^2(\theta)$$

\$\$\$nn,2nn

$\theta =$

✓

Solution or Explanation

[Click to View Solution](#)

11. 1/1 points | [Previous Answers](#)SCalcET8 4.1.043.

Find the critical numbers of the function. (Enter your answers as a comma-separated list. If an answer does not exist, enter DNE.)

$$f(x) = x^6 e^{-7x}$$

\$\$\$0,67

$x =$

✓

Solution or Explanation

[Click to View Solution](#)

12. 1/1 points | [Previous Answers](#)SCalcET8 4.1.047.

Find the absolute maximum and absolute minimum values of f on the given interval.

$$f(x) = 17 + 2x - x^2, \quad [0, 5]$$

\$\$\$2

absolute minimum value

✓

\$\$\$18

absolute maximum value

✓

Solution or Explanation

$f(x) = 17 + 2x - x^2, \quad [0, 5]. \quad f'(x) = 2 - 2x = 0 \Leftrightarrow x = 1. \quad f(0) = 17, \quad f(1) = 18, \quad \text{and} \quad f(5) = 2. \quad \text{So} \quad f(1) = 18 \text{ is the absolute maximum value and } f(5) = 2 \text{ is the absolute minimum value.}$

13.2/2 points | [Previous Answers](#)SCalcET8 4.1.054.Find the absolute maximum and absolute minimum values of f on the given interval.

$$f(x) = \frac{x}{x^2 - x + 1}, \quad [0, 3]$$

\$\$\$0

absolute minimum value

✓

\$\$\$1

absolute maximum value

✓

Solution or Explanation

$$f(x) = \frac{x}{x^2 - x + 1}, \quad [0, 3]. \quad f'(x) = \frac{(x^2 - x + 1) - x(2x - 1)}{(x^2 - x + 1)^2} = \frac{x^2 - x + 1 - 2x^2 + x}{(x^2 - x + 1)^2} = \frac{1 - x^2}{(x^2 - x + 1)^2} = 0 \Leftrightarrow x = \pm 1, \text{ but } x = -1$$

is not in the given interval, $[0, 3]$. $f(0) = 0$, $f(1) = 1$, and $f(3) = \frac{3}{7}$. So $f(1) = 1$ is the absolute maximum value and $f(0) = 0$ is the absolute minimum value.

14.2/2 points | [Previous Answers](#)SCalcET8 4.1.057.Find the absolute maximum and absolute minimum values of f on the given interval.

$$f(t) = 2 \cos(t) + \sin(2t), \quad [0, \pi/2]$$

\$\$\$0

absolute minimum value

✓ \$\$\$ $\sqrt{3} + \sqrt{3}2$

absolute maximum value

✓

Solution or Explanation

$$f(t) = 2 \cos(t) + \sin(2t), \quad [0, \pi/2].$$

$$f'(t) = -2 \sin(t) + \cos(2t) \cdot 2 = -2 \sin(t) + 2(1 - 2 \sin^2(t)) = -2(2 \sin^2(t) + \sin(t) - 1) = -2(2 \sin(t) - 1)(\sin(t) + 1).$$

$$f'(t) = 0 \Rightarrow \sin(t) = \frac{1}{2} \text{ or } \sin(t) = -1 \Rightarrow t = \frac{\pi}{6}. \quad f(0) = 2, \quad f\left(\frac{\pi}{6}\right) = \sqrt{3} + \frac{1}{2}\sqrt{3} = \frac{3}{2}\sqrt{3} \approx 2.60, \text{ and } f\left(\frac{\pi}{2}\right) = 0.$$

So $f\left(\frac{\pi}{6}\right) = \frac{3}{2}\sqrt{3}$ is the absolute maximum value and $f\left(\frac{\pi}{2}\right) = 0$ is the absolute minimum value.

15.1/1 points | [Previous Answers](#)SCalcET8 4.1.053.Find the absolute maximum and absolute minimum values of f on the given interval.

$$f(x) = x + \frac{25}{x}, [0.2, 20]$$

absolute minimum value

10

10

absolute maximum value

125.2

125.2

Solution or Explanation

$$f(x) = x + \frac{25}{x}, [0.2, 20]. \quad f'(x) = 1 - \frac{25}{x^2} = \frac{(x+5)(x-5)}{x^2} = 0 \Leftrightarrow x = \pm 5, \text{ but } x = -5 \text{ is not in the given interval, } [0.2, 20].$$

$f'(x)$ does not exist when $x = 0$, but 0 is not in the given interval, so 5 is the only critical number.

$f(0.2) = 125.2$, $f(5) = 10$, and $f(20) = 21.25$. So $f(0.2) = 125.2$ is the maximum value and $f(5) = 10$ is the absolute minimum value.

16.1/0 points | [Previous Answers](#)SCalcET8 4.1.061.Find the absolute maximum and absolute minimum values of f on the given interval.

$$f(x) = \ln(x^2 + 3x + 12), [-2, 1]$$

absolute minimum value

$\ln\left(\frac{39}{4}\right)$

$\ln\left(\frac{39}{4}\right)$

absolute maximum value

$\ln(16)$

$\ln(16)$

Solution or Explanation

$$f(x) = \ln(x^2 + 3x + 12), [-2, 1]. \quad f'(x) = \frac{1}{x^2 + 3x + 12} \cdot (2x + 3) = 0 \Leftrightarrow x = -\frac{3}{2}. \text{ Since } x^2 + 3x + 12 > 0 \text{ for all } x, \text{ the}$$

domain of f and f' is \mathbb{R} . $f(-2) = \ln 10 \approx 2.303$, $f\left(-\frac{3}{2}\right) = \ln \frac{39}{4} \approx 2.277$, and $f(1) = \ln 16 \approx 2.773$. So $f(1) = \ln 16 \approx 2.773$ is the absolute maximum value and $f\left(-\frac{3}{2}\right) = \ln \frac{39}{4} \approx 2.277$ is the absolute minimum value.

17.1/1 points | [Previous Answers](#)SCalcET8 4.1.503.XP.

Find the critical numbers of the function. (Enter your answers as a comma-separated list. If an answer does not exist, enter DNE.)

$$g(x) = x^{1/9} - x^{-8/9}$$

-8

x =

-8

Solution or Explanation

[Click to View Solution](#)

18.2/2 points | [Previous Answers](#)SCalcET8 4.1.505.XP.Find the absolute maximum and absolute minimum values of f on the given interval.

$$f(t) = t\sqrt{16 - t^2}, \quad [-1, 4]$$

absolute minimum value

absolute maximum value

Solution or Explanation

$$f(t) = t\sqrt{16 - t^2}, \quad [-1, 4].$$

$$f'(t) = t \cdot \frac{1}{2}(16 - t^2)^{-1/2}(-2t) + (16 - t^2)^{1/2} \cdot 1 = \frac{-t^2}{\sqrt{16 - t^2}} + \sqrt{16 - t^2} = \frac{-t^2 + (16 - t^2)}{\sqrt{16 - t^2}} = \frac{16 - 2t^2}{\sqrt{16 - t^2}}.$$

$$f'(t) = 0 \Rightarrow 16 - 2t^2 = 0 \Rightarrow t^2 = 8 \Rightarrow t = \pm\sqrt{8}, \text{ but } t = -\sqrt{8} \text{ is not in the given interval, } [-1, 4].$$

$f'(t)$ does not exist if $16 - t^2 = 0 \Rightarrow t = \pm 4$, but -4 is not in the given interval. $f(-1) = -\sqrt{15}$, $f(\sqrt{8}) = 8$, and $f(4) = 0$. So $f(\sqrt{8}) = 8$ is the absolute maximum value and $f(-1) = -\sqrt{15}$ is the absolute minimum value.

19.2/2 points | [Previous Answers](#)SCalcET8 4.1.510.XP.Find the absolute minimum and absolute maximum values of f on the given interval.

$$f(x) = x - \ln(5x), \quad \left[\frac{1}{2}, 2\right]$$

absolute minimum

absolute maximum

Solution or Explanation

[Click to View Solution](#)