

The due date for this assignment is past. Your work can be viewed below, but no changes can be made.

Important! Before you view the answer key, decide whether or not you plan to request an extension. Your Instructor may *not* grant you an extension if you have viewed the answer key. Automatic extensions are not granted if you have viewed the answer key.

[Request Extension](#)

1. 2/2 points | [Previous Answers](#)SCalcET8 3.4.003.

Write the composite function in the form $f(g(x))$. [Identify the inner function $u = g(x)$ and the outer function $y = f(u)$.]

$$y = \tan(\pi x)$$

$$(g(x), f(u)) = ($$

$$\pi x, \tan(u)$$

$$\pi x, \tan(u))$$

Find the derivative dy/dx .

$$\frac{dy}{dx} =$$

$$\sec^2(\pi x)(\pi)$$

$$\pi \sec^2(\pi x)$$

Solution or Explanation

Let $u = g(x) = \pi x$ and $y = f(u) = \tan(u)$. Then $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = (\sec^2(u))(\pi) = \pi \sec^2(\pi x)$.

2. 1/1 points | [Previous Answers](#)SCalcET8 3.4.007.

Find the derivative of the function.

$$F(x) = (3x^6 + 4x^3)^4$$

$$F'(x) =$$

$$4(3x^6 + 4x^3)^3(18x^5 + 12x^2)$$

$$24x^{11} (3x^3 + 2) (3x^3 + 4)^3$$

Solution or Explanation

$F(x) = (3x^6 + 4x^3)^4 \Rightarrow F'(x) = 4(3x^6 + 4x^3)^3 \cdot \frac{d}{dx}(3x^6 + 4x^3) = 4(3x^6 + 4x^3)^3(18x^5 + 12x^2)$. We can factor as follows:
 $4(x^3)^3(3x^3 + 4)^3 6x^2(3x^3 + 2) = 24x^{11}(3x^3 + 4)^3(3x^3 + 2)$.

3. 1/1 points | [Previous Answers](#)SCalcET8 3.4.011.

Find the derivative of the function.

$$f(\theta) = \cos(\theta^2)$$

$$f'(\theta) =$$

$$-2\theta \sin(\theta^2)$$

$$-2\theta \sin(\theta^2)$$

Solution or Explanation

$$f(\theta) = \cos(\theta^2) \Rightarrow f'(\theta) = -\sin(\theta^2) \cdot \frac{d}{d\theta}(\theta^2) = -\sin(\theta^2) \cdot (2\theta) = -2\theta \sin(\theta^2)$$

4. 1/1 points | [Previous Answers](#)SCalcET8 3.4.021.

Find the derivative of the function.

$$y = \sqrt{\frac{x}{x+9}}$$

$$y' =$$

$$\frac{9}{2\sqrt{x}(x+9)^{3/2}}$$

$$\frac{9}{2\sqrt{x}(x+9)^{3/2}}$$

Solution or Explanation

$$y = \sqrt{\frac{x}{x+9}} = \left(\frac{x}{x+9}\right)^{1/2} \Rightarrow$$

$$y' = \frac{1}{2} \left(\frac{x}{x+9}\right)^{-1/2} \frac{d}{dx} \left(\frac{x}{x+9}\right) = \frac{1}{2} \left(\frac{x^{-1/2}}{(x+9)^{-1/2}}\right) \left(\frac{(x+9)(1) - x(1)}{(x+9)^2}\right)$$

$$= \frac{1}{2} \left(\frac{(x+9)^{1/2}}{x^{1/2}}\right) \left(\frac{9}{(x+9)^2}\right) = \frac{9}{2\sqrt{x}(x+9)^{3/2}}$$

5. 1/1 points | [Previous Answers](#)SCalcET8 3.4.023.

Find the derivative of the function.

$$y = e^{\tan(\theta)}$$

$$y' =$$

$$e^{\tan(\theta)} \sec^2(\theta)$$

$$e^{\tan(\theta)} \sec^2(\theta)$$

Solution or Explanation

$$y = e^{\tan(\theta)} \Rightarrow y' = e^{\tan(\theta)} \cdot \frac{d}{d\theta}(\tan(\theta)) = (\sec^2(\theta))e^{\tan(\theta)}$$

6. 1/1 points | [Previous Answers](#)SCalcET8 3.4.031.

Find the derivative of the function.

$$F(t) = e^{3t \sin(2t)}$$

$$F'(t) =$$

$$e^{3t \sin(2t)} (3 \sin(2t) + 6t \cos(2t))$$



$$e^{3t \sin(2t)} (6t \cos(2t) + 3 \sin(2t))$$

Solution or Explanation

By the [example](#), $F(t) = e^{3t \sin(2t)} \Rightarrow$

$$F'(t) = e^{3t \sin(2t)} (3t \sin(2t))' = e^{3t \sin(2t)} (3t \cdot 2 \cos(2t) + \sin(2t) \cdot 3) = e^{3t \sin(2t)} (6t \cos(2t) + 3 \sin(2t))$$

7. 1/1 points | [Previous Answers](#)SCalcET8 3.4.041.

Find the derivative of the function.

$$f(t) = \cos^2(e^{\cos^2(t)})$$

$$f'(t) =$$

$$-2 \cos(\cos^2(t)) (-\sin(\cos^2(t))) (\cos^2(t)) (2 \cos(t)) (-\sin(t))$$



$$4 \cos(e^{\cos^2(t)}) \sin(e^{\cos^2(t)}) e^{\cos^2(t)} \cos(t) \sin(t)$$

Solution or Explanation

[Click to View Solution](#)8. 1/0 points | [Previous Answers](#)SCalcET8 3.4.042.

Find the derivative of the function.

$$y = \sqrt{7x + \sqrt{7x + \sqrt{7x}}}$$

$$y' =$$

$$\frac{1}{2} \sqrt{7x + \sqrt{7x + \sqrt{7x}}} \left\{ 7 + 7 + 7 \cdot \frac{1}{2} \sqrt{7x + \sqrt{7x}} \right\}$$



$$\frac{\frac{1}{2} \sqrt{7x + \sqrt{7x + \sqrt{7x}}} + 7}{2 \sqrt{7x + \sqrt{7x + \sqrt{7x}}}}$$

Solution or Explanation

[Click to View Solution](#)

9. 2/2 points | [Previous Answers](#)SCalcET8 3.4.050.

Find y' and y'' .

$$y = e^{4e^x}$$

$$y' =$$

$$4e^{x+4e^x}$$

$$y'' =$$

$$4e^{x+4e^x} (4e^x + 1)$$

Solution or Explanation

[Click to View Solution](#)

10. 1/1 points | [Previous Answers](#)SCalcET8 3.4.053.

Find an equation of the tangent line to the curve at the given point.

$$y = \sin(\sin(x)), \quad (\pi, 0)$$

$$y = -1(x - \pi)$$

$$y = \pi - x$$

Solution or Explanation

[Click to View Solution](#)

11. 1/1 points | [Previous Answers](#)SCalcET8 3.4.062.

If $h(x) = \sqrt{7 + 6f(x)}$, where $f(4) = 7$ and $f'(4) = 3$, find $h'(4)$.

$$h'(4) = \frac{9}{7}$$


Solution or Explanation

[Click to View Solution](#)


12.2/2 points | [Previous Answers](#)SCalcET8 3.4.069.

Suppose f is differentiable on \mathbb{R} . Let $F(x) = f(e^x)$ and $G(x) = e^{f(x)}$. Find expressions for the following.

(a) $F'(x)$

- ☐ $f'(e^x)$
☐ $f(e^x)e^x$
☐ $f'(e^x)xe^x$
☐ $f(e^x)xe^x$
☒  $f'(e^x)e^x$

(b) $G'(x)$

- ☐ $e^{f(x)}f(x)$
☐ $e^{f'(x)}$
☒  $e^{f(x)}f'(x)$
☐ $e^{f'(x)}f(x)$
☐ $e^{f'(x)}f'(x)$



Solution or Explanation

$$(a) \quad F(x) = f(e^x) \Rightarrow F'(x) = f'(e^x) \frac{d}{dx}(e^x) = f'(e^x)e^x$$

$$(b) \quad G(x) = e^{f(x)} \Rightarrow G'(x) = e^{f(x)} \frac{d}{dx}f(x) = e^{f(x)}f'(x)$$

13.1/1 points | [Previous Answers](#)SCalcET8 3.4.073.

If $F(x) = f(6f(4f(x)))$, where $f(0) = 0$ and $f'(0) = 1$, find $F'(0)$.

$$F'(0) = \boxed{24} \quad \text{✓} \quad \text{24}$$

Solution or Explanation

[Click to View Solution](#)14.1/0 points | [Previous Answers](#)SCalcET8 3.4.079.

The displacement of a particle on a vibrating string is given by the equation $s(t) = 3 + \frac{1}{4} \sin(3\pi t)$ where s is measured in centimeters and t in seconds. Find the velocity of the particle after t seconds.

$$v(t) =$$

$$14 \cos(3\pi t) [3\pi]$$

$$\frac{3\pi}{4} \cos(3\pi t) \text{ cm/s}$$

Solution or Explanation

[Click to View Solution](#)

15.1/0 points | [Previous Answers](#)SCalcET8 3.4.071.

Let $r(x) = f(g(h(x)))$, where $h(1) = 3$, $g(3) = 5$, $h'(1) = 3$, $g'(3) = 3$, and $f'(5) = 5$. Find $r'(1)$.

$r'(1) =$  

Solution or Explanation

[Click to View Solution](#)

16.3/3 points | [Previous Answers](#)SCalcET8 3.4.083.MI.

The motion of a spring that is subject to a frictional force or a damping force (such as a shock absorber in a car) is often modeled by the product of an exponential function and a sine or cosine function. Suppose the equation of motion of a point on such a spring is

$$s(t) = 4e^{-1.6t} \sin(2\pi t)$$





where s is measured in centimeters and t in seconds. Find the velocity after t seconds.

$$v(t) =$$



$$4[e^{-1.6t}(-1.6)(\sin(2\pi t)) + e^{-1.6t}(\cos(2\pi t)(2\pi))]$$



$$4e^{-1.6t} (2\pi \cos(2\pi t) - 1.6 \sin(2\pi t))$$

Graph the position function for $0 \leq t \leq 2$.



Graph the velocity function for $0 \leq t \leq 2$.





Solution or Explanation

[Click to View Solution](#)

17.1/1 points | [Previous Answers](#)SCalcET8 3.4.505.XP.

Find the derivative of the function.

$$y = \left(\frac{x^2 + 2}{x^2 - 2} \right)^6$$

$y' =$

$$-6 \frac{(x^2 + 2)^5 [(2x)(x^2 - 2) - (x^2 + 2)(2x)]}{(x^2 - 2)^7}$$

$$-48x \frac{(x^2 + 2)^5}{(x^2 - 2)^7}$$

Solution or Explanation

[Click to View Solution](#)

18.2/2 points | [Previous Answers](#)SCalcET8 3.4.515.XP.

Find the derivative of the function.

$$y = \cos(\cos(\cos(x)))$$

$y' =$

$$- \sin(\cos(\cos(x))) [-\sin(\cos(x))] [-\sin(x)]$$

$$- \sin(\cos(\cos(x))) \sin(\cos(x)) \sin(x)$$

Solution or Explanation

[Click to View Solution](#)

19.1/0 points | [Previous Answers](#)SCalcET8 3.4.516.XP.

Find the derivative of the function.

$$y = 5^{4^{x^2}}$$

 $y' =$

$$5^{4^{x^2}} \ln(5) 4^{x^2} \ln(4) (2x)$$

$$5^{4^{x^2}} \ln(5) 4^{x^2} \ln(4) (2x)$$

Solution or Explanation

[Click to View Solution](#)20.2/2 points | [Previous Answers](#)SCalcET8 3.5.001.

Consider the following equation.

$$4x^2 - y^2 = 3$$

(a) Find y' by implicit differentiation. $y' =$

$$-8x - 2y$$

$$\frac{4x}{y}$$

(b) Solve the equation explicitly for y and differentiate to get y' in terms of x . $y' = \pm$

$$\frac{4x}{\sqrt{4x^2 - 3}}$$

$$\frac{4x}{\sqrt{4x^2 - 3}}$$

Solution or Explanation

$$(a) \quad \frac{d}{dx}(4x^2 - y^2) = \frac{d}{dx}(3) \Rightarrow 8x - 2yy' = 0 \Rightarrow 2yy' = 8x \Rightarrow y' = \frac{4x}{y}$$

$$(b) \quad 4x^2 - y^2 = 3 \Rightarrow y^2 = 4x^2 - 3 \Rightarrow y = \pm \sqrt{4x^2 - 3}, \text{ so } y' = \pm \frac{1}{2}(4x^2 - 3)^{-1/2}(8x) = \pm \frac{4x}{\sqrt{4x^2 - 3}}.$$

$$\text{From part (a), } y' = \frac{4x}{y} = \frac{4x}{\pm \sqrt{4x^2 - 3}}, \text{ which agrees with part (b).}$$

21.1/1 points | [Previous Answers](#)SCalcET8 3.5.005.Find dy/dx by implicit differentiation.

$$x^2 - 8xy + y^2 = 8$$

 $y' =$

$$8y - 2x - 8x + 2y$$

$$\frac{4y - x}{y - 4x}$$



Solution or Explanation

$$\begin{aligned} \frac{d}{dx}(x^2 - 8xy + y^2) &= \frac{d}{dx}(8) \Rightarrow 2x - 8[xy' + y(1)] + 2yy' = 0 \\ &\Rightarrow 2yy' - 8xy' = 8y - 2x \\ &\Rightarrow y'(2y - 8x) = 8y - 2x \\ &\Rightarrow y' = \frac{4y - x}{y - 4x} \end{aligned}$$

22.1/1 points | [Previous Answers](#)SCalcET8 3.5.011.Find dy/dx by implicit differentiation.

$$y \cos(x) = 3x^2 + 2y^2$$

 $y' =$

$$6x + y \sin(x) \cos(x) - 4y$$

$$\frac{y \sin(x) + 6x}{\cos(x) - 4y}$$



Solution or Explanation

$$\begin{aligned} \frac{d}{dx}(y \cos(x)) &= \frac{d}{dx}(3x^2 + 2y^2) \Rightarrow y(-\sin(x)) + \cos(x) \cdot y' = 6x + 4yy' \Rightarrow \cos(x) \cdot y' - 4yy' = 6x + y \sin(x) \Rightarrow \\ y'(\cos(x) - 4y) &= 6x + y \sin(x) \Rightarrow y' = \frac{6x + y \sin(x)}{\cos(x) - 4y} \end{aligned}$$

23.2/0 points | [Previous Answers](#)SCalcET8 3.5.020.Find dy/dx by implicit differentiation.

$$\tan(x - y) = \frac{y}{9 + x^2}$$

 $y' =$

$$4 \sec^2(x - y) + 18x^2 \sec^2(x - y) + 81 \sec^2(x - y) + 2xyx^2 + 9 + x^4 \sec^2(x - y) + 18x^2 \sec^2(x - y) + 81 \sec^2(x - y)$$

$$\frac{(9 + x^2) \sec^2(x - y) + 2x \tan(x - y)}{1 + (9 + x^2) \sec^2(x - y)}$$



Solution or Explanation

[Click to View Solution](#)

24.1/1 points | [Previous Answers](#)SCalcET8 3.5.025.

Use implicit differentiation to find an equation of the tangent line to the curve at the given point.

$$y \sin(12x) = x \cos(2y), \quad (\pi/2, \pi/4)$$

y =

$$-3(x - \pi/2) + \pi/4$$

$$-3x + \frac{7\pi}{4}$$



Solution or Explanation

$y \sin(12x) = x \cos(2y) \Rightarrow y \cdot \cos(12x) \cdot 12 + \sin(12x) \cdot y' = x(-\sin(2y) \cdot 2y') + \cos(2y) \cdot 1 \Rightarrow$
 $\sin(12x) \cdot y' + 2x \sin(2y) \cdot y' = -12y \cos(12x) + \cos(2y) \Rightarrow y'(\sin(12x) + 2x \sin(2y)) = -12y \cos(12x) + \cos(2y) \Rightarrow$
 $y' = \frac{-12y \cos(12x) + \cos(2y)}{\sin(12x) + 2x \sin(2y)}$. When $x = \frac{\pi}{2}$ and $y = \frac{\pi}{4}$, we have $y' = \frac{(-3\pi)(1) + 0}{0 + \pi \cdot 1} = \frac{-3\pi}{\pi} = -3$, so an equation of the
 tangent line is $y - \frac{\pi}{4} = -3\left(x - \frac{\pi}{2}\right)$, or $y = -3x + \frac{7\pi}{4}$.

25.1/1 points | [Previous Answers](#)SCalcET8 3.5.029.

Use implicit differentiation to find an equation of the tangent line to the curve at the given point.


$$x^2 + y^2 = (4x^2 + 2y^2 - x)^2$$

(0, 0.5)
(cardioid)

y =

$$x + 0.5$$

$$x + \frac{1}{2}$$



Solution or Explanation

[Click to View Solution](#)

26.1/1 points | [Previous Answers](#)SCalcET8 3.5.030.

Use implicit differentiation to find an equation of the tangent line to the curve at the given point.

$$x^{2/3} + y^{2/3} = 4$$


$$(-3\sqrt{3}, 1)$$

(astroid)

y =

$$(-1^3\sqrt{-3\sqrt{3}})(x+3\sqrt{3})+1$$

$$\left(\frac{1}{\sqrt{3}}\right)x + 4$$



Solution or Explanation

[Click to View Solution](#)27.1/1 points | [Previous Answers](#)SCalcET8 3.5.035.Find y'' by implicit differentiation.

$$x^2 + 4y^2 = 4$$

y'' =

$$-14y^3$$

$$-\frac{1}{4y^3}$$

Solution or Explanation

$$x^2 + 4y^2 = 4 \Rightarrow 2x + 8yy' = 0$$

$$\Rightarrow y' = -x/(4y)$$

$$\Rightarrow y'' = -\frac{\frac{1}{4} \cdot y \cdot 1 - x \cdot y'}{y^2} = -\frac{\frac{1}{4}y - x[-x/(4y)]}{y^2} = -\frac{\frac{1}{4} \cdot 4y^2 + x^2}{4y^3} = -\frac{1}{4} \cdot \frac{4}{4y^3}$$

[since x and y must satisfy the
original equation $x^2 + 4y^2 = 4$]

$$\text{Thus, } y'' = -\frac{1}{4y^3}.$$

28.0/0 points | [Previous Answers](#)SCalcET8 3.5.040.If $x^2 + xy + y^3 = 1$, find the value of y''' at the point where $x = 1$. $2(27y^4 - 81y^3 + 6y^2 + 12y - 1)(3y^2 + 1)^4$

✗ 42

Solution or Explanation

If $x = 1$ in $x^2 + xy + y^3 = 1$, then we get $1 + y + y^3 = 1 \Rightarrow y^3 + y = 0 \Rightarrow y(y^2 + 1) \Rightarrow y = 0$, so the point where $x = 1$ is $(1, 0)$. Differentiating implicitly with respect to x gives us $2x + xy' + y \cdot 1 + 3y^2 \cdot y' = 0$. Substituting 1 for x and 0 for y gives us $2 + y' + 0 + 0 = 0 \Rightarrow y' = -2$. Differentiating $2x + xy' + y + 3y^2y' = 0$ implicitly with respect to x gives us $2 + xy'' + y' \cdot 1 + y' + 3(y^2y'' + y' \cdot 2yy') = 0$. Now substitute 1 for x , 0 for y , and -2 for y' . $2 + y'' + (-2) + (-2) + 3(0 + 0) = 0 \Rightarrow y'' = 2$. Differentiating $2 + xy'' + 2y' + 3y^2y'' + 6y(y')^2 = 0$ implicitly with respect to x gives us $xy''' + y'' \cdot 1 + 2y'' + 3(y^2y''' + y'' \cdot 2yy') + 6[y \cdot 2y'y'' + (y')^2y'] = 0$. Now substitute 1 for x , 0 for y , -2 for y' , and 2 for y'' . $y''' + 2 + 4 + 3(0 + 0) + 6[0 + (-8)] = 0 \Rightarrow y''' = -2 - 4 + 48 = 42$.

29.3/3 points | [Previous Answers](#)SCalcET8 3.5.502.XP.

Consider the following.

$$\cos(x) + \sqrt{y} = 6$$

(a) Find y' by implicit differentiation. $y' =$ $2\sin(x)12(y)-12$ ✓ $2\sqrt{y} \sin(x)$ (b) Solve the equation explicitly for y and differentiate to get y' in terms of x . $y' =$ $2(6-\cos(x))(\sin(x))$ ✓ $2\sin(x)(6 - \cos(x))$ (c) Check that your solutions to parts (a) and (b) are consistent by substituting the expression for y into your solution for part (a). $y' =$ $2\sin(x)12((6-\cos(x))^2)-12$ ✓ $2\sin(x)(6 - \cos(x))$

Solution or Explanation

[Click to View Solution](#)

30.1/1 points | [Previous Answers](#)SCalcET8 3.5.508.XP.

Find dy/dx by implicit differentiation.

$$5 \cos(x) \sin(y) = 1$$

$y' =$

$$5 \sin(x) \sin(y) 5 \cos(x) \cos(y)$$



$$\tan(x) \tan(y)$$

Solution or Explanation

[Click to View Solution](#)