

Module 6

Ann: Test 2 today
└ Hope to grade by wednesday

Preview: 1) Let $f(x) = 3x^3 + 2$; $g(x) = \sqrt{7-x}$

a) find $h = f - g$; Domain?

b) $h = \frac{2}{f}$; Domain?

c) Domains of $g \circ f$ and $f \circ g$?

2) Graph $g(x) = \begin{cases} 1 & -1 \leq x < 0 \\ 3 & 0 \leq x \leq 2 \\ -1 & 3 < x < 4 \end{cases}$

3) Find f^{-1} where $f(x) = 2 - \sqrt{x}$

$$\text{Dom}(f) = [0, \infty)$$

$$\text{Range}(f) = (-\infty, 2]$$

Note

$$x = f(f^{-1}(x)) = 2 - \sqrt{f^{-1}(x)}.$$

$$S_0, \quad X-2 = -\sqrt{f^{-1}(X)}$$

$$-X+2 = \sqrt{f^{-1}(X)}$$

$f^{-1}(x) = (2-x)^2$ is the rule for f^{-1} .

$$f^{-1}: (-\infty, 2] \longrightarrow [0, \infty) \quad \text{is } f^{-1}.$$

$$x \mapsto (2-x)^2$$

(Content: Inverses of functions.

Notation: $f \circ g$ is the function $f(g(x))$; $\text{id}(x) = x$.

Def: $g: B \rightarrow A$ is the inverse for \sim function $f: A \rightarrow B$ if $g \circ f = id_A$ and $f \circ g = id_B$.

$$A \xrightarrow{f} B \xrightarrow{f^{-1}} A$$

Denote g by f^{-1} .

Fact: Inverses are unique.

Def : A function is invertible if it has an inverse.

Fact: A function f is invertible iff it is bijective, that is surjective and injective.

Recall: Surjective means every element of the codomain is achieved and f is injective if whenever $f(a) = f(b)$, we have $a = b$.

Horiz. Test: A function $f: U \rightarrow V$, $U, V \subseteq \mathbb{R}$, $\text{im}(f) = V$ is invertible iff for every $c \in V$, the line $y=c$ intersects the graph of f only once. (this is a way to check injectivity visually)

Composing functions

Fact: for $f: A \rightarrow B$, $g: B \rightarrow C$, the domain of $g \circ f$ is A .

In this class, we give functions implicitly by their rules (grumble grumble), so here $\text{Dom}(g \circ f)$ is $g^{-1}(\text{Dom}(f))$.

Also, by $g \circ f$, we really mean $(g|_{\text{im}(f)} \circ f)$.
restrict g 's domain.

Restrict g 's down to

BB: 6.1, etc.


W6.W1

Preview 1) $f(x) = \sqrt{16-x^2}$; $g(x) = x+4$

a) Domain of f/g ?

b) Domain of $g \circ f$?

c) Domain of $f \circ g$?



Find $f \circ g$ and $g \circ f$.

3) Let $f(x) = \frac{7}{\sqrt{3x^3-9}}$; Find f^{-1} .

~~4) tires cost 45 USD per tire.
To put n wheels on an n -wheeler, it costs 35 times square root n plus tire costs. You need to find how many wheels each transaction had on an order.
Find such a formula.~~

5) A really cool spherical bubble is growing at a rate of $3 \frac{\text{cm}}{\text{min}}$ from a vat of liquid of Volume 1 cm^3 . If the bubble needs a thickness of at least 1 mm , how long till the bubble bursts? Surface area of a sphere is $4\pi r^2$. How large did the bubble get?

Content: BB 6.2, etc.

W6F 1

Ann i-Test 4 Aug 84 i/.

- Wiki 6 today

Content: 6, 3, wiki

Review: 1) Suppose $Ax - By = C$ with
 $A > B > 0 > C$ graph this line.

$$-By = C - Ax$$



$$y = \frac{C - Ax}{-B} = \frac{C}{-B} + \frac{-Ax}{-B}$$

$$= -\frac{C}{B} + \frac{Ax}{B}$$

$A, B > 0$, so $A/B > 0$

$C < 0$ and $B > 0$,

so $-\frac{C}{B} \rightarrow$ signs $- \overline{+} = +$



what are $g(w)$'s


w-ints and z-ints?

w-int: $(-2, 0), (1, 0), (2, 0)$


z-int: $(0, 1.5)$

3) Find AROC for h between

$x=2$ and $x=3$.




4) $h(x)$ is ; graph $3h(\frac{x}{2} - 1) + 1$.



5) $f(x)$ is a transformation of x^2 ,

while for function $f(x)$.



Module 7

W7M1

Preview

① An isosceles right triangle and a square are cut out of m meters of metal. What side lengths minimize area?

m is given
 b is variable we want
and x

$$A_{\Delta} = \frac{1}{2}b^2, \quad A_{\square}(l) = l^2$$

$$\text{also, } 4l + 2b + \sqrt{2}b = m$$

$$\text{so } l = \frac{m - (2 + \sqrt{2})b}{4}$$

$$\begin{aligned} A_{\text{tot}} &= \frac{1}{2}b^2 + \left(\frac{m - (2 + \sqrt{2})b}{4} \right)^2 \\ &= \frac{1}{2}b^2 + \left(\frac{m}{4} \right)^2 - 2m \cdot \frac{(2 + \sqrt{2})b}{4} + \left(\frac{(2 + \sqrt{2})}{4} \right)^2 b^2 \end{aligned}$$

Fact: min/max of poly is @ vertex:

Formula: For $ax^2 + bx + c$, vertex is at $\left(-\frac{b}{2a}, f\left(-\frac{b}{2a}\right) \right)$

Here we have poly in b

$$\left(\frac{1}{2} + \left(\frac{(2 + \sqrt{2})}{4} \right)^2 \right) b^2 - 2m \left(\frac{(2 + \sqrt{2})}{4} \right) b + \left(\frac{m}{4} \right)^2$$

$$\text{so vertex is at } \frac{-2m \left(\frac{(2 + \sqrt{2})}{4} \right)}{-2 \left(\frac{1}{2} + \left(\frac{(2 + \sqrt{2})}{4} \right)^2 \right)} = b_{\min}$$

Hence the base and length that minimize area are respectively

$$\frac{m \left(\frac{(2 + \sqrt{2})}{4} \right)}{\frac{1}{2} + \left(\frac{(2 + \sqrt{2})}{4} \right)^2} \quad \text{and} \quad \frac{m - (2 + \sqrt{2}) \cdot \left(\frac{m \left(\frac{(2 + \sqrt{2})}{4} \right)}{\frac{1}{2} + \left(\frac{(2 + \sqrt{2})}{4} \right)^2} \right)}{4}$$

Content:

Def: A function f is a quadratic if f is of the form $f(x) = ax^2 + bx + c$.

ex) 1) $4x^2 + 3x + 2$

2) $xy^4 + zy^2 + 1$; here note
 $= x(y^2)^2 + z(y^2) + 1$

Def: A root/zero of f is a $x \in \mathbb{R}$ such that $f(x) = 0$

Formula: for $f(x) = ax^2 + bx + c$,

the roots of $f(x)$ are

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b}{2a} \pm \sqrt{\left(\frac{-b}{2a} \right)^2 - \frac{c}{a}}$$

Proof: $0 = ax^2 + bx + c$

$$-c = ax^2 + bx$$

$$\frac{-c}{a} = x^2 + \frac{b}{a}x$$

$$\frac{-c}{a} = x^2 + \frac{b}{a}x + \left(\frac{b}{2a} \right)^2 - \left(\frac{b}{2a} \right)^2$$

$$\frac{-c}{a} + \left(\frac{b}{2a} \right)^2 = x^2 + \frac{b}{a}x + \left(\frac{b}{2a} \right)^2$$

$$\frac{-c}{a} + \left(\frac{b}{2a} \right)^2 = \left(x + \frac{b}{2a} \right)^2$$

$$\pm \sqrt{\left(\frac{b}{2a} \right)^2 - \frac{c}{a}} = x + \frac{b}{2a}$$

$$x = \frac{-b}{2a} \pm \sqrt{\left(\frac{b}{2a} \right)^2 - \frac{c}{a}}$$

Note: Vertex on graph of parabola is symmetric about roots, hence vertex's x -coord is $-\frac{b}{2a}$

Now BB 7.1, etc.


Contn: BB $d^2 + d^3$

Thm: Alek's walk around and help.

Ans: wiki is graded

Preview: 1) Dale needs to add a privacy fence to his rectangular resort. A cliff covers one side of his property. If he bought 1 km of fence, how much area can he cover?

2) $f(x)$ has plot



What's formula for f ?

3) $f(x) = x^2 + x$; $g(x) = \sqrt{x-2}$
find $f \circ g$ domain:

$$f \circ g = \sqrt{x-2}^2 + \sqrt{x-2}$$


$$= x-2 + \sqrt{x-2}$$

$$\text{dom}(f \circ g) = \text{dom}(g) = [2, \infty)$$

$\text{dom}(f \circ g) \subseteq \text{dom}(g)$
 $\text{dom}(f) = \mathbb{R}$, so no restrictions from here.

Preview for end goal of class

4) $f(x)$ has plot




Preview for next module

5) plot $f(x) = \frac{1}{3}(x-1)^2(x+3)(x+1)^3$

LC: pos

deg: $2+1+3=6$ even



Wiki

W8M1

review) 1) M2 - 63%

$3x + 2y = 3$; find line \perp through $(7, 2)$.

2) M3 - 90%

$$\text{is } (y+3)^2 = x - y^2 + 5y$$

a function of x ?

3) M4 - 50%

Domain of $\frac{\sqrt{x+7}}{3x-10}$

4) M6 - 90%
 $f(x) = \frac{x+2}{3x-1}$; $g(x) = 2x-1$; Find $f \circ g$.

Preview

5) $f(x)$ has graph



$\text{sgn}(f(x))$

roots w/ mult?

pts of inf

inc/dec?

EB?

Content: M8. *