

0.1 Landscape Simulation: Different Autocorrelation Scenarios




Figure 1: Simulated landscapes under different levels of autocorrelation (5, 10, 20; from top to bottom). Autocorrelation only affected the layers elev and forest, which were both simulated using a Gaussian random field neutral landscape model (Schlather et al., 2015) using the R-package NLMR (Sciaini et al., 2018). Simulations were repeated 100 times for each autocorrelation scenario, thus resulting in 300 unique landscape configurations.

0.2 Dynamic Tentative Distribution Parameters




Figure 2: Tentative parameter estimates for the von Mises distribution (top row) and gamma distribution (bottom row) fitted to steps with different durations. The von Mises distribution requires one parameter, namely a concentration parameter (κ). The gamma distribution requires a shape parameter (k) and a scale parameter (θ). The subscript ₀ is used to indicate that these are tentative distribution parameters (sensu Avgar et al., 2016 and Fieberg et al., 2021).

0.3 Distribution of Relative Turning Angles following Different Step Durations




Figure 3: Density of relative turning angles associated with steps of $\Delta t = 1$ following steps with different durations for all three autocorrelation scenarios (5, 10, and 20). To generate this figure, we assumed a missingness of 0.5 and forgiveness of 5.

0.4 Model Estimates across all Scenarios




Figure 4: Parameter estimates across different autocorrelation scenarios (5, 10, 20; panels a, b, and c) and missingness levels (0% - 50%; from left to right). True simulation parameters are indicated by the solid black lines. Parameter estimates from the different approaches are given by the colored symbols, and their bootstrap 95% CIs across 100 replicates by the colored lines.

0.5 Case Study Covariates




Figure 5: Covariates used for the case study, overlaid with the GPS data of a spotted hyena called “Apollo” (lines in black). Apollo was originally collared in 2007 in northern Botswana and monitored until 2011. The depicted area is part of the Okavango Delta, which is a massive wetland area. Data was projected to a local projection (EPSG:32734).

0.6 Case Study Model Output

Table 1: Model results from the case study using GPS data collected on Apollo. In F1, forgiveness was set to one (only 2-hour steps were considered), whereas in F3-S and F3-SH a forgiveness of three was employed (allowing for step durations of up to 6 hours). In model F3-S, the step duration was interacted with step descriptors. In model F3-SH, step duration was interacted with step descriptors and habitat covariates.

Coefficient	F1	F3-S	F3-SH
sl	0.00002 (0.00001)	0.00001 (0.00001)	0.00001 (0.00001)
log_sl	-0.02251 (0.01862)	-0.01805 (0.0157)	-0.01846 (0.0157)
cos_ta	0.03166 (0.03055)	-0.0085 (0.02557)	-0.009 (0.02558)
Water	-1.71418*** (0.20628)	-1.56915*** (0.13995)	-1.61501*** (0.17121)
DistanceToWater	-0.00005*** (0.00001)	-0.00005*** (0.00001)	-0.00005*** (0.00001)
Trees	1.51764* (0.88816)	-0.2178 (0.62517)	0.60764 (0.75259)
sl:duration4		-0.00001 (0.00002)	-0.00001 (0.00002)
sl:duration6		-0.00004** (0.00002)	-0.00004** (0.00002)
log_sl:duration4		0.08122 (0.05075)	0.07978 (0.0508)
log_sl:duration6		0.02867 (0.02471)	0.03058 (0.02474)
cos_ta:duration4		-0.07526 (0.05784)	-0.07635 (0.05788)
cos_ta:duration6		-0.16358*** (0.06055)	-0.16105*** (0.06059)
Water:duration4			0.08548 (0.34111)
Water:duration6			0.20782 (0.46194)
DistanceToWater:duration4			0.00002 (0.00002)
DistanceToWater:duration6			-0.00001 (0.00003)
Trees:duration4			-0.04696 (1.61148)
Trees:duration6			-6.73474*** (1.97442)
Steps	2,179	4,505	4,505
AIC	-	47,565	47,564

Significance codes: * $p < 0.10$ ** $p < 0.05$ *** $p < 0.01$

Bibliography

- Avgar, T., Potts, J. R., Lewis, M. A., & Boyce, M. S. (2016). Integrated Step Selection Analysis: Bridging the Gap Between Resource Selection and Animal Movement. *Methods in Ecology and Evolution*, 7(5), 619–630. <https://doi.org/10.1111/2041-210X.12528>
- Fieberg, J., Signer, J., Smith, B., & Avgar, T. (2021). A 'How to' Guide for Interpreting Parameters in Habitat-Selection Analyses. *Journal of Animal Ecology*, 90(5), 1027–1043. <https://doi.org/10.1111/1365-2656.13441>
- Schlather, M., Malinowski, A., Menck, P. J., Oesting, M., & Stokorb, K. (2015). Analysis, Simulation and Prediction of Multivariate Random Fields with Package RandomFields. *Journal of Statistical Software*, 63, 1–25. <https://doi.org/10.18637/jss.v063.i08>
- Sciaini, M., Fritsch, M., Scherer, C., & Simpkins, C. E. (2018). NLMR and landscapetools: An Integrated Environment for Simulating and Modifying Neutral Landscape Models in R. *Methods in Ecology and Evolution*, 9(11), 2240–2248. <https://doi.org/10.1111/2041-210X.13076>