

5DGSPTSBL: Geometric Fast-Slow Stable Boundary Layer Model

5DGSPTSBL Contributors

2026-08-03

Abstract

We present a five-dimensional geometric singular perturbation model for the stable boundary layer with state variables $(\tilde{e}, q_\theta, S, T_s, T_g)$. The model combines desingularized turbulence dynamics, slaved thermodynamic closures, and slow thermal coupling to support fold-locus diagnostics and data-driven parameter identification.

1 Closures and Smooth Regularization

To maintain C^r -smoothness near neutral stratification and support robust implicit integration with autodiff Jacobians, the stratification closure uses a hyperbolic embedding of $\max(0, x)$:

$$\text{smooth_max}(x; \epsilon) = \frac{x + \sqrt{x^2 + \epsilon^2}}{2}. \quad (1)$$

With $x = (\theta_0 - T_s)/h_{\text{sbl}}(\tilde{e})$, the closure becomes

$$\theta_z(T_s, \tilde{e}) = \text{smooth_max}\left(\frac{\theta_0 - T_s}{h_{\text{sbl}}(\tilde{e})}; \epsilon_{\text{strat}}\right), \quad (2)$$

where ϵ_{strat} is a small smoothing scale exposed in `SBLParams`.

The bulk transfer coefficients are

$$C_D = \left[\frac{\kappa}{\ln((z_1 - d)/z_{0m})} \right]^2, \quad (3)$$

$$C_H = \frac{\kappa^2}{\ln((z_1 - d)/z_{0m}) \ln((z_1 - d)/z_{0h})}. \quad (4)$$

1.1 Transcritical Ignition and Surface-Flux Parameterization

Breakout from the laminar floor $\tilde{e} \approx \sqrt{\delta}$ into the active turbulent manifold occurs when the shear S exceeds a local ignition threshold S_{trans} . Linearizing the desingularized fast equation $\frac{d\tilde{e}}{d\tau} = f_1(\tilde{e}, q_\theta)$ about the laminar state $(\tilde{e}_0, q_{\theta,0}) = (\sqrt{\delta}, 0)$ gives

$$\left. \frac{\partial f_1}{\partial \tilde{e}} \right|_{(\sqrt{\delta}, 0)} = \frac{1}{2} c_m \ell S^2 - \frac{3}{2\ell} \delta. \quad (5)$$

Setting this growth rate to zero yields the explicit transcritical threshold

$$S_{\text{trans}} = \frac{1}{\ell} \sqrt{\frac{3\delta}{c_m}}. \quad (6)$$

For $S < S_{\text{trans}}$, the laminar floor remains locally exponentially stable, while for $S > S_{\text{trans}}$ the principal fast eigenvalue becomes positive and trajectories are ejected toward the upper turbulent branch of the fast manifold.

To compare buffered and unbuffered surface-cooling regimes without introducing dimensional control parameters, we write the surface energy balance in terms of the nondimensional ground-flux ratio

$$\Pi_G = \frac{G}{R_{\text{net}}} = \frac{\frac{k_g}{d_g}(T_s - T_g)}{R_{\text{net}}(T_s)}. \quad (7)$$

Using Π_G makes regime boundaries invariant to soil conductivity and depth through (k_g, d_g) and to differing atmospheric radiative forcing profiles through R_{net} , while preserving the buffered versus runaway cooling interpretation.

1.2 Projection and Unfolding Proof Sketches

Let $\pi_{Ri}(\tilde{e}, q_\theta, S, T_s, T_g) = \frac{g\theta_z(T_s, \tilde{e})}{\theta_0 S^2}$ denote the scalar Richardson-number projection. Its differential is the nonzero row functional

$$D\pi_{Ri} = \left(\frac{g}{\theta_0 S^2} \frac{\partial \theta_z}{\partial \tilde{e}} \quad 0 \quad -\frac{2g\theta_z}{\theta_0 S^3} \quad \frac{g}{\theta_0 S^2} \frac{\partial \theta_z}{\partial T_s} \quad 0 \right), \quad (8)$$

so $\text{rank}(D\pi_{Ri}) = 1$ whenever $S > 0$ and $\theta_z > 0$. Since the fold set $\mathcal{C}_{\text{fold}} \subset \mathbb{R}^5$ is three-dimensional, the dimension count

$$\dim(T_p \mathcal{C}_{\text{fold}} \cap \ker D\pi_{Ri}(p)) \geq 3 + 4 - 5 = 2 \quad (9)$$

shows that the kernel of the restricted map $D\pi_{Ri}|_{T_p \mathcal{C}_{\text{fold}}}$ is necessarily nontrivial. Hence Richardson-number level sets intersect the folded fast manifold along two-dimensional fibers rather than at isolated points, which is the geometric source of non-universal projected critical Ri thresholds.

For the cusp unfolding, let (η_1, η_2) be the center-manifold control coordinates induced by geostrophic shear and deep-soil temperature. The physical unfolding matrix is

$$M = \begin{pmatrix} \partial_{S_g} \eta_1 & \partial_{T_{\text{deep}}} \eta_1 \\ \partial_{S_g} \eta_2 & \partial_{T_{\text{deep}}} \eta_2 \end{pmatrix}, \quad (10)$$

with leading entries $\partial_{S_g} \eta_1 = \epsilon_1/(\tilde{e}\tau_S)$ and $\partial_{T_{\text{deep}}} \eta_2 = \epsilon_1\epsilon_2 k_g/(\tilde{e}C_s d_g)$. Therefore

$$\det(M) = \frac{\epsilon_1^2 \epsilon_2 k_g}{\tilde{e}^2 C_s d_g \tau_S} \left(1 - \frac{\partial R_{\text{net}}}{\partial T_s} \frac{d_g}{k_g} \right), \quad (11)$$

which remains positive for the nocturnal linearized radiation law used here because $\partial R_{\text{net}}/\partial T_s < 0$. The unfolding is thus transverse and structurally stable in the (S_g, T_{deep}) control plane.

2 Governing Equations

The desingularized fast-slow system in fast time τ is

$$\frac{d\tilde{e}}{d\tau} = \frac{1}{2}c_m\ell S^2\tilde{e} - \frac{g}{2\theta_0}q_\theta - \frac{1}{2\ell}\tilde{e}^3, \quad (12)$$

$$\frac{dq_\theta}{d\tau} = -c_w\theta_z(T_s, \tilde{e})\tilde{e}^3 - \frac{g}{\theta_0}c_\theta\ell q_\theta^2 - \frac{C_\theta}{\ell}\tilde{e}^2q_\theta, \quad (13)$$

$$\frac{dS}{d\tau} = \frac{\epsilon_1}{\max(\tilde{e}, \sqrt{\delta})} \mathcal{F}_S, \quad (14)$$

$$\frac{dT_s}{d\tau} = \frac{\epsilon_1}{\max(\tilde{e}, \sqrt{\delta})} \frac{1}{C_s} \left(R_{\text{net}} + \rho c_p q_\theta + \frac{k_g}{d_g}(T_g - T_s) \right), \quad (15)$$

$$\frac{dT_g}{d\tau} = \frac{\epsilon_1\epsilon_2}{\max(\tilde{e}, \sqrt{\delta})} \frac{\kappa_g}{d_g^2}(T_s - T_g). \quad (16)$$

Remark on Asymptotic Normal Form: The dissipation term $-\frac{1}{2\ell}\tilde{e}^3$ represents the leading-order C^r -smooth normal form of Kolmogorov dissipation $\frac{e^{3/2}}{\ell}$ in fast time τ . Omitted $\mathcal{O}(\delta)$ terms contribute a bounded linear damping correction $-\frac{3\delta}{4\ell}\tilde{e}$ near the laminar floor $\tilde{e} \sim \sqrt{\delta}$, which is uniformly subdominant across the active turbulent manifold for $\delta \leq 10^{-4}$.

Writing the omitted remainder as $\mathcal{R}_\delta(\tilde{e}) = \frac{3\delta}{4\ell}\tilde{e} + \mathcal{O}(\delta^2)$, the desingularized dissipation error obeys

$$\max_{\tilde{e} \in [\sqrt{\delta}, \tilde{e}_{\text{max}}]} |\mathcal{R}_\delta(\tilde{e})| \leq \frac{3\delta^{3/2}}{4\ell}. \quad (17)$$

For the representative manuscript regime $\delta = 10^{-5}$ and $\ell = 10$ m, this gives $\max|\mathcal{R}_\delta| \lesssim 2.4 \times 10^{-9} \text{ m s}^{-2}$, which is below $10^{-5}\%$ of the active cubic dissipation magnitude at $\tilde{e} = 0.2$. The cubic fast-time normal form therefore preserves the physical dissipation scale to manuscript accuracy.

The interior mixing length is

$$\ell(z_{\text{eff}}) = \frac{\kappa z_{\text{eff}}}{1 + \kappa z_{\text{eff}}/\ell_\infty}, \quad z_{\text{eff}} = z - d + z_{0m}. \quad (18)$$

3 Notes Traceability

The manuscript templates were aligned with project notes emphasizing:

- desingularization using $\tilde{e} = \sqrt{e + \delta}$,
- geometric fast-slow interpretation of the closure manifold,
- smooth stratification regularization for Fenichel-compatible dynamics.

Primary note sources:

- `notes/Governing Equations and Turbulence Closure.md`
- `notes/Closures.md`
- `notes/Types.md`




Figure 1: CASES-99 benchmark simulation showing (a) fast state dynamics, (b) surface and ground temperatures, (c) phase space trajectory with fold locus contour, and (d) fast subsystem Jacobian determinant stability.




Figure 2: Phase space trajectory in $(\tilde{e}, q_\theta)$ space demonstrating convergence to the slow manifold.

4 Numerical Results and Phase Manifold

5 Reproducibility

The manuscript was assembled from repository templates and notes using `julia -project=. scripts/assemble_manuscript.jl`. The primary benchmark figure is generated by `scripts/run_cases99.jl` and written to `paper/figures/cases99_manifold.png`. lection —

References

- [Fenichel(1979)] Neil Fenichel. Geometric singular perturbation theory for ordinary differential equations. *Journal of Differential Equations*, 31(1):53–98, 1979. doi: 10.1016/0022-0396(79)90152-9.
- [Jones(1995)] Christopher K. R. T. Jones. Geometric singular perturbation theory. In Italy Montecatini Terme, editor, *Lecture Notes in Mathematics*, volume 1609 of *Lecture Notes in Mathematics*, pages 44–118. Springer Berlin Heidelberg, Berlin, Heidelberg, 1995. doi: 10.1007/BFB0095239.
- [Krupa and Szmolyan(2001)] Martin Krupa and Peter Szmolyan. Extending geometric singular perturbation theory to nonhyperbolic points—Fold and canard points in two dimensions. *SIAM Journal on Mathematical Analysis*, 33(2):286–314, 2001. doi: 10.1137/S0036141099360919.
- [Berglund and Gentz(2002)] Nils Berglund and Barbara Gentz. Geometric singular perturbation theory for stochastic differential equations. *Journal of Differential Equations*, 191(1):1–54, 2002. doi: 10.1016/S0022-0396(03)00020-2.
- [Lorenz(1986)] Edward N. Lorenz. On the existence of a slow manifold. *Journal of the Atmospheric Sciences*, 43(15):1547–1558, 1986. doi: 10.1175/1520-0469(1986)043<1547:OTEOAS>2.0.CO;2.
- [Maeschalck and Desroches(2013)] P. Maeschalck and M. Desroches. Numerical continuation techniques for planar slow-fast systems. *SIAM Journal on Applied Dynamical Systems*, 12(3):1159–1180, 2013. doi: 10.1137/120877386.
- [Kuehn(2015)] Christian Kuehn. *Multiple Time Scale Dynamics: An Introduction Specializing to Differential Equations*, volume 191 of *Applied Mathematical Sciences*. Springer International Publishing, Cham, Switzerland, 2015. ISBN 978-3-319-12315-8. doi: 10.1007/978-3-319-12315-8.
- [Jelbart and Kuehn(2022)] Samuel Jelbart and Christian Kuehn. Discrete geometric singular perturbation theory. *Discrete and Continuous Dynamical Systems - Series A*, 43(1):57–120, 2022. doi: 10.3934/dcds.2022142.
- [Kristiansen(2023)] Kristian Uldall Kristiansen. A review of multiple-time-scale dynamics: Fundamental phenomena and mathematical methods. In Bernhelm Booß-Bavnbek, Jens Christensen, Katherine Richardson, and Oriol Codina, editors, *Mathematics Online First Collections*, pages 309–363. Springer Nature Switzerland, Cham, 2023. ISBN 978-3-031-45105-8. doi: 10.1007/16618_2023_75.
- [McNider and Pielke(1981)] Richard T. McNider and Roger A. Pielke. Diurnal boundary-layer development over sloping terrain. *Journal of the Atmospheric Sciences*, 38(10):2198–2212, 1981. doi: 10.1175/1520-0469(1981)038<2198:DBLDOS>2.0.CO;2.
- [Mellor and Yamada(1982)] George L. Mellor and Tetsuji Yamada. Development of a turbulence closure model for geophysical fluid problems. *Reviews of Geophysics*, 20(4):851–875, 1982. doi: 10.1029/RG020i004p00851.
- [Nieuwstadt(1984)] F. T. M. Nieuwstadt. The turbulent structure of the stable atmospheric boundary layer. *Journal of the Atmospheric Sciences*, 41(14):2202–2216, 1984. doi: 10.1175/1520-0469(1984)041<2202:TTSOTS>2.0.CO;2.

- [England and McNider(1995)] David E. England and Richard T. McNider. Stability functions based upon shear functions. *Oceanographic Literature Review*, 74(1-2):113–130, 1995. doi: 10.1007/BF00715713.
- [McNider et al.(1995)McNider, England, Friedman, and Shi] R. McNider, David E. England, M. Friedman, and X. Shi. Predictability of the stable atmospheric boundary layer. *Journal of the Atmospheric Sciences*, 52(10):1602–1614, 1995. doi: 10.1175/1520-0469(1995)052<1602:POTSAB>2.0.CO;2.
- [Shi et al.(2005)Shi, McNider, Singh, England, Friedman, Lapenta, and Norris] X. Shi, R. McNider, M. P. Singh, David E. England, M. Friedman, W. Lapenta, and W. Norris. On the behavior of the stable boundary layer and role of initial conditions. *Pure and Applied Geophysics*, 162(10):1811–1829, 2005. doi: 10.1007/S00024-005-2694-7.
- [Cuxart et al.(2004)Cuxart, Holtslag, Beare, Bazile, Beljaars, Cheng, Conangla, Ek, Freedman, Hamdi, et al.] J. Cuxart, A. Holtslag, R. Beare, E. Bazile, A. Beljaars, A. Cheng, L. Conangla, M. Ek, F. Freedman, R. Hamdi, et al. Single-column model intercomparison for a stably stratified atmospheric boundary layer. *Boundary-Layer Meteorology*, 118(2):273–303, 2004. doi: 10.1007/s10546-005-3780-1.
- [Zilitinkevich et al.(2006)Zilitinkevich, Elperin, Kleerorin, and Rogachevskii] Sergej Zilitinkevich, Tov Elperin, Nathan Kleerorin, and Igor Rogachevskii. Energy- and flux-budget (EFB) turbulence closure model for stably stratified flows. Part I: Steady-state, homogeneous regimes. *Atmospheric Boundary Layers*, 125(2):11–35, 2006. doi: 10.1007/s10546-007-9189-2.
- [Jiménez and Cuxart(2006)] M. A. Jiménez and J. Cuxart. Study of the probability density functions from a large-eddy simulation for a stably stratified boundary layer. *Boundary-Layer Meteorology*, 128(1):117–135, 2006. doi: 10.1007/S10546-005-9009-5.
- [Wiel et al.(2012)Wiel, Moene, Jonker, Baas, Basu, Donda, Sun, and Holtslag] V. Wiel, A. Moene, H. Jonker, P. Baas, S. Basu, Jmm Judith Donda, Jielun Sun, and A. Holtslag. The minimum wind speed for sustainable turbulence in the nocturnal boundary layer. *Journal of the Atmospheric Sciences*, 69(11):3116–3127, 2012. doi: 10.1175/JAS-D-12-0107.1.
- [Grachev et al.(2013)Grachev, Andreas, et al.] Andrey A. Grachev, Edgar L. Andreas, et al. The critical Richardson number and turbulence in stably stratified boundary layers. *Boundary-Layer Meteorology*, 147(1):23–42, 2013. doi: 10.1007/s10546-012-9772-8.
- [Mahrt(2014)] Larry Mahrt. Stably stratified atmospheric boundary layers. *Annual Review of Fluid Mechanics*, 46(1):23–45, 2014. doi: 10.1146/annurev-fluid-010313-141354.
- [Monahan et al.(2015)Monahan, Rees, He, and McFarlane] Adam H. Monahan, T. Rees, Yanping He, and Norman A. McFarlane. Multiple regimes of wind, stratification, and turbulence in the stable boundary layer. *Journal of the Atmospheric Sciences*, 72(8):3178–3198, 2015. doi: 10.1175/JAS-D-14-0311.1.
- [Wiel et al.(2017)Wiel, Vignon, Baas, Hooijdonk, Linden, Hooft, Bosveld, Roode, Moene, and Genthon] B. V. D. Wiel, É. Vignon, P. Baas, Ivo G. S. van Hooijdonk, S. V. D. Linden, J. Hooft, F. Bosveld, S. R. D. Roode, A. Moene, and C. Genthon. Regime transitions in near-surface temperature inversions: A conceptual model. *Journal of the Atmospheric Sciences*, 74(4):1057–1073, 2017. doi: 10.1175/JAS-D-16-0180.1.

- [Hooft et al.(2019)Hooft, Baas, Tiggelen, Ansorge, and Wiel] J. Hooft, P. Baas, M. V. Tiggelen, C. Ansorge, and B. J. H. Wiel. An idealized description for the diurnal cycle of the dry atmospheric boundary layer. *Journal of the Atmospheric Sciences*, 76(12):3717–3736, 2019. doi: 10.1175/JAS-D-19-0023.1.
- [Ramsey and Monahan(2022)] Elizabeth Ramsey and Adam H. Monahan. Empirical low-dimensional dynamics of atmospheric stable boundary layer temperature inversions. *Journal of the Atmospheric Sciences*, 79(7):1965–1984, 2022. doi: 10.1175/JAS-D-21-0205.1.
- [Shaw et al.(2022)Shaw, Berg, Debnath, Deskos, Draxl, Ghate, Hasager, Kotamarthi, Mirocha, Muradyan, et al.] W. Shaw, L. Berg, M. Debnath, G. Deskos, C. Draxl, V. Ghate, C. Hasager, R. Kotamarthi, J. Mirocha, P. Muradyan, et al. Scientific challenges to characterizing the wind resource in the marine atmospheric boundary layer. *Wind Energy Science*, 7(6):2307–2334, 2022. doi: 10.5194/wes-2021-156.
- [He and Li(2025)] Jun-Yi He and Qiu-Sheng Li. *Winds in the Atmospheric Boundary Layer*. Springer Nature Singapore, Singapore, 2025. ISBN 978-981-95-1516-5. doi: 10.1007/978-981-95-1517-2.
- [Acevedo and Mahrt(2010)] Otávio Acevedo and Larry Mahrt. Systematic vertical variation of mesoscale fluxes in the nocturnal boundary layer. *Boundary-Layer Meteorology*, 135(1):19–30, 2010. doi: 10.1007/s10546-010-9465-4.
- [Poulos et al.(2002)Poulos, Blumen, Fritts, Lundquist, Sun, Burns, Nappo, Banta, Newsom, Cuxart, et al.] G. Poulos, W. Blumen, D. Fritts, J. Lundquist, Jielun Sun, S. Burns, C. Nappo, R. Banta, R. Newsom, J. Cuxart, et al. CASES-99: A comprehensive investigation of the stable atmospheric boundary layer. *Bulletin of the American Meteorological Society*, 83(4):555–581, 2002. doi: 10.1175/1520-0477(2002)083<0555:CACIOT>2.3.CO;2.
- [Grachev et al.(2005)Grachev, Fairall, Persson, Andreas, and Guest] A. Grachev, C. Fairall, P. Persson, Edgar L. Andreas, and P. Guest. Stable boundary-layer scaling regimes: The SHEBA data. *Boundary-Layer Meteorology*, 116(2):201–235, 2005. doi: 10.1007/s10546-004-2729-0.
- [Vignon et al.(2017)Vignon, Hourdin, Genthon, Gallee, Bazile, Lefebvre, Madeleine, and Wiel] É. Vignon, F. Hourdin, C. Genthon, H. Gallee, E. Bazile, M. Lefebvre, J. Madeleine, and Bas J. H. van de Wiel. Antarctic boundary layer parametrization in a general circulation model: 1-D simulations facing summer observations at Dome C. *Journal of Geophysical Research: Atmospheres*, 122(13):6818–6843, 2017. doi: 10.1002/2017JD026802.
- [Su et al.(2024)Su, Lu, Yuan, Wang, He, and Xia] Lian Su, Chunsong Lu, Jinlong Yuan, Xiaofei Wang, Qing He, and Haiyun Xia. Measurement report: The promotion of the low-level jet and thermal effects on the development of the deep convective boundary layer at the southern edge of the Taklimakan Desert. *Atmospheric Chemistry and Physics*, 24(18):10947–10963, 2024. doi: 10.5194/acp-24-10947-2024.
- [Zhang et al.(2024)Zhang, Zhang, Cai, Song, and Zhang] Lu Zhang, Hongsheng Zhang, Xuhui Cai, Yu Song, and Xiaoye Zhang. Characteristics of turbulence intermittency, fine structure, and flux correction in the Taklimakan desert. *Journal of the Atmospheric Sciences*, 81(2):459–475, 2024. doi: 10.1175/JAS-D-23-0107.1.

- [Hairer and Wanner(2010)] Ernst Hairer and Gerhard Wanner. *Solving Ordinary Differential Equations II: Stiff and Differential-Algebraic Problems*. Springer Series in Computational Mathematics. Springer-Verlag, Berlin, 2nd edition, 2010. doi: 10.1007/978-3-642-05221-7.
- [Krauskopf et al.(2007)Krauskopf, Osinga, and Galán-Vioque] B. Krauskopf, H. Osinga, and J. Galán-Vioque. *Numerical Continuation Methods for Dynamical Systems: Path-Following and Boundary Value Problems*. Springer, 2007. doi: 10.1007/978-1-4020-6356-5.
- [Rackauckas and Nie(2017)] Christopher Rackauckas and Qing Nie. DifferentialEquations.jl – A performant and feature-rich ecosystem for solving differential equations in Julia. *Journal of Open Research Software*, 5(1):15, 2017. doi: 10.5334/jors.151.
- [Steinebach(2023)] Gerd Steinebach. Construction of Rosenbrock–Wanner method Rodas5P and numerical benchmarks within the Julia Differential Equations package. *BIT Numerical Mathematics*, 63(2):27, 2023. doi: 10.1007/s10543-023-00967-x.