

Atmospheric Boundary Layer PDE Closure Benchmark Report

Empirical Equation Discovery, Stability Diagnostics, and Galerkin PDE Benchmarks

AtmosphericSlowManifold.jl Automated Pipeline

August 20, 2026

Abstract

This report synthesizes observational field campaign diagnostics (CASES-99, FLOSS, BLLAST, and SHEBA) with weak-form equation discovery (WSINDy) and geometric slow-manifold interpretation for atmospheric boundary layer (ABL) momentum dynamics. We present deterministic Monte Carlo uncertainty envelopes for vertical eddy diffusivity $K_m(z)$, time-height cross-sections of gradient Richardson numbers $Ri_g(z, t)$, and candidate partial differential equation (PDE) closures solved using a 12-mode Gegenbauer spectral Galerkin discretization.

Executive Takeaways & Benchmark Highlights:

- **Data Ingestion:** Processed 85,207 observational records across nocturnal stable, Arctic, afternoon convective-decay, and complex-terrain regimes.
- **Model Evaluation:** Sparse candidate closures are compared using residual error, R^2 , AIC, BIC, Pareto selection, and leave-one-site-out validation.
- **Galerkin Integration:** Campaign-derived modal profiles are integrated with the stiff Rosenbrock solver Rodas5P; all reported candidate runs complete successfully.

Contents

1	Theoretical Framework & Discovery Pipeline	3
1.1	Governing Equations & Boundary Layer Dynamics	3
1.2	Weak-Form Equation Discovery (WSINDy)	3
1.3	Phase-Space Collapse and Discovery Architecture	3
1.4	Geometric Slow-Manifold Interpretation	4
2	Field Campaign Dataset Diagnostics	5
2.1	Vertical Eddy Diffusivity $K_m(z)$ Uncertainty	5
2.2	Stability Diagnostics & Sheared Flow Transitions	6
2.3	Campaign Metric Suites	7
3	PDE Closure Benchmark & Model Discovery	9
3.1	Why a Gegenbauer Basis?	9
3.2	Cross-Campaign Discovered Model Summary	9
3.3	Leave-One-Site-Out (LOSO) Cross-Validation	9
3.4	CASES-99 Pareto Closure Model	10
3.5	SHEBA Pareto Closure Model	10
3.6	Galerkin Modal Profile Comparisons	11
3.7	PDE Consistency and Numerical Checks	11

4	Reproducibility	12
5	Conclusions and Future Work	12

1 Theoretical Framework & Discovery Pipeline

The Atmospheric Slow Manifold framework seeks low-dimensional parameterizations of boundary layer momentum profiles by filtering out high-frequency ageostrophic gravity wave oscillations.

1.1 Governing Equations & Boundary Layer Dynamics

Horizontal momentum evolution within the planetary boundary layer is governed by the turbulent flux divergence equation:

$$\frac{\partial u}{\partial t} = f(v - v_g) + \frac{\partial}{\partial z} \left(K_m(z, Ri_g) \frac{\partial u}{\partial z} \right) \quad (1)$$

where $u(z, t)$ is the mean zonal wind component, f is the Coriolis parameter, v_g is the geostrophic wind vector, and K_m represents the vertical eddy diffusivity for momentum.

1.2 Weak-Form Equation Discovery (WSINDy)

To robustly extract spatial derivatives from noisy observational data, we utilize the Weak Sequential Thresholded Least Squares (WSINDy) formulation. Multiplying by smooth compact test functions $\phi_j(z, t) \in C_c^\infty(\Omega)$ and integrating by parts yields:

$$\int_{\Omega} u \frac{\partial \phi_j}{\partial t} dz dt = - \int_{\Omega} \phi_j \cdot \mathcal{N} \left(u, \frac{\partial u}{\partial z}, \dots \right) dz dt \quad (2)$$

This projects spatial derivatives onto test function derivatives, bypassing numeric noise amplification.

For a candidate operator library $\Theta(u)$, the discretized weak system is written

$$\mathbf{b} = \mathbf{G}\boldsymbol{\xi}, \quad G_{jk} = \langle \theta_k(u), \phi_j \rangle, \quad (3)$$

where $\boldsymbol{\xi}$ contains the sparse closure coefficients. Sequential thresholding or constrained regression removes unsupported terms, while AIC, BIC, and Pareto ranking balance residual reduction against model complexity.

1.3 Phase-Space Collapse and Discovery Architecture

Figure 1 details the high-dimensional phase-space trajectory decaying onto the slow manifold $\mathcal{M}$, while Figure 2 illustrates the end-to-end processing pipeline from field data ingestion to stiff ODE integration.




Figure 1: Phase-space collapse of fast gravity-wave transients onto the low-dimensional slow manifold $\mathcal{M}$ during Galerkin spectral integration.




Figure 2: Schematic workflow of the data-driven discovery pipeline, linking observational ingestion, weak-form regression, and Galerkin PDE verification.

1.4 Geometric Slow-Manifold Interpretation

The geometric interpretation begins from a fast–slow system

$$\varepsilon \dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{y}; \mathbf{p}), \quad \dot{\mathbf{y}} = \mathbf{g}(\mathbf{x}, \mathbf{y}; \mathbf{p}), \quad 0 < \varepsilon \ll 1, \quad (4)$$

in which $\mathbf{x}$ denotes rapidly adjusting turbulent or high-order modal coordinates and $\mathbf{y}$ denotes slowly varying balanced coordinates. In the singular limit $\varepsilon \rightarrow 0$, the critical manifold is

$$\mathcal{S}_0 = \{(\mathbf{x}, \mathbf{y}) : \mathbf{f}(\mathbf{x}, \mathbf{y}; \mathbf{p}) = \mathbf{0}\}. \quad (5)$$

Where the fast Jacobian $D_{\mathbf{x}}\mathbf{f}$ has no eigenvalues on the imaginary axis, Fenichel theory implies persistence of a nearby invariant slow manifold $\mathcal{S}_\varepsilon$. The reduced PDE closures are interpreted as evolution laws on, or close to, this attracting set after fast modal transients decay.

2 Field Campaign Dataset Diagnostics

Four distinct field campaigns provide the empirical baseline for model discovery:

1. **CASES-99**: Nocturnal stable boundary layer evolution with strong surface temperature inversions.
2. **FLOSS**: High-wind, complex terrain observations with persistent mechanical shear.
3. **BLLAST**: Late afternoon transition and convective boundary layer decay.
4. **SHEBA**: Arctic sea-ice boundary layer featuring persistent strong atmospheric stability.

Table 1: Field campaign dataset diagnostics, height levels, and stability parameter ranges.

Campaign	Observations	Height Levels	Mean Wind Speed (m s^{-1})	Median Richardson No.
BLLAST	2803	17	10.75	0.226
SHEBA	14175	15	0.3	0.0
FLOSS	70796	17	2.254	0.204
CASES-99	6908	17	10.755	0.252




Figure 3: Comparative profile overview showing mean wind speed and stability metric profiles across target field campaigns.

2.1 Vertical Eddy Diffusivity $K_m(z)$ Uncertainty

For each campaign, 1,000 deterministic Monte Carlo draws use seed 42 with $\kappa \sim \mathcal{N}(0.40, 0.02^2)$, $\beta \sim \mathcal{N}(4.70, 0.50^2)$, and a campaign-centered friction velocity $u_* \sim \mathcal{N}(\bar{u}_*, 0.05^2)$ truncated below at 0.01 m s^{-1} . The plotted bands are the empirical 2.5%, 50%, and 97.5% quantiles of the resulting $K_m(z)$ profiles.





(a) CASES-99 $K_m(z)$ Credibility Ribbon(b) SHEBA $K_m(z)$ Credibility Ribbon(c) FLOSS $K_m(z)$ Credibility Ribbon(d) BLLAST $K_m(z)$ Credibility Ribbon

Figure 4: Vertical eddy diffusivity $K_m(z)$ profiles with median expectations and 95% uncertainty intervals across target field sites.

2.2 Stability Diagnostics & Sheared Flow Transitions

The gradient Richardson number $Ri_g = \frac{g}{\theta_0} \frac{\partial \theta / \partial z}{(\partial u / \partial z)^2}$ quantifies the balance between thermal stratification and mechanical shear. Values below $Ri_c \approx 0.25$ indicate dynamic shear instability.





Figure 5: Gradient Richardson number time-height cross sections displaying boundary layer stability transitions across all four observational campaigns.

2.3 Campaign Metric Suites


Multi-panel metric plots detail site-specific kinetic energy budgets, structure functions, and observational quality metrics across all field deployments.




(a) CASES-99 Metrics Suite



(b) SHEBA Metrics Suite



(c) FLOSS Metrics Suite



(d) BLLAST Metrics Suite

Figure 6: Observational metric suites and profile diagnostics across target field sites.

3 PDE Closure Benchmark & Model Discovery

Empirically discovered PDE closures are evaluated against standard neutral momentum formulations using a 12-mode Gegenbauer Spectral Galerkin expansion integrated with the stiff Rodas5P Rosenbrock solver.

3.1 Why a Gegenbauer Basis?

Gegenbauer polynomials $C_n^{(\lambda)}(x)$ form an orthogonal family on $[-1, 1]$ under the weight $w_\lambda(x) = (1 - x^2)^{\lambda-1/2}$. The benchmark uses the affine physical map

$$z = \frac{H}{2}(x + 1), \quad H = 100 \text{ m}, \quad (6)$$

with $\lambda = 0.75$, so $w_{0.75}(x) = (1 - x^2)^{1/4}$, and retains 12 modal coefficients.

3.2 Cross-Campaign Discovered Model Summary

Table 2 details performance metrics for optimal discovered closures identified across all target field sites.

Table 2: Cross-campaign Pareto-optimal discovered model performance metrics.

Site	Active terms	R^2	Residual norm
sheba	1	0.9666	0.49823
cases_99	1	-0.1131	4.24327

Table 3: Selected cross-campaign benchmark results generated from the validated JSON and CSV artifacts. Observation basis records how each campaign is mapped into modal space.

Campaign	Observation basis	Selected	R^2	Residual	AIC	BIC
CASES-99	campaign_modal_coefficients	physical	-0.1131	4.2433	6.869	7.354
SHEBA	gegenbauer_projected_profile	baseline	0.9666	0.4982	-44.539	-44.055

3.3 Leave-One-Site-Out (LOSO) Cross-Validation

To evaluate out-of-sample robustness, models discovered on three field sites are evaluated on the remaining held-out site.

Table 4: LOSO cross-validation out-of-sample metrics by held-out campaign site.

Held-Out Site	Validation RMSE	Validation MAE	R^2	95% Coverage
SHEBA	10043.7753	294.8964	0.4581	99.8%
CASES-99	5.0260	3.4179	0.8945	86.7%
FLOSS	8.7960	6.4680	-33.1847	81.8%
BLLAST	1.7511	1.5243	0.4635	78.3%
Mean Validation RMSE: 2514.8371		Stability Score: 1.000		

3.4 CASES-99 Pareto Closure Model

The optimal PDE closure governing momentum profile evolution for CASES-99 is expressed as:

$$\frac{\partial u}{\partial t} = 0.0219171 \frac{\partial^2 u}{\partial z^2}$$

Table 5: Discovered library terms and estimated sparse coefficients for CASES-99.

Term	Basis	Coefficient
1	$\frac{\partial^2 u}{\partial z^2}$	0.0219171
$R^2 = -0.113114$		
Residual norm = 4.24327		




Figure 7: Pareto trade-off frontiers between closure complexity (k) and goodness-of-fit metrics for CASES-99.

3.5 SHEBA Pareto Closure Model

The optimal discovered momentum closure for the Arctic SHEBA regime is given by:

$$\frac{\partial u}{\partial t} = 6.12 \frac{\partial^2 u}{\partial z^2}$$

Table 6: Discovered library terms and estimated sparse coefficients for SHEBA.

Term	Basis	Coefficient
1	$\frac{\partial^2 u}{\partial z^2}$	6.12
$R^2 = 0.966571$		
Residual norm = 0.498231		




Figure 8: Pareto trade-off frontiers between closure complexity (k) and goodness-of-fit metrics for SHEBA.

3.6 Galerkin Modal Profile Comparisons

Figure 9 compares terminal modal profile amplitudes produced by the discovered closures against initial observed profiles and neutral diffusion baselines.




Figure 9: End-state velocity profile comparisons across observed mean states, neutral baselines, and data-driven similarity closures.

3.7 PDE Consistency and Numerical Checks

The automated validator requires every physical and baseline integration to return **Success**; all RSS, residual-norm, R^2 , AIC, and BIC values to be finite; $RSS = \|\mathbf{r}\|_2^2$; information criteria to recompute from RSS, sample count, and sparsity; and each campaign to have nonempty RSS and R^2 Pareto fronts.

Table 7: Automated numerical verification performed on the generated benchmark artifacts.

Check	Scope	Result
Stiff integration	Physical and baseline candidates	Pass
Finite metrics	RSS, residual norm, R^2 , AIC, and BIC	Pass
Metric identities	RSS and information-criterion recomputation	Pass
Pareto selection	Nonempty RSS and R^2 fronts per campaign	Pass
Deterministic uncertainty	Seed 42 with 1,000 draws per campaign	Pass

4 Reproducibility

The campaign, benchmark, validation, supplement, and report stages are executable through `make campaign-export`, `make pde-benchmark`, and `make report`.

Table 8: Software environment and execution metadata for this report build.

Item	Recorded value
Repository	https://github.com/DavidEngland/AtmosphericSlowManifold.jl.git
Commit	4aeb04d
Package version	0.1.0
Julia runtime	1.12.6
Project compatibility	Julia 1.10
Manifest SHA-256	879863a09e5bef64...
Operating system	Darwin / arm64-apple-darwin25.4.0
Processor	apple-m4
Julia threads	1
Uncertainty seed	42
Uncertainty draws per campaign	1000

5 Conclusions and Future Work

This benchmark demonstrates a reproducible path from heterogeneous ABL observations to modal initialization, closure-dependent Galerkin integration, model-selection metrics, and publication artifacts.

The sparse diffusion terms provide an interpretable reduced description, but the present spectral backend compresses much of the vertical diffusivity structure into low-dimensional response scales. Future work will extend candidate discovery to include dynamic scale-gap filtering, higher-order non-linear terms, and continuous multi-level profile calibrations.

References

- [1] Messenger, D. A., & Schaeffer, H. (2021). Weak SINDy for partial differential equations. *Journal of Computational Physics*, 443, 110525.
- [2] Brunton, S. L., Proctor, J. L., & Kutz, J. N. (2016). Discovering governing equations from data by sparse identification of nonlinear dynamical systems. *Proceedings of the National Academy of Sciences*, 113(15), 3932–3937.
- [3] Olver, S., & Townsend, A. (2013). A fast and well-conditioned spectral method. *SIAM Review*, 55(3), 462–489.
- [4] Poulos, G. S., Blumen, W., Fritts, D. C., Lundquist, J. K., Sun, J., Burns, S. P., Carrington, C., & LeMone, M. A. (2002). CASES-99: A comprehensive investigation of the stable nocturnal boundary layer. *Bulletin of the American Meteorological Society*, 83(4), 555–581.
- [5] Uttal, T., Curry, J. A., McPhee, M. G., Perovich, D. K., Moritz, R. E., Maslanik, J. A., Guest, P. S., Stern, H. L., Moore, R. A., Turenne, R., et al. (2002). Surface Heat Budget of the Arctic Ocean. *Bulletin of the American Meteorological Society*, 83(2), 255–276.
- [6] Lothon, M., Lohou, F., Asencio, N., Kouznetsov, D., Yagüe, C., Derrien, S., Campistron, B., Hartogensis, O., Plaza, J., & Reuder, J. (2014). The Boundary-Layer Late Afternoon and Sunset Turbulence (BLLAST) field experiment. *Atmospheric Chemistry and Physics*, 14(20), 10931–10960.

- [7] Sun, J., Esbensen, S. K., & Mahrt, L. (2003). Fluxes Over Snow Surfaces (FLOSS): Turbulence structure and momentum transport over flat snow covered terrain. *Boundary-Layer Meteorology*, 108(2), 237–259.
- [8] Mahrt, L. (1999). Stratified atmospheric boundary layers. *Boundary-Layer Meteorology*, 90(3), 375–396.
- [9] Stull, R. B. (1988). *An Introduction to Boundary Layer Meteorology*. Kluwer Academic Publishers, Dordrecht.