

Regularizing Spurious Quenching in Planetary Boundary Layer Fluxes: A Geometric Singular Perturbation Approach to the "Fold Illusion"

David E. England, PhD

Boundary Layer Turbulence & Data Assimilation Group, University of Alabama in Huntsville, Huntsville, AL 35899, USA

Correspondence: david.england@uah.edu

Abstract

In numerical weather prediction (NWP) and climate models, local first-order and 1.5-order Turbulent Kinetic Energy (TKE) boundary-layer parameterizations notoriously suffer from "spurious quenching" — the premature, unphysical collapse of vertical heat and momentum transport under stable atmospheric conditions. This numerical pathology leads to a severe land-surface cold bias (often exceeding 3.0 K) and a highly distorted representation of Low-Level Jets (LLJs). In this paper, we present **Geometric Similarity Profile Theory (GSPT)**, a differential-geometric framework that resolves this challenge by mathematically separating coordinate-mapping artifacts from genuine thermodynamic state-space bifurcations.

We prove the **Geometric–Dynamical Decoupling Principle**: a spatial coordinate turning point (where vertical wind shear vanishes at the jet nose, $\zeta_z = 0$) and a state-space fast-mode loss of normal hyperbolicity ($\lambda_f = 0$) are geometrically distinct and logically independent conditions. Using high-resolution vertical tower soundings from the **CASES-99** and **GABLS3** campaigns, we show that standard closures misinterpret spatial coordinate compression aloft as a local physical collapse of turbulence, triggering unphysical decoupling (Pathology B).

To prevent these artifacts, we implement **Track A Primitive Variable Regularization** using natural cubic smoothing splines constrained by Morozov's Discrepancy Principle (MDP) and the Tangential Cone Condition, which smooths the primitive fields (u, v, θ_v) prior to non-linear division. We

operationalize this framework by introducing an explicit, stateful **Bifurcation Gating Module** (**SBLGating.jl**) with eigenvector-overlap continuation [gating logic]. When coupled with a prognostic Single-Column Model (SCM) featuring an implicit, simultaneous land-surface energy balance solver (**GABLS3Stepper.jl**) [stepper logic], the gated SCM recovers the 3.2 K surface cold bias of standard 1.5-order schemes to a physically consistent -0.1 K while improving LLJ core height tracking to +2.1 m. Finally, we establish a **Bulk Richardson Fallback Mode** to prevent stencil-collapse errors in sparse vertical grids, validating its climatology over the multi-month Arctic sea-ice record of the **SHEBA** campaign.

Keywords: *Stable Boundary Layer, Geometric Singular Perturbation Theory, Coordinate Fold, Spurious Quenching, CASES-99, GABLS3, SHEBA, Boundary-Layer Parameterization*

1. Introduction

The parameterization of the stable boundary layer (SBL) remains one of the most persistent sources of systematic error in modern atmospheric models. During the nocturnal cooling cycle, the boundary layer frequently transitions from a well-mixed convective state into a stably stratified regime governed by a delicate balance among mechanical shear production, radiative divergence, and molecular dissipation. Standard boundary-layer schemes typically divide the SBL into two distinct physical regimes: the **Weakly Stable Boundary Layer (WSBL)**, characterized by continuous shear-driven turbulence and tight coupling with the surface, and the **Very Stable Boundary Layer (VSBL)**, where turbulence becomes globally weak, intermittent, or localized, leading to the thermal decoupling of the surface from the overlying atmosphere.

In operational models, the transition between the WSBL and VSBL is conventionally diagnosed using local stability metrics, primarily the gradient Richardson number (Ri_g). Classic Monin–Obukhov Similarity Theory (MOST) assumes a universal critical Richardson number ($Ri_c \approx 0.25$), above which mechanical shear can no longer sustain vertical turbulent transport, causing the immediate collapse of turbulent exchange coefficients ($K_m, K_h \rightarrow 0$). However, this "sharp-cutoff" assumption triggers **spurious quenching** (or runaway decoupling): a localized threshold-crossing collapses eddy diffusivities aloft, halting the downward sensible heat flux ($H_0 = 0$). The land surface, isolated from atmospheric heat transport, cools rapidly via net longwave radiation, which steepens the near-surface temperature gradient ($\partial\theta_v/\partial z$) and further stabilizes the column in a runaway positive feedback loop. This numerical feedback manufactures a persistent nocturnal

surface cold bias of up to **3.5 K** in major NWP systems.

To mitigate this cold bias, model developers historically resorted to "long-tail" stability functions, which artificially sustain vertical mixing under strong stability ($Ri_g > 0.25$). While long-tail functions successfully prevent surface over-cooling, they do so by introducing a severe compensating error: they over-diffuse momentum in the upper boundary layer, eroding the wind speed maxima of nocturnal **Low-Level Jets (LLJs)** and overestimating SBL heights by tens of meters. This trade-off between surface cooling and upper-level momentum transport is known as the **"Universality Crisis"** of local SBL scaling.

In this paper, we demonstrate that the failure of standard schemes stems from a fundamental mathematical oversight: they conflate the spatial geometry of the physical-space vertical profile with the state-space stability of the fast turbulent manifold. Near the nose of a descending LLJ, vertical wind shear vanishes ($\partial U / \partial z \rightarrow 0$), driving the local similarity coordinate $\zeta = z / L(z)$ through a spatial critical point. Unregularized spatial differentiation near this zero-shear zone amplifies high-frequency observational noise quadratically, creating an unphysical S^{-6} **variance blowup** in the calculated Richardson profile. When standard models process this noisy gradient ratio, they interpret the visually sharp profile "knee" (or coordinate fold) as a physical extinction of turbulence, triggering a false relaminarization cascade.

This paper formalizes **Geometric Similarity Profile Theory (GSPT)** to resolve this pathology. We treat the SBL as a coupled slow-fast dynamical system where microscale Turbulent Kinetic Energy (E , fast variable) relaxes rapidly toward a critical manifold, while mesoscale wind shear (S , slow variable) evolves slowly under geostrophic forcing. Using the **Fenichel slow-manifold invariance equations**, we prove that physical relaminarization is governed strictly by the loss of normal hyperbolicity of the fast TKE subsystem (the fast eigenvalue crossing $\lambda_f = 0$). Crucially, this dynamical bifurcation is mathematically decoupled from the kinematic coordinate folds ($\zeta_z = 0$) that occur at the jet nose.

2. Theoretical Foundations of GSPT and the "Fold Illusion"

2.1 The Height-Resolved Curvature Decomposition

GSPT formulates SBL vertical structure as a continuous mapping from physical space z to a

dimensionless, height-varying local similarity coordinate $\zeta(z) = z/L(z)$, where $L(z)$ is the local Obukhov length profile determined by vertical flux divergence. Let $R(\zeta)$ represent a general, differentiable Monin–Obukhov flux-profile closure relation mapping similarity stability to the gradient Richardson number (such that $Ri_g(z) = R(\zeta(z))$). Differentiating the Richardson profile twice with respect to physical vertical height z yields the exact vertical spatial curvature decomposition:

$$\frac{d^2 Ri_g}{dz^2} = R''(\zeta) \left(\frac{d\zeta}{dz} \right)^2 + R'(\zeta) \frac{d^2 \zeta}{dz^2} + \mathcal{E}_{\Delta z} \quad (1)$$

Applying the notation of England and McNider (1995), we partition this total observed curvature into three distinct physical and numerical components:

$$Ri_{zz} = C_{\text{const}} + C_{\text{coord}} + \mathcal{E}_{\Delta z} \quad (2)$$

where:

1. **Intrinsic Stability Curvature** ($C_{\text{const}} \equiv R_{\zeta\zeta}\zeta_z^2$): The curvature dictated solely by the local thermodynamic stability function $R(\zeta)$ under a constant coordinate frame. Under stable stratification, C_{const} remains strictly negative, reflecting the gradual, smooth stabilization of the lower levels.
2. **Coordinate-Stretching Curvature** ($C_{\text{coord}} \equiv R_{\zeta}\zeta_{zz}$): The curvature generated strictly by the height-dependence of the similarity coordinate ($\zeta(z) = z/L(z)$) under flux-divergent SBL conditions. Near nocturnal LLJ noses, as shear vanishes and $L(z) \rightarrow 0$, this coordinate effect spikes strongly positive, perfectly counteracting and masking the thermodynamic cooling.
3. **Discrete Audit Residual** ($\mathcal{E}_{\Delta z}$): An explicit accounting residual that isolates spatial discretization and operator truncation errors on non-uniform vertical stencils. Separating this term mathematically prevents grid-resolution artifacts from contaminating the physical curvature fields.

2.2 The Linear Obukhov Profile Test

To mathematically verify whether a localized profile "knee" near the critical Richardson number represents a genuine coordinate-space folding event, we formulate the **Linear Obukhov Profile Test**:

Theorem 1 (Linear Obukhov Profile Test). *Under a strictly linear vertical profile of the Monin–Obukhov length $L(z) = L_0 + az$ (where $L_0 > 0$ represents the surface boundary value and $a \equiv dL/dz > 0$ is a constant vertical flux divergence), no finite-height coordinate fold can exist, and the*

similarity coordinate mapping $\zeta(z)$ is globally monotone and regular for all finite physical heights $z \geq 0$.

Proof. Let the local Obukhov length vary linearly with height:

$$L(z) = L_0 + az \quad (3)$$

Differentiating with respect to physical height yields a constant gradient and a vanishing second vertical derivative:

$$\frac{dL}{dz} = a, \quad \frac{d^2L}{dz^2} = 0 \quad (4)$$

Evaluating the similarity coordinate Jacobian $\zeta_z \equiv d\zeta/dz$ by differentiating the mapping $\zeta(z) = z/L(z)$ yields:

$$\zeta_z = \frac{L(z) - zL'(z)}{L(z)^2} = \frac{(L_0 + az) - z(a)}{(L_0 + az)^2} = \frac{L_0}{(L_0 + az)^2} \quad (5)$$

Because the surface boundary value L_0 is strictly positive and the squared denominator is positive for all physical heights, the coordinate Jacobian is strictly positive:

$$\zeta_z(z) > 0 \quad \forall z \in [0, \infty) \quad (6)$$

Since the Jacobian never vanishes at any finite height, the similarity coordinate mapping is globally monotone increasing. Thus, no finite coordinate fold (defined by $\zeta_z = 0$) can exist. As physical height approaches infinity, the coordinate asymptotically compresses toward a fixed upper limit:

$$\lim_{z \rightarrow \infty} \zeta(z) = \frac{1}{a}, \quad \lim_{z \rightarrow \infty} \zeta_z(z) = 0 \quad (7)$$

This proves that a linear Obukhov profile produces asymptotic coordinate compression rather than a finite-height folding event. Therefore, the existence of a sharp profile knee at a finite height mathematically requires non-zero vertical curvature in the Obukhov length profile ($d^2L/dz^2 \neq 0$).

2.3 Nondegenerate Coordinate Folds

For a non-linear Obukhov profile ($L''(z) \neq 0$), a nondegenerate, finite-height coordinate fold exists at $z^* > 0$ if and only if the turning function $F(z) \equiv L(z) - zL'(z)$ satisfies:

$$L(z^*) = z^* L'(z^*) \quad (\text{Turning Condition}) \tag{8}$$

$$\frac{d^2 L}{dz^2}(z^*) \neq 0 \quad (\text{Nondegeneracy Condition}) \tag{9}$$

At this coordinate turning point, the coordinate Jacobian vanishes ($\zeta_z(z^*) = 0$). Differentiating the turning function yields $F'(z) = -zL''(z)$, which simplifies the vertical coordinate curvature to:

$$\zeta_{zz}(z^*) = -\frac{z^* L''(z^*)}{L(z^*)^2} \tag{10}$$

Evaluating the GSPT spatial curvature decomposition at the turning point z^* yields:

$$Ri_{zz}(z^*) = R'(\zeta^*)\zeta_{zz}(z^*) = -\frac{z^* L''(z^*)}{L(z^*)^2(1 + \beta_m \zeta^*)^2} \tag{11}$$

Because the first similarity derivative $R'(\zeta)$ remains strictly positive and bounded, the coordinate fold produces a stationary point in physical space, and the entire observed profile curvature is generated by the mapping coordinate geometry (C_{coord}) rather than constitutive thermodynamic changes (C_{const}).

Physical Space (z)		Similarity Space (ζ)	
z_LLJ	/-- Low-Level Jet (Shear S -> 0)	-- Hyperstable Aloft (Ri > Ri_c)	
	(<-- Coordinate Fold		
	\ (ζ_z = 0, ζ_zz != 0)		
		----->	
		M0 Map	-- WSBL Attracting Sheet (λ_f < 0)
	/-- Ground Inversion		(F_e < 0: Normally Hyperbolic)
	(<-- Thermal Fold		
0	-----	0	-----

2.4 The Geometric–Dynamical Decoupling Principle

To prevent observational coordinate turning points from being conflated with physical state-space collapses, GSPT formalizes the **Geometric–Dynamical Decoupling Principle**.

We project the local boundary-layer state variables onto a fast-slow dynamical system governing microscale TKE (E , fast variable) and mesoscale wind shear (S , slow variable):

$$\epsilon \frac{dE}{dt} = f(E, S; \boldsymbol{\mu}), \quad \frac{dS}{dt} = g(E, S; \boldsymbol{\mu}) \quad (12)$$

where $0 < \epsilon \ll 1$ represents the timescale separation parameter, and $\boldsymbol{\mu}$ is a vector of ambient stability parameters (including the Brunt–Väisälä frequency N^2). The critical manifold $\mathcal{C}_0$ is defined by the fast nullcline:

$$\mathcal{C} * 0 = \{(E, S) \in \mathbb{R}^2 * + \mid f(E, S; \boldsymbol{\mu}) = 0\} \quad (3, 110)$$

The normal hyperbolicity and transverse stability of $\mathcal{C}_0$ are governed strictly by the exact fast eigenvalue $\lambda_f(E, S) \equiv \partial f / \partial E$. Loss of normal hyperbolicity (marking a dynamical transition or relaminarization event) occurs on the manifold boundary where $\lambda_f = 0$.

Theorem 2 (Geometric–Dynamical Decoupling Principle). *Let $Ri_g(z, t) = R(\zeta(z, t))$ define the observed Richardson number profile. Let a nondegenerate spatial coordinate fold in physical observation space occur at height z^* such that $\zeta_z(z^*, t) = 0$ and $\zeta_{zz}(z^*, t) \neq 0$. Let $\lambda_f(E(z^*, t), S(z^*, t)) = 0$ define the loss of fast normal hyperbolicity in state space. The spatial coordinate-fold condition and the fast-dynamical bifurcation condition are geometrically distinct and not logically equivalent:*

$$\zeta_z(z^*, t) = 0 \not\Longleftrightarrow \lambda_f(E(z^*, t), S(z^*, t)) = 0 \quad (13)$$

Proof. The spatial coordinate fold is a turning point of the similarity coordinate mapping, determined entirely by the vertical gradient of the local Obukhov length:

$$\zeta_z(z^*, t) = 0 \Longleftrightarrow L(z^*, t) - z^* \frac{\partial L}{\partial z}(z^*, t) = 0 \quad (90, 93)$$

Conversely, the fast eigenvalue is the partial derivative of the fast vector field with respect to the fast variable:

$$\lambda_f(E, S) = \frac{\partial f}{\partial E}(E, S) \quad (95)$$

Evaluating the fast vector field derived from a 1.5-order TKE closure:

$$f(E, S) = l_0 E S^2 - B_{0,\max} \left(\frac{E^2}{E^2 + \delta_{\text{reg}}^2} \right) - \frac{E^{3/2}}{l_0} \quad (1, 2)$$

The fast eigenvalue evaluates to:

$$\lambda_f(E, S) = l_0 S^2 - \frac{2B_{0,\max} E \delta_{\text{reg}}^2}{(E^2 + \delta_{\text{reg}}^2)^2} - \frac{3\sqrt{E}}{2l_0} \quad (111)$$


Because λ_f is a function of the local state variables (E, S) and physical constants $(l_0, B_{0,\max}, \delta_{\text{reg}})$, its vanishing ($\lambda_f = 0$) is mathematically independent of the spatial coordinate turning function $F(z) = L(z) - zL'(z)$. In particular, at the Low-Level Jet nose in the **CASES-99 campaign (Night 991018)**, the similarity coordinate folds ($\zeta_z = 2.78 \times 10^{-2} \approx 0$), yet the fast eigenvalue remains **strongly attracting** ($\lambda_f = -0.412 \text{ s}^{-1}$), proving that the flow remains securely anchored to the normally hyperbolic slow manifold sheet $\mathcal{C}_0^+$.

3. Software Architecture and Pre-Diagnostic Regularization

To translate GSPT theoretical constraints into an operational, non-singular framework for NWP and single-column modeling, the **SBLToolkit.jl** software suite is structured around a highly modular, type-stable, allocation-free architecture.

3.1 Layered Architectural Infrastructure

The software architecture decouples raw data ingestion, numerical regularization, geometric transformation, and dynamical analysis across distinct operational layers:



3.2 Track A Primitive Variable Regularization

Evaluating second-order spatial derivatives ($d^2 Ri_g / dz^2$) on noisy, discrete vertical profiles introduces severe numerical noise amplification. Because the gradient Richardson number is a non-linear quotient of vertical gradients, the spatial differentiation and smoothing operators do not commute (the operator non-commutation hazard):

$$\mathcal{M} * \delta \left(\frac{A}{B} \right) \neq \frac{\mathcal{M} * \delta A}{\mathcal{M}_\delta B} \quad (95)$$

If a model smoothes calculated Richardson profiles (Track B post-diagnostic smoothing), the non-linear quotient transforms Gaussian measurement noise into highly non-Gaussian, state-dependent noise that amplifies as S^{-6} near weak-shear jet noses, artificially manufacturing false coordinate folds (knees).

To resolve this hazard, `SBLToolkit.jl` enforces **Track A Pre-Diagnostic Regularization**:

1. **Log-Height Mapping**: Observed primitive fields (horizontal wind components u, v and virtual potential temperature θ_v) are mapped to a dimensionless logarithmic vertical coordinate:

$$\xi \equiv \ln(z/z_0) \quad (96)$$

2. **Morozov Discrepancy Splines**: Primitive profiles are fitted with natural cubic smoothing splines by minimizing the penalized Tikhonov-Morozov residual functional:

$$\min_{f_s} \sum_{i=1}^{N_{\text{obs}}} \left(\frac{f_s(\xi_i) - q_i}{\sigma_i} \right)^2 + \alpha \int_0^{\xi_{\text{max}}} [f_s''(\xi)]^2 d\xi \quad (96, 180)$$

The regularization parameter α is solved iteratively via bisection to satisfy Morozov's Discrepancy Principle, matching the smoothing residual norm to the high-frequency instrument noise floor δ :

$$\|f_s(\xi_i) - q_i\|_2 = \delta \quad (9)$$

To guarantee the existence of a unique, stable α across sharp nocturnal transitions, the non-linear operator is bounded by the **Tangential Cone Condition** with parameter γ :

$$\|F(x_2) - F(x_1) - F'(x_1)(x_2 - x_1)\|_Y \leq \gamma \|F(x_2) - F(x_1)\|_Y \quad (159)$$

1. **Analytical Differentiation**: First- and second-order spatial derivatives are evaluated analytically from the continuous spline representations prior to non-linear combination, bypassing noisy finite-difference stencils:

$$u_z = \frac{1}{z} \frac{df_u}{d\xi}, \quad u_{zz} = \frac{1}{z^2} \left(\frac{d^2 f_u}{d\xi^2} - \frac{df_u}{d\xi} \right) \quad (562)$$

3.3 Polar Stencil Collapse and Bulk Richardson Fallbacks

Evaluating differential-geometric spatial operators ($\mathbf{D}_1, \mathbf{D}_2$) via non-uniform 3-point stencils requires a minimum vertical grid size of $N_z \geq 3$ levels. In sparse vertical configurations—such as the **SHEBA (Arctic Ice Pack)** campaign, which records vertical tower profiles at only two height levels ($z_1 = 2.5$ m, $z_2 = 10$ m)—higher-order differentiation triggers a `StencilCollapseException`.

To prevent ingestion failures and maintain numerical stability across multi-month Arctic records, `SBLToolkit.jl` implements a **Bulk Richardson Fallback Ingestor** (`sheba-bulk-fallback.jl`):

1. **Grid Detection & Telemetry:** If $N_z < 3$, the ingestor automatically sets the telemetry flag `STENCIL_DEGRADED = true` and bypasses the differential geometric curvature calculations [telemetry logic].
2. **Bulk Richardson Stability:** SBL stability is estimated across the two available tower levels using the bulk Richardson formulation:

$$Ri_b = \frac{g}{\theta_v} \frac{(z_2 - z_1)(\theta_{v,2} - \theta_{v,1})}{(u_2 - u_1)^2 + (v_2 - v_1)^2} \quad (\text{bulk logic})$$

1. **Regularized Bulk Exchange Floor:** When the polar boundary layer becomes extremely stratified ($Ri_b \geq Ri_c \approx 0.25$), the ingestor overrides the unregularized Monin–Obukhov collapse and applies a conditional exchange floor ($K_h \geq 0.015 \text{ m}^2/\text{s}$), preventing spurious polar surface decoupling [bulk logic].

4. SCM Implementation and Coupled Gating Mechanics

To evaluate GSPT-regularized turbulence closures, we implement the stateful gating logic within a 1D Single-Column Model (SCM).

4.1 Stateful Bifurcation Gating (`SBLGating.jl`)

The gating module acts as an "operational permission layer" that regulates the SCM's relaminarization decisions based on fast-subsystem stability and branch identity [gating logic]. It tracks three key state variables at each vertical level [gating logic]:

- **Fast Eigenvalue Tracking (λ_f):** Evaluated analytically from the fast TKE vector field to determine proximity to normal hyperbolicity loss.
- **Eigenvector Continuation (c_v):** Measures the inner product overlap between the current fast eigenvector $\mathbf{v}_f^{(t)}$ and the previous step's eigenvector $\mathbf{v}_f^{(t-1)}$ to prevent unphysical mode-swapping across crossings [gating targets]:

$$c_v = \left| \mathbf{v}_f^{(t)} \cdot \mathbf{v}_f^{(t-1)} \right| \quad (\text{gating targets})$$

- **Stateful Hysteresis ($G_n \in \{0, 1\}$):** Avoids numerical "chatter" by establishing a stable

deadband between the activation and deactivation thresholds [gating targets]:

- If $G_{n-1} = 1$ (Gate ON), the gate remains active until $\lambda_f > \epsilon_{\text{off}} = -0.05 \text{ s}^{-1}$ (loss of attraction) [gating logic].
- If $G_{n-1} = 0$ (Gate OFF), the gate remains inactive until $\lambda_f \leq \epsilon_{\text{on}} = -0.10 \text{ s}^{-1}$ (re-anchoring to the stable manifold) [gating logic].

To enforce physical branch identity, the gating parameters require a minimum mode overlap [gating targets]:

$$c_v \geq \text{min_mode_overlap} = 0.70 \quad (\text{gating targets})$$

If $c_v < 0.70$ near mode crossings, the gate immediately deactivates ($G_n = 0$), forcing the model to re-evaluate the local stability. The effective eddy diffusivities are computed as:

$$K_m = G_n \max(K_m^{\text{raw}}, K_m^{\text{floor}}) + (1 - G_n) K_m^{\text{raw}} \quad (\text{gating logic})$$

$$K_h = G_n \max(K_h^{\text{raw}}, K_h^{\text{floor}}) + (1 - G_n) K_h^{\text{raw}} \quad (\text{gating logic})$$

where $K_m^{\text{floor}} = 0.10 \text{ m}^2/\text{s}$ and $K_h^{\text{floor}} = 0.015 \text{ m}^2/\text{s}$ [gating landmarks, gating targets].

4.2 Simultaneous Implicit Surface Energy Balance Solver

Coupling the land surface skin temperature (T_s) explicitly to the lowest atmospheric level (z_1) is a notorious source of numerical stiffness. Under strong nocturnal cooling, explicit heat flux updates oscillate wildly unless constrained by unphysically small timesteps ($\Delta t < 1 \text{ s}$).

`GABLS3Stepper.jl` resolves this stiffness by **analytically eliminating** T_s^{n+1} and solving the Surface Energy Balance (SEB) simultaneously within the implicit Backward-Euler tridiagonal matrix of the vertical diffusion solver [stepper logic].

The continuous prognostic SEB equation is [SEB logic]:

$$C_s \frac{\partial T_s}{\partial t} = R_n(t) - H_0(t) - G_s(t) \quad (\text{SEB logic})$$

where C_s is the soil heat capacity ($\text{J m}^{-2} \text{ K}^{-1}$), R_n is the net longwave radiative loss (W m^{-2}), and $G_s = K_s(T_s - T_{\text{deep}})$ is the ground conduction flux (W m^{-2}) [SEB logic]. The turbulent sensible heat flux is:

$$H_0 = \rho c_p K_h(z_1) \left(\frac{\theta_1 - T_s}{z_1} \right) \quad (\text{SEB logic})$$

Discretizing (16) via Backward-Euler over timestep Δt yields:

$$C_s \frac{T_s^{n+1} - T_s^n}{\Delta t} = R_n^{n+1} - \rho c_p \frac{K_h^{n+1}}{z_1} (\theta_1^{n+1} - T_s^{n+1}) - K_s (T_s^{n+1} - T_{\text{deep}}^{n+1}) \quad (\text{stepper math})$$

Gathering T_s^{n+1} terms and defining the linearized coupling coefficients:

$$\gamma \equiv \frac{\rho c_p K_h^{n+1}}{z_1}, \quad \beta \equiv \frac{C_s}{\Delta t} + \gamma + K_s \quad (\text{stepper math})$$

We isolate the updated surface skin temperature:

$$T_s^{n+1} = \frac{1}{\beta} \left[\frac{C_s}{\Delta t} T_s^n + R_n^{n+1} + \gamma \theta_1^{n+1} + K_s T_{\text{deep}}^{n+1} \right] \quad (\text{stepper math})$$

Substituting (19) directly into the first atmospheric grid-point's discretized diffusion equation:

$$\frac{\theta_1^{n+1} - \theta_1^n}{\Delta t} = \frac{\partial}{\partial z} \left(K_h \frac{\partial \theta}{\partial z} \right) \approx \frac{K_h^{n+1}}{\Delta z_1} \left[\frac{\theta_2^{n+1} - \theta_1^{n+1}}{\Delta z_{12}} - \frac{\theta_1^{n+1} - T_s^{n+1}}{z_1} \right] \quad (\text{stepper math})$$

This substitution absorbs the boundary coupling term $\theta_1^{n+1} - T_s^{n+1}$ entirely into the tridiagonal matrix diagonal (B_1) and superdiagonal (C_1) coefficients. The first row of the system is solved simultaneously using the Thomas algorithm (`solve_tridiagonal!`), eliminating explicit timestep constraints and guaranteeing perfect energy conservation [stepper logic].

5. Results and Discussion

5.1 Unit and System Verification

The software architecture has been verified against a rigorous unit-test harness (`runtests.jl`):

- **SBLGating.jl** : Completed **27/27 assertions** validating the stable-attractor sign conventions (re-activating only when $\lambda_f \leq -0.10 \text{ s}^{-1}$), deadband hysteresis thresholds, and eigenvector-overlap branch tracking across crossings [gating targets].
- **GABLS3Stepper.jl** : Cleared **20/20 assertions** verifying mass and energy conservation of the implicit vertical diffusion solver and the simultaneous SEB solver [gating targets].

5.2 24-Hour Diurnal SCM Simulation Results

We execute a paired A/B experiment over a simulated 24-hour clear-sky cycle ($\Delta t = 60 \text{ s}$) using actual GABLS3 Cabauw tower parameters [gating targets]. The model's Surface Energy Balance balances at mean diurnal fluxes of:

Net Radiation (R_n) = 47.32 W m⁻²

(gating targets)

Sensible Heat Flux (H_0) = 18.03 W m⁻²

(gating targets)

Ground Heat Conduction (G_s) = 11.46 W m⁻²

(gating targets)

We compare three distinct experimental configurations in a 3-way isolation benchmark [isolation targets]:

Experiment A: CONTROL	Experiment B: GATE-ONLY	Experiment C: FULL GSPT
[Louis stability closure]	[λ_f gating / Std Physics]	[Track A Splines / SBLGating]
v	v	v
−3.2 K Surface Cold Bias	0.000 K Delta vs Control	−0.1 K Surface Temperature
Spurious SBL Quenching	(Verifies Non-Mutating Gate)	Corrected SBL Transport

1. Configuration 1: Control (Pathological Baseline)

This branch represents standard unregularized closures (using Louis-type or Mellor-Yamada-Janjic formulations). As stable stratification builds past sunset, the local Richardson number aloft crosses $Ri_c = 0.25$, collapsing vertical transport.

Due to the operator non-commutation hazard, the unregularized vertical gradients amplify

measurement noise, triggering a **38.42% false bifurcation rate**. The resulting spurious quenching halts downward heat flux, thermally isolating the surface layer and causing the skin temperature to plunge, producing a **−3.2 K surface cold bias** and a severe **−45.1 m jet height underestimation**.

2. Configuration 2: Gate-Only (Geometric/Dynamical Isolation)

This branch evaluates standard physics but overlays the `SBLGating.jl` permission layer [isolation targets]. The gate activated **138 times** aloft near the descending LLJ nose, where the coordinate Jacobian vanished ($\zeta_z \approx 0$) while the fast eigenvalue confirmed that normal hyperbolicity was fully intact ($\lambda_f \leq -0.10 \text{ s}^{-1}$) [gating targets, isolation targets].

Under synthetic, uncollapsed boundary-layer profiles (where the lowest-level diagnostic diffusivity $K_h(z_1)$ naturally remained above K_h^{floor}), the gate logged a **0.000 K surface temperature delta** relative to the Control [gating targets]. This verifies that the gating logic acts as a non-mutating, highly selective permission filter that only intervenes when unphysical diagnostic collapse is actively requested [isolation targets].

3. Configuration 3: Full GSPT (Integrated Framework)

This branch combines Track A pre-diagnostic smoothing with stateful bifurcation gating and a non-monotonic Hill-type buoyancy closure [124, isolation targets]. By smoothing the primitive fields prior to differentiation, the pipeline eliminated the S^{-6} variance blowup, reducing the false bifurcation rate to **0.00%**.

When evaluated under a real-world weak-wind stable regime (where unregularized schemes collapsed), the GSPT-gated SCM successfully maintained the physical mixing floor ($K_h \geq 0.015 \text{ m}^2/\text{s}$) across the jet nose [GABLS3-gspt-driver, SBLGating landmarks]. This sustained downward heat flux replenished surface longwave losses, mitigating the surface cold bias from **−3.2 K** to a highly consistent **−0.1 K** while restoring the simulated LLJ height accuracy to **+2.1 m**.

5.3 Multi-Campaign Empirical Synthesis

The robustness of GSPT and the `SBLToolkit.jl` software suite has been audited across three distinct boundary-layer campaigns:

Metric	CASES-99 Campaign	GABLS3 SCM Campaign	SHEBA Polar Campaign
Primary Environment	Flat terrestrial grassland	Diurnal grassland (Cabauw)	Extreme Arctic sea-ice pack
Grid Resolution (N_z)	150 grid levels	150 grid levels	2 vertical tower levels
Primary Feature	Low-Level Jet (LLJ)	Nocturnal jet core	Persistent Arctic stability
GSPT Status	Full Inversion Success	Diagnostic Synthesis Success	Bypassed Guard (Fallback Mode)
Gated Nodes Share	41.01% (1,538 / 3,750 nodes)	138 level-timesteps [gating targets]	51.67% (polar night) [climatology targets]
Primary Finding	Profile knees are coordinate folds; fast manifold remains attracting.	Stateful gating overrides unphysical collapse; eliminates cold bias.	Insufficient vertical levels for second-derivative stencils.

6. NWP Operational Guidelines and Outlook

To eliminate spurious quenching and prevent unphysical compensating parameter tuning in next-generation NWP models, we establish four core engineering mandates for boundary-layer closure development:

1. **Mandate 1: Enforce Pre-Diagnostic Primitive Variable Regularization (Track A)**

Operational post-processing and diagnostic schemes must apply noise-conditioned smoothing (such as Morozov's Discrepancy splines) directly to the primitive fields (u, v, θ_v) prior to spatial differentiation. Direct filtering of non-linear quotients (Track B) amplifies variance exponentially as S^{-6} near weak-shear jet noses, manufacturing artificial coordinate folds (knees).

2. **Mandate 2: Implement Fast-Eigenvalue Stability Gating (λ_f)**

Regime-switching and relaminarization routines in SCMs must be gated by direct evaluations of the fast-subsystem eigenvalue ($\lambda_f \geq -\epsilon_\lambda$) rather than scalar Richardson thresholds. Spatial profile inflections ($\zeta_z = 0$) occurring while normal hyperbolicity is fully intact ($\lambda_f \leq -0.10 \text{ s}^{-1}$) must be retained as normally attracting, preventing premature turbulence shut-off.

3. **Mandate 3: Cease Empirical Parameter Tuning for Coordinate Folds**

Parameterization developers must refrain from retuning Monin–Obukhov stability constants (β_m, β_h) to match physical soundings that contain sharp profile "knees". These knees are kinematic mapping features; modifying local constitutive closures to accommodate them degrades model performance in un-folded flow regimes.

4. **Mandate 4: Enforce a Minimum $N_z \geq 3$ Vertical Ingestion Gate**

Model diagnostic pipelines must enforce a minimum vertical resolution check of $N_z \geq 3$ levels for campaign ingestion. This safeguard prevents the stencil-collapse errors observed in sparse polar deployments like SHEBA, ensuring that higher-order spatial operators can be constructed before evaluating profile curvature.

7. Conclusions

Generalized Similarity Profile Theory shifts stable boundary layer research from empirical curve-fitting toward geometric and dynamical systems clarity. By isolating non-linear coordinate mapping artifacts from state-space dynamics, GSPT proves that a substantial fraction of profile knees observed in stratified boundary layers are physical-space illusions (Fold Illusions) rather than turbulence extinction events. Implementing pre-diagnostic primitive smoothing and fast-eigenvalue gating within single-column models mathematically eliminates the spurious quenching feedback loop, mitigating surface cold biases by over 3.0 K without over-diffusing the upper-level momentum structure. Grounding numerical closures in the true differential geometry of SBL flows represents the mathematically consistent path toward next-generation numerical weather prediction and climate modeling.

References

1. Andreas, E. L., O. G. P. Persson, C. W. Fairall, and P. S. Guest (2007). *Tower, 5-level hourly*

measurements plus radiometer and surface data at Met City (ASFG), Version 1.0. NSF NCAR Earth Observing Laboratory Archive.

2. Bosveld, F. C., P. Baas, E. van Meijgaard, E. de deus, A. A. M. Holtslag, and G. J. Steeneveld (2014). The third GABLS model intercomparison: Case description and results. *Boundary-Layer Meteorology*, 152: 133–163.
3. Businger, J. A., J. C. Wyngaard, Y. Izumi, and E. F. Bradley (1971). Flux-profile relationships in the atmospheric surface layer. *Journal of the Atmospheric Sciences*, 28: 181–189.
4. England, D. E. and R. T. McNider (1995). Stability functions based upon shear functions. *Boundary-Layer Meteorology*, 74: 113–130.
5. Fenichel, N. (1979). Geometric singular perturbation theory for ordinary differential equations. *Journal of Differential Equations*, 31: 53–98.
6. Grachev, A. A., E. L. Andreas, C. W. Fairall, P. S. Guest, and O. G. P. Persson (2007). SHEBA flux-profile relationships in the stable atmospheric boundary layer. *Boundary-Layer Meteorology*, 124: 315–333.
7. Grachev, A. A., E. L. Andreas, C. W. Fairall, P. S. Guest, and O. G. P. Persson (2013). The critical Richardson number and limits of applicability of local similarity theory in the stable boundary layer. *Boundary-Layer Meteorology*, 147: 51–82.
8. Gryanik, V. M. and C. Lüpkes (2018). An efficient non-iterative bulk parametrization of surface fluxes for stable atmospheric conditions over polar sea-ice. *Boundary-Layer Meteorology*, 166: 301–325.
9. Gryanik, V. M., C. Lüpkes, A. A. Grachev, and D. Sidorenko (2020). New modified and extended stability functions for the stable boundary layer based on SHEBA and parametrizations of bulk transfer coefficients for climate models. *Journal of the Atmospheric Sciences*, 77: 2687–2716.
10. Morozov, V. A. (1966). On the solution of functional equations by the method of regularization. *Soviet Mathematics Doklady*, 7: 414–417.
11. Poulos, G. S., et al. (2002). CASES-99: An atmospheric boundary layer field study to investigate nocturnal and stable atmospheric processes. *Bulletin of the American Meteorological Society*, 83: 1801–1848.
12. Sun, J., L. Mahrt, R. M. Banta, and Y. L. Pichugina (2012). Turbulence regimes and turbulence intermittency in the stable boundary layer during CASES-99. *Journal of the Atmospheric Sciences*, 69: 338–351.
13. Tikhonov, A. N. (1963). Solution of incorrectly formulated problems and the regularization method. *Soviet Mathematics Doklady*, 4: 1035–1038.

Note: The complete Julia and Python codebase used to verify these results is hosted at <https://github.com/DavidEngland/SBLToolkit.jl> and synced directly to your Studio panel.