

classNumber $Q(\sqrt{-143}) = 10$ **Lean 4 Formal Proof Certificate**

Issued: June 23, 2026

1. Project and Proof Summary

This certificate documents the Lean 4 formal proof of $\text{classNumber } Q(\sqrt{-143}) = 10$ developed under the Morningstar Project (Theorema Aureum 143, Volume I).

The number field $K = Q(\sqrt{-143}) = \text{AdjoinRoot}(X^2 + 143)$ has discriminant -143 , ring of integers $Z[w]$ where $w = (-1 + \sqrt{-143})/2$ (satisfying $w^2 - w + 36 = 0$), Minkowski bound < 8 , and class number $h(K) = 10$.

Lean toolchain: Lean 4, Mathlib v4.12.0. No sorry. No axioms beyond {propext, Classical.choice, Quot.sound}.

2. Proved Bricks (0 sorry, classical trio)

All theorems below are sorry-free with classical trio axiom footprint only.

[PROVED]

BSD_classNumber_lower_bound

 $10 \leq \text{classNumber } K$ (unconditional).Proof: $\text{orderOf}([p_2]) = 10 \leq \text{Fintype.card}(\text{ClassGroup}) = \text{classNumber } K$.**[PROVED]**

span_gen_OK_eq_p2_pow_10

 $\text{span}\{-28+3w\} = p_2^{10}$ in $Z[w]$. A 7-step ideal-arithmetic chain:(1) gen_OK has norm $1024 = 2^{10}$ (norm_num).(2) $p_2 = \text{span}\{2, w\}$ is prime ($Z \bmod 2$ ring map).(3) $p_2 * p_2' = \text{span}\{2\}$ (explicit Z -combination).(4) $\text{Norm}(p_2') = 2$ (map_pow + absNorm_p2_eq_2).(5) $\text{span}\{\text{gen_OK}\} + p_2' = \text{top}$ (IsMaximal + coprimeness).(6) $p_2^{10} \leq \text{span}\{\text{gen_OK}\}$ (Euclid step via binomial).(7) $J = \text{top}$ from $\text{absNorm}(J) = 1$ (Ideal.absNorm_eq_one_iff).**[PROVED]**

BSD_p2_pow_10_principal

BSD_p2_pow_10_principal_hyp discharged.

 p_2^{10} is principal; generator = $-28+3w$, norm = $2^{10} = 1024$.**[PROVED]**

BSD_orderOf_p2_eq_10

 $\text{orderOf}([p_2]) = 10$ in $\text{ClassGroup}(Z[w])$.Divides 10: p_2^{10} is principal $\Rightarrow \text{mk0}(p_2)^{10} = 1$ in ClassGroup .Exceeds 9: master_not_principal_1_to_9 shows p_2^k non-principal for $k=1..9$.**[PROVED]**

BSD_classGroup_gen_by_p2_of_card_10

 $\text{classNumber } K = 10 \wedge p_2^{10} \text{ principal} \Rightarrow \text{ClassGroup}(Z[w]) = \langle [p_2] \rangle$.Proof: $\text{Fin } 10 \rightarrow \text{ClassGroup}, k \mapsto [p_2]^k$, injective ($\text{orderOf}=10 \Rightarrow \text{distinct}$).Domain and codomain have equal card (both 10) $\Rightarrow$ surjective.Hence every x in ClassGroup is a power of $[p_2]$.**[CONDITIONAL]**

BSD_classGroup_gen_by_p2_of_bqf

 $\text{BSD_BQF_ClassNumber_bridge} \Rightarrow \text{ClassGroup}(Z[w]) = \langle [p_2] \rangle$.**[CONDITIONAL]**

BSD_classNumber_eq_10_one_gate

 $\text{BSD_BQF_ClassNumber_bridge} \Rightarrow \text{classNumber } K = 10$ (ideal route).Chain: $\text{bridge} \rightarrow \text{classNumber } K = 10 \rightarrow \text{ClassGroup generated by } [p_2] \rightarrow \text{upper bound}$.Combined with $\text{BSD_classNumber_lower_bound}$: equality.**[CONDITIONAL]**

BSD_classNumber_via_bqf_bridge

 $\text{BSD_BQF_ClassNumber_bridge} \Rightarrow \text{classNumber } K = 10$ (BQF count route).

Directly: bridge says $\text{classNumber } K = \text{length}(\text{reducedForms143}) = 10$.

3. Named Open Surface

One named open surface remains. It is a def Prop (not proved, not an axiom):

[OPEN]

BSD_BQF_ClassNumber_bridge

$\text{NumberField.classNumber } K = \text{reducedForms143.length} (= 10)$.

Mathematical content: the Gauss-Dirichlet bijection between $SL_2(\mathbb{Z})$ -equivalence classes of primitive binary quadratic forms of discriminant -143 and ideal classes in $\text{ClassGroup}(\mathbb{Z}[w])$.

Lean gap: $\text{BinaryQuadraticForm.classGroupEquiv}$ is absent from Mathlib v4.12.0.

What IS proved: the 10 reduced BQFs are fully enumerated and proved complete ($\text{BSD_allReducedForms143}$, $\text{BSD_numReducedForms143} = 10$). Only the formal bijection to ClassGroup is missing.

4. Proof Chain Architecture

UNCONDITIONAL (0 sorry, 0 named open surfaces):

```
norm_form_gen_1024_BSD      : (-28)^2 + (-28)*3 + 36*3^2 = 1024
master_not_principal_1_to_9 : p2^k non-principal for k = 1..9
span_gen_0K_eq_p2_pow_10    : span{-28+3w} = p2^10 [7-step chain]
BSD_p2_pow_10_principal     : BSD_p2_pow_10_principal_hyp discharged
BSD_orderOf_p2_eq_10        : orderOf([p2]) = 10
BSD_classNumber_lower_bound : 10 <= classNumber K
```

CONDITIONAL (0 sorry, 1 named open surface: BSD_BQF_ClassNumber_bridge):

```
BSD_classGroup_gen_by_p2_of_bqf : bridge => ClassGroup = <[p2]>
BSD_classNumber_eq_10_one_gate  : bridge => classNumber K = 10 [ideal route]
BSD_classNumber_via_bqf_bridge  : bridge => classNumber K = 10 [BQF route]
```

5. Arithmetic Evidence (proved by norm_num / decide)

Splitting of 2: $w^2 - w + 36 = 0 \pmod{2} \rightarrow 2$ splits; $p_2 = \text{span}\{2, w\}$, $p_2' = \text{span}\{2, w-1\}$

Splitting of 3: $w^2 - w + 36 = 0 \pmod{3} \rightarrow 3$ splits; $p_3 = \text{span}\{3, w\}$

Splitting of 7: $w^2 - w + 36 = 0 \pmod{7} \rightarrow 7$ splits

Inertness of 5: $w^2 - w + 36 = 0 \pmod{5}$ has no solution $\rightarrow 5$ is inert

Norm form: $a^2 + ab + 36b^2 = 1$ has no integer solution (p_2 is non-principal)

Generator norm: $(-28)^2 + (-28)*3 + 36*3^2 = 1024 = 2^{10}$

Minkowski bound: $(2/\pi)*\sqrt{143} < 8 \rightarrow$ every class has representative of norm ≤ 7

absNorm(p2): $\text{Ideal.absNorm}(p_2) = 2$ (proved via ZMod ring map)

6. GitHub Repositories

ClassNumber-143: [DavidFox998/ClassNumber-143](https://github.com/DavidFox998/ClassNumber-143) -- Self-contained class number proof (BSD namespace)

BSD Tower: [DavidFox998/Birch-and-Swinnerton-Dyer](https://github.com/DavidFox998/Birch-and-Swinnerton-Dyer) Full BSD tower mirror (Towers.BSD namespace)

7. Axiom Footprint

Every proved brick satisfies:

- Axioms = {propext, Classical.choice, Quot.sound} (classical trio only).
- No sorry / admit / sorryAx in any landed brick.
- No new research-grade axioms.

The conditional combinators (BSD_classNumber_eq_10_one_gate, etc.) are also sorry-free; they thread the single named open surface as an explicit hypothesis, never as an axiom.