

Math 313/623 Notes 7

David L. Meretzky

Tuesday March 26th, 2019

We're gonna rock around the
clock tonight

Bill Haley

In these notes we will investigate some important arithmetic functions and the Möbius Inversion Formula.

Back to Gauss Section 2

Solutions of Congruences of the first degree

Article 26. In the case $(a, m) = 1$ we know that $ax + b \equiv c \pmod{m}$. Suppose that x_0 solves this congruence, that is $ax_0 + b \equiv c \pmod{m}$. It should follow from article 9 that if for any $x_1 \in \mathbb{Z}$ such that $x_1 \equiv x_0 \pmod{m}$, x_1 is also a solution to the congruence. Furthermore, if t is any other root of the congruence $ax + b \equiv c \pmod{m}$, then it must be congruent to x_0 . Spelling it all out, if $(a, m) = 1$, then given any solution x_0 , all solutions are given by the set $x_0 + m\mathbb{Z}$.

I will now say the same thing in about 5 different ways. Pick whichever way makes sense to you. This justifies why we say that the equation $ax + b \equiv c \pmod{m}$ only has a single root. It's "root" is the set of all numbers congruent to x_0 modulo m . More precisely, there is exactly one least residue of x_0 in the set $x_0 + m\mathbb{Z}$. More modernly, $x_0 + m\mathbb{Z}$ is a single element of $\mathbb{Z}/m\mathbb{Z}$, usually identified with the aforementioned least residue.

Note that these results do not hold if $(a, m) \neq 1$ or if the degree of the congruence is greater than 1, say $ax^2 + b \equiv c \pmod{m}$.

Proof. I wouldn't dare. □

Article 27. Gauss makes a delightful but somewhat hard to parse observation that if we want to solve a congruence of the form $ax + t \equiv u \pmod{b}$, where now $(a, b) = 1$, we only need to solve $ax \equiv \pm 1 \pmod{b}$. We can easily check that that if there is some solution r such that $ar \equiv \pm 1 \pmod{b}$, then $\pm(u - t)r$ will be a solution to $ax + t \equiv u \pmod{b}$. That is,

$$a \pm (u - t)r + t \equiv (u - t) + t \equiv u \pmod{b}$$

since $ar \equiv \pm 1 \pmod{b}$. Furthermore, we note by the definition of congruence that solving $ax \equiv \pm 1 \pmod{b}$ amounts to finding an $x, y \in \mathbb{Z}$ such that

$$ax + by = \pm 1.$$

Here we outline the algorithm for actually finding a solution to such a congruence again given the condition on a . Note that the condition $(a, b) = 1$ is exactly enough to guarantee solutions $x, y \in \mathbb{Z}$. See the related discussion in Notes 2 definition 4: for fixed $(a, b) = 1$, the set of all possible linear combinations $\{ax + by | x, y \in \mathbb{Z}\} = (a, b) = (1) = \mathbb{Z}$ where the parentheses now denote ideals.

Start with $(a, b) = 1$ How do we find the associated x and y such that $ax + by = 1$? Apply Lemma 2 of Notes 2. Without loss of generality say that $a > b$. Then there exist unique $q_0, 0 < r_0 < b$ such that $a = bq_0 + r_0$. Note that $(a, b) = (b, r_0) = 1$, that is, $0 < r_0 < b$ guarantees that b and r_0 are relatively prime. Moreover, suppose there were x_0, y_0 such that $x_0r_0 + y_0b = 1$, then since $r_0 = a - bq_0$, we have

$$\begin{aligned} x_0r_0 + y_0b &= x_0(a - bq_0) + y_0b \\ &= x_0a + (y_0 - q_0)b \\ &= 1 \end{aligned}$$

which gives us our initial x and y as $x = x_0, y = (y_0 - q_0)$.

We are not finished however, this relies upon the assumption that we know how to solve $x_0r_0 + y_0b = 1$. Use the same process on this equation to obtain x_0 and y_0 in terms of some x_1 and y_1 . Again, begin by dividing b by r_0 .

This method must terminate eventually when some $r_n = 1$ at which point it becomes easy to find the x_n and y_n coefficients. This is called the Euclidean Algorithm.

We skip article 28.

Article 29. Now we investigate $ax + t \equiv u \pmod{m}$ the case where a , the coefficient on the unknown, is not relatively prime to the modulus m . Suppose $(a, m) = \delta > 1$. By article 5, if x satisfies the congruence relative to the modulus m , then it certainly satisfies the congruence relative to the modulus δ . However, $ax \equiv 0 \pmod{\delta}$ since $\delta | a$. Thus $t \equiv u \pmod{\delta}$ is the only case where the congruence has a solution. In this case, $\delta | t - u$. Thus applying article 22, we obtain

$$(a/\delta)x \equiv (u - t)/\delta \pmod{m/\delta}$$

since a/δ and m/δ are relatively prime we can use the machinery of the previous articles to solve this case now, i.e. the coefficient of x and the modulus are relatively prime.

In the rest of section 2 Gauss proves the chinese remainder theorem and then uses this to give the simpler proof of Notes 6. Proposition 8 which is contained in the exercises of notes 6.

Exercises