





Appendix A: Supplementary data

Supplementary Table 1. *Ex vivo* expression of differentiation markers and inhibitory receptors across age groups.

<i>Ex vivo</i>		Median	Mean $\pm$ SD	One-way ANOVA
(% in V 2 T cells)		(IQR)	(n=13-28)	*Kruskal-Wallis
CD45RA	Neonate	61.5 (54.33 - 71.43)	62.81 $\pm$ 9.891	p=0.0024
	Infant	59.48 (45.3 - 68.48)	58.16 $\pm$ 16.39	
	Adult	35.4 (14.5 - 52.95)	39.18 $\pm$ 28.46	
CD27	Neonate	90.66 (86.49 - 93.97)	88.47 $\pm$ 8.927	p=0.0001*
	Infant	89.1 (84.05 - 92.85)	86.11 $\pm$ 10.87	
	Adult	55.2 (39.38 - 79.01)	55.73 $\pm$ 25.32	
NKG2A	Neonate	5.05 (3.34 - 6.84)	5.4 $\pm$ 3.19	p < 0.0001
	Infant	45.71 (31.93 - 48.48)	41.83 $\pm$ 15.94	
	Adult	72.94 (59.75 - 83.72)	70.2 $\pm$ 16.81	
CD16	Neonate	10.38 (9.555 - 29.05)	17.46 $\pm$ 11.9	p < 0.0001
	Infant	11.35 (5.395 - 21.81)	15.16 $\pm$ 12.07	
	Adult	34.82 (10.03 - 67.2)	40.02 $\pm$ 29.99	
PD-1	Neonate	27.8 (21.08 - 36.05)	29.89 $\pm$ 11.68	p = 0.0002*
	Infant	17.86 (13.33 - 21.54)	18.52 $\pm$ 8.767	
	Adult	10.9 (3.9 - 24.45)	14.75 $\pm$ 13.22	



Supplementary Figure 1. The proportion of V 9 chains in the 990-996 nucleotide range after stimulation mirrors the frequency of total V 2 T cells. Two weeks after stimulation, CBMC were lysed for RNA extraction and the CDR3 length distribution was determined by run off analysis for TCR gamma chains using a V9 segment. The bar graph shows the relative proportion of V 9 chains of different length (mean + SD) after each treatment, with a focus on the proportion of V 9 chains in the 990-996 nucleotide range. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.



Supplementary Figure 3. Perforin⁺ and granulysin⁺ V 2 T cells are enriched in the PD-1⁻ and CD16⁺ subsets. Sixteen days after stimulation with ZOL or live BCG, the frequency of cells storing cytotoxic effector molecules was assessed by intracellular staining. **A)** The scatter plot depicts the frequency of perforin⁺ V 2 T cells in the PD-1⁺ subset, the PD-1⁻ subset and the total V 2 population after BCG stimulation. **B)** The scatter plot shows the frequency of perforin⁺ V 2 T cells in the CD16⁺ subset, the CD16⁻ subset and the total V 2 population after BCG stimulation. **C)** The scatter plot compares the frequency of granulysin⁺ V 2 T cells in the CD16⁺ subset, the CD16⁻ subset and the total V 2 population after ZOL stimulation. * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001.