



1 Supplemental data




- 11 displays the proportion of granulysin<sup>+</sup> V 2 cells across the four subsets identified by CD56 and
- 12 PD-1, showing mean, standard deviation and individual values for each subset.
- 13


14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24




**Supplemental Figure 2. The subsets identified by CD56 and PD-1 differentially express CD4 and CD8.** After expansion, CB V 2 cells were assessed for expression of CD4 and CD8 relative to CD56 and PD-1. A) The pseudocolor plots show the expression of CD4 and CD8 in the PD-1<sup>+</sup>CD56<sup>-</sup> (left) and PD-1<sup>-</sup>CD56<sup>+</sup> (right) subset of total V 2 cells (middle dot plot). B) The scatter plots summarize the results for 5 specimens, displaying mean, SD and individual values of the proportion of CD8<sup>+</sup> (left), CD4<sup>+</sup> (center) and CD8<sup>-</sup>CD4<sup>-</sup> (right) V 2 cells 16 days after ZOL stimulation.



27 **Supplemental figure 3. The proportion of perforin<sup>+</sup> cells is enriched in the NKG2A<sup>+</sup> and**  
28 **CD16<sup>+</sup> subset of V 2 lymphocytes *ex vivo* and after expansion.** Before and after stimulation,  
29 CBMC were assessed for expression of cytotoxic mediators relative to NK receptors. In A) the plots  
30 show the proportion of perforin<sup>+</sup> V 2 cells in the NKG2A<sup>+</sup> and NKG2A<sup>-</sup> subset *ex vivo* (left) and 16  
31 days after ZOL stimulation (right). Paired values for individual specimens and means are  
32 displayed. B) The scatter plot shows the proportion of perforin<sup>+</sup> V 2 cells in the CD16<sup>+</sup> and CD16<sup>-</sup>  
33 subsets 16 days after ZOL stimulation, displaying mean, SD and individual values. C) The  
34 pseudocolor plots highlight the expression of CD16 relative to perforin in total V 2 cells, in  
35 PD-1<sup>+</sup>CD56<sup>-</sup> cells and in PD-1<sup>-</sup>CD56<sup>+</sup> cells (from left to right).



**Supplemental figure 4. The proportion of CD107a<sup>+</sup> V 2 cells after expansion with BCG is higher in female than male neonates.** CBMC were cultured for 16 days after ZOL or BCG stimulation, then restimulated for 6 hours with plastic-immobilized anti- TCR. The box plots show mean, median, IQR and outliers for the proportion of CD107a<sup>+</sup> V 2 cells after restimulation in cultures expanded with ZOL (left) or BCG (right), comparing female and male neonates.



55

56 **Supplemental figure 5. Sorting of neonatal V2 cell subsets based on CD56 and PD-1**  
 57 **expression yields highly purified fractions.** Fourteen days after ZOL stimulation CBMC were  
 58 stained with anti V2 (clone B6), anti-CD56 and anti-PD-1 for sorting. Cells for bulk RNA  
 59 sequencing were sorted on a FACS Aria II directly in trizol. A small aliquot of cells was sorted in  
 60 FBS-coated tubes for purity assessment. The pseudocolor plots show the pre-sorting V2  
 61 population (first row) and the purity of PD-1<sup>+</sup>CD56<sup>-</sup> (second row) and PD-1<sup>-</sup>CD56<sup>+</sup> (third row)  
 62 fractions after sorting for a representative specimen.

