

# Data Science: Unsupervised Learning

Adriano Z. Zambom

Unsupervised Learning

# Unsupervised Learning: Principal Components Analysis

## Unsupervised Learning: Principal Components Analysis

# Principal Components Analysis

## PCA

- Objective:

# Principal Components Analysis

## PCA

- Objective: Perform dimension reduction when we have many variables

## PCA

- Objective: Perform dimension reduction when we have many variables
- That is, explain the variance-covariance structure of the data through a few linear combinations

## PCA

- Objective: Perform dimension reduction when we have many variables
- That is, explain the variance-covariance structure of the data through a few linear combinations
- With 12 variables, for example, there will be more than 200 three-dimensional scatterplots to be studied

## PCA

- Objective: Perform dimension reduction when we have many variables
- That is, explain the variance-covariance structure of the data through a few linear combinations
- With 12 variables, for example, there will be more than 200 three-dimensional scatterplots to be studied
- Summary: Dimension reduction + interpretation

# Principal Components Analysis

## PCA

- Helpful when?



# Principal Components Analysis

## PCA

- Helpful when? we have multicollinearity: correlations between the x-variables

# Principal Components Analysis

## PCA

- Helpful when? we have multicollinearity: correlations between the x-variables
- Ex: What would you see in a plot of two x-variables that have a nearly perfect correlation?

# Principal Components Analysis

## PCA

- Helpful when? we have multicollinearity: correlations between the x-variables
- Ex: What would you see in a plot of two x-variables that have a nearly perfect correlation? around a line

# Principal Components Analysis

## PCA

- Helpful when? we have multicollinearity: correlations between the x-variables
- Ex: What would you see in a plot of two x-variables that have a nearly perfect correlation? around a line
- That line could be used as a new (one-dimensional) axis to represent the variation among data points.

# Principal Components Analysis

## PCA

- Helpful when? we have multicollinearity: correlations between the x-variables
- Ex: What would you see in a plot of two x-variables that have a nearly perfect correlation? around a line
- That line could be used as a new (one-dimensional) axis to represent the variation among data points.
- Ex: Suppose that we have verbal, math, and total SAT scores. Note that (at most) two dimensions of the data are necessary because  $\text{total} = \text{verbal} + \text{math}$

# Principal Components Analysis

## PCA

- Helpful when? we have multicollinearity: correlations between the x-variables
- Ex: What would you see in a plot of two x-variables that have a nearly perfect correlation? around a line
- That line could be used as a new (one-dimensional) axis to represent the variation among data points.
- Ex: Suppose that we have verbal, math, and total SAT scores. Note that (at most) two dimensions of the data are necessary because  $\text{total} = \text{verbal} + \text{math}$   
'at most' two dimensions and not exactly two: if there is a strong correlation between verbal and math, then it may be possible that there is only one true dimension to the data.

# Principal Components Analysis

## PCA

- The total variability is given by all variables, say we have  $p$  variables.
- However, sometimes we can find just a few ( less than  $p$ ) that explain almost all variability contained in all these  $p$  variables.

# Principal Components Analysis

## PCA

- The total variability is given by all variables, say we have  $p$  variables.
- However, sometimes we can find just a few (less than  $p$ ) that explain almost all variability contained in all these  $p$  variables.
- For example: if you have a large number of variables, say  $p$ , for your regression analysis, it may be difficult to fit the model, or you may not want to use all these variables. Instead, you can use only a few components  $k < p$ , and the prediction of your response variable  $Y$  may be satisfactory.



# Principal Components Analysis

**PCA** Suppose we have random variables  $X_1, \dots, X_p$ .

# Principal Components Analysis

**PCA** Suppose we have random variables  $X_1, \dots, X_p$ . Assume these random variables together  $\mathbf{X}' = [X_1, X_2, \dots, X_p]$  have the covariance matrix  $\Sigma$

# Principal Components Analysis

**PCA** Suppose we have random variables  $X_1, \dots, X_p$ . Assume these random variables together  $\mathbf{X}' = [X_1, X_2, \dots, X_p]$  have the covariance matrix  $\Sigma$  with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p \geq 0$ .

# Principal Components Analysis

**PCA** Suppose we have random variables  $X_1, \dots, X_p$ . Assume these random variables together  $\mathbf{X}' = [X_1, X_2, \dots, X_p]$  have the covariance matrix  $\Sigma$  with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p \geq 0$ . Consider the linear combinations

$$Y_1 = \mathbf{a}'_1 \mathbf{X} = a_{11}X_1 + a_{12}X_2 + \dots + a_{1p}X_p$$

# Principal Components Analysis

**PCA** Suppose we have random variables  $X_1, \dots, X_p$ . Assume these random variables together  $\mathbf{X}' = [X_1, X_2, \dots, X_p]$  have the covariance matrix  $\Sigma$  with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p \geq 0$ . Consider the linear combinations

$$Y_1 = \mathbf{a}'_1 \mathbf{X} = a_{11}X_1 + a_{12}X_2 + \dots + a_{1p}X_p$$

$$Y_2 = \mathbf{a}'_2 \mathbf{X} = a_{21}X_1 + a_{22}X_2 + \dots + a_{2p}X_p$$

# Principal Components Analysis

**PCA** Suppose we have random variables  $X_1, \dots, X_p$ . Assume these random variables together  $\mathbf{X}' = [X_1, X_2, \dots, X_p]$  have the covariance matrix  $\Sigma$  with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p \geq 0$ . Consider the linear combinations

$$Y_1 = \mathbf{a}'_1 \mathbf{X} = a_{11}X_1 + a_{12}X_2 + \dots + a_{1p}X_p$$

$$Y_2 = \mathbf{a}'_2 \mathbf{X} = a_{21}X_1 + a_{22}X_2 + \dots + a_{2p}X_p$$

$$\vdots$$

$$Y_p = \mathbf{a}'_p \mathbf{X} = a_{p1}X_1 + a_{p2}X_2 + \dots + a_{pp}X_p$$

Then

$$\text{Var}(Y_i) =$$

# Principal Components Analysis

**PCA** Suppose we have random variables  $X_1, \dots, X_p$ . Assume these random variables together  $\mathbf{X}' = [X_1, X_2, \dots, X_p]$  have the covariance matrix  $\Sigma$  with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p \geq 0$ . Consider the linear combinations

$$Y_1 = \mathbf{a}'_1 \mathbf{X} = a_{11}X_1 + a_{12}X_2 + \dots + a_{1p}X_p$$

$$Y_2 = \mathbf{a}'_2 \mathbf{X} = a_{21}X_1 + a_{22}X_2 + \dots + a_{2p}X_p$$

$$\vdots$$

$$Y_p = \mathbf{a}'_p \mathbf{X} = a_{p1}X_1 + a_{p2}X_2 + \dots + a_{pp}X_p$$

Then

$$\text{Var}(Y_i) = \mathbf{a}'_i \Sigma \mathbf{a}_i, i = 1, 2, \dots, p$$

$$\text{Cov}(Y_i, Y_k) =$$

# Principal Components Analysis

**PCA** Suppose we have random variables  $X_1, \dots, X_p$ . Assume these random variables together  $\mathbf{X}' = [X_1, X_2, \dots, X_p]$  have the covariance matrix  $\Sigma$  with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p \geq 0$ . Consider the linear combinations

$$Y_1 = \mathbf{a}'_1 \mathbf{X} = a_{11}X_1 + a_{12}X_2 + \dots + a_{1p}X_p$$

$$Y_2 = \mathbf{a}'_2 \mathbf{X} = a_{21}X_1 + a_{22}X_2 + \dots + a_{2p}X_p$$

$$\vdots$$

$$Y_p = \mathbf{a}'_p \mathbf{X} = a_{p1}X_1 + a_{p2}X_2 + \dots + a_{pp}X_p$$

Then

$$\text{Var}(Y_i) = \mathbf{a}'_i \Sigma \mathbf{a}_i, i = 1, 2, \dots, p$$

$$\text{Cov}(Y_i, Y_k) = \mathbf{a}'_i \Sigma \mathbf{a}_k, i, k = 1, 2, \dots, p$$



# Principal Components Analysis

**PCA** Suppose we have random variables  $X_1, \dots, X_p$ . Assume these random variables together  $\mathbf{X}' = [X_1, X_2, \dots, X_p]$  have the covariance matrix  $\Sigma$  with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p \geq 0$ . Consider the linear combinations

$$Y_1 = \mathbf{a}'_1 \mathbf{X} = a_{11}X_1 + a_{12}X_2 + \dots + a_{1p}X_p$$

$$Y_2 = \mathbf{a}'_2 \mathbf{X} = a_{21}X_1 + a_{22}X_2 + \dots + a_{2p}X_p$$

$$\vdots$$

$$Y_p = \mathbf{a}'_p \mathbf{X} = a_{p1}X_1 + a_{p2}X_2 + \dots + a_{pp}X_p$$

Then

$$\text{Var}(Y_i) = \mathbf{a}'_i \Sigma \mathbf{a}_i, i = 1, 2, \dots, p$$

$$\text{Cov}(Y_i, Y_k) = \mathbf{a}'_i \Sigma \mathbf{a}_k, i, k = 1, 2, \dots, p$$

**The principal components** are those *uncorrelated* linear combinations  $Y_1, \dots, Y_p$  whose variances are as large as possible.

# Principal Components Analysis

## PCA

- **First principle component** = linear combination  $\mathbf{a}'_1 \mathbf{X} = a_{11}X_1 + a_{12}X_2 + \dots + a_{1p}X_p$  that maximizes  $Var(\mathbf{a}'_1 \mathbf{X})$  subject to  $\mathbf{a}'_1 \mathbf{a}_1 = 1$ .

# Principal Components Analysis

## PCA

- **First principle component** = linear combination  $\mathbf{a}'_1 \mathbf{X} = a_{11}X_1 + a_{12}X_2 + \dots + a_{1p}X_p$  that maximizes  $Var(\mathbf{a}'_1 \mathbf{X})$  subject to  $\mathbf{a}'_1 \mathbf{a}_1 = 1$ .
- **Second principle component** = linear combination  $\mathbf{a}'_2 \mathbf{X}$  that maximizes  $Var(\mathbf{a}'_2 \mathbf{X})$  subject to  $\mathbf{a}'_2 \mathbf{a}_2 = 1$  and  $Cov(\mathbf{a}'_1 \mathbf{X}, \mathbf{a}'_2 \mathbf{X}) = 0$ .

# Principal Components Analysis

## PCA

- **First principle component** = linear combination  $\mathbf{a}'_1 \mathbf{X} = a_{11}X_1 + a_{12}X_2 + \dots + a_{1p}X_p$  that maximizes  $Var(\mathbf{a}'_1 \mathbf{X})$  subject to  $\mathbf{a}'_1 \mathbf{a}_1 = 1$ .
- **Second principle component** = linear combination  $\mathbf{a}'_2 \mathbf{X}$  that maximizes  $Var(\mathbf{a}'_2 \mathbf{X})$  subject to  $\mathbf{a}'_2 \mathbf{a}_2 = 1$  and  $Cov(\mathbf{a}'_1 \mathbf{X}, \mathbf{a}'_2 \mathbf{X}) = 0$ .
- ...
- **i-th principle component** = linear combination  $\mathbf{a}'_i \mathbf{X}$  that maximizes  $Var(\mathbf{a}'_i \mathbf{X})$  subject to  $\mathbf{a}'_i \mathbf{a}_i = 1$  and  $Cov(\mathbf{a}'_i \mathbf{X}, \mathbf{a}'_k \mathbf{X}) = 0 \forall k < i$ .

# Principal Components Analysis

## PCA

**Result** The variances of the Principal components have the following property

$$\begin{aligned}\sigma_{11} + \sigma_{22} + \dots + \sigma_{pp} &= \sum_{i=1}^p \text{Var}(X_i) = \lambda_1 + \lambda_2 + \dots + \lambda_p \\ &= \sum_{i=1}^p \text{Var}(Y_i)\end{aligned}$$

# Principal Components Analysis

## PCA

**Result** The variances of the Principal components have the following property

$$\begin{aligned}\sigma_{11} + \sigma_{22} + \dots + \sigma_{pp} &= \sum_{i=1}^p \text{Var}(X_i) = \lambda_1 + \lambda_2 + \dots + \lambda_p \\ &= \sum_{i=1}^p \text{Var}(Y_i)\end{aligned}$$

That is, total population variance is equal to the sum of the eigen-values. Hence, The proportion of total variance explained by the k-th principal component is

$$\frac{\lambda_k}{\lambda_1 + \lambda_2 + \dots + \lambda_p}$$

# Principal Components Analysis

## PCA

**Result** The vectors  $\mathbf{a}_i$  that define the principal components are actually the eigen-vectors associated with the matrix  $\Sigma$

# Principal Components Analysis

## PCA

**Example 8.1 (Calculating the population principal components)** Suppose that random variables  $X_1$ ,  $X_2$  and  $X_3$  have the covariance matrix

$$\Sigma = \begin{bmatrix} 1 & -2 & 0 \\ -2 & 5 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

It may be verified that the eigenvalue–eigenvector pairs are

$$\lambda_1 = 5.83, \quad \mathbf{e}'_1 = [.383, -.924, 0]$$

$$\lambda_2 = 2.00, \quad \mathbf{e}'_2 = [0, 0, 1]$$

$$\lambda_3 = 0.17, \quad \mathbf{e}'_3 = [.924, .383, 0]$$

Therefore, the principal components become

$$Y_1 = \mathbf{e}'_1 \mathbf{X} = .383X_1 - .924X_2$$

$$Y_2 = \mathbf{e}'_2 \mathbf{X} = X_3$$

$$Y_3 = \mathbf{e}'_3 \mathbf{X} = .924X_1 + .383X_2$$



# Principal Components Analysis

## PCA

In this example we have


$$\sigma_1 + \sigma_2 + \sigma_3 = 1 + 5 + 2 = \lambda_1 + \lambda_2 + \lambda_3 = 5.83 + 2 + 0.17$$

So the proportion of variability explained by the first principal component is

$$\frac{5.83}{5.83 + 2 + 0.17} = 0.728$$

# Principal Components Analysis

## PCA



**Figure 8.1** The constant density ellipse  $\mathbf{x}'\Sigma^{-1}\mathbf{x} = c^2$  and the principal components  $y_1, y_2$  for a bivariate normal random vector  $\mathbf{X}$  having mean  $\mathbf{0}$ .

# Principal Components Analysis

## PCA

Example: (Summarizing sample variability with two sample principal components) A census provided information, by tract, on five socioeconomic variables for Madison, Wisconsin, area.

# Principal Components Analysis

## PCA

Example: (Summarizing sample variability with two sample principal components) A census provided information, by tract, on five socioeconomic variables for Madison, Wisconsin, area.

Coefficients for the Principal Components  
(Correlation Coefficients in Parentheses)

Variable	$\hat{e}_1 (r_{\hat{y}_1, x_k})$	$\hat{e}_2 (r_{\hat{y}_2, x_k})$	$\hat{e}_3$	$\hat{e}_4$	$\hat{e}_5$
Total population	-0.039(-.22)	0.071(.24)	0.188	0.977	-0.058
Profession	0.105(.35)	0.130(.26)	-0.961	0.171	-0.139
Employment (%)	-0.492(-.68)	0.864(.73)	0.046	-0.091	0.005
Government employment (%)	0.863(.95)	0.480(.32)	0.153	-0.030	0.007
Medium home value	0.009(.16)	0.015(.17)	-0.125	0.082	0.989
Variance ( $\hat{\lambda}_i$ ):	107.02	39.67	8.37	2.87	0.15
Cumulative percentage of total variance	67.7	92.8	98.1	99.9	1.000

# Principal Components Analysis

PCA in R:

# Principal Components Analysis

PCA in R:

Let's use the dataset:  
?USArrests

# Principal Components Analysis

PCA in R:

Let's use the dataset:

?USArrests

There are a few functions to run PCA in R: ?

?prcomp (stats)

?princomp (stats)

?PCA (FactoMineR)

?dudi.pca (ade4)

?acp (amap).

# Principal Components Analysis

PCA in R:

Let's use the dataset:

?USArrests

There are a few functions to run PCA in R: ?

?prcomp (stats)

?princomp (stats)

?PCA (FactoMineR)

?dudi.pca (ade4)

?acp (amap).

Let's focus on ?prcomp and ?princomp



# Principal Components Analysis

PCA in R:

```
fit = prcomp(USArrests, scale. = TRUE)
```

# Principal Components Analysis

PCA in R:

```
fit = prcomp(USArrests, scale. = TRUE)
```

Standard deviations (1, ..., p=4):

```
[1] 1.5748783 0.9948694 0.5971291 0.4164494
```

Rotation (n x k) = (4 x 4):

	PC1	PC2	PC3	PC4
Murder	-0.5358995	0.4181809	-0.3412327	0.64922780
Assault	-0.5831836	0.1879856	-0.2681484	-0.74340748
UrbanPop	-0.2781909	-0.8728062	-0.3780158	0.13387773
Rape	-0.5434321	-0.1673186	0.8177779	0.08902432

# Principal Components Analysis

```
fit2 = princomp(USArrests, cor = TRUE)  
pca2$sdev  
unclass(pca2$loadings)
```

# Principal Components Analysis

## PCA

### The number of Principal Components

# Principal Components Analysis

## PCA

### The number of Principal Components

There is no definitive answer to this question. Consider:

- the amount of total sample variance explained

# Principal Components Analysis

## PCA

### The number of Principal Components

There is no definitive answer to this question. Consider:

- the amount of total sample variance explained
- the relative sizes of the eigenvalues (the variances of the sample components)

# Principal Components Analysis

## PCA

### The number of Principal Components

There is no definitive answer to this question. Consider:

- the amount of total sample variance explained
- the relative sizes of the eigenvalues (the variances of the sample components)
- the subject-matter interpretation of the components

# Principal Components Analysis

## PCA

### The number of Principal Components

There is no definitive answer to this question. Consider:

- the amount of total sample variance explained
- the relative sizes of the eigenvalues (the variances of the sample components)
- the subject-matter interpretation of the components
- A component associated with an eigenvalue near zero and, hence deemed unimportant, may indicate an unsuspected linear dependency in the data.



# Principal Components Analysis

## PCA

### The number of Principal Components

There is no definitive answer to this question. Consider:

- the amount of total sample variance explained
- the relative sizes of the eigenvalues (the variances of the sample components)
- the subject-matter interpretation of the components
- A component associated with an eigenvalue near zero and, hence deemed unimportant, may indicate an unsuspected linear dependency in the data.
- A useful visual aid to determining an appropriate number of principal components is a scree plot. With the eigenvalues ordered from largest to smallest, a scree plot is a plot of  $\hat{\lambda}_i$  versus  $i$ -the magnitude of an eigenvalue versus its number. **look for the elbow!**

# Principal Components Analysis

PCA




Figure 8.2 A scree plot.

# Principal Components Analysis

## **PCA**


Use *screeplot* function in R

```
screeplot(fit)
```

```
screeplot(fit2)
```

# Principal Components Analysis

## PCA



**Figure 8.4** Sample principal components and ellipses of constant distance.

# Principal Components Analysis

## Videos:

[https://www.youtube.com/watch?v=\\_UVHneBUBW0](https://www.youtube.com/watch?v=_UVHneBUBW0)

[https://www.youtube.com/watch?v=HMOI\\_lkzW08](https://www.youtube.com/watch?v=HMOI_lkzW08)

<https://www.youtube.com/watch?v=kw9R0nD69OU>

<https://www.youtube.com/watch?v=NLrb41ls4qo>

## Exercises:

### 1. Apply PCA on the wine data:

```
wine <- read.table("http://archive.ics.uci.edu/ml/machine-learning-databases/wine/wine.data",  
sep=",")
```

### 2. <https://learnche.org/pid/latent-variable-modelling/principal-component-analysis/pca-exercises>

Remember to take a look at the loadings, plots, variability explained, choosing the number of components, etc