

MA1E01: Chapter 3 Summary

Differentiation

Definitions

- **The Derivative Function:** The function $f'(x)$ defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

is called the derivative of f with respect to x . The domain of the derivative function consists of all x in the domain of f for which the limit above exists.

- **Differentiability:** A function f is said to be differentiable at x_0 if the limit

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$$

exists.

- **Equation of tangent:** The equation of the tangent line to a curve $y = f(x)$ at the point $x = x_0$ is

$$y = f(x_0) + f'(x_0)(x - x_0).$$

- **Higher derivatives:** The n^{th} derivative function is obtained by differentiating the $(n-1)^{\text{th}}$ derivative function,

$$f^{(n)}(x) = \frac{d^n f}{dx^n} = \lim_{h \rightarrow 0} \frac{f^{(n-1)}(x+h) - f^{(n-1)}(x)}{h}.$$

- **Local Linear Approximation:** The local linear approximation to the curve $y = f(x)$ in the vicinity of the point x_0 is given by the equation of the tangent line to the curve at that point, i.e.,

$$f(x) \approx f(x_0) + f'(x_0)(x - x_0).$$

- **Differentials:** The differential dy gives the change in the tangent line of $f(x)$ at x as x runs over the interval from x to $x + dx$. This is different to Δy which gives the change in the value of the function $f(x)$ over this interval.
- **Measurement/Propagated Error:** If x_0 is the exact value of a certain quantity and x is the measured value, then the measurement error is

$$dx = \Delta x = x - x_0.$$

Any quantity that depends on this measurement will have an associated propagated error. If $y_0 = f(x_0)$ is the exact value of a quantity being computed, then the value based on the measurement x is $y = f(x)$. Hence the propagated error is

$$\Delta y = y - y_0 = f(x) - f(x_0).$$

If the measurement error of x_0 is small, then we can approximate the propagated error by the differential

$$\Delta y \approx dy = f'(x)dx.$$

Relative/Percentage Error: The relative error associated with measuring a quantity is $\Delta y/y$. Again, if $\Delta x = dx$ is small enough, we can approximate this by the differential

$$\frac{\Delta y}{y} \approx \frac{dy}{y} = \frac{f'(x)dx}{f(x)}.$$

Theorems

- **Differentiability and continuity:** If a function f is differentiable at x_0 , then f is continuous at x_0 . The converse is false, differentiability does not necessarily imply continuity, for example, the absolute value function. As a corollary, if a function f is not continuous at x_0 , then it is not differentiable at x_0 .

Miscellaneous Results

- **Techniques of Differentiation:**

1. Product Rule:

$$\frac{d}{dx}(f \cdot g) = \frac{df}{dx}g + f\frac{dg}{dx}.$$

2. Quotient Rule:

$$\frac{d}{dx}\left(\frac{f}{g}\right) = \frac{\frac{df}{dx}g - f\frac{dg}{dx}}{g^2}.$$

3. Chain Rule: For compositions $y = f(g(x))$, we set $u = g(x)$, which implies $y = f(u)$, and then

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \frac{df(u)}{du} \frac{dg(x)}{dx}.$$

4. Implicit Differentiation: For functions defined implicitly by some equation, we can differentiate both sides of the equation using the rules above and solve for dy/dx in terms of x and y .

- **Related Rates:** For quantities related by a particular formula, then the rate of change at a particular time requires knowledge of the quantities and their derivatives at that time. For example, the volume of a cone is $V = \frac{\pi}{3}r^2h$, then the rate of change of the volume as r and h are allowed to vary is

$$\frac{dV}{dt} = \frac{\pi}{3} \left[2rh \frac{dr}{dt} + r^2 \frac{dh}{dt} \right].$$