
Algorithm Design and Analysis (ECS 122A)

Study Guide

Davis Computer Science Club
Tutoring Committee

Spring Quarter 2015

Contents

| | | |
|----------|---|----------|
| 1 | Asymptotic Notation | 2 |
| 1.1 | O-Notation (Big O) | 2 |
| 1.1.1 | Example | 3 |
| 1.2 | o-Notation (Little O) | 4 |
| 1.2.1 | Example | 5 |
| 1.2.2 | Example | 5 |
| 1.3 | Ω -Notation (Big Omega) | 6 |
| 1.3.1 | Example | 7 |
| 1.4 | ω -Notation (Little Omega) | 8 |
| 1.5 | Θ -notation | 9 |

1 Asymptotic Notation

1.1 O-Notation (Big O)

Notation:

$$f(n) \in O(g(n))$$

Formal Definition:

For a given function $g(n)$, $O(g(n))$ is the set of functions for which there exists positive constants c and n_0 such that $0 \leq f(n) \leq c \cdot g(n)$ for all $n \geq n_0$.

$$O(g(n)) = \{f(n) : \exists c, n_0 \text{ s.t. } 0 \leq f(n) \leq c \cdot g(n) \forall n \geq n_0\}$$

Informal Definition:

The function $g(n)$ is an asymptotic upper bound for the function $f(n)$ if there exists constants c and n_0 such that $0 \leq f(n) \leq c \cdot g(n)$ for $n \geq n_0$.

Another way to perceive Big O notation is that for $f(n) \in O(g(n))$, the function f 's asymptotic¹ growth is no faster than that of function g 's.

Limit Definition:

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} < \infty$$

¹Asymptotic: As given variable approaches infinity.

1.1.1 Example

Prove that asymptotic upper bound of $f(n) = 2n + 10$ is $g(n) = n^2$.

$$\begin{aligned} 0 \leq f(n) &\leq c \cdot g(n) \text{ for } n \geq n_0 \\ 0 \leq 2n + 10 &\leq c \cdot n^2 \text{ for } n \geq n_0 \end{aligned}$$

Arbitrarily choose c and n_0 values. Simplest is to turn one of the variables into the value 1 and solve. For this example, we will assign the value 1 to n_0 .

$$\begin{aligned} 0 \leq 2n + 10 &\leq c \cdot n^2 \text{ for } n \geq 1 \\ 2(1) + 10 &\leq c \cdot (1)^2 \\ 12 &\leq c \end{aligned}$$

By picking $n_0 = 1$ and $c = 12$, the inequality of $2n + 10 \leq 12n^2$ will hold true for all $n \geq 1$. Since there exists a constant c and n_0 that fulfill this inequality, we have proven that $f(n) = 2n + 10 = O(n^2)$.

1.2 o-Notation (Little O)

Notation:

$$f(n) \in o(g(n))$$

Formal Definition:

For a given function $g(n)$, $o(g(n))$ is the set of functions for which every positive constant $c > 0$, there exists a constant $n_0 > 0$ such that $0 \leq f(n) \leq c \cdot g(n)$ for all $n \geq n_0$.

$$o(g(n)) = \{f(n) : \exists n_0 \text{ s.t. } 0 \leq f(n) \leq c \cdot g(n) \forall n \geq n_0, c \geq 0\}$$

Informal Definition:

The function $g(n)$ is an upper bound that is not asymptotically tight. For all positive constant values of c , there must exist a constant n_0 such that $0 \leq f(n) \leq c \cdot g(n)$ for all $n \geq n_0$. The value of n_0 may not depend on n , but may depend on c .

Another way to perceive Little O notation is that for $f(n) \in o(g(n))$, the function f 's asymptotic growth is strictly less than that of the function g 's. In this sense, Little O can be seen as a “stronger” bound in comparison to Big O. By proving that a function is an element of Little O, it also proves that the function is an element of Big O.

Limit Definition:

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

1.2.1 Example

Prove that $f(n) = 2n$ has an upper bound $o(n^2)$.

$$\begin{aligned}0 \leq c \cdot g(n) &\leq f(n) \text{ for } n \geq n_0 \\0 \leq c \cdot 2n &\leq n^2 \text{ for } n \geq n_0 \\2c &\leq n \text{ for } n \geq n_0 \\2c &\leq n_0\end{aligned}$$

For Little O to hold true, the inequality needs to hold true for all $c > 0$ and for all $n > n_0$. From simplifying the inequality, we assert that the inequality will hold true as long as the value of n_0 is twice the value of c . Given that they are both constants, then there exists a constant value of n_0 for all positive constant c that fulfill this inequality.

Another method to solve this problem is to use the limit definition.

$$\begin{aligned}\lim_{n \rightarrow \infty} \frac{2n}{n^2} \\ \lim_{n \rightarrow \infty} \frac{2}{n} = 0\end{aligned}$$

1.2.2 Example

Prove that $f(n) = 2n^2$ does not have the upper bound $o(n^2)$.

$$\begin{aligned}0 \leq c \cdot g(n) &\leq f(n) \text{ for } n \geq n_0 \\0 \leq c \cdot 2n^2 &\leq n^2 \text{ for } n \geq n_0 \\2c &\leq 1 \text{ for } n \geq n_0\end{aligned}$$

For a function to have the Little O bound, the inequality must hold true for all positive c . However, simplification of the inequality asserts that the inequality will only hold true for all $c < \frac{1}{2}$. Therefore, $f(n) = 2n^2$ does not have the upper bound $o(n^2)$.

1.3 Ω -Notation (Big Omega)

Notation:

$$f(n) \in \Omega(g(n))$$

Formal Definition:

For a given function $g(n)$, $\Omega(g(n))$ is the set of functions for which there exists positive constants c and n_0 such that $0 \leq c \cdot g(n) \leq f(n)$ for all $n \geq n_0$.

$$\Omega(g(n)) = \{f(n) : \exists c, n_0 \text{ s.t. } 0 \leq c \cdot g(n) \leq f(n) \forall n \geq n_0\}$$

Informal Definition:

The function $g(n)$ is an asymptotic lower bound for the function $f(n)$ if there exists constants c and n_0 such that $0 \leq c \cdot g(n) \leq f(n)$ for $n \geq n_0$.

Limit Definition:

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} > 0$$

1.3.1 Example

Prove that the asymptotic lower bound of $f(n) = 8n^2$ is $g(n) = n$.

$$\begin{aligned} 0 \leq c \cdot g(n) &\leq f(n) \text{ for } n \geq n_0 \\ 0 \leq c \cdot n &\leq 8n^2 \text{ for } n \geq n_0 \end{aligned}$$

Arbitrarily choose c and n_0 values. Simplest is to turn one of the variables into the value 1 and solve. For this example, we will assign the value 1 to c .

$$\begin{aligned} 0 \leq n &\leq 8n^2 \text{ for } n \geq n_0 \\ (n_0) &\leq 8(n_0)^2 \\ 1 &\leq 8n_0 \\ \frac{1}{8} &\leq n_0 \end{aligned}$$

By picking $n_0 = 1$ and $c = 1$, the inequality of $n \leq 8n^2$ will hold true for all $n \geq 1$. Since there exists a constant c and n_0 that fulfill this inequality, we have proven that $f(n) = n^2 = \Omega(n)$.

1.4 ω -Notation (Little Omega)

Notation:

$$f(n) \in \omega(g(n))$$

Formal Definition:

For a given function $g(n)$, $\omega(g(n))$ is the set of functions for which every positive constant $c > 0$, there exists a constant $n_0 > 0$ such that $0 \leq c \cdot g(n) \leq f(n)$ for all $n \geq n_0$.

$$\omega(g(n)) = \{f(n) : \exists n_0 \text{ s.t. } 0 \leq c \cdot g(n) \leq f(n) \forall n \geq n_0, c \geq 0\}$$

Informal Definition:

The function $g(n)$ is a lower bound that is not asymptotically tight. For all positive constant values of c , there must exist a constant n_0 such that $0 \leq c \cdot g(n) \leq f(n)$ for all $n \geq n_0$. The value of n_0 may not depend on n , but may depend on c .

Another way to perceive Little ω notation is that for $f(n) \in \omega(g(n))$, the function f 's asymptotic growth is strictly greater than that of the function g 's. In this sense, Little ω can be seen as a “stronger” bound in comparison to Big Ω . By proving that a function is an element of Little ω , it also proves that the function is an element of Big Ω .

Limit Definition:

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

1.5 Θ -notation

Definition: The function $g(n)$ is an asymptotic