

7.2) Trigonometric Integrals

Exercise: Integrate $\int \sin^4(x) \cos^3(x) dx$.

$$\int \sin^4(x) \cos^3(x) dx = \int \sin^4(x) (1 - \sin^2(x)) \cos(x) dx$$

$$\int \sin^4(x) \cos^3(x) dx = \int (\sin^4(x) - \sin^6(x)) \cos(x) dx$$

$$\int \sin^4(x) \cos^3(x) dx = \int (u^4 - u^6) du$$

$$\int \sin^4(x) \cos^3(x) dx = \frac{u^5}{5} - \frac{u^7}{7} + C$$

$$\int \sin^4(x) \cos^3(x) dx = \frac{\sin^5(x)}{5} - \frac{\sin^7(x)}{7} + C$$

Exercise: Integrate the following:

a) $\int \cos^4(x) \sin^2(x) dx$

b) $\int \sin^5(x) \cos^3(x) dx$.

a) $\int \cos^4(x) \sin^2(x) dx = \int (\cos^4(x) - \cos^6(x)) dx$

$$\int \cos^4(x) \sin^2(x) dx = \int \left(\frac{(1+\cos(2x))^2}{4} - \frac{(1+\cos(2x))^3}{8} \right) dx$$

$$\int \cos^4(x) \sin^2(x) dx = \int \left(\frac{1}{4} + \frac{\cos(2x)}{2} + \frac{\cos^2(2x)}{4} - \frac{1}{8} - \frac{3\cos(2x)}{8} - \frac{3\cos^2(2x)}{8} - \frac{\cos^3(2x)}{8} \right) dx$$

$$\int \cos^4(x) \sin^2(x) dx = \int \left(\frac{1}{8} + \frac{\cos(2x)}{8} - \frac{\cos^2(2x)}{8} - \frac{\cos(2x)(1-\sin^2(2x))}{8} \right) dx$$

$$\int \cos^4(x) \sin^2(x) dx = \int \left(\frac{1}{8} + \frac{\cos(2x)}{8} - \frac{1+\cos(4x)}{16} - \frac{\cos(2x)}{8} + \frac{\cos(2x)\sin^2(2x)}{8} \right) dx$$

$$\int \cos^4(x) \sin^2(x) dx = \int \left(\frac{1}{16} - \frac{\cos(4x)}{16} + \frac{\cos(2x)\sin^2(2x)}{8} \right) dx$$

$$\int \cos^4(x) \sin^2(x) dx = \frac{1}{16}x - \frac{\sin(4x)}{64} + \frac{\sin^3(2x)}{48} + C$$

b) $\int \sin^5(x) \cos^3(x) dx = \int \sin^5(x) (1 - \sin^2(x)) \cos(x) dx$

$$\int \sin^5(x) \cos^3(x) dx = \int (\sin^5(x) - \sin^7(x)) \cos(x) dx$$

$$\int \sin^5(x) \cos^3(x) dx = \int (u^5 - u^7) du$$

$$\int \sin^5(x) \cos^3(x) dx = \frac{u^6}{6} - \frac{u^8}{8} + C$$

$$\int \sin^5(x) \cos^3(x) dx = \frac{\sin^6(x)}{6} - \frac{\sin^8(x)}{8} + C$$

Exercises: Integrate the following:

a) $\int \csc^4(x) \cot^3(x) dx$

b) $\int \tan^5(x) dx$

c) $\int \cos(5x) \sin(3x) dx$

a) $\int \csc^4(x) \cot^3(x) dx = \int \csc^2(x) (\cot^3(x) + \cot^5(x)) dx$

$$\int \csc^4(x) \cot^3(x) dx = - \int (u^3 + u^5) du$$

$$\int \csc^4(x) \cot^3(x) dx = - \frac{u^4}{4} - \frac{u^6}{6} + C$$

$$\int \csc^4(x) \cot^3(x) dx = - \frac{\cot^4(x)}{4} - \frac{\cot^6(x)}{6} + C$$

b) $\int \tan^5(x) dx = \int \tan(x) (\sec^2(x) - 1)^2 dx$

$$\int \tan^5(x) dx = \int \tan(x) (1 - 2\sec^2(x) + \sec^4(x)) dx$$

$$\int \tan^5(x) dx = \int \tan(x) (1 - 2\sec^2(x) + \sec^2(x) + \sec^2(x) \tan^2(x)) dx$$

$$\int \tan^5(x) dx = \int \tan(x) (1 - \sec^2(x) + \sec^2(x) \tan^2(x)) dx$$

$$\int \tan^5(x) dx = \int (\tan(x) - \tan(x) \sec^2(x) + \sec^2(x) \tan^3(x)) dx$$

$$\int \tan^5(x) dx = \ln |\sec(x)| - \frac{1}{2} \tan^2(x) + \frac{1}{4} \tan^4(x) + C$$

c) $\int \cos(5x) \sin(3x) dx = \int \frac{1}{2} (\sin(8x) - \sin(2x)) dx$

$$\int \cos(5x) \sin(3x) dx = - \frac{1}{16} \cos(8x) + \frac{1}{4} \cos(2x) + C$$

7.3)

Exercise: Integrate $\int \frac{\sqrt{4-x^2}}{x^2} dx$

Let $x = 2\sin(\theta)$, and $dx = 2\cos(\theta)d\theta$, and $\theta = \sin^{-1}\left(\frac{x}{2}\right)$

$$\int \frac{\sqrt{4-x^2}}{x^2} dx = \int \frac{\sqrt{4-4\sin^2(\theta)}}{(4\sin^2(\theta))} (2\cos(\theta)) d\theta$$

$$\int \frac{\sqrt{4-x^2}}{x^2} dx = \int \frac{2\sqrt{1-\sin^2(\theta)}}{(4\sin^2(\theta))} (2\cos(\theta)) d\theta$$

$$\int \frac{\sqrt{4-x^2}}{x^2} dx = \int \frac{\cos(\theta)}{\sin^2(\theta)} (\cos(\theta)) d\theta$$

$$\int \frac{\sqrt{4-x^2}}{x^2} dx = \int \frac{\cos^2(\theta)}{\sin^2(\theta)} d\theta = \int \cot^2(\theta) d\theta = \int (\csc^2(\theta) - 1) d\theta$$

$$\int \frac{\sqrt{4-x^2}}{x^2} dx = -\cot(\theta) - \theta + C = -\frac{\sqrt{4-x^2}}{x} - \sin^{-1}\left(\frac{x}{2}\right) + C$$

Exercise: Integrate $\int \frac{dx}{\sqrt{x^2-4}}$

Let $x = 2\sec(\theta)$, and $dx = \sec(\theta)\tan(\theta)d\theta$, and $\theta = \sec^{-1}\left(\frac{x}{2}\right)$

$$\int \frac{2\tan(\theta)\sec(\theta)d\theta}{\sqrt{4\sec^2(\theta)-4}} = \int \frac{2\tan(\theta)\sec(\theta)d\theta}{2\tan(\theta)} = \int \sec(\theta)d\theta = \ln|\sec(\theta) + \tan(\theta)| + C$$

$$\ln|\sec(\theta) + \tan(\theta)| + C = \ln\left|\frac{x}{2} + \frac{\sqrt{x^2-4}}{2}\right| + C$$

Exercises: Integrate the following:

a) $\int \frac{x^3}{\sqrt{9x^2+4}} dx$

b) $\int \frac{x}{\sqrt{x^2+4x+5}} dx$

a)

Let $x = \frac{2}{3}\tan(\theta)$, and $dx = \frac{2}{3}\sec^2(\theta)d\theta$, and $\theta = \tan^{-1}\left(\frac{3x}{2}\right)$

$$\int \frac{x^3}{\sqrt{9x^2+4}} dx = \int \frac{\frac{8}{27}\tan^3(\theta)}{\sqrt{4\tan^2(\theta)+4}} \left(\frac{2}{3}\sec^2(\theta)\right) d\theta$$

$$\int \frac{x^3}{\sqrt{9x^2+4}} dx = \frac{16}{81} \int \frac{\tan^3(\theta)}{2\sec(\theta)} \sec^2(\theta) d\theta$$

$$\int \frac{x^3}{\sqrt{9x^2+4}} dx = \frac{8}{81} \int \tan^3(\theta) \sec(\theta) d\theta = \frac{8}{81} \int \tan(\theta) (\sec^2(\theta) - 1) \sec(\theta) d\theta$$

$$\int \frac{x^3}{\sqrt{9x^2+4}} dx = \frac{8}{81} \int (\sec^2(\theta) - 1) \tan(\theta) \sec(\theta) d\theta$$

$$\int \frac{x^3}{\sqrt{9x^2+4}} dx = \frac{8}{81} \int (u^2 - 1) du$$

$$\int \frac{x^3}{\sqrt{9x^2+4}} dx = \frac{8}{81} \left(\frac{u^3}{3} - u \right) + C = \frac{8 \sec^3(\theta)}{243} - \frac{8}{81} \sec(\theta) + C$$

$$\text{Let } \frac{3x}{2} = \tan(\theta), \text{ so } \sec(\theta) = \frac{\sqrt{9x^2+4}}{2}$$

$$\int \frac{x^3}{\sqrt{9x^2+4}} dx = \frac{(9x^2+4)^{3/2}}{243} - \frac{4(9x^2+4)^{1/2}}{81} + C$$

b)

$$\text{Let } x+2 = \tan(\theta), \text{ and } dx = \sec^2(\theta) d\theta, \text{ and } \theta = \tan^{-1}(x+2)$$

$$\int \frac{x}{\sqrt{x^2+4x+5}} dx = \int \frac{x}{\sqrt{(x+2)^2+1}} dx = \int \frac{\tan(\theta)-2}{\sqrt{\tan^2(\theta)+1}} \sec^2(\theta) d\theta$$

$$\int \frac{x}{\sqrt{x^2+4x+5}} dx = \int \left(\frac{\tan(\theta)}{\sec(\theta)} - \frac{2}{\sec(\theta)} \right) \sec^2(\theta) d\theta$$

$$\int \frac{x}{\sqrt{x^2+4x+5}} dx = \int \left(\frac{\tan(\theta)}{\sec(\theta)} - \frac{2}{\sec(\theta)} \right) \sec^2(\theta) d\theta$$

$$\int \frac{x}{\sqrt{x^2+4x+5}} dx = \int (\tan(\theta) \sec(\theta) - 2 \sec(\theta)) d\theta$$

$$\int \frac{x}{\sqrt{x^2+4x+5}} dx = \sec(\theta) - 2 \ln |\sec(\theta) + \tan(\theta)| + C$$

$$\int \frac{x}{\sqrt{x^2+4x+5}} dx = \sqrt{x^2+4x+5} - 2 \ln \left| \sqrt{x^2+4x+5} + x+2 \right| + C$$

7.4) **Exercise:** Integrate $\int \frac{x^3-x^2-3x-1}{x^2-2x-3} dx$

$$\frac{x^3-x^2-3x-1}{x^2-2x-3} = x + 1 + \frac{2x+2}{x^2-2x-3} = x + 1 + \frac{2}{x-3}$$

$$\int \frac{x^3-x^2-3x-1}{x^2-2x-3} dx = \int \left(x + 1 + \frac{2}{x-3} \right) dx = \frac{1}{2}x^2 + x + 2\ln|x-3| + C$$

Exercise: Integrate $\int \frac{5x^2-40x-36}{x^3-4x^2-12x} dx$

$$x^3 - 4x^2 - 12x = x(x+2)(x-6)$$

$$\frac{5x^2-40x-36}{x^3-4x^2-12x} = \frac{A}{x} + \frac{B}{x+2} + \frac{C}{x-6}$$

$$5x^2 - 40x - 36 = A(x+2)(x-6) + Bx(x-6) + Cx(x+2)$$

$$5x^2 - 40x - 36 = A(x^2 - 4x - 12) + B(x^2 - 6x) + C(x^2 + 2x)$$

$$5x^2 - 40x - 36 = x^2(A+B+C) + x(-4A-6B+2C) + (-12A)$$

$$-12A = -36, \text{ so } A = 3.$$

$$3 + B + C = 5$$

$$-12 - 6B + 2C = -40$$

$$B + C = 2$$

$$-6B + 2C = -28$$

$$2B + 2C = 4$$

$$6B - 2C = 28$$

$$8B = 32, \text{ so } B = 4, \text{ and } C = -2$$

$$\int \frac{5x^2-40x-36}{x^3-4x^2-12x} dx = \int \left(\frac{3}{x} + \frac{4}{x+2} - \frac{2}{x-6} \right) dx = 3\ln|x| + 4\ln|x+2| - 2\ln|x-6| + C$$

Exercises: Integrate the following:

$$\text{a) } \int \frac{10x^2-7x+9}{x^3-x^2-x+1} dx$$

$$\text{b) } \int \frac{x^3+4x^2+5x+14}{x^4+4x^2-5} dx$$

$$\text{a) } x^3 - x^2 - x + 1 = x^2(x-1) - 1(x-1) = (x^2-1)(x-1) = (x+1)(x-1)^2$$

$$\frac{10x^2-7x+9}{x^3-x^2-x+1} = \frac{A}{x+1} + \frac{B}{x-1} + \frac{C}{(x-1)^2}$$

$$10x^2 - 7x + 9 = A(x-1)^2 + B(x-1)(x+1) + C(x+1)$$

$$10x^2 - 7x + 9 = A(x^2 - 2x + 1) + B(x^2 - 1) + C(x + 1)$$

$$10x^2 - 7x + 9 = x^2(A + B) + x(-2A + C) + 1(A - B + C)$$

$$10 = A + B$$

$$-7 = -2A + C$$

$$9 = A - B + C$$

Add them all up to get

$$12 = 2C$$

$$C = 6.$$

$$-7 = -2A + 6$$

$$A = 6.5, \text{ and } B = 3.5.$$

$$\int \frac{10x^2 - 7x + 9}{x^3 - x^2 - x + 1} dx = \int \left(\frac{6.5}{x+1} + \frac{3.5}{x-1} + \frac{6}{(x-1)^2} \right) dx = 6.5 \ln |x+1| + 3.5 \ln |x-1| - \frac{6}{x-1} + C$$

$$\text{b) } x^4 + 4x^2 - 5 = (x^2 - 1)(x^2 + 5) = (x - 1)(x + 1)(x^2 + 5)$$

$$\frac{x^3 + 4x^2 + 5x + 14}{x^4 + 4x^2 - 5} = \frac{A}{x+1} + \frac{B}{x-1} + \frac{Cx+D}{x^2+5}$$

$$x^3 + 4x^2 + 5x + 14 = A(x - 1)(x^2 + 5) + B(x + 1)(x^2 + 5) + (Cx + D)(x^2 - 1)$$

$$x^3 + 4x^2 + 5x + 14 = A(x^3 - x^2 + 5x - 5) + B(x^3 + x^2 + 5x + 5) + C(x^3 - x) + D(x^2 - 1)$$

$$x^3 + 4x^2 + 5x + 14 = x^3(A + B + C) + x^2(-A + B + D) + x(5A + 5B - C) + 1(-5A + 5B - D)$$

$$1 = A + B + C$$

$$4 = -A + B + D$$

$$5 = 5A + 5B - C$$

$$14 = -5A + 5B - D$$

Add the first and third equations together and add the second and fourth equations together:

$$6 = 6A + 6B$$

$$18 = -6A + 6B$$

$$1 = A + B$$

$$3 = -A + B$$

$$B = 2, A = -1, C = 0, \text{ and } D = 1.$$

$$\int \frac{x^3 + 4x^2 + 5x + 14}{x^4 + 4x^2 - 5} dx = \int \left(\frac{-1}{x+1} + \frac{2}{x-1} + \frac{1}{x^2+5} \right) dx = -\ln |x+1| + 2\ln |x-1| + \frac{1}{\sqrt{5}} \tan^{-1} \left(\frac{1}{\sqrt{5}} x \right) + C$$