

CSC236 Midterm Test Solutions

A correlation between the midterm test and exam has been confirmed. Please use this resource to study well!

Written by AlexTheMango, with pleasure [^_-]/

Prepared for December 19, 2024

Question #1

Let \mathcal{F} be the collection of all functions with domain \mathbb{N} and co-domain \mathbb{R} .

Given $A, B \in \mathcal{P}(\mathcal{F})$, define addition on $\mathcal{P}(\mathcal{F})$ by

$$A + B := \{f + g : f \in A, g \in B\}.$$

Recall that $f + g$ is the function with domain \mathbb{N} and co-domain \mathbb{R} such that

$$(f + g)(n) = f(n) + g(n).$$

Also recall that, if $h \in \mathcal{F}$, then

$$O(h) = \{q \in \mathcal{F} : (\exists n_0, c \in \mathbb{N})(\forall n \geq n_0)[|q(n)| \leq c|h(n)|]\}.$$

Claim: For arbitrary nonnegative $u, v \in \mathcal{F}$, it follows that $O(u) + O(v) = O(u + v)$.

Proof.

This proof demonstrates a double-subset inclusion to show equality.

Forward Inclusion — $O(u) + O(v) \subseteq O(u + v)$:

Let $h \in [O(u) + O(v)]$. Then, $h = f + g$, where $f \in O(u)$ and $g \in O(v)$.

By definition, there exists $c_1, c_2, n_1, n_2 > 0$ such that

- $|f(n)| \leq c_1|u(n)|$ for all $n \geq n_1$;
- $|g(n)| \leq c_2|v(n)|$ for all $n \geq n_2$.

Choose $n_0 = \max(n_1, n_2)$ and $c = \max(c_1, c_2)$.

Using the definition, triangle inequality, and assumption that u, v are nonnegative functions,

it follows that

$$\begin{aligned} |h(n)| &= |f(n) + g(n)| \leq |f(n)| + |g(n)| \leq c_1|u(n)| + c_2|v(n)| \\ &\leq c|u(n)| + c|v(n)| \\ &\leq c(|u(n)| + |v(n)|) \\ &= c(u(n) + v(n)) = c(|u(n) + v(n)|) = c|(u + v)(n)|. \end{aligned}$$

Thus, $|h(n)| \leq c|(u + v)(n)|$.

By definition, $h \in O(u + v)$. Therefore, $O(u) + O(v) \subseteq O(u + v)$.

Backward Inclusion — $O(u) + O(v) \supseteq O(u + v)$:

Let $h \in O(u + v)$.

By definition, there exists $c, n_0 > 0$ such that $|h(n)| \leq c|(u + v)(n)|$ for all $n \geq n_0$.

It follows that $|h(n)| \leq c|u(n) + v(n)| = c(u(n) + v(n))$, as u, v are nonnegative functions.

Let $w(n) = h(n) - cu(n)$. Consider the following cases for $w(n)$.

Case — $w(n) > 0$:

Notice that $w(n) > 0 \implies h(n) - cu(n) > 0 \implies h(n) > cu(n)$.

Since u is a nonnegative function, then $h(n)$ must be positive.

Recall $|h(n)| \leq c|(u + v)(n)| = c|u(n) + v(n)|$, and both u, v are nonnegative functions.

It follows that

$$\begin{aligned} |w(n)| &= |h(n) - cu(n)| = h(n) - cu(n) \\ &= |h(n)| - cu(n) \leq |cu(n) + cv(n)| - cu(n) \\ &= \cancel{cu(n)} + cv(n) - \cancel{cu(n)} = cv(n) = c|v(n)|. \end{aligned}$$

Thus, $|w(n)| \leq c|v(n)|$. This means $w(n) \in O(v)$.

Write $h(n) = cu(n) + w(n)$. It is obvious that $cu(n) \in O(u)$, and recall that $w(n) \in O(v)$.

Thus, $h(n) \in O(u + v)$.

Case — $w(n) \leq 0$:

Notice that $w(n) \leq 0 \implies h(n) - cu(n) \leq 0 \implies h(n) \leq cu(n)$. So, choose $h(n) = cu(n)$.

Then, $w(n) = h(n) - cu(n) = \cancel{cu(n)} - \cancel{cu(n)} = 0$.

Notice that $|w(n)| = |0| = 0 \leq c|v(n)|$, in fact, for any $c > 0$.

Clearly, $w(n) \in O(v)$.

Write $h(n) = cu(n) + w(n)$. It is obvious that $cu(n) \in O(u)$, and recall that $w(n) \in O(v)$.

Thus, $h(n) \in O(u + v)$.

Conclusion of Cases:

In all cases, $h(n) \in O(u + v)$ has been demonstrated.

Therefore, $O(u) + O(v) \subseteq O(u + v)$.

Conclusion:

Since both inclusions hold, $O(u) + O(v) = O(u + v)$.

□

Question #2

Let \mathcal{F} be as in *Question #1*. Let \mathcal{G} be the collection of all functions with domain $\mathcal{N} \times \mathcal{N}$ and co-domain \mathcal{R} . Let $V \in \mathcal{G}$.

For every $i \in \mathbb{N}$, let $g_i(n) = \sum_{j=1}^i V(j, n)$, and let $f_i(n) = V(i, n)$.

(a)

Claim: For all $i \in \mathbb{N}$, it follows that $O(g_i) = \sum_{j=0}^i O(f_j)$.

Proof.

Denote the predicate:

$$P(i) := O(g_i) = \sum_{j=0}^i O(f_j)$$

Proceed using the principle of simple induction over $P(i)$ for all $i \in \mathbb{N}$.

Base Case:

Let $i = 0$.

Then,

$$\begin{aligned} O(g_i) &= O(g_0) \\ &= O\left(\sum_{j=0}^0 V(j, n)\right) \\ &= O(V(0, n)) \\ &= O(f_0) \\ &= \sum_{j=0}^0 O(f_j) \\ &= \sum_{j=0}^i O(f_j). \end{aligned}$$

Thus, $P(0)$.

Induction Hypothesis:

Assume for some $k \in \mathbb{N}$, $P(k)$.

This means $O(g_k) = \sum_{j=0}^k O(f_j)$.

Induction Step:

Notice that

$$\begin{aligned} O(g_{k+1}) &= O\left(\sum_{j=0}^{k+1} V(j, n)\right) \\ &= O\left(\sum_{j=0}^{k+1} f_j\right) \\ &= O\left(\sum_{j=0}^k f_j + f_{k+1}\right) \\ &= O\left(\sum_{j=0}^k f_j\right) + O(f_{k+1}), \text{ by Question \#1} \\ &= \sum_{j=0}^k O(f_j) + O(f_{k+1}), \text{ by the Induction Hypothesis} \\ &= \sum_{j=0}^{k+1} O(f_j). \end{aligned}$$

Thus, $P(k) \implies P(k+1)$.

Conclusion:

Therefore, by the principle of simple induction, $P(i)$ holds for all $i \in \mathbb{N}$.

□

(b)

Claim: If $g(n) = g_n(n)$, then $O(g) = \sum_{j=0}^n O(f_j)$ does **not** necessarily hold.

(b)

Claim: If $g(n) = g_n(n)$, then $O(g) = \sum_{j=0}^n O(f_j)$ does **not** necessarily hold.

Proof.

To show that the equivalence in the claim does not necessarily hold, consider a counterexample.

Fix n_0 . Define $f_j(n)$ as follows:

$$f_j(n) = \begin{cases} n^2 & \text{if } j = n, \\ 1 & \text{if } j \neq n. \end{cases}$$

Consider the function $g_n(n) = \sum_{j=0}^n f_j(n)$.

For $n > n_0$, compute $g_n(n)$ as follows:

$$g_n(n) = \sum_{j=0}^n f_j(n) = \sum_{j=0}^{n-1} f_j(n) + f_n(n).$$

Substituting the definition of $f_j(n)$, this leads to:

$$\sum_{j=0}^{n-1} f_j(n) = \sum_{j=0}^{n-1} 1 = n.$$

Since $f_n(n) = n^2$, it follows that:

$$g_n(n) = n + n^2.$$

Therefore, $O(g_n) = O(n + n^2) = O(n^2)$. Let this be the left-hand side (LHS).

On the other hand, consider $\sum_{j=0}^n O(f_j)$:

$$f_j(n) = 1 \text{ for all } j \neq n.$$

Hence, $O(f_j) = O(1)$. There are n terms where $f_j(n) = 1$, so:

$$\sum_{j=0}^n O(f_j) = \sum_{j=0}^n O(1) = (n+1)O(1).$$

This simplifies to $O(n+1) = O(n)$. Let this be the right-hand side (RHS).

Clearly, $LHS = O(n^2) \neq O(n) = RHS$.

Note that this analysis holds for $n > n_0$, as n_0 is fixed and n can grow arbitrarily large. Fixing n_0 ensures a concrete starting point, while allowing $n > n_0$ provides generality for the counterexample. The counterexample demonstrates that $O(g) = \sum_{j=0}^n O(f_j)$ does not necessarily hold in general.

Thus, the equivalence in the claim is disproved.

□

Question #3

Claim: $f(n) = \lceil \sqrt{n} \rceil - \lfloor \sqrt{n} - 4 \rfloor$ is asymptotically constant (i.e. $f(n) \in \Theta(1)$).

Proof.

By definition, if x and y are arbitrary real numbers, then

$$(x \leq \lceil x \rceil < x + 1)$$

and

$$(y - 1 < \lfloor y \rfloor \leq y).$$

Rewrite the second inequality as $-y \leq -\lfloor y \rfloor < -(y - 1)$.

By adding the two inequalities, it follows that $x - y \leq \lceil x \rceil - \lfloor y \rfloor < x + 1 - (y - 1) = x - y + 2$.

Let $x = \sqrt{n}$ and $y = \sqrt{n} - 4$, for arbitrary natural n .

Then, $\lceil x \rceil - \lfloor y \rfloor = \lceil \sqrt{n} \rceil - \lfloor \sqrt{n} - 4 \rfloor = f(n)$. As well, $x - y = \sqrt{n} - (\sqrt{n} - 4) = 4$.

This means $x - y \leq \lceil x \rceil - \lfloor y \rfloor < x - y + 2 \implies 4 \leq f(n) < 4 + 2 \implies 4 \leq f(n) < 6$.

Let $n_0 = 0, c = 4, d = 6$. Let $g(n) = 1$.

Notice that $4 \leq f(n) < 6 \implies cg(n) \leq f(n) \leq dg(n)$, for all $n \geq n_0 = 0$ with $c = 4, d = 6$.

Therefore, $f(n) \in \Theta(g(n)) \implies f(n) \in \Theta(1)$. Indeed, $f(n)$ is asymptotically constant.

□

Question #4

Claim: The recurrence, $T(n) = 3T(\frac{n}{3}) + n^2 - n$, can be solved using the master theorem, and there exists a function $g(n)$ such that $T \in \Theta(g(n))$.

Proof.

The recurrence $T(n) = 3T(\frac{n}{3}) + n^2 - n$ has the form $T(n) = aT(\frac{n}{b}) + f(n)$, where $a = 3$, $b = 3$, and $f(n) = n^2 - n$. Since $f(n) = n^2 - n$ asymptotically behaves like n^2 , it follows that $f(n) \in \Theta(n^2)$, implying $k = 2$.

Master theorem applies to recurrences of this form, provided $a > 0$, $b > 1$, and $f(n)$ is non-negative for sufficiently large n . Here, $a = 3$, $b = 3$, and $f(n) = n^2 - n$ satisfies all these conditions since n^2 dominates n as $n \rightarrow \infty$.

Next, compute $\log_b a$:

$$\log_b a = \log_3 3 = 1.$$

Compare $\log_b a$ with k :

$$k = 2 > \log_3 3 = 1.$$

By the master theorem, when $k > \log_b a$, this leads to $T(n) \in \Theta(n^k)$.

Thus:

$$T(n) \in \Theta(n^2).$$

Therefore, there exists a function $g(n) = n^2$ such that $T(n) \in \Theta(g(n))$.

□

Question #5

Claim: Every regex without the Kleene star $*$ represents a finite language.

Proof.

Let r be a regular expression without the Kleene star $*$.

Define the predicate:

$$P(r) := \mathcal{L}(r) \text{ is a finite language.}$$

Proceed using the principle of structural induction over $P(r)$ for all regular expressions r without the Kleene star $*$.

Base Case:

By the definition of regular expressions,

- $\mathcal{L}(\emptyset) = \emptyset$
- $\mathcal{L}(\epsilon) = \{\epsilon\}$
- $\mathcal{L}(a) = \{a\}$, where $a \in \Sigma$ is an arbitrary symbol

Clearly, all three languages as denoted above are finite.

Thus, $P(\emptyset), P(\epsilon), P(a)$ all hold.

Induction Hypothesis:

Assume that for some regular expressions r_1, r_2 without the Kleene star $*$, $P(r_1), P(r_2)$ hold.

This means the languages $\mathcal{L}(r_1), \mathcal{L}(r_2)$ are finite.

Induction Step:

Consider that every language without the Kleene star $*$ can be obtained by the union or concatenation of languages.

By the Induction Hypothesis, $\mathcal{L}(r_1)$ and $\mathcal{L}(r_2)$ are finite languages. Recall that languages are sets of elements, where the elements are symbols of some alphabet Σ .

By definition, $\mathcal{L}(r_1 + r_2) = \mathcal{L}(r_1) \cup \mathcal{L}(r_2)$, which is finite as the union of finite sets (languages) is a finite set.

By definition, $\mathcal{L}(r_1 r_2) = \mathcal{L}(r_1) \frown \mathcal{L}(r_2)$, which is finite as the concatenation of two finite languages remains finite.

Conclusion:

By the principle of structural induction, every regular expression without the Kleene star $*$ represents a finite language.

□

Question #6

wordsgo here

Proof.

proofgoes here

□

Question #7

wordsgo here

Proof.

proofgoes here

□