

## Homework 7

**Liu Huihang**

SA18017026

QQ: 184050473

MAIL: huihang@mail.ustc.edu.cn

**Problem 1 (Slide 7 page 6)** Consider Bart Simpson density as following

$$f(x) = \frac{1}{2}\phi(x; 0, 1) + \frac{1}{10} \sum_{j=0}^4 \phi(x; (j/2) - 1, 1/10)$$

where  $\phi(x; \mu, \sigma)$  denotes a Normal density with mean  $\mu$  and standard deviation  $\sigma$ .

Generate 1000 sample from Bart Simpson distribution, estimate its density by histogram and compare different interval segmentation method. Estimate density use naive density estimator with image and compare the effect of different bandwidth  $h$ .

**Problem 2 (Slide 7 page 46)** Prove the asymptotic normality of KDE  $\hat{f}_h(x)$  if  $f''$  exists and  $h = cn^{-1/5}$ .

**Solution**

We have

$$E[\hat{f}_h(x) - f(x)] = \frac{1}{2}h^2 f''(x)\kappa_{21} + o(h^2)$$

$$Var[\hat{f}_h(x)] = \frac{1}{nh} f(x)\kappa_{02} + o\left(\frac{1}{nh}\right)$$

$$\frac{\hat{f}_h(x) - E[\hat{f}_h(x)]}{\sqrt{Var[\hat{f}_h(x)]}} \rightsquigarrow N(0, 1)$$

Thus,  $h = cn^{-1/5}$

$$\frac{\hat{f}_h(x) - f(x) - \frac{1}{2}h^2 f''(x)\kappa_{21}}{\sqrt{\frac{1}{nh} f(x)\kappa_{02}}} \rightsquigarrow N(0, 1)$$

$$\frac{\hat{f}_h(x) - f(x)}{\sqrt{\frac{n^{-4/5}}{c} f(x)\kappa_{02}}} - \frac{\frac{1}{2}c^2 n^{-2/5} f''(x)\kappa_{21}}{\sqrt{\frac{n^{-4/5}}{c} f(x)\kappa_{02}}} = n^{2/5} \frac{\hat{f}_h(x) - f(x)}{\sqrt{\frac{1}{c} f(x)\kappa_{02}}} - \frac{\frac{1}{2}c^2 f''(x)\kappa_{21}}{\sqrt{\frac{1}{c} f(x)\kappa_{02}}}$$

this leads to

$$n^{2/5} \left( \hat{f}_h(x) - f(x) \right) \rightsquigarrow N \left( \frac{1}{2}c^2 f''(x)\kappa_{21}, \frac{1}{c} f(x)\kappa_{02} \right)$$

■

**Problem 3** Generate 1000 samples from distribution  $0.3N(0, 1) + 0.7N(1, 0.32)$ , draw one figure with different estimated density functions with different bandwidth  $h$  where  $h$  is given by kedd.

**Problem 4 (Slide 7 page 44)** Suppose now that we have estimated an unknown density  $f$  using some kernel  $K_A$  and bandwidth  $h_A$ , what bandwidth  $h_B$  should we use in the estimation with kernel  $K_B$  when we want to get approximately the same degree of smoothness as we had in the case of  $K_A$  and  $h_A$ ?

Prove that the answer is given by the following formula:

$$h_B = h_A \frac{\delta_0^B}{\delta_0^A}.$$

**Solution**

$\delta_0$  is calculated to minimize AMSE, thus we take  $h$  to minimize the AMSE,

$$\text{AMSE} = \frac{1}{4}h^4(f''(x))^2\kappa_{21}^2 + \frac{1}{nh}f(x)\kappa_{02},$$

so this leads to

$$h(x) = \left( \frac{f(x)\kappa_{02}}{(f''(x))^2\kappa_{21}^2} \right)^{1/5} n^{-1/5} = \delta_0 \left( \frac{f(x)}{(f''(x))^2} \right)^{1/5} n^{-1/5}$$

Thus, to get the same degree of smoothness as we had in the case of  $K_A$  and  $h_A$ , we should choose

$$h_B = h_A \frac{\delta_0^B}{\delta_0^A}.$$

■

## Homework 8

**Liu Huihang**

SA18017026

QQ: 184050473

MAIL: huihang@mail.ustc.edu.cn

**Problem 1** Let  $X_1, \dots, X_n$ , i.i.d from density  $f(x)$ , denote  $r$  derivative of density function by  $\hat{f}_h^{(r)}$ , please find the asymptotic distribution of  $\hat{f}_h^{(r)}$ .

**Solution**

We have

$$\begin{aligned} E \left[ \hat{f}_h^{(r)}(x) \right] &= \frac{1}{h} \int K \left( \frac{x-z}{h} \right) f^{(r)}(z) dz \\ &= f^{(r)}(x) + \frac{1}{2} f^{(r+2)}(x) \kappa_{21} h^2 + o(h^2), \end{aligned}$$

and

$$\text{Var} \left[ \hat{f}_h^{(r)}(x) \right] = \frac{f(x)}{nh^{2r+1}} \int \left[ K^{(r)}(u) \right]^2 du + o \left( \frac{1}{nh^{2r+1}} \right).$$

Choose  $h(x) = cn^{-1/(2r+5)}$ . Thus,

$$\frac{\hat{f}_h^{(r)}(x) - E \left[ \hat{f}_h^{(r)}(x) \right]}{\sqrt{\text{Var} \left[ \hat{f}_h^{(r)}(x) \right]}} \rightsquigarrow N(0, 1),$$

Because

$$\begin{aligned} & \frac{\hat{f}_h^{(r)}(x) - f^{(r)}(x) - \frac{1}{2} f^{(r+2)}(x) \kappa_{21} h^2}{\sqrt{\frac{f(x)}{nh^{2r+1}} \int \left[ K^{(r)}(u) \right]^2 du}} \\ &= \frac{\hat{f}_h^{(r)}(x) - f^{(r)}(x) - \frac{1}{2} f^{(r+2)}(x) \kappa_{21} c^2 n^{-2/(2r+5)}}{\sqrt{n^{-4/(2r+5)} c^{-2r-1} f(x) \int \left[ K^{(r)}(u) \right]^2 du}} \\ &= n^{2/(2r+5)} \frac{\hat{f}_h^{(r)}(x) - f^{(r)}(x)}{\sqrt{c^{-2r-1} f(x) \int \left[ K^{(r)}(u) \right]^2 du}} - \frac{\frac{1}{2} c^2 f^{(r+2)}(x) \kappa_{21}}{\sqrt{c^{-2r-1} f(x) \int \left[ K^{(r)}(u) \right]^2 du}} \end{aligned}$$

so we obtain

$$n^{2/(2r+5)} \left( \hat{f}_h^{(r)}(x) - f^{(r)}(x) \right) \rightsquigarrow N \left( \frac{1}{2} c^2 f^{(r+2)}(x) \kappa_{21}, c^{-2r-1} f(x) \int \left[ K^{(r)}(u) \right]^2 du \right).$$

■

**Problem 2** *Generate 1000 samples from distribution  $0.3N(0, 1) + 0.7N(1, 0.32)$ , draw one figure with different estimated derivative of density functions with different bandwidth  $h$  where  $h$  is given by kedd.*

**Problem 3 (Adaptive KDE)** *Write program to draw the figure of 21 pages of courseware. (Referring algorithm steps in reading materials)*

**Problem 4 (Kernel Density Estimation with Boundary Correction)** *Reade accessory materials. Write program to realize the result in 39 pages of slide. (Reading material, Figure 3)*