



MODULE 3: Potato pest and disease management




C I P C A M E R O O N

Module 3 Outline

- 3.1. The disease triangle
- 3.2. Controlling late blight
- 3.3. Controlling bacterial wilt
- 3.4. Controlling soft rot (or blackleg)
- 3.5. Controlling viral diseases
- 3.6. Controlling potato tuber moth
- 3.7. Controlling nematodes
- 3.8. Safe use of chemicals

3.1. The disease triangle



The disease triangle

- **Pests and diseases** in potato reduce tuber quality causing up to 100% yield losses.
- A nexus of three factors are conducive to pests and diseases in the field and in storage:
 - ✓ a crop attacked by pests and diseases,
 - ✓ the **pathogen** or pest (organism causing disease or damage), and
 - ✓ the **environment** (conditions that favor the pathogen or pest).

Cont'd...

The disease triangle

- A pathogen can be lodged in the plant or tuber for a long time without symptoms (**latent infection** – e.g., bacterial wilt).
- Late blight, bacterial wilt, soft rot (or blackleg), viral diseases, potato tuber moth, and nematodes are some harmful diseases and pests.
- Pest/disease control strategy: Break the harmful nexus.

3.2. Controlling late blight



Causal agent and symptoms

- Caused by the fungus *Phytophthora infestans*, can lead to crop failure.
- Tell-tale signs:
 - ✓ On leaves: Large, irregular brown spots that dry out,
 - ✓ On stems: Weakened stems, and
 - ✓ On tubers: Brown patches of dry rot around the edges.
- Severe infestation occurs at high relative humidity (>90%) and cool temperature (<22°C).
- Spreads swiftly in the field.


Steps to control late blight

- Timely spraying of two major groups of chemicals to prevent infection and kill the pathogen after infection:
 - **Contact fungicides** kill by contact. Examples: Mancozeb, Pencozeb, Balear (Tropic), Plantineb, Manesam, Mancobex, and Mancozan.
 - **Systemic fungicides** enter the plant and kill the pathogen. Examples: Ridomil (Gold and Plus), Fungi-Pro, Fungicur, Monchamp, Metalm 75 WP, Metrostar and Parastar.
- Planting less susceptible varieties.

Spraying schedule

- Use a contact fungicide after plant emergence.
- Spray around 3 g/liter (45 – 50 g/sprayer) of a systemic fungicide 2 weeks later; dosage can vary.
- Subsequent sprays: Contact fungicides at 2-week intervals until the canopy turns yellow (maturity). If disease symptoms are visible in the field, spray a systemic fungicide (a maximum of twice per season for less susceptible varieties).

3.3 .Controlling bacterial wilt



Causal agent and mode of infection

- Bacterial wilt is both a soil and seedborne disease caused by *Ralstonia solanacearum*.
- Infects the crop at all stages causing crop failure.
- Can also spread via infected seed, water, roots, soil, farming tools, livestock and people.
- Also affects plants such as chili, tomato, tobacco, black nightshade and eggplant, and several weed species.
- Not every wilt is caused by bacterial wilt.



Classic symptoms

- On immersing a small bit of the stem's base in a glass of water, bacteria oozes from the stem and a downward movement in the water will occur.
- Black or brown rings when a tuber is cut in half.
- Bacteria oozing from tuber eyes, revealed by soil sticking to tuber eyes when the crop is harvested.

Controlling bacterial wilt

- There is no commercial chemical to control bacterial wilt.
- Cultural practices are the answer:
 - ✓ Plant bacterial wilt-free seed,
 - ✓ Crop rotation with crops that don't belong to the potato family, such as cereals (maize as first rotation crop)
 - ✓ Uproot wilting plants and soil around roots,
 - ✓ Clean tools before and after use,
 - ✓ In serious infestations, plant non-host plants for more than 5 years, and
 - ✓ Remove potato volunteers.

3.4. Controlling soft rot or blackleg



Causes and symptoms

- Caused by a unique bacterium (*Pectobacterium atrosepticum*) that changes tuber tissue into liquid; high humidity aids infection.
- Infected tuber: Decay in one spot spreads rapidly, resulting in rotting tissue that is mushy, slimy and water soaked.
- Infected tubers rot either in the field or in storage and produce a foul smell.
- **Black lesions** at the base of the stem; affected tissue becomes soft and water soaked under humid conditions and then the plant collapses.

Controlling soft rot (blackleg)

- Control is through a recommended integrated approach.

3.5. Controlling viral diseases



Viruses are tough to spot

- Viral diseases are difficult to recognize in the field.
- Mild infections can show no signs of disease at all.
- More than one type of virus can infect the field or plant.
- Potato leafroll virus: Infected plants have leaflets curling upward, turn pale yellow and feel brittle and fragile.
- Good fertilization / irrigation increases resistance to viruses.


What to watch out for

- Differentiate between a healthy and diseased plant.
- Viruses result in smaller potato tubers which farmers unknowingly mistake for seed, leading to reduced yields during subsequent seasons (**seed degeneration**).

Controlling viral diseases

- A mix of practices can control viral diseases in potatoes:
 - ✓ Using clean/certified seed,
 - ✓ Controlling insects that spread viral diseases (insecticides, traps, predators, etc.),
 - ✓ Planting varieties that are less susceptible to viral diseases, and
 - ✓ Uprooting and destroying non-potato plants which can attract and host vectors of viral diseases.

3.6. Controlling potato tuber moth



Symptoms to look out for

- Potato tuber moths (potato tuber worms) infest the crop in the field and move with tubers to the store.
- Moth larvae penetrate tubers through the eyes and create twisting **tunnels** in the tubers.
- They reproduce continuously in stored potatoes causing huge losses.
- They display fewer characteristic tunnels in stores but they have clearly visible **excreta**, mainly at the eyes. Often, these tubers become drier than healthy ones.

Preventing infestation

- Avoid planting in too light and loose soil.
- Deep planting or high hilling up protect the tubers.
- Decontaminate storage rooms prior to storing.
- Inspect the tubers carefully before and during storage and remove tubers showing openings/galleries and/or excreta.
- Use natural repellent plants such as **Lantana**.

Controlling the potato tuber moth

- ✓ Spraying the field with appropriate insecticides
 - The first treatment starts right after plant emergence.
 - Subsequent treatments are done at regular intervals of 2-3 weeks.
 - Combining fungicides and insecticides is more economical.
 - In dry weather, irrigation is a good preventative measure against these worms.

3.7. Controlling nematodes



Types, symptoms and marketability

- Potato cyst nematodes:
 - ✓ Impair plant growth, reducing it to dwarf,
 - ✓ May turn yellow and show signs of wilting, and
 - ✓ In the long run, tubers decrease in size causing reduction in yield.
- Root-knot nematodes (widespread):
 - ✓ Attack tubers causing blemishes, and
 - ✓ Make tubers unmarketable.
- Tubers containing nematodes are edible.
- Infested potatoes are more susceptible to bacterial wilt.

3.8. Safe use of chemicals



When to use chemicals

- Maintain pest populations and disease severity at acceptable levels
- Keep pesticides to levels that are economically justified and safe for human/environmental health.
- Use chemicals after exhausting cultural or biological control:
 - ✓ Natural predators,
 - ✓ Varieties with pest/disease resistance,
 - ✓ Healthy seed potatoes,
 - ✓ Rotation with other crops, and
 - ✓ Organic pesticides (e.g., neem).

Safe use of chemicals

- Strictly follow the manufacturer's instructions on dosage, spraying and frequency of application.
- Wear recommended protective gear while spraying.
- Misuse of pesticides can lead to poisoning.
- Strictly avoid spraying crops before harvest. **No fungicides or insecticides should be applied during the last 2-3 weeks of the crop.**
- Ensure the safe disposal of chemical containers.



CIP is a research-for-development organization with a focus on potato, sweetpotato and Andean roots and tubers. It delivers innovative science-based solutions to enhance access to affordable nutritious food, foster inclusive sustainable business and employment growth, and drive the climate resilience of root and tuber agri-food systems. Headquartered in Lima, Peru, CIP has a research presence in more than 20 countries in Africa, Asia and Latin America.

www.cipotato.org



CIP is a CGIAR research center

CGIAR is a global research partnership for a food-secure future. Its science is carried out by 15 research centers in close collaboration with hundreds of partners across the globe.

www.cgiar.org

CIP thanks all donors and organizations which globally support its work through their contributions to the CGIAR Trust Fund. <https://www.cgiar.org/funders/>

© 2021. International Potato Center. All rights reserved.



This work by the International Potato Center is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0).

To view a copy of this license, visit <https://creativecommons.org/licenses/by/4.0/>. Permissions beyond the scope of this license may be available at: <http://www.cipotato.org/contact/>