

# Chapter 8: Hypothesis Testing with Two Samples

DSCC 462

Computational Introduction to Statistics

Anson Kahng

Fall 2022

# Comparison of Two Means

# Comparison of Two Means

- We used a hypothesis test to compare the unknown mean of a single population to some fixed, known value,  $\mu_0$

# Comparison of Two Means

- We used a hypothesis test to compare the unknown mean of a single population to some fixed, known value,  $\mu_0$
- Often, we want to compare the means of two separate populations, where *both* means are unknown

# Comparison of Two Means

- We used a hypothesis test to compare the unknown mean of a single population to some fixed, known value,  $\mu_0$
- Often, we want to compare the means of two separate populations, where *both* means are unknown
  - E.g., is the average height of Americans equal to the average height of Canadians?

# Comparison of Two Means

- We used a hypothesis test to compare the unknown mean of a single population to some fixed, known value,  $\mu_0$
- Often, we want to compare the means of two separate populations, where *both* means are unknown
  - E.g., is the average height of Americans equal to the average height of Canadians?
- Must determine whether the two samples are paired or independent

# Comparison of Two Means

- We used a hypothesis test to compare the unknown mean of a single population to some fixed, known value,  $\mu_0$
- Often, we want to compare the means of two separate populations, where *both* means are unknown
  - E.g., is the average height of Americans equal to the average height of Canadians?
- Must determine whether the two samples are paired or independent
  - Paired: Weight before and after surgery for a group of men

# Comparison of Two Means

- We used a hypothesis test to compare the unknown mean of a single population to some fixed, known value,  $\mu_0$
- Often, we want to compare the means of two separate populations, where *both* means are unknown
  - E.g., is the average height of Americans equal to the average height of Canadians?
- Must determine whether the two samples are paired or independent
  - Paired: Weight before and after surgery for a group of men
  - Independent: Height of Americans compared to height of Canadians



# Paired Samples

# Paired Samples

- For each observation in the first group, there is a corresponding observation in the second group

# Paired Samples

- For each observation in the first group, there is a corresponding observation in the second group
- *Self-pairing*: Measurements are taken on a single subject at two distinct time points (before and after)

# Paired Samples

- For each observation in the first group, there is a corresponding observation in the second group
- *Self-pairing*: Measurements are taken on a single subject at two distinct time points (before and after)
- *Matched pairing*: Match two individuals with similar demographics / characteristics and compare their differences in response

# Paired Samples

- For each observation in the first group, there is a corresponding observation in the second group
- *Self-pairing*: Measurements are taken on a single subject at two distinct time points (before and after)
- *Matched pairing*: Match two individuals with similar demographics / characteristics and compare their differences in response
  - Make the pair as similar as possible with respect to important characteristics (e.g., age, gender, socioeconomic status, etc.) depending on the setting

# Paired Samples

# Paired Samples

- Use pairing to control for extraneous sources of variation that may otherwise influence results

# Paired Samples

- Use pairing to control for extraneous sources of variation that may otherwise influence results
- By measuring on the same sample, we remove natural biological variability between people



# Paired Samples

- Use pairing to control for extraneous sources of variation that may otherwise influence results
- By measuring on the same sample, we remove natural biological variability between people
- If we pair based on a certain characteristic (e.g., age), then we do not have to worry about that characteristic (age) influencing the results

# Paired Samples

- Use pairing to control for extraneous sources of variation that may otherwise influence results
- By measuring on the same sample, we remove natural biological variability between people
- If we pair based on a certain characteristic (e.g., age), then we do not have to worry about that characteristic (age) influencing the results
- In general, pairing makes comparisons more *precise*

# Paired Samples

# Paired Samples

- In the situation of paired samples, our data often is the difference between elements in each pair

# Paired Samples

- In the situation of paired samples, our data often is the difference between elements in each pair
  - Sample 1:  $x_{11}, x_{21}, \dots, x_{n1}$

# Paired Samples

- In the situation of paired samples, our data often is the difference between elements in each pair
  - Sample 1:  $x_{11}, x_{21}, \dots, x_{n1}$
  - Sample 2:  $x_{12}, x_{22}, \dots, x_{n2}$

# Paired Samples

- In the situation of paired samples, our data often is the difference between elements in each pair
  - Sample 1:  $x_{11}, x_{21}, \dots, x_{n1}$
  - Sample 2:  $x_{12}, x_{22}, \dots, x_{n2}$
  - Difference:  $d_1 = x_{11} - x_{12}, d_2 = x_{21} - x_{22}, \dots, d_n = x_{n1} - x_{n2}$

# Paired Samples: Example



# Paired Samples: Example

- We measure the sitting heart rate for 40 people after they have been sitting for 5 minutes

# Paired Samples: Example

- We measure the sitting heart rate for 40 people after they have been sitting for 5 minutes
- These 40 people then run on a treadmill for 30 minutes, sit for 5 minutes, and then have their heart rate measured

# Paired Samples: Example

- We measure the sitting heart rate for 40 people after they have been sitting for 5 minutes
- These 40 people then run on a treadmill for 30 minutes, sit for 5 minutes, and then have their heart rate measured
- Sample 1: "before" heart rate

# Paired Samples: Example

- We measure the sitting heart rate for 40 people after they have been sitting for 5 minutes
- These 40 people then run on a treadmill for 30 minutes, sit for 5 minutes, and then have their heart rate measured
- Sample 1: "before" heart rate
- Sample 2: "after" heart rate

# Paired Samples: Example

- We measure the sitting heart rate for 40 people after they have been sitting for 5 minutes
- These 40 people then run on a treadmill for 30 minutes, sit for 5 minutes, and then have their heart rate measured
- Sample 1: "before" heart rate
- Sample 2: "after" heart rate
- Goal: We are interested in how heart rate changed after running; we care about *before* – *after*

# Paired Samples: Example

# Paired Samples: Example

- The data is paired, so we can use the differences  $d_i = b_i - a_i$  as the data

# Paired Samples: Example

- The data is paired, so we can use the differences  $d_i = b_i - a_i$  as the data
  - Reduces to the one-sample problem: compare differences to 0



# Paired Samples: Example

- The data is paired, so we can use the differences  $d_i = b_i - a_i$  as the data
  - Reduces to the one-sample problem: compare differences to 0
- We need the mean of the  $d_i$ 's and its sampling distribution

# Paired Samples: Example

- The data is paired, so we can use the differences  $d_i = b_i - a_i$  as the data
  - Reduces to the one-sample problem: compare differences to 0
- We need the mean of the  $d_i$ 's and its sampling distribution
- Mean:  $\bar{x}_d$ , sample standard deviation  $s_d$  (unknown true  $\sigma_d$ )

# Paired Samples: Example

- The data is paired, so we can use the differences  $d_i = b_i - a_i$  as the data
  - Reduces to the one-sample problem: compare differences to 0
- We need the mean of the  $d_i$ 's and its sampling distribution
- Mean:  $\bar{x}_d$ , sample standard deviation  $s_d$  (unknown true  $\sigma_d$ )
- Standard error:  $\frac{s_d}{\sqrt{n}}$

# Paired Samples: Example

- The data is paired, so we can use the differences  $d_i = b_i - a_i$  as the data
  - Reduces to the one-sample problem: compare differences to 0
- We need the mean of the  $d_i$ 's and its sampling distribution
- Mean:  $\bar{x}_d$ , sample standard deviation  $s_d$  (unknown true  $\sigma_d$ )
- Standard error:  $\frac{s_d}{\sqrt{n}}$
- Assumption:  $\bar{x}_d \sim N\left(\mu_d, \sigma_d/\sqrt{n}\right)$

$$n = 40$$

# Paired Samples: Example

# Paired Samples: Example

- Let the true difference in population mean be  $\mu_d$

# Paired Samples: Example

- Let the true difference in population mean be  $\mu_d$
- Then, our hypotheses are:

# Paired Samples: Example

- Let the true difference in population mean be  $\mu_d$
- Then, our hypotheses are:
  - $H_0 : \mu_d = 0$  and  $H_0 : \mu_d \neq 0$



# Paired Samples: Example

- Let the true difference in population mean be  $\mu_d$
- Then, our hypotheses are:
  - $H_0 : \mu_d = 0$  and  $H_1 : \mu_d \neq 0$
- Test statistic:  $t = \frac{\bar{x}_d - \mu_d}{s_d / \sqrt{n}}$

# Paired Samples: Example

- Let the true difference in population mean be  $\mu_d$
- Then, our hypotheses are:
  - $H_0 : \mu_d = 0$  and  $H_a : \mu_d \neq 0$
- Test statistic:  $t = \frac{\bar{x}_d - \mu_d}{s_d / \sqrt{n}}$
- Under the null hypothesis,  $t \sim t_{n-1}$  ( $n - 1$  degrees of freedom)

# Paired Samples: Example

- Let the true difference in population mean be  $\mu_d$
- Then, our hypotheses are:
  - $H_0 : \mu_d = 0$  and  $H_a : \mu_d \neq 0$
- Test statistic:  $t = \frac{\bar{x}_d - \mu_d}{s_d / \sqrt{n}}$
- Under the null hypothesis,  $t \sim t_{n-1}$  ( $n - 1$  degrees of freedom)
- Calculate p-value,  $p$ , by finding the probability of observing a mean difference at least as extreme as  $\bar{x}_d$ , given that  $\mu_d = 0$

# Paired Samples: Example

- Let the true difference in population mean be  $\mu_d$
- Then, our hypotheses are:
  - $H_0 : \mu_d = 0$  and  $H_a : \mu_d \neq 0$
- Test statistic:  $t = \frac{\bar{x}_d - \mu_d}{s_d / \sqrt{n}}$
- Under the null hypothesis,  $t \sim t_{n-1}$  ( $n - 1$  degrees of freedom)
- Calculate p-value,  $p$ , by finding the probability of observing a mean difference at least as extreme as  $\bar{x}_d$ , given that  $\mu_d = 0$
- Reject  $H_0$  if  $p \leq \alpha$

# Paired Samples: Example

# Paired Samples: Example

- Consider the heart rates before and after running on a treadmill example

# Paired Samples: Example

- Consider the heart rates before and after running on a treadmill example

*d*

Before	After	Difference
55	60	-5
62	75	-13
61	65	-4
72	89	-17
57	70	-13

# Paired Samples: Example

- Consider the heart rates before and after running on a treadmill example

Before	After	Difference
55	60	-5
62	75	-13
61	65	-4
72	89	-17
57	70	-13

- Does heart rate significantly change after running on a treadmill?



# Paired Samples: Example

- Consider the heart rates before and after running on a treadmill example

Before	After	Difference
55	60	-5
62	75	-13
61	65	-4
72	89	-17
57	70	-13

- Does heart rate significantly change after running on a treadmill?
- Test at the  $\alpha = 0.01$  significance level

# Paired Samples: Example

Before	After	Difference
55	60	-5
62	75	-13
61	65	-4
72	89	-17
57	70	-13

# Paired Samples: Example

Before	After	Difference
55	60	-5
62	75	-13
61	65	-4
72	89	-17
57	70	-13

- $$\bar{x}_d = \frac{-5 - 13 - 4 - 17 - 13}{5} = -10.4$$

# Paired Samples: Example

Before	After	Difference
55	60	-5
62	75	-13
61	65	-4
72	89	-17
57	70	-13

- $\bar{x}_d = \frac{-5 - 13 - 4 - 17 - 13}{5} = -10.4$

- $s_d^2 = \frac{(-5 + 10.4)^2 + (-13 + 10.4)^2 + (-4 + 10.4)^2 + (-17 + 10.4)^2 + (-13 + 10.4)^2}{5 - 1} = 31.8$

# Paired Samples: Example

$$d = b - a$$

Before	After	Difference
55	60	-5
62	75	-13
61	65	-4
72	89	-17
57	70	-13

$$\bullet \bar{x}_d = \frac{-5 - 13 - 4 - 17 - 13}{5} = -10.4$$

$$\bullet s_d^2 = \frac{(-5 + 10.4)^2 + (-13 + 10.4)^2 + (-4 + 10.4)^2 + (-17 + 10.4)^2 + (-13 + 10.4)^2}{5 - 1} = 31.8$$

$$\bullet s_d = \sqrt{31.8} = 5.64$$

# Paired Samples: Example

# Paired Samples: Example

- $H_0 : \mu_d = 0$  vs.  $H_1 : \mu_d \neq 0$

# Paired Samples: Example

- $H_0 : \mu_d = 0$  vs.  $H_1 : \mu_d \neq 0$
- Calculate t statistic:  $t = \frac{\bar{x}_d - \mu_d}{s_d/\sqrt{n}} = \frac{-10.4 - 0}{5.64/\sqrt{5}} = -4.12$



# Paired Samples: Example

- $H_0 : \mu_d = 0$  vs.  $H_1 : \mu_d \neq 0$
- Calculate t statistic:  $t = \frac{\bar{x}_d - \mu_d}{s_d/\sqrt{n}} = \frac{-10.4 - 0}{5.64/\sqrt{5}} = -4.12$
- Degrees of freedom:  $df = 5 - 1 = 4$

# Paired Samples: Example

- $H_0 : \mu_d = 0$  vs.  $H_1 : \mu_d \neq 0$
- Calculate t statistic:  $t = \frac{\bar{x}_d - \mu_d}{s_d/\sqrt{n}} = \frac{-10.4 - 0}{5.64/\sqrt{5}} = -4.12$
- Degrees of freedom:  $df = 5 - 1 = 4$
- In R: `2*pt(-4.12, 4) = 0.0146`

# Paired Samples: Example

- $H_0 : \mu_d = 0$  vs.  $H_1 : \mu_d \neq 0$
- Calculate t statistic:  $t = \frac{\bar{x}_d - \mu_d}{s_d/\sqrt{n}} = \frac{-10.4 - 0}{5.64/\sqrt{5}} = -4.12$
- Degrees of freedom:  $df = 5 - 1 = 4$
- In R: `2*pt(-4.12, 4) = 0.0146`
- Since  $0.0146 > \alpha = 0.01$ , we fail to reject  $H_0$

# Paired Samples: Example

- $H_0 : \mu_d = 0$  vs.  $H_1 : \mu_d \neq 0$  ← 2-tailed

$$H_0 : \mu_d \geq 0$$

$$H_1 : \mu_d < 0$$


- Calculate t statistic:  $t = \frac{\bar{x}_d - \mu_d}{s_d / \sqrt{n}} = \frac{-10.4 - 0}{5.64 / \sqrt{5}} = -4.12$

- Degrees of freedom:  $df = 5 - 1 = 4$

- In R:  $2 * pt(-4.12, 4) = 0.0146$

- Since  $0.0146 > \alpha = 0.01$ , we fail to reject  $H_0$

- There is insufficient evidence to conclude that the average heart rate after running on a treadmill for 30 minutes is significantly different than the average sitting heart rate at the  $\alpha = 0.01$  significance level



$$n = 5$$

# Paired Samples: Confidence Intervals

# Paired Samples: Confidence Intervals

- We can also calculate confidence intervals for paired differences

# Paired Samples: Confidence Intervals

- We can also calculate confidence intervals for paired differences
- Our (two-sided) interval for unknown  $\sigma$  is  $\bar{x}_d \pm t_{\alpha/2} \frac{s_d}{\sqrt{n}}$


# Paired Samples: Confidence Intervals

- We can also calculate confidence intervals for paired differences
- Our (two-sided) interval for unknown  $\sigma$  is  $\bar{x}_d \pm t_{\alpha/2} \frac{s_d}{\sqrt{n}}$
- Continuing with the heart rates example, we can get the following 99% confidence interval:



# Paired Samples: Confidence Intervals

- We can also calculate confidence intervals for paired differences
- Our (two-sided) interval for unknown  $\sigma$  is  $\bar{x}_d \pm t_{\alpha/2} \frac{s_d}{\sqrt{n}}$
- Continuing with the heart rates example, we can get the following 99% confidence interval:

$$\begin{aligned} \bar{x}_d \pm t_{\alpha/2} \frac{s_d}{\sqrt{n}} &= -10.4 \pm \overbrace{4.60}^{t_{0.01/2, 4}} \cdot \frac{5.65}{\sqrt{5}} \\ &= (-21.62, 1.62) \end{aligned}$$


# Paired Samples: Confidence Intervals

- We can also calculate confidence intervals for paired differences
- Our (two-sided) interval for unknown  $\sigma$  is  $\bar{x}_d \pm t_{\alpha/2} \frac{s_d}{\sqrt{n}}$
- Continuing with the heart rates example, we can get the following 99% confidence interval:

$$\begin{aligned}\bar{x}_d \pm t_{\alpha/2} \frac{s_d}{\sqrt{n}} &= -10.4 \pm 4.60 \cdot \frac{5.65}{\sqrt{5}} \\ &= (-21.62, 1.62)\end{aligned}$$

- I am 99% confident that the interval (-21.62, 1.62) captures the true mean before-after difference in heart rate for subjects running on a treadmill

# Paired Samples: R Code

```
> before <- c(55, 62, 61, 72, 57)
> after <- c(60, 75, 65, 89, 70)
> d <- before-after
> t.test(d)
```

One Sample t-test

```
data: d
t = -4.1239, df = 4, p-value = 0.01457
alternative hypothesis: true mean is not equal to 0
95 percent confidence interval:
 -17.401928 -3.398072
sample estimates:
mean of x
 -10.4
```

```
> t.test(before, after, paired=T)
```

Paired t-test

```
data: before and after
t = -4.1239, df = 4, p-value = 0.01457
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 -17.401928 -3.398072
sample estimates:
mean of the differences
 -10.4
```

# One-sided Paired t-test Example

# One-sided Paired t-test Example

- Setup: Trace metals in drinking water affect the flavor, and an unusually high concentration can pose a health hazard


# One-sided Paired t-test Example

- Setup: Trace metals in drinking water affect the flavor, and an unusually high concentration can pose a health hazard
- Ten pairs of data were taken measuring zinc concentration in bottom water and surface water (two layers of water from same sources)

# One-sided Paired t-test Example

- Setup: Trace metals in drinking water affect the flavor, and an unusually high concentration can pose a health hazard
- Ten pairs of data were taken measuring zinc concentration in bottom water and surface water (two layers of water from same sources)
- Based on this sample, the mean difference in zinc concentrations between the bottom water and surface water was  $\bar{x}_d = 0.0804$  and the standard deviation was  $s_d = 0.9023$

# One-sided Paired t-test Example

- Setup: Trace metals in drinking water affect the flavor, and an unusually high concentration can pose a health hazard
- Ten pairs of data were taken measuring zinc concentration in bottom water and surface water (two layers of water from same sources)
- Based on this sample, the mean difference in zinc concentrations between the bottom water and surface water was  $\bar{x}_d = 0.0804$  and the standard deviation was  $s_d = 0.9023$   

- Q: Does the true average zinc concentration in the bottom water exceed that of surface water?



# One-sided Paired t-test Example

# One-sided Paired t-test Example

- Let  $\alpha = 0.05$

# One-sided Paired t-test Example

- Let  $\alpha = 0.05$

↙ let more than top.

- $H_0 : \mu_d \leq 0$  vs.  $H_1 : \mu_d > 0$

# One-sided Paired t-test Example

- Let  $\alpha = 0.05$
- $H_0 : \mu_d \leq 0$  vs.  $H_1 : \mu_d > 0$
- Calculate t:  $t = \frac{\bar{x}_d - \mu_d}{s_d/\sqrt{n}} = \frac{0.0804 - 0}{0.0923/\sqrt{10}} = \underline{2.88}$

# One-sided Paired t-test Example

- Let  $\alpha = 0.05$
- $H_0 : \mu_d \leq 0$  vs.  $H_1 : \mu_d > 0$
- Calculate t:  $t = \frac{\bar{x}_d - \mu_d}{s_d/\sqrt{n}} = \frac{0.0804 - 0}{0.0923/\sqrt{10}} = 2.88$
- Degrees of freedom:  $df = 10 - 1 = 9$

# One-sided Paired t-test Example

- Let  $\alpha = 0.05$
- $H_0 : \mu_d \leq 0$  vs.  $H_1 : \mu_d > 0$
- Calculate t:  $t = \frac{\bar{x}_d - \mu_d}{s_d/\sqrt{n}} = \frac{0.0804 - 0}{0.0923/\sqrt{10}} = 2.88$
- Degrees of freedom:  $df = 10 - 1 = 9$
- R: `1-pt (2.88, 9) = 0.0091`

# One-sided Paired t-test Example

- Let  $\alpha = 0.05$
- $H_0 : \mu_d \leq 0$  vs.  $H_1 : \mu_d > 0$
- Calculate t:  $t = \frac{\bar{x}_d - \mu_d}{s_d/\sqrt{n}} = \frac{0.0804 - 0}{0.0923/\sqrt{10}} = 2.88$
- Degrees of freedom:  $df = 10 - 1 = 9$
- R: `1-pt (2.88, 9) = 0.0091`
- Since  $0.0091 < \alpha = 0.05$ , we reject  $H_0$

# One-sided Paired t-test Example

- Let  $\alpha = 0.05$
- $H_0 : \mu_d \leq 0$  vs.  $H_1 : \mu_d > 0$
- Calculate t:  $t = \frac{\bar{x}_d - \mu_d}{s_d/\sqrt{n}} = \frac{0.0804 - 0}{0.0923/\sqrt{10}} = 2.88$
- Degrees of freedom:  $df = 10 - 1 = 9$
- R: `1-pt(2.88, 9) = 0.0091`
- Since  $0.0091 < \alpha = 0.05$ , we reject  $H_0$
- There is sufficient evidence to conclude that, on average, the bottom zinc concentration is higher than the surface zinc concentration



# Independent Samples

# Independent Samples

- Suppose we have measurements on two samples of subjects

# Independent Samples

- Suppose we have measurements on two samples of subjects
  - Heart rates for a group of people who have been sitting

# Independent Samples

- Suppose we have measurements on two samples of subjects
  - Heart rates for a group of people who have been sitting
  - Heart rates for a group of people who have been running

# Independent Samples

- Suppose we have measurements on two samples of subjects
  - Heart rates for a group of people who have been sitting
  - Heart rates for a group of people who have been running
- The two underlying populations are independent and normally distributed

# Independent Samples

- Suppose we have measurements on two samples of subjects
  - Heart rates for a group of people who have been sitting
  - Heart rates for a group of people who have been running
- The two underlying populations are independent and normally distributed
- The first population has mean  $\mu_1$  and the second population has mean  $\mu_2$

# Independent Samples

- Suppose we have measurements on two samples of subjects
  - Heart rates for a group of people who have been sitting
  - Heart rates for a group of people who have been running
- The two underlying populations are independent and normally distributed
- The first population has mean  $\mu_1$  and the second population has mean  $\mu_2$
- Test whether the two populations are identical

# Independent Samples

- Suppose we have measurements on two samples of subjects
  - Heart rates for a group of people who have been sitting
  - Heart rates for a group of people who have been running
- The two underlying populations are independent and normally distributed
- The first population has mean  $\mu_1$  and the second population has mean  $\mu_2$
- Test whether the two populations are identical
  - $H_0 : \mu_1 = \mu_2$  vs.  $H_1 : \mu_1 \neq \mu_2$



# Independent Samples

$$\bar{x} \sim \bar{X}$$

- Suppose we have measurements on two samples of subjects
  - Heart rates for a group of people who have been sitting
  - Heart rates for a group of people who have been running
- The two underlying populations are independent and normally distributed
- The first population has mean  $\mu_1$  and the second population has mean  $\mu_2$
- Test whether the two populations are identical

$$H_0 : \mu_1 = \mu_2 \text{ vs. } H_1 : \mu_1 \neq \mu_2 \quad \rightarrow$$

$$\bar{x}_1 \sim \bar{X}_1 = \bar{x}_2 \sim \bar{X}_2$$

$$N(\mu, \sigma)$$

$$\text{Or, equivalently, } H_0 : \mu_1 - \mu_2 = 0 \text{ vs. } H_1 : \mu_1 - \mu_2 \neq 0$$

$$\bar{x}_1 - \bar{x}_2 \sim \bar{X}_1 - \bar{X}_2$$

# Independent Samples

# Independent Samples

- Population 1: mean  $\mu_1$  and variance  $\sigma_1^2$

# Independent Samples

- Population 1: mean  $\mu_1$  and variance  $\sigma_1^2$ 
  - Draw sample of size  $n_1$  with sample mean  $\bar{x}_1$  and sample variance  $s_1^2$

# Independent Samples

- Population 1: mean  $\mu_1$  and variance  $\sigma_1^2$ 
  - Draw sample of size  $n_1$  with sample mean  $\bar{x}_1$  and sample variance  $s_1^2$
- Population 2: mean  $\mu_2$  and variance  $\sigma_2^2$

# Independent Samples

- Population 1: mean  $\mu_1$  and variance  $\sigma_1^2$ 
  - Draw sample of size  $n_1$  with sample mean  $\bar{x}_1$  and sample variance  $s_1^2$
- Population 2: mean  $\mu_2$  and variance  $\sigma_2^2$ 
  - Draw sample of size  $n_2$  with sample mean  $\bar{x}_2$  and sample variance  $s_2^2$

# Independent Samples

- Population 1: mean  $\mu_1$  and variance  $\sigma_1^2$ 
  - Draw sample of size  $n_1$  with sample mean  $\bar{x}_1$  and sample variance  $s_1^2$
- Population 2: mean  $\mu_2$  and variance  $\sigma_2^2$ 
  - Draw sample of size  $n_2$  with sample mean  $\bar{x}_2$  and sample variance  $s_2^2$
- We can compare the means of these two populations in two different ways

# Independent Samples

- Population 1: mean  $\mu_1$  and variance  $\sigma_1^2$ 
  - Draw sample of size  $n_1$  with sample mean  $\bar{x}_1$  and sample variance  $s_1^2$
- Population 2: mean  $\mu_2$  and variance  $\sigma_2^2$ 
  - Draw sample of size  $n_2$  with sample mean  $\bar{x}_2$  and sample variance  $s_2^2$
- We can compare the means of these two populations in two different ways
  - Equal population variances: known ( $\sigma_1^2 = \sigma_2^2$ ) or unknown (t-test)



# Independent Samples

- Population 1: mean  $\mu_1$  and variance  $\sigma_1^2$ 
  - Draw sample of size  $n_1$  with sample mean  $\bar{x}_1$  and sample variance  $s_1^2$
- Population 2: mean  $\mu_2$  and variance  $\sigma_2^2$ 
  - Draw sample of size  $n_2$  with sample mean  $\bar{x}_2$  and sample variance  $s_2^2$
- We can compare the means of these two populations in two different ways
  - Equal population variances: known ( $\sigma_1^2 = \sigma_2^2$ ) or unknown (t-test)
  - Unequal population variances:  $\sigma_1^2 \neq \sigma_2^2$

# Equal, Known Variances

# Equal, Known Variances

- Consider situations where it is either known or reasonable to assume that the two population variances are the same, and we know  $\sigma^2 = \sigma_1^2 = \sigma_2^2$

# Equal, Known Variances

- Consider situations where it is either known or reasonable to assume that the two population variances are the same, and we know  $\sigma^2 = \sigma_1^2 = \sigma_2^2$
- Suppose we are interested in the difference between the two population means

# Equal, Known Variances

- Consider situations where it is either known or reasonable to assume that the two population variances are the same, and we know  $\sigma^2 = \sigma_1^2 = \sigma_2^2$
- Suppose we are interested in the difference between the two population means
- By extension of the CLT,  $\bar{X}_1 - \bar{X}_2$  is approximately normal with mean  $\mu_1 - \mu_2$  and standard

error  $\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$

$$\text{var}(\bar{X}_1 - \bar{X}_2) = \text{var}(\bar{X}_1) + \text{var}(\bar{X}_2)$$

$$\sigma^2 = \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}$$

$$\sigma = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

# Equal, Known Variances

- Consider situations where it is either known or reasonable to assume that the two population variances are the same, and we know  $\sigma^2 = \sigma_1^2 = \sigma_2^2$
- Suppose we are interested in the difference between the two population means
- By extension of the CLT,  $\bar{X}_1 - \bar{X}_2$  is approximately normal with mean  $\mu_1 - \mu_2$  and standard

error  $\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$

$\sqrt{\frac{\sigma^2}{n_1} + \frac{\sigma^2}{n_2}}$

- Because we assume  $\sigma^2 = \sigma_1^2 = \sigma_2^2$ , the standard error is  $\sqrt{\sigma^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)} = \sigma \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$

# Equal, Known Variances

- Consider situations where it is either known or reasonable to assume that the two population variances are the same, and we know  $\sigma^2 = \sigma_1^2 = \sigma_2^2$
- Suppose we are interested in the difference between the two population means
- By extension of the CLT,  $\bar{X}_1 - \bar{X}_2$  is approximately normal with mean  $\mu_1 - \mu_2$  and standard error  $\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$
- Because we assume  $\sigma^2 = \sigma_1^2 = \sigma_2^2$ , the standard error is  $\sqrt{\sigma^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}$
- Then, we can use a z-test with  $z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\sigma^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}}$  as our test statistic

# Equal, Unknown Variances



# Equal, Unknown Variances

- When  $\sigma^2$  is unknown, we use a sample estimate (t-test)

# Equal, Unknown Variances

- When  $\sigma^2$  is unknown, we use a sample estimate (t-test)
- We assume that both populations have the same variance, so we want to consider both samples in our calculation of the sample variance,  $s_p^2$

# Equal, Unknown Variances

- When  $\sigma^2$  is unknown, we use a sample estimate (t-test)
- We assume that both populations have the same variance, so we want to consider both samples in our calculation of the sample variance,  $s_p^2$
- We can think of  $s_p^2$  as the *pooled* estimate of common variance  $\sigma^2$ :

# Equal, Unknown Variances

- When  $\sigma^2$  is unknown, we use a sample estimate (t-test)
- We assume that both populations have the same variance, so we want to consider both samples in our calculation of the sample variance,  $s_p^2$
- We can think of  $s_p^2$  as the *pooled* estimate of common variance  $\sigma^2$ :


- $$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{(n_1 - 1) + (n_2 - 1)}$$

# Equal, Unknown Variances

- When  $\sigma^2$  is unknown, we use a sample estimate (t-test)
- We assume that both populations have the same variance, so we want to consider both samples in our calculation of the sample variance,  $s_p^2$
- We can think of  $s_p^2$  as the *pooled* estimate of common variance  $\sigma^2$ :
  - $$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{(n_1 - 1) + (n_2 - 1)}$$
- In other words,  $s_p^2$  is the weighted average of the two sample variances,  $s_1^2$  and  $s_2^2$ , where the weights are the degrees of freedom for each sample

# Equal, Unknown Variances

# Equal, Unknown Variances

- We can use a t-test statistic:  $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}}$  

# Equal, Unknown Variances

- We can use a t-test statistic:  $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}}$
- The test statistic  $t$  has a t distribution with  $(n_1 - 1) + (n_2 - 1) = n_1 + n_2 - 2$  degrees of freedom



# Equal, Unknown Variances

- We can use a t-test statistic:  $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}}$
- The test statistic  $t$  has a t distribution with  $(n_1 - 1) + (n_2 - 1) = n_1 + n_2 - 2$  degrees of freedom
- Determine the p-value,  $p$ , based on this t distribution

# Equal, Unknown Variances

- We can use a t-test statistic:  $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}}$
- The test statistic  $t$  has a t distribution with  $(n_1 - 1) + (n_2 - 1) = n_1 + n_2 - 2$  degrees of freedom
- Determine the p-value,  $p$ , based on this t distribution
- Reject  $H_0$  if  $p \leq \alpha$

# Equal, Unknown Variances: Example

# Equal, Unknown Variances: Example

- Setup: A study examines whether targeted political ads makes potential voters more likely to donate money to candidates

# Equal, Unknown Variances: Example

- Setup: A study examines whether targeted political ads makes potential voters more likely to donate money to candidates
- A sample of 54 potential voters who saw targeted political ads had mean donation amount  $\bar{x}_1 = 240$  USD with standard deviation  $s_1 = 30$  USD

# Equal, Unknown Variances: Example

- Setup: A study examines whether targeted political ads makes potential voters more likely to donate money to candidates
  - A sample of 54 potential voters who saw targeted political ads had mean donation amount  $\bar{x}_1 = 240$  USD with standard deviation  $s_1 = 30$  USD
  - A sample of 49 potential voters who did not see targeted political ads had mean donation amount  $\bar{x}_2 = 220$  USD with standard deviation  $s_2 = 48$  USD

# Equal, Unknown Variances: Example

- Setup: A study examines whether targeted political ads makes potential voters more likely to donate money to candidates
- A sample of 54 potential voters who saw targeted political ads had mean donation amount  $\bar{x}_1 = 240$  USD with standard deviation  $s_1 = 30$  USD
- A sample of 49 potential voters who did not see targeted political ads had mean donation amount  $\bar{x}_2 = 220$  USD with standard deviation  $s_2 = 48$  USD
- Is there a significant difference (at the  $\alpha = 0.05$  significance level) in average donations between potential voters who saw targeted political ads and those who did not, assuming equal variances?

Q

# Equal, Unknown Variances: Example



# Equal, Unknown Variances: Example

- Hypotheses:  $H_0 : \mu_1 - \mu_2 = 0$ ,  $H_1 : \mu_1 - \mu_2 \neq 0$ , significance  $\alpha = 0.05$

# Equal, Unknown Variances: Example

- Hypotheses:  $H_0 : \mu_1 - \mu_2 = 0$ ,  $H_1 : \mu_1 - \mu_2 \neq 0$ , significance  $\alpha = 0.05$
- Calculate the pooled (sample) variance:

$$s_p^2 = \frac{(n_1 - 1) s_1^2 + (n_2 - 1) s_2^2}{(n_1 - 1) + (n_2 - 1)} = \frac{(54 - 1) 30^2 + (49 - 1) 48^2}{54 - 1 + 49 - 1}$$
$$= 1567$$

# Equal, Unknown Variances: Example

- Hypotheses:  $H_0 : \mu_1 - \mu_2 = 0$ ,  $H_1 : \mu_1 - \mu_2 \neq 0$ , significance  $\alpha = 0.05$
- Calculate the pooled (sample) variance:

# Equal, Unknown Variances: Example

- Hypotheses:  $H_0 : \mu_1 - \mu_2 = 0$ ,  $H_1 : \mu_1 - \mu_2 \neq 0$ , significance  $\alpha = 0.05$
- Calculate the pooled (sample) variance:

1567

- Next, calculate the t-statistic:

$$t = \frac{(\bar{x}_2 - \bar{x}_1) - (\mu_2 - \mu_1)}{\sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}} = \frac{(220 - 240) - 0}{\sqrt{1567 \left( \frac{1}{59} + \frac{1}{48} \right)}} = \boxed{2.56}$$

# Equal, Unknown Variances: Example

- Hypotheses:  $H_0 : \mu_1 - \mu_2 = 0$ ,  $H_1 : \mu_1 - \mu_2 \neq 0$ , significance  $\alpha = 0.05$
- Calculate the pooled (sample) variance:
- Next, calculate the t-statistic:

# Equal, Unknown Variances: Example

# Equal, Unknown Variances: Example

- We compare this t-value with a t distribution with  $\overset{n_1}{(54 - 1)} + \overset{n_2}{(49 - 1)} = 101$  df

$$t = 2.561$$

# Equal, Unknown Variances: Example

- We compare this t-value with a t distribution with  $(54 - 1) + (49 - 1) = 101$  df
- In R:  $2 * (1 - pt(2.561, 101)) = \underline{0.012}$

$t$   $\uparrow$   $\uparrow$   
 $df$



# Equal, Unknown Variances: Example

- We compare this t-value with a t distribution with  $(54 - 1) + (49 - 1) = 101$  df
- In R: `2 * (1 - pt (2.561, 101) ) = 0.012`
- Since the p-value is  $0.012 < 0.05$ , we reject the null hypothesis

# Equal, Unknown Variances: Example

- We compare this t-value with a t distribution with  $(\underline{54 - 1}) + (\underline{49 - 1}) = 101$  df
- In R:  $2 * (1 - pt(2.561, 101)) = 0.012$
- Since the p-value is  $0.012 < 0.05$ , we reject the null hypothesis
- There is significant evidence to conclude that the average donation amount from potential voters who saw targeted political ads is significantly different from the average donation amount from potential voters who did not see targeted political ads

# Equal, Unknown Variances: Example

# Equal, Unknown Variances: Example

- What if we want a  $(1 - \alpha)\%$  confidence interval for  $\mu_1 - \mu_2$ ?

# Equal, Unknown Variances: Example

- What if we want a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$ ?

- We can write this as  $(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}$

# Equal, Unknown Variances: Example

- What if we want a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$ ?
- We can write this as  $(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}$
- We are  $(1 - \alpha) \%$  confident that this interval contains the true mean difference  $\mu_1 - \mu_2$

# Equal, Unknown Variances: Example

- What if we want a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$ ?
- We can write this as  $(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}$
- We are  $(1 - \alpha) \%$  confident that this interval contains the true mean difference  $\mu_1 - \mu_2$
- In our voter example, a 95% confidence interval for the mean difference in donations is as follows:

# Equal, Unknown Variances: Example

- What if we want a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$ ?
- We can write this as  $(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}$
- We are  $(1 - \alpha) \%$  confident that this interval contains the true mean difference  $\mu_1 - \mu_2$
- In our voter example, a 95% confidence interval for the mean difference in donations is as follows:

$$\begin{aligned} (\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)} &= (240 - 220) \pm 1.984 \sqrt{1567.248 \left( \frac{1}{54} + \frac{1}{49} \right)} \\ &= (4.50, 35.50) \end{aligned}$$



# Equal, Unknown Variances: Example

- What if we want a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$ ? ↖ ad ↗ no ed
- We can write this as  $(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}$
- We are  $(1 - \alpha) \%$  confident that this interval contains the true mean difference  $\mu_1 - \mu_2$
- In our voter example, a 95% confidence interval for the mean difference in donations is as follows:

$$\begin{aligned} (\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)} &= (240 - 220) \pm 1.984 \sqrt{1567.248 \left( \frac{1}{54} + \frac{1}{49} \right)} \\ &= (4.50, 35.50) \end{aligned}$$

- We are 95% confident that the interval (4.50, 35.50) captures the true difference in average donation amounts

# Unequal Variance (Welch t-test)

# Unequal Variance (Welch t-test)

- Let's now consider the situation where the variances of the two populations are not assumed to be equal

# Unequal Variance (Welch t-test)

- Let's now consider the situation where the variances of the two populations are not assumed to be equal
- Need to modify the two-sample t-test

# Unequal Variance (Welch t-test)

- Let's now consider the situation where the variances of the two populations are not assumed to be equal
- Need to modify the two-sample t-test
- Instead of used a pooled estimate of the variance, we substitute  $s_1^2$  for  $\sigma_1^2$  and  $s_2^2$  for  $\sigma_2^2$

# Unequal Variance (Welch t-test)

- Let's now consider the situation where the variances of the two populations are not assumed to be equal
- Need to modify the two-sample t-test
- Instead of used a pooled estimate of the variance, we substitute  $s_1^2$  for  $\sigma_1^2$  and  $s_2^2$  for  $\sigma_2^2$

• Test statistic:  $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$

$$\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

# Unequal Variance (Welch t-test)

- Let's now consider the situation where the variances of the two populations are not assumed to be equal
- Need to modify the two-sample t-test
- Instead of used a pooled estimate of the variance, we substitute  $s_1^2$  for  $\sigma_1^2$  and  $s_2^2$  for  $\sigma_2^2$

- Test statistic:  $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$
- The exact distribution for this t-statistic is difficult to derive

# Unequal Variance (Welch t-test)

- Let's now consider the situation where the variances of the two populations are not assumed to be equal
- Need to modify the two-sample t-test
- Instead of used a pooled estimate of the variance, we substitute  $s_1^2$  for  $\sigma_1^2$  and  $s_2^2$  for  $\sigma_2^2$

Test statistic:  $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$

- The exact distribution for this t-statistic is difficult to derive
  - Use an approximation

$t, df$




# Unequal Variance (Welch t-test)

# Unequal Variance (Welch t-test)

- Need to come up with a value for the degrees of freedom (Welch-Satterthwaite equation)

# Unequal Variance (Welch t-test)

- Need to come up with a value for the degrees of freedom (Welch-Satterthwaite equation)

$$\nu = \frac{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2}{\left( \frac{\left( \frac{s_1^2}{n_1} \right)^2}{n_1 - 1} + \frac{\left( \frac{s_2^2}{n_2} \right)^2}{n_2 - 1} \right)}$$


# Unequal Variance (Welch t-test)

- Need to come up with a value for the degrees of freedom (Welch-Satterthwaite equation)

$$\nu = \frac{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2}{\left( \frac{\left( \frac{s_1^2}{n_1} \right)^2}{n_1 - 1} + \frac{\left( \frac{s_2^2}{n_2} \right)^2}{n_2 - 1} \right)}$$

- Under  $H_0 : \mu_1 = \mu_2$ , we can approximate the distribution of  $t$  by a t distribution with  $\nu$  degrees of freedom = nu

# Unequal Variance (Welch t-test)

- Need to come up with a value for the degrees of freedom (Welch-Satterthwaite equation)

$$\nu = \frac{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2}{\left( \frac{\left( \frac{s_1^2}{n_1} \right)^2}{n_1 - 1} + \frac{\left( \frac{s_2^2}{n_2} \right)^2}{n_2 - 1} \right)}$$

- Under  $H_0 : \mu_1 = \mu_2$ , we can approximate the distribution of  $t$  by a t distribution with  $\nu$  degrees of freedom
- Calculate the p-value,  $p$ , based on this t distribution

$p_t(t, \nu)$

# Unequal Variance (Welch t-test)

- Need to come up with a value for the degrees of freedom (Welch-Satterthwaite equation)

$$\nu = \frac{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2}{\left( \frac{\left( \frac{s_1^2}{n_1} \right)^2}{n_1 - 1} + \frac{\left( \frac{s_2^2}{n_2} \right)^2}{n_2 - 1} \right)}$$

- Under  $H_0 : \mu_1 = \mu_2$ , we can approximate the distribution of  $t$  by a t distribution with  $\nu$  degrees of freedom
- Calculate the p-value,  $p$ , based on this t distribution
- Reject  $H_0$  if  $p \leq \alpha$

# Unequal Variance (Welch t-test): Example

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?



# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3
  - Sample standard deviation: 1.86

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3
  - Sample standard deviation: 1.86
- Take a sample of 7 people between 10 and 30 years old

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3
  - Sample standard deviation: 1.86
- Take a sample of 7 people between 10 and 30 years old
  - Sample mean blood calcium level: 10.6

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3
  - Sample standard deviation: 1.86
- Take a sample of 7 people between 10 and 30 years old
  - Sample mean blood calcium level: 10.6
  - Sample standard deviation: 0.92

# Unequal Variance (Welch t-test): Example

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?



# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3
  - Sample standard deviation: 1.86

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3
  - Sample standard deviation: 1.86
- Take a sample of 7 people between 10 and 30 years old

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3
  - Sample standard deviation: 1.86
- Take a sample of 7 people between 10 and 30 years old
  - Sample mean blood calcium level: 10.6

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3
  - Sample standard deviation: 1.86
- Take a sample of 7 people between 10 and 30 years old
  - Sample mean blood calcium level: 10.6
  - Sample standard deviation: 0.92

# Unequal Variance (Welch t-test): Example

- Setup: Are average blood calcium levels different for people over 60 years old compared to 10-30 year olds?
- Take a sample of 15 people who are over 60 years old
  - Sample mean blood calcium level: 9.3
  - Sample standard deviation: 1.86
- Take a sample of 7 people between 10 and 30 years old
  - Sample mean blood calcium level: 10.6
  - Sample standard deviation: 0.92
- Test at  $\alpha = 0.05$  significance level

# Unequal Variance (Welch t-test): Example



# Unequal Variance (Welch t-test): Example

- Test at  $\alpha = 0.05$  significance level

# Unequal Variance (Welch t-test): Example

- Test at  $\alpha = 0.05$  significance level
- Let  $\mu_1$  be the average blood calcium level for people over 60 years old

# Unequal Variance (Welch t-test): Example

- Test at  $\alpha = 0.05$  significance level
- Let  $\mu_1$  be the average blood calcium level for people over 60 years old
- Let  $\mu_2$  be the average blood calcium level for people between 10-30 years old

# Unequal Variance (Welch t-test): Example

- Test at  $\alpha = 0.05$  significance level
- Let  $\mu_1$  be the average blood calcium level for people over 60 years old
- Let  $\mu_2$  be the average blood calcium level for people between 10-30 years old
- Hypotheses:  $H_0 : \mu_1 - \mu_2 = 0$  vs.  $H_1 : \mu_1 - \mu_2 \neq 0$

# Unequal Variance (Welch t-test): Example

# Unequal Variance (Welch t-test): Example

- Calculate t-statistic:

# Unequal Variance (Welch t-test): Example

- Calculate t-statistic:

# Unequal Variance (Welch t-test): Example

- Calculate t-statistic:
- Calculate degrees of freedom ( $\nu$ ):



# Unequal Variance (Welch t-test): Example

- Calculate t-statistic:
- Calculate degrees of freedom ( $\nu$ ):

# Unequal Variance (Welch t-test): Example

- Calculate t-statistic: 
$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}} = \frac{9.3 - 10.6}{\sqrt{\frac{1.86^2}{15} + \frac{0.92^2}{7}}} = -2.19$$

- Calculate degrees of freedom ( $\nu$ ):

- R: 
$$\nu = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)^2}{\left[\frac{\left(\frac{s_1^2}{n_1}\right)^2}{n_1 - 1} + \frac{\left(\frac{s_2^2}{n_2}\right)^2}{n_2 - 1}\right]} = \frac{\left(\frac{1.86^2}{15} + \frac{0.92^2}{7}\right)^2}{\left[\frac{\left(\frac{1.86^2}{15}\right)^2}{14} + \frac{\left(\frac{0.92^2}{7}\right)^2}{6}\right]} = 19.82$$
  

$$p = 2 \times pt(-2.19, 19.82) = 0.0407$$

# Unequal Variance (Welch t-test): Example

- Calculate t-statistic:
- Calculate degrees of freedom ( $\nu$ ):
- R:
- Since the p-value is less than  $\alpha = 0.05$ , we reject the null hypothesis and conclude a difference in average blood calcium level between the two age groups

# Unequal Variance (Welch t-test): Example

# Unequal Variance (Welch t-test): Example

- Similarly, we can calculate a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$  as

$$\underline{(\bar{x}_1 - \bar{x}_2)} \pm \underline{t_{\alpha/2}} \sqrt{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)} \quad \text{SE}$$

# Unequal Variance (Welch t-test): Example

- Similarly, we can calculate a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$  as

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)}$$

- We are  $(1 - \alpha) \%$  confident that this interval contains the true mean difference  $\mu_1 - \mu_2$

# Unequal Variance (Welch t-test): Example

- Similarly, we can calculate a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$  as

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)}$$

- We are  $(1 - \alpha) \%$  confident that this interval contains the true mean difference  $\mu_1 - \mu_2$
- For our blood calcium example, a 95% confidence interval for the difference in average blood calcium levels for 60+ year olds and 10-30 year olds is

# Unequal Variance (Welch t-test): Example

- Similarly, we can calculate a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$  as

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)}$$

- We are  $(1 - \alpha) \%$  confident that this interval contains the true mean difference  $\mu_1 - \mu_2$
- For our blood calcium example, a 95% confidence interval for the difference in average blood calcium levels for 60+ year olds and 10-30 year olds is

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)} = (9.3 - 10.6) \pm 2.09 \sqrt{\left( \frac{1.86^2}{15} + \frac{0.92^2}{7} \right)} = (-2.54, -0.06)$$



# Unequal Variance (Welch t-test): Example

- Similarly, we can calculate a  $(1 - \alpha) \%$  confidence interval for  $\mu_1 - \mu_2$  as

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)}$$

- We are  $(1 - \alpha) \%$  confident that this interval contains the true mean difference  $\mu_1 - \mu_2$
- For our blood calcium example, a 95% confidence interval for the difference in average blood calcium levels for 60+ year olds and 10-30 year olds is

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)} = (9.3 - 10.6) \pm 2.09 \sqrt{\left( \frac{1.86^2}{15} + \frac{0.92^2}{7} \right)} = (-2.54, -0.06)$$

- I am 95% confident that the interval  $(-2.54, -0.06)$  captures the true difference in average blood calcium levels between 60+ year olds and 10-30 year olds

# Unequal Variance (Welch t-test): Example

# Unequal Variance (Welch t-test): Example

- Setup: Do babies born in Rochester have larger birthweights than babies born in Buffalo?

# Unequal Variance (Welch t-test): Example

- Setup: Do babies born in Rochester have larger birthweights than babies born in Buffalo?
- Let  $\mu_1$  be the mean birthweight of babies born in Rochester

# Unequal Variance (Welch t-test): Example

- Setup: Do babies born in Rochester have larger birthweights than babies born in Buffalo?
- Let  $\mu_1$  be the mean birthweight of babies born in Rochester
- Let  $\mu_2$  be the mean birthweight of babies born in Buffalo

# Unequal Variance (Welch t-test): Example

- Setup: Do babies born in Rochester have larger birthweights than babies born in Buffalo?
- Let  $\mu_1$  be the mean birthweight of babies born in Rochester
- Let  $\mu_2$  be the mean birthweight of babies born in Buffalo
- Assume unequal variances

# Unequal Variance (Welch t-test): Example

- Setup: Do babies born in Rochester have larger birthweights than babies born in Buffalo?
- Let  $\mu_1$  be the mean birthweight of babies born in Rochester
- Let  $\mu_2$  be the mean birthweight of babies born in Buffalo
- Assume unequal variances
- Let  $\alpha = 0.05$  be our significance level

# Unequal Variance (Welch t-test): Example



# Unequal Variance (Welch t-test): Example

- Hypotheses:  $H_0 : \mu_1 \leq \mu_2$  vs.  $H_1 : \mu_1 > \mu_2$ , samples of sizes  $n_1 = 140$  and  $n_2 = 172$

# Unequal Variance (Welch t-test): Example

- Hypotheses:  $H_0 : \mu_1 \leq \mu_2$  vs.  $H_1 : \mu_1 > \mu_2$ , samples of sizes  $n_1 = 140$  and  $n_2 = 172$
- $\bar{x}_1 = 8.2$  lbs,  $\bar{x}_2 = 7.9$  lbs,  $s_1^2 = 1.4$  lbs<sup>2</sup>, and  $s_2^2 = 1.1$  lbs<sup>2</sup>

# Unequal Variance (Welch t-test): Example

- Hypotheses:  $H_0 : \mu_1 \leq \mu_2$  vs.  $H_1 : \mu_1 > \mu_2$ , samples of sizes  $n_1 = 140$  and  $n_2 = 172$
- $\bar{x}_1 = 8.2$  lbs,  $\bar{x}_2 = 7.9$  lbs,  $s_1^2 = 1.4$  lbs<sup>2</sup>, and  $s_2^2 = 1.1$  lbs<sup>2</sup>
- Calculating  $t =$

# Unequal Variance (Welch t-test): Example

- Hypotheses:  $H_0 : \mu_1 \leq \mu_2$  vs.  $H_1 : \mu_1 > \mu_2$ , samples of sizes  $n_1 = 140$  and  $n_2 = 172$
- $\bar{x}_1 = 8.2$  lbs,  $\bar{x}_2 = 7.9$  lbs,  $s_1^2 = 1.4$  lbs<sup>2</sup>, and  $s_2^2 = 1.1$  lbs<sup>2</sup>
- Calculating  $t =$
  
- Calculating  $\nu =$

# Unequal Variance (Welch t-test): Example

- Hypotheses:  $H_0 : \mu_1 \leq \mu_2$  vs.  $H_1 : \mu_1 > \mu_2$ , samples of sizes  $n_1 = 140$  and  $n_2 = 172$

- $\bar{x}_1 = 8.2$  lbs,  $\bar{x}_2 = 7.9$  lbs,  $s_1^2 = 1.4$  lbs<sup>2</sup>, and  $s_2^2 = 1.1$  lbs<sup>2</sup>

- Calculating  $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}} = \frac{0.3}{\sqrt{\frac{1.4^2}{140} + \frac{1.1^2}{172}}} = 2.1$

- Calculating  $\nu =$


$$n_1 = 140$$
$$n_2 = 172$$

$$\nu = 260.353$$

- In R:  $p = 1 - pt(2.1, 260.363) = \underline{0.018} < \alpha = 0.05$

$$t > 0$$

$$t < 0$$



$$1 - pt(t)$$

$$pt(t)$$

# Unequal Variance (Welch t-test): Example


- Hypotheses:  $H_0 : \mu_1 \leq \mu_2$  vs.  $H_1 : \mu_1 > \mu_2$ , samples of sizes  $n_1 = 140$  and  $n_2 = 172$

- $\bar{x}_1 = 8.2$  lbs,  $\bar{x}_2 = 7.9$  lbs,  $s_1^2 = 1.4$  lbs<sup>2</sup>, and  $s_2^2 = 1.1$  lbs<sup>2</sup>

- Calculating  $t =$

$$z = \frac{\bar{x} - \mu}{\sigma / \sqrt{n}}$$

- Calculating  $\nu =$



- In R:

- Conclusion:

# Unequal Variance (Welch t-test): Example

- Hypotheses:  $H_0 : \mu_1 \leq \mu_2$  vs.  $H_1 : \mu_1 > \mu_2$ , samples of sizes  $n_1 = 140$  and  $n_2 = 172$
- $\bar{x}_1 = 8.2$  lbs,  $\bar{x}_2 = 7.9$  lbs,  $s_1^2 = 1.4$  lbs<sup>2</sup>, and  $s_2^2 = 1.1$  lbs<sup>2</sup>
- Calculating  $t =$

$$\Pr(\text{Dell} \mid \text{batt} > 8.5) = \frac{\Pr(\text{batt} > 8.5 \mid \text{Dell}) \Pr(\text{Dell})}{\Pr(\text{batt} > 8.5)}$$

- Calculating  $\nu =$

$$\Pr(\text{batt} > 8.5 \mid \text{Dell}) \Pr(\text{Dell}) + \Pr(\text{batt} > 8.5 \mid \text{Mac}) \Pr(\text{Mac})$$

- In R:
- Conclusion: