

# Chapter 4: Probability and Combinatorics

DSCC 462

Computational Introduction to Statistics

Anson Kahng

Fall 2022

# Probability

# Probability

# Probability

- The outcome that we will observe is often uncertain

# Probability

- The outcome that we will observe is often uncertain
  - Flip a coin

# Probability

- The outcome that we will observe is often uncertain
  - Flip a coin
  - Draw a card

# Probability

- The outcome that we will observe is often uncertain
  - Flip a coin
  - Draw a card
  - Roll a die

# Probability

- The outcome that we will observe is often uncertain
  - Flip a coin
  - Draw a card
  - Roll a die
  - Income of a selected individual



# Probability

- The outcome that we will observe is often uncertain
  - Flip a coin
  - Draw a card
  - Roll a die
  - Income of a selected individual
- We want to find the *probability* of each event happening

# Probability

- The outcome that we will observe is often uncertain
  - Flip a coin
  - Draw a card
  - Roll a die
  - Income of a selected individual
- We want to find the *probability* of each event happening
- Probability is the mathematics of random occurrences

# Events

# Events

- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$

# Events

- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$ 
  - Example: Flip of a coin,  $S = \{\text{Heads}, \text{Tails}\}$

# Events

- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$ 
  - Example: Flip of a coin,  $S = \{\text{Heads}, \text{Tails}\}$
- *A random experiment* occurs when an element of  $S$  is randomly selected

# Events

- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$ 
  - Example: Flip of a coin,  $S = \{\text{Heads}, \text{Tails}\}$
- *A random experiment* occurs when an element of  $S$  is randomly selected
- **Event:** The basic element to which probability can be applied

# Events

- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$ 
  - Example: Flip of a coin,  $S = \{\text{Heads}, \text{Tails}\}$
- *A random experiment* occurs when an element of  $S$  is randomly selected
- **Event:** The basic element to which probability can be applied
  - “Probability of an event happening”



# Events

- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$ 
  - Example: Flip of a coin,  $S = \{\text{Heads}, \text{Tails}\}$
- *A random experiment* occurs when an element of  $S$  is randomly selected
- **Event:** The basic element to which probability can be applied
  - “Probability of an event happening”
  - Events can be possible outcomes or observed values

# Events

- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$ 
  - Example: Flip of a coin,  $S = \{\text{Heads}, \text{Tails}\}$
- A *random experiment* occurs when an element of  $S$  is randomly selected
- **Event:** The basic element to which probability can be applied
  - “Probability of an event happening”
  - Events can be possible outcomes or observed values
  - Either happens or it does not

# Events

- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$ 
  - Example: Flip of a coin,  $S = \{\text{Heads}, \text{Tails}\}$
- A *random experiment* occurs when an element of  $S$  is randomly selected
- **Event:** The basic element to which probability can be applied
  - “Probability of an event happening”
  - Events can be possible outcomes or observed values
  - Either happens or it does not
- Events are represented by uppercase letters:  $A, B, C, \dots$

# Events


- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$ 
  - Example: Flip of a coin,  $S = \{\text{Heads}, \text{Tails}\}$
- *A random experiment* occurs when an element of  $S$  is randomly selected
- **Event:** The basic element to which probability can be applied
  - “Probability of an event happening”
  - Events can be possible outcomes or observed values
  - Either happens or it does not
- Events are represented by uppercase letters:  $A, B, C, \dots$
- List the event in  $\{ \}$  brackets

# Events

- **Sample space:** All possible outcomes that can be observed in a given situation, denoted  $S$ 
  - Example: Flip of a coin,  $S = \{\text{Heads}, \text{Tails}\}$
- A *random experiment* occurs when an element of  $S$  is randomly selected
- **Event:** The basic element to which probability can be applied
  - “Probability of an event happening”
  - Events can be possible outcomes or observed values
  - Either happens or it does not
- Events are represented by uppercase letters:  $A, B, C, \dots$
- List the event in  $\{ \}$  brackets
- Example:  $A = \{\text{roll an even number on a six-sided die}\} = \{2, 4, 6\}$


# Operations on Events

- Let  $A$  and  $B$  be events, or subsets of  $S$ , where  $A \subset S$  and  $B \subset S$




# Intersection

- Intersection ( $A \cap B$ ): The event "both  $A$  and  $B$ ", or all elements in  $S$  in both  $A$  and  $B$



# Union


- Union ( $A \cup B$ ): The event "either  $A$  or  $B$ ", or all elements in  $S$  in either  $A$  or  $B$





# Complement

- Complement ( $A^c$ ,  $\bar{A}$ , or  $A'$ ): The event "not  $A$ ", or all elements in  $S$  not in  $A$



# Operations Example

# Operations Example

- Suppose we have the following, where  $A \subset S$ ,  $B \subset S$ , and  $C \subset S$ :

$$S = \{1,2,3,4,5,6,7,8\}$$

$$A = \{1,2,3,4\}$$

$$B = \{2,4,6,8\}$$

$$C = \{7,8\}$$

# Operations Example

- Suppose we have the following, where  $A \subset S$ ,  $B \subset S$ , and  $C \subset S$ :

$$S = \{1,2,3,4,5,6,7,8\}$$

$$A = \{1,2,3,4\}$$

$$B = \{2,4,6,8\}$$

$$C = \{7,8\}$$

- Evaluate the following expressions:

$$A \cap B = \{2,4\}$$

$$(A \cup C) \cap B = \{2,4,8\}$$

$$A^c \cap C = \{7,8\}$$

$$(A \cap B^c) \cup C = \{1,3,7,8\}$$

# Operations on Events: De Morgan's Laws

# Operations on Events: De Morgan's Laws


- De Morgan's Laws:

# Operations on Events: De Morgan's Laws

- De Morgan's Laws:
  - $(A \cup B)^c = A^c \cap B^c$


# Operations on Events: De Morgan's Laws

- De Morgan's Laws:
  - $(A \cup B)^c = A^c \cap B^c$
  - $(A \cap B)^c = A^c \cup B^c$






# Events




# Events

- Null events are events that can never occur, represented as  $\emptyset$



# Events

- Null events are events that can never occur, represented as  $\emptyset$
- Disjoint or mutually exclusive events are events that cannot occur simultaneously;  
 $A$  and  $B$  are disjoint if and only if  $A \cap B = \emptyset$



# Cardinality

# Cardinality

- The *cardinality* of  $A$  is the number of elements in the set, denoted  $|A|$

# Cardinality

- The *cardinality* of  $A$  is the number of elements in the set, denoted  $|A|$
- Three types of cardinality:
  - Finite:  $|A| < \infty$
  - Countable:  $|A| = \infty$  but elements can be listed as  $x_1, x_2, \dots$
  - Uncountable:  $|A| = \infty$  and elements cannot be listed as  $x_1, x_2, \dots$

# Probability

# Probability

- **Probability:** If an experiment is repeated  $n$  times under identical conditions, and if event  $A$  occurs  $m$  times, then as  $n$  grows large, the ratio  $m/n$  approaches a fixed limit that is the probability of event  $A$ :  $\Pr(A) = \frac{m}{n}$




# Probability

- **Probability:** If an experiment is repeated  $n$  times under identical conditions, and if event  $A$  occurs  $m$  times, then as  $n$  grows large, the ratio  $m/n$  approaches a fixed limit that is the probability of event  $A$ :  $\Pr(A) = \frac{m}{n}$
- Relative frequency of occurrence of an event when repeated many times

# Probability

- **Probability:** If an experiment is repeated  $n$  times under identical conditions, and if event  $A$  occurs  $m$  times, then as  $n$  grows large, the ratio  $m/n$  approaches a fixed limit that is the probability of event  $A$ :  $\Pr(A) = \frac{m}{n}$
- Relative frequency of occurrence of an event when repeated many times
- $\Pr(A) = \frac{\text{\# of times } A \text{ occurs}}{\text{total \# of trials}}$

# Probability



# Probability Rules

# Probability Rules

- $0 \leq \Pr(A) \leq 1$

# Probability Rules

- $0 \leq \Pr(A) \leq 1$
- $\Pr(S) = 1$

# Probability Rules

- $0 \leq \Pr(A) \leq 1$
- $\Pr(S) = 1$
- $\Pr(\emptyset) = 0$


# Probability Rules

- $0 \leq \Pr(A) \leq 1$
- $\Pr(S) = 1$
- $\Pr(\emptyset) = 0$
- $\Pr(A^c) = 1 - \Pr(A)$



# Probability Rules

- $0 \leq \Pr(A) \leq 1$
- $\Pr(S) = 1$
- $\Pr(\emptyset) = 0$
- $\Pr(A^c) = 1 - \Pr(A)$
- If  $A \subset B$ , then  $\Pr(A) \leq \Pr(B)$




# Mutual Exclusivity and Exhaustiveness

# Mutual Exclusivity and Exhaustiveness

- When the probabilities of mutually exclusive events sum to 1, the events are *exhaustive* (i.e., no other possible outcomes)

# Mutual Exclusivity and Exhaustiveness

- When the probabilities of mutually exclusive events sum to 1, the events are *exhaustive* (i.e., no other possible outcomes)




# Addition Rule: Mutually Exclusive Events

# Addition Rule: Mutually Exclusive Events

- If  $A$  and  $B$  are mutually exclusive, we have  $\Pr(A \cup B) = \Pr(A) + \Pr(B)$

# Addition Rule: Mutually Exclusive Events

- If  $A$  and  $B$  are mutually exclusive, we have  $\Pr(A \cup B) = \Pr(A) + \Pr(B)$



# Addition Rule: General




# Addition Rule: General

- In general, we have  $\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$

# Addition Rule: General

- In general, we have  $\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$



# Probability Example

# Probability Example

- Suppose that 55% of cancer patients are female, 20% of cancer patients have previously undergone chemotherapy, and 15% of cancer patients are both female and have undergone chemotherapy

# Probability Example

↙ A

↙ B

- Suppose that 55% of cancer patients are female, 20% of cancer patients have previously undergone chemotherapy, and 15% of cancer patients are both female and have undergone chemotherapy
- What is the probability that a patient is female or has undergone chemotherapy?

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$55 + 20 - 15 = \underline{60\%}$$

# Conditional Probability

# Conditional Probability

- Often, we are interested in determining the probability that an event will occur given that we already know the outcome of another event
  - Example: What is the probability that it rains tomorrow given that it rained today?

# Conditional Probability

- Often, we are interested in determining the probability that an event will occur given that we already know the outcome of another event
  - Example: What is the probability that it rains tomorrow given that it rained today?
- **Conditional Probability:** The probability that event  $A$  will occur given that we already know the outcome of event  $B$



# Conditional Probability

- Often, we are interested in determining the probability that an event will occur given that we already know the outcome of another event
  - Example: What is the probability that it rains tomorrow given that it rained today?
- **Conditional Probability:** The probability that event  $A$  will occur given that we already know the outcome of event  $B$
- $\Pr(A | B) =$  probability of  $A$  given  $B$

# Multiplicative Rule

# Multiplicative Rule

- The *multiplicative rule of probability* tells us the following:

$$\Pr(A \cap B) = \Pr(A) \cdot \Pr(B | A)$$

$$\Pr(A \cap B) = \Pr(B) \cdot \Pr(A | B)$$

# Multiplicative Rule

- The *multiplicative rule of probability* tells us the following:


$$\Pr(A \cap B) = \Pr(A) \cdot \Pr(B | A)$$

$$\Pr(A \cap B) = \Pr(B) \cdot \Pr(A | B)$$

- Rearranging yields *conditional probability expressions*:

$$\rightarrow \underline{\Pr(B | A)} = \frac{\Pr(A \cap B)}{\Pr(A)}$$

$$\Pr(A | B) = \frac{\Pr(A \cap B)}{\Pr(B)}$$



# Conditional Probability Example

# Conditional Probability Example


- Setup:
  - Suppose 10,000 students enter college
  - 450 students changed majors
  - 300 students who changed majors were males
  - 3000 students were males

# Conditional Probability Example

- Setup:
  - Suppose 10,000 students enter college
  - 450 students changed majors
  - 300 students who changed majors were males
  - 3000 students were males
- Q1: What is the probability of changing majors given that you are a male?

# Conditional Probability Example

- Setup:
  - Suppose 10,000 students enter college
  - 450 students changed majors
  - 300 students who changed majors were males
  - 3000 students were males



- Q1: What is the probability of  $\overbrace{\text{changing majors}}^A$  given that you are a  $\overbrace{\text{male}}^B$ ?

$$P(A|B) = \frac{Pr(A \cap B)}{Pr(B)} = \frac{300 / 10K}{3000 / 10K} = 1/10$$

- Q2: What is the probability of changing majors given that you are not a male?

$$Pr(A|B^c) = \frac{Pr(A \cap B^c)}{Pr(B^c)} = \frac{150}{7000} \approx 2\%$$



# Multiplicative Rule Example

# Multiplicative Rule Example

- Setup:
  - The probability that you will be sick tomorrow is 0.6
  - If you are sick tomorrow, the probability that you will be sick the next day is 0.7
  - If you are not sick tomorrow, the probability that you will be sick the next day is 0.2

# Multiplicative Rule Example

- Setup:
  - The probability that you will be sick tomorrow is 0.6
  - If you are sick tomorrow, the probability that you will be sick the next day is 0.7
  - If you are not sick tomorrow, the probability that you will be sick the next day is 0.2
- Q1: What is the probability that you are sick tomorrow and the next day?

# Multiplicative Rule Example

- Setup:


- The probability that you will be sick tomorrow is 0.6
- If you are sick tomorrow, the probability that you will be sick the next day is 0.7
- If you are not sick tomorrow, the probability that you will be sick the next day is 0.2
- Q1: What is the probability that you are sick tomorrow and the next day?

$$P(A \cap B) = P(B|A) \cdot P(A) = 0.7 \times 0.6 = \underline{0.42}$$

- Q2: What is the probability that you are not sick tomorrow but sick the following day?


$$Pr(A^c \cap B) = Pr(B|A^c) \cdot P(A^c) = (1 - 0.6)(0.2) = \underline{0.08}$$

# Conditional Probability




# Conditional Probability

- Note,  $\Pr(B | A) \neq 1 - \Pr(A | B)$



# Conditional Probability


- Note,  $\Pr(B | A) \neq 1 - \Pr(A | B)$
- Similarly,  $\Pr(B | A) \neq 1 - \Pr(B | A^c)$



# Conditional Probability

- Note,  $\Pr(B | A) \neq 1 - \Pr(A | B)$
- Similarly,  $\Pr(B | A) \neq 1 - \Pr(B | A^c)$
- But,  $\Pr(B | A) = 1 - \Pr(B^c | A)$

$$\Pr(B) = 1 - \Pr(B^c)$$





# Conditional Probability Example

# Conditional Probability Example

- Setup:
  - Consider a random experiment where 3 balls are randomly selected (without replacement) from 5 balls labeled 1, 2, 3, 4, 5. Sample space:

123, 124, 125, 134, 135, 145

234, 235, 245

345

# Conditional Probability Example

- Setup:
  - Consider a random experiment where 3 balls are randomly selected (without replacement) from 5 balls labeled 1, 2, 3, 4, 5. Sample space:

123, 124, 125, 134, 135, 145  
234, 235, 245  
345

- Let  $A = \{1 \text{ is selected}\}$  and  $B = \{5 \text{ is selected}\}$ . What is  $\Pr(A | B)$ ?

$$\Pr(A | B) = \frac{\Pr(A \cap B)}{\Pr(B)} = \frac{3}{6} = \frac{1}{2}$$

# Independence

# Independence

- **Independence:** The outcome of one event has no effect on the outcome of another event
  - If  $A$  and  $B$  are independent, then  $\Pr(A | B) = \Pr(A)$  (and  $\Pr(B | A) = \Pr(B)$ )

# Independence

- **Independence:** The outcome of one event has no effect on the outcome of another event
  - If  $A$  and  $B$  are independent, then  $\Pr(A | B) = \Pr(A)$  (and  $\Pr(B | A) = \Pr(B)$ )
  - This is because intersection is decomposable:  $\Pr(B|A) \cdot \Pr(A) = \Pr(A) \cdot \Pr(B)$
  - If  $A$  and  $B$  are independent, then  $\Pr(A \cap B) = \Pr(A) \cdot \Pr(B)$
  - From this, we see that  $\Pr(A | B) = \frac{\Pr(A \cap B)}{\Pr(B)} = \frac{\Pr(A) \cdot \Pr(B)}{\Pr(B)} = \Pr(A)$

# Independence Example

# Independence Example

- Setup:
  - Suppose we flip a coin twice; tosses are independent
  - Let  $A = \{\text{first flip is heads}\}$  and  $B = \{\text{second flip is heads}\}$
  - $\Pr(A) = \Pr(B) = 1/2$



# Independence Example

- Setup:
  - Suppose we flip a coin twice; tosses are independent
  - Let  $A = \{\text{first flip is heads}\}$  and  $B = \{\text{second flip is heads}\}$
  - $\Pr(A) = \Pr(B) = 1/2$
- What is  $\Pr(A \cap B)$  (probability that both flips are heads)?

$1/4$

$1/2^2$

# Mutual Independence

# Mutual Independence

- Suppose we have  $n$  events,  $N$ . These  $n$  events are **mutually independent** iff, for every subset of events  $M \subseteq N$ , we have

$$\Pr \left( \bigcap_{i \in M} A_i \right) = \prod_{i \in M} \Pr(A_i)$$

# Mutual Independence

- Suppose we have  $n$  events,  $N$ . These  $n$  events are **mutually independent** iff, for every subset of events  $M \subseteq N$ , we have

$$\Pr\left(\bigcap_{i \in M} A_i\right) = \prod_{i \in M} \Pr(A_i)$$

- Consider the case of  $n = 3$ . Events  $A_1, A_2, A_3$  are independent iff the following hold:

$$\Pr(A_1 \cap A_2) = \Pr(A_1) \cdot \Pr(A_2)$$

$$\Pr(A_1 \cap A_3) = \Pr(A_1) \cdot \Pr(A_3)$$

$$\Pr(A_2 \cap A_3) = \Pr(A_2) \cdot \Pr(A_3)$$

$$\Pr(A_1 \cap A_2 \cap A_3) = \Pr(A_1) \cdot \Pr(A_2) \cdot \Pr(A_3)$$

# Mutual Independence

- Suppose we have  $n$  events,  $N$ . These  $n$  events are **mutually independent** iff, for every subset of events  $M \subseteq N$ , we have

$$\Pr\left(\bigcap_{i \in M} A_i\right) = \prod_{i \in M} \Pr(A_i)$$

- Consider the case of  $n = 3$ . Events  $A_1, A_2, A_3$  are independent iff the following hold:

$$\Pr(A_1 \cap A_2) = \Pr(A_1) \cdot \Pr(A_2)$$

$$\Pr(A_1 \cap A_3) = \Pr(A_1) \cdot \Pr(A_3)$$

$$\Pr(A_2 \cap A_3) = \Pr(A_2) \cdot \Pr(A_3)$$

$$\Pr(A_1 \cap A_2 \cap A_3) = \Pr(A_1) \cdot \Pr(A_2) \cdot \Pr(A_3)$$

- If all but the last equality hold,  $A_1, A_2, A_3$  are *pairwise independent*, but not mutually independent

# Pairwise Independence: Example

# Pairwise Independence: Example

- Setup: Consider rolling a fair ~~six~~<sup>4</sup>-sided die. Consider the events  $A = \{\underline{1}, 2\}$ ,  $B = \{\underline{1}, 3\}$ , and  $C = \{2, 3\}$
- $\Pr(A) = \Pr(B) = \Pr(C) = \frac{1}{2}$
- $\Pr(\underline{A \cap B}) = \frac{1}{4} = \Pr(A) \cdot \Pr(B)$
- $\Pr(A \cap C) = \frac{1}{4}$       -1
- $\Pr(B \cap C) = \frac{1}{4}$       //
- $\Pr(A \cap B \cap C) = 0 \neq (\frac{1}{2})^3$

# Pairwise Independence: Example

- Setup: Consider rolling a fair six-sided die. Consider the events  $A = \{1,2\}$ ,  $B = \{1,3\}$ , and  $C = \{2,3\}$ 
  - $\Pr(A) = \Pr(B) = \Pr(C) =$
  - $\Pr(A \cap B) =$
  - $\Pr(A \cap C) =$
  - $\Pr(B \cap C) =$
  - $\Pr(A \cap B \cap C) =$
- These events are pairwise independent but not mutually independent



# Independence vs. Mutual Exclusivity

# Independence vs. Mutual Exclusivity

- Independence and mutual exclusivity are not the same thing

# Independence vs. Mutual Exclusivity

- Independence and mutual exclusivity are not the same thing
- If  $A$  and  $B$  are mutually exclusive, then  $\Pr(A | B) = 0$  and  $\Pr(B | A) = 0$

# Independence vs. Mutual Exclusivity

- Independence and mutual exclusivity are not the same thing
- If  $A$  and  $B$  are mutually exclusive, then  $\Pr(A | B) = 0$  and  $\Pr(B | A) = 0$
- This is not the same thing as independence, where  $\Pr(A | B) = \Pr(A)$  and  $\Pr(B | A) = \Pr(B)$

# Independence vs. Mutual Exclusivity

$A$        $A^c$



- Independence and mutual exclusivity are not the same thing
- If  $A$  and  $B$  are mutually exclusive, then  $\Pr(A | B) = 0$  and  $\Pr(B | A) = 0$
- This is not the same thing as independence, where  $\Pr(A | B) = \Pr(A)$  and  $\Pr(B | A) = \Pr(B)$
- Independence: the other event still may occur; its probability is unaffected

# Law of Total Probability

# Law of Total Probability


- Consider a collection of mutually exclusive and exhaustive events  $A_1, A_2, \dots, A_n$  that *partitions* the sample space  $S$

# Law of Total Probability





- Consider a collection of mutually exclusive and exhaustive events  $A_1, A_2, \dots, A_n$  that *partitions* the sample space  $S$

- Then, for any event  $E$ , the law of total probability states the following:



$$\Pr(E) = \Pr(E \cap A_1) + \Pr(E \cap A_2) + \dots + \Pr(E \cap A_n)$$

$$= \Pr(E | A_1) \cdot \Pr(A_1) + \Pr(E | A_2) \cdot \Pr(A_2) + \dots + \Pr(E | A_n) \cdot \Pr(A_n)$$



$$\Pr(A|B) = \Pr(A)$$

$$\Pr(B|A) = \Pr(B)$$




# Law of Total Probability

$$P(A \cap B) = P(A|B) P(B)$$

- Consider a collection of mutually exclusive and exhaustive events  $A_1, A_2, \dots, A_n$  that *partitions* the sample space  $S$
- Then, for any event  $E$ , the law of total probability states the following:

$$\begin{aligned} \Pr(E) &= \Pr(E \cap A_1) + \Pr(E \cap A_2) + \dots + \Pr(E \cap A_n) \\ &= \Pr(E|A_1) \cdot \Pr(A_1) + \Pr(E|A_2) \cdot \Pr(A_2) + \dots + \Pr(E|A_n) \cdot \Pr(A_n) \end{aligned}$$



# Bayes' Theorem

# Bayes' Theorem

- Let's say you have an idea of  $\Pr(B | A)$  but want to know about  $\Pr(A | B)$

# Bayes' Theorem

- Let's say you have an idea of  $\Pr(B | A)$  but want to know about  $\Pr(A | B)$
- Recall that  $\Pr(A | B) \cdot \Pr(B) = \Pr(B | A) \cdot \Pr(A) = \Pr(A \cap B)$


# Bayes' Theorem

- Let's say you have an idea of  $\Pr(B | A)$  but want to know about  $\Pr(A | B)$
- Recall that  $\Pr(A | B) \cdot \Pr(B) = \Pr(B | A) \cdot \Pr(A) = \Pr(A \cap B)$
- Rearranging yields Bayes' Theorem:

$$\Pr(A | B) = \frac{\Pr(B | A) \cdot \Pr(A)}{\Pr(B)} = \frac{\Pr(B | A) \cdot \Pr(A)}{\Pr(B | A) \cdot \Pr(A) + \Pr(B | A^c) \cdot \Pr(A^c)}$$

# Bayes' Theorem

3 Blue 1 Brown



- Let's say you have an idea of  $\Pr(B | A)$  but want to know about  $\Pr(A | B)$

- Recall that  $\Pr(A | B) \cdot \Pr(B) = \Pr(B | A) \cdot \Pr(A) = \Pr(A \cap B)$


- Rearranging yields Bayes' Theorem:

$$\Pr(A | B) = \frac{\Pr(B | A) \cdot \Pr(A)}{\Pr(B)} = \frac{\Pr(B | A) \cdot \Pr(A)}{\Pr(B | A) \cdot \Pr(A) + \Pr(B | A^c) \cdot \Pr(A^c)}$$

Posterior

Likelihood

Prior



# Bayes' Theorem: Example

# Bayes' Theorem: Example

- Setup:
  - Given that you have diabetes, there is a 70% chance you are also overweight
  - Given that you do not have diabetes, there is a 35% chance you are overweight
  - 10% of people have diabetes



# Bayes' Theorem: Example

- Setup:


- Given that you have diabetes, there is a 70% chance you are also overweight
- Given that you do not have diabetes, there is a 35% chance you are overweight
- 10% of people have diabetes

- Q: Given that a randomly selected person is overweight, what is the probability that he has diabetes?

$$P(D|W) = \frac{P(W|D) P(D)}{P(W)}$$

$$= \frac{.7 \times .1}{P(W|D) P(D) + P(W|D^c) P(D^c)}$$

$.70 \times .10$ 
 $.35$ 
 $.40$



Diagnostic Tests

(Bayes' rule)

# Diagnostic Tests

- Apply Bayes' theorem to diagnostic testing and screening

# Diagnostic Tests

- Apply Bayes' theorem to diagnostic testing and screening
- Assume there are two mutually exclusive and exhaustive states of health:
  - $D_1$ : the event that a subject has the disease
  - $D_2$ : the event that a subject does not have the disease

# Diagnostic Tests

- Apply Bayes' theorem to diagnostic testing and screening
- Assume there are two mutually exclusive and exhaustive states of health:
  - $D_1$ : the event that a subject has the disease
  - $D_2$ : the event that a subject does not have the disease
- Assume that we run a screening test on a patient to determine if they have the disease, with two mutually exclusive and exhaustive outcomes:
  - $T^+$ : the test is positive
  - $T^-$ : the test is negative

# Diagnostic Tests

- Apply Bayes' theorem to diagnostic testing and screening
- Assume there are two mutually exclusive and exhaustive states of health:
  - $D_1$ : the event that a subject has the disease
  - $D_2$ : the event that a subject does not have the disease
- Assume that we run a screening test on a patient to determine if they have the disease, with two mutually exclusive and exhaustive outcomes:
  - $T^+$ : the test is positive
  - $T^-$ : the test is negative
- Typically, we are interested in  $\Pr(D_1 | T^+)$  (true positive rate of a test)

# Diagnostic Tests

# Diagnostic Tests

- **Sensitivity:** Probability of a positive test result given that the individual tested actually has the disease (true positive):
  - $\Pr(T^+ | D_1)$



# Diagnostic Tests

- **Sensitivity:** Probability of a positive test result given that the individual tested actually has the disease (true positive):
  - $\Pr(T^+ | D_1)$
- **False negative probability:** Probability of a negative test result given that the individual tested actually has the disease (false negative):
  - $\Pr(T^- | D_1) = 1 - \text{Sensitivity}$

# Diagnostic Tests

- **Sensitivity:** Probability of a positive test result given that the individual tested actually has the disease (true positive):
  - $\Pr(T^+ | D_1)$
- **False negative probability:** Probability of a negative test result given that the individual tested actually has the disease (false negative):
  - $\Pr(T^- | D_1) = 1 - \text{Sensitivity}$
- **Specificity:** Probability of a negative test result given that the individual tested does not have the disease (true negative):
  - $\Pr(T^- | D_2)$

# Diagnostic Tests

- **Sensitivity:** Probability of a positive test result given that the individual tested actually has the disease (true positive):
  - $\Pr(T^+ | D_1)$
- **False negative probability:** Probability of a negative test result given that the individual tested actually has the disease (false negative):
  - $\Pr(T^- | D_1) = 1 - \text{Sensitivity}$
- **Specificity:** Probability of a negative test result given that the individual tested does not have the disease (true negative):
  - $\Pr(T^- | D_2)$
- **False positive probability:** Probability of a positive test result given that the individual tested does not have the disease (false positive):
  - $\Pr(T^+ | D_2) = 1 - \text{Specificity}$

# Positive Predictive Value (PPV)

# Positive Predictive Value (PPV)

- **Positive predictive value (PPV):** The probability that a person with a positive test result actually has the disease
  - $\Pr(D_1 | T^+)$

# Positive Predictive Value (PPV)

- **Positive predictive value (PPV):** The probability that a person with a positive test result actually has the disease
  - $\Pr(D_1 | T^+)$
- Using Bayes' Rule, sensitivity, and specificity:

$$\begin{aligned}\Pr(D_1 | T^+) &= \frac{\Pr(D_1 \cap T^+)}{\Pr(T^+)} \\ &= \frac{\Pr(T^+ | D_1) \cdot \Pr(D_1)}{\Pr(T^+ | D_1) \cdot \Pr(D_1) + \Pr(T^+ | D_2) \cdot \Pr(D_2)}\end{aligned}$$

# Positive Predictive Value (PPV)

- **Positive predictive value (PPV):** The probability that a person with a positive test result actually has the disease

- $\Pr(D_1 | T^+)$

- Using Bayes' Rule, sensitivity, and specificity:

$$\begin{aligned}\Pr(D_1 | T^+) &= \frac{\Pr(D_1 \cap T^+)}{\Pr(T^+)} \\ &= \frac{\Pr(T^+ | D_1) \cdot \Pr(D_1)}{\Pr(T^+ | D_1) \cdot \Pr(D_1) + \Pr(T^+ | D_2) \cdot \Pr(D_2)}\end{aligned}$$

- What are  $\Pr(D_1)$  and  $\Pr(D_2)$ ?
  - $\Pr(D_1)$ : probability of having the disease, or prevalence of the disease
  - $\Pr(D_2) = 1 - \Pr(D_1)$

# Negative Predictive Value (NPV)



# Negative Predictive Value (NPV)

- **Negative predictive value (NPV):** The probability that a person with a negative test result actually does not have the disease
  - $\Pr(D_2 | T^-)$

# Negative Predictive Value (NPV)

- **Negative predictive value (NPV):** The probability that a person with a negative test result actually does not have the disease
- $\Pr(D_2 | T^-)$
- Using Bayes' Rule, sensitivity, and specificity:

$$\begin{aligned}\Pr(D_2 | T^-) &= \frac{\Pr(D_2 \cap T^-)}{\Pr(T^-)} \\ &= \frac{\Pr(T^- | D_2) \cdot \Pr(D_2)}{\Pr(T^- | D_2) \cdot \Pr(D_2) + \Pr(T^- | D_1) \cdot \Pr(D_1)}\end{aligned}$$

# Diagnostic Tests: Example

# Diagnostic Tests: Example

- Cancer test has the following properties:
  - The test gives a positive result 95% of the time when the patient has cancer
  - The test gives a negative result 90% of the time when the patient does not have cancer
  - About 12% of patients have cancer

# Diagnostic Tests: Example

- Cancer test has the following properties:
  - The test gives a positive result 95% of the time when the patient has cancer
  - The test gives a negative result 90% of the time when the patient does not have cancer
  - About 12% of patients have cancer
- Q: A patient tested positive for cancer. What is the probability that they have cancer?

# Diagnostic Tests: Example

# Diagnostic Tests: Example

- Cancer test has the following properties:
  - The test gives a positive result 95% of the time when the patient has cancer
  - The test gives a negative result 90% of the time when the patient does not have cancer
  - About 12% of patients have cancer

# Diagnostic Tests: Example

- Cancer test has the following properties:
  - The test gives a positive result 95% of the time when the patient has cancer
  - The test gives a negative result 90% of the time when the patient does not have cancer
  - About 12% of patients have cancer
- Q: A patient tested positive for cancer. What is the probability that they have cancer?



# Diagnostic Tests: Example

- Cancer test has the following properties:
  - The test gives a positive result 95% of the time when the patient has cancer
  - The test gives a negative result 90% of the time when the patient does not have cancer
  - About 12% of patients have cancer
- Q: A patient tested positive for cancer. What is the probability that they have cancer?

$$\begin{aligned}\Pr(C | pos) &= \frac{\Pr(C \cap pos)}{\Pr(pos)} \\&= \frac{\Pr(pos | C) \cdot \Pr(C)}{\Pr(pos | C) \cdot \Pr(C) + \Pr(pos | C^c) \cdot \Pr(C^c)} \\&= \frac{0.95 \cdot 0.12}{0.95 \cdot 0.12 + (1 - 0.90) \cdot (1 - 0.12)} \\&= 0.5644\end{aligned}$$

# Combinatorics

# Counting Outcomes

# Counting Outcomes

- If each outcome in the sample space is equally likely, then computing probabilities is an exercise in counting

# Counting Outcomes

- If each outcome in the sample space is equally likely, then computing probabilities is an exercise in counting
- For a sample space  $S$  and an event  $E \subseteq S$ , the probability of  $E$  (under an equiprobable model) is  $\Pr(E) = \frac{N}{D}$
- Where  $N$  is the total number of outcomes in  $E$  and  $D$  is the total number of outcomes in  $S$

# Counting Outcomes

- If each outcome in the sample space is equally likely, then computing probabilities is an exercise in counting
- For a sample space  $S$  and an event  $E \subseteq S$ , the probability of  $E$  (under an equiprobable model) is  $\Pr(E) = \frac{N}{D}$ 
  - Where  $N$  is the total number of outcomes in  $E$  and  $D$  is the total number of outcomes in  $S$
- We're going to learn how to count the number of outcomes

# Ordered vs. Unordered Selection

# Ordered vs. Unordered Selection

- **Ordered selection** of size  $n$  from sample space  $S$ : select  $n$  distinct objects from  $S$  where order of selection matters
  - Care about the names and order of choices



# Ordered vs. Unordered Selection

- **Ordered selection** of size  $n$  from sample space  $S$ : select  $n$  distinct objects from  $S$  where order of selection matters
  - Care about the names and order of choices
- **Unordered selection** of size  $n$  from sample space  $S$ : select  $n$  distinct objects from  $S$  where order of selection does not matter
  - Care about the names of choices (think of it as a set)

# Rule of Product

# Rule of Product

- Suppose a procedure can be broken down into  $m$  tasks

# Rule of Product

- Suppose a procedure can be broken down into  $m$  tasks
- There are  $n_i$  distinct ways to perform the  $i^{th}$  task, for  $i = 1, \dots, m$

# Rule of Product

- Suppose a procedure can be broken down into  $m$  tasks
- There are  $n_i$  distinct ways to perform the  $i^{th}$  task, for  $i = 1, \dots, m$
- Then, there are  $n_1 \cdot n_2 \cdot \dots \cdot n_m$  distinct ways to perform the entire procedure

# Rule of Product: Example

# Rule of Product: Example

- How many valid three-digit numbers (i.e., between 100 and 999, inclusive) have three different digits and only a single odd number in the middle?

# Rule of Product: Example



# Rule of Product: Example

- How many valid three-digit numbers (i.e., between 100 and 999, inclusive) have three different digits and only a single odd digit in the center?

Break this down into  $m = 3$  tasks

Task 1: Select an odd (center) digit,  $n_1 = 5$

Task 2: Select a first (even) digit that is not 0,  $n_2 = 4$

Task 3: Select a last (even) digit,  $n_3 = 4$

Total:  $n_1 \cdot n_2 \cdot n_3 = 5 \cdot 4 \cdot 4 = 80$

# Tree Method (Rule of Sum)

# Tree Method (Rule of Sum)

- Suppose a procedure can be broken down into  $m$  disjoint and exhaustive cases

# Tree Method (Rule of Sum)

- Suppose a procedure can be broken down into  $m$  disjoint and exhaustive cases
- There are  $n_i$  distinct ways to get the  $i^{th}$  case, for  $i = 1, \dots, m$


# Tree Method (Rule of Sum)

- Suppose a procedure can be broken down into  $m$  disjoint and exhaustive cases
- There are  $n_i$  distinct ways to get the  $i^{th}$  case, for  $i = 1, \dots, m$
- Then, there are  $n_1 + n_2 + \dots + n_m$  distinct ways to perform the entire procedure

# Tree Method (Rule of Sum)

- Suppose a procedure can be broken down into  $m$  disjoint and exhaustive cases
- There are  $n_i$  distinct ways to get the  $i^{th}$  case, for  $i = 1, \dots, m$
- Then, there are  $n_1 + n_2 + \dots + n_m$  distinct ways to perform the entire procedure
- Often, use the rule of sum (tree method) and the rule of product together

# Rule of Sum (OR) and Rule of Product (AND)



# Factorials



# Factorials

- Factorial:  $n!$  is the product of all positive integers less than or equal to  $n$ 
  - $n! = n \cdot (n - 1) \cdot \dots \cdot 1$

# Factorials

- Factorial:  $n!$  is the product of all positive integers less than or equal to  $n$ 
  - $n! = n \cdot (n - 1) \cdot \dots \cdot 1$
- Allows us to calculate the number of ways in which  $n$  objects can be ordered

# Factorials

- Factorial:  $n!$  is the product of all positive integers less than or equal to  $n$ 
  - $n! = n \cdot (n - 1) \cdot \dots \cdot 1$
- Allows us to calculate the number of ways in which  $n$  objects can be ordered
- By convention,  $0! = 1$  (there is one way of ordering zero things)

# Factorials

- Factorial:  $n!$  is the product of all positive integers less than or equal to  $n$ 
  - $n! = n \cdot (n - 1) \cdot \dots \cdot 1$
- Allows us to calculate the number of ways in which  $n$  objects can be ordered
- By convention,  $0! = 1$  (there is one way of ordering zero things)
- In R: use `factorial(x)`

# Permutation

# Permutation

- Suppose we want to select and order  $k$  objects from a total of  $n$  objects
  - Ordered selection

# Permutation

- Suppose we want to select and order  $k$  objects from a total of  $n$  objects
  - Ordered selection
- There are  $n$  ways to select the first object,  $n - 1$  ways to select the second object, and so on until we have  $n - k + 1$  ways to select the final object

$$\begin{aligned} P(n, k) &= n \cdot (n - 1) \cdot \dots \cdot (n - k + 1) \\ &= \frac{n!}{(n - k)!} \end{aligned}$$

# Permutation: Example



# Permutation: Example

- Q1: How many four-letter “words” are there where each letter is distinct?

# Permutation: Example

- Q1: How many four-letter "words" are there where each letter is distinct?
- Q2: How many ways are there of assigning three students to seven orientation groups, where each student must go to a different group?

# Combination

# Combination

- Suppose we want to select  $k$  objects from  $n$  objects (unordered selection)

# Combination

- Suppose we want to select  $k$  objects from  $n$  objects (unordered selection)
- There are  $P(n, k)$  ways to select and order  $k$  out of  $n$  objects

# Combination

- Suppose we want to select  $k$  objects from  $n$  objects (unordered selection)
- There are  $P(n, k)$  ways to select and order  $k$  out of  $n$  objects
- There are  $k!$  ways to order  $k$  distinct objects

# Combination

- Suppose we want to select  $k$  objects from  $n$  objects (unordered selection)
- There are  $P(n, k)$  ways to select and order  $k$  out of  $n$  objects
- There are  $k!$  ways to order  $k$  distinct objects
- Therefore, we have  $C(n, k) = \frac{P(n, k)}{k!} = \frac{n!}{k!(n - k)!} = \binom{n}{k}$

# Combination

- Suppose we want to select  $k$  objects from  $n$  objects (unordered selection)
- There are  $P(n, k)$  ways to select and order  $k$  out of  $n$  objects
- There are  $k!$  ways to order  $k$  distinct objects
- Therefore, we have  $C(n, k) = \frac{P(n, k)}{k!} = \frac{n!}{k!(n - k)!} = \binom{n}{k}$
- Interpretation:  $C(n, k)$  is the number of ways in which  $k$  objects can be selected from a total of  $n$  objects (without replacement) without regard to order



# Combination

- Suppose we want to select  $k$  objects from  $n$  objects (unordered selection)
- There are  $P(n, k)$  ways to select and order  $k$  out of  $n$  objects
- There are  $k!$  ways to order  $k$  distinct objects
- Therefore, we have  $C(n, k) = \frac{P(n, k)}{k!} = \frac{n!}{k!(n - k)!} = \binom{n}{k}$
- Interpretation:  $C(n, k)$  is the number of ways in which  $k$  objects can be selected from a total of  $n$  objects (without replacement) without regard to order
- In R, use `choose(n, k)`

# Combination

- Suppose we want to select  $k$  objects from  $n$  objects (unordered selection)
- There are  $P(n, k)$  ways to select and order  $k$  out of  $n$  objects
- There are  $k!$  ways to order  $k$  distinct objects
- Therefore, we have  $C(n, k) = \frac{P(n, k)}{k!} = \frac{n!}{k!(n - k)!} = \binom{n}{k}$
- Interpretation:  $C(n, k)$  is the number of ways in which  $k$  objects can be selected from a total of  $n$  objects (without replacement) without regard to order
- In R, use `choose(n, k)`
- *Binomial coefficient*

# Combination: Example (Poker Hands)

# Combination: Example (Poker Hands)

- Setting: A poker hand consists of five cards dealt from a standard deck of 52 cards (4 suits of 13 values)

# Combination: Example (Poker Hands)

- Setting: A poker hand consists of five cards dealt from a standard deck of 52 cards (4 suits of 13 values)
- Q1: How many different five-card hands are there?

# Combination: Example (Poker Hands)

- Setting: A poker hand consists of five cards dealt from a standard deck of 52 cards (4 suits of 13 values)
- Q1: How many different five-card hands are there?
- Q2: What is the probability of getting four of the same kind?

# Combination: Example (Urn)

# Combination: Example (Urn)

- Setting: An urn contains 35 yellow balls (numbered 1-35) and 35 pink balls (numbered 1-35). Four balls are chosen at random



# Combination: Example (Urn)

- Setting: An urn contains 35 yellow balls (numbered 1-35) and 35 pink balls (numbered 1-35). Four balls are chosen at random
- Q1: What is the probability that there are two pairs of balls which have the same number?

# Combination: Example (Urn)

- Setting: An urn contains 35 yellow balls (numbered 1-35) and 35 pink balls (numbered 1-35). Four balls are chosen at random
- Q1: What is the probability that there are two pairs of balls which have the same number?
- Q2: What is the probability that there is exactly one pair of balls with matching numbers?

# Combination: Example (Urn)

# Combination: Example (Urn)

- Setting: An urn contains 35 yellow balls (numbered 1-35) and 35 pink balls (numbered 1-35). Four balls are chosen at random

# Combination: Example (Urn)

- Setting: An urn contains 35 yellow balls (numbered 1-35) and 35 pink balls (numbered 1-35). Four balls are chosen at random
- Q3: What is the probability that the balls are all the same color and consecutively numbered?

# Stars and Bars: Intuition

# Stars and Bars: Intuition

- How many ways are there of choosing three *positive* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?

# Stars and Bars: Intuition


- How many ways are there of choosing three *positive* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?






# Stars and Bars: Intuition

- How many ways are there of choosing three *positive* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?




# Stars and Bars: Intuition

- How many ways are there of choosing three *positive* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?

- $\binom{6-1}{3-1} = \binom{5}{2}$ : 

# Stars and Bars: Intuition


- How many ways are there of choosing three *positive* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?

- $\binom{6-1}{3-1} = \binom{5}{2}$ : 

- How many ways are there of choosing three *nonnegative* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?

# Stars and Bars: Intuition

- How many ways are there of choosing three *positive* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?


- $\binom{6-1}{3-1} = \binom{5}{2}$ : A stars and bars diagram representing the equation x1 + x2 + x3 = 6 with positive integers. It consists of five stars arranged in a horizontal line. There are two vertical bars placed between the stars: one between the first and second star, and another between the third and fourth star. This configuration divides the five stars into three groups: one star in the first group, two stars in the second group, and two stars in the third group, which corresponds to the solution (1, 2, 2).

- How many ways are there of choosing three *nonnegative* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?




# Stars and Bars: Intuition

- How many ways are there of choosing three *positive* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?


- $\binom{6-1}{3-1} = \binom{5}{2}$ : A stars and bars diagram representing the equation  $x_1 + x_2 + x_3 = 6$  with positive integers. It consists of five stars and two vertical bars. The stars are arranged in three groups: one star before the first bar, two stars between the first and second bars, and two stars after the second bar. This represents the solution  $x_1=1, x_2=2, x_3=2$ .

- How many ways are there of choosing three *nonnegative* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?




# Stars and Bars: Intuition

- How many ways are there of choosing three *positive* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?

- $\left( \begin{matrix} 6-1 \\ 3-1 \end{matrix} \right) = \left( \begin{matrix} 5 \\ 2 \end{matrix} \right)$ :
 
 A diagram showing 6 stars arranged in a horizontal line. There are two vertical bars placed between the stars, dividing the 6 stars into three groups of 2 stars each. This represents the solution (2, 2, 2) for the equation x1 + x2 + x3 = 6.

- How many ways are there of choosing three *nonnegative* numbers,  $x_1, x_2, x_3$ , such that  $x_1 + x_2 + x_3 = 6$ ?

- $\left( \begin{matrix} 6+3-1 \\ 3-1 \end{matrix} \right) = \left( \begin{matrix} 8 \\ 2 \end{matrix} \right)$ :
 
 A diagram showing 8 stars arranged in a horizontal line. There are two vertical bars placed between the stars, dividing the 8 stars into three groups of 2 stars each. This represents the solution (2, 2, 2) for the equation x1 + x2 + x3 = 6, where each variable is nonnegative.

# Stars and Bars: More Formally

# Stars and Bars: More Formally

- Suppose there are  $n$  objects and  $k$  bins. Bins are distinguishable, but objects are not. The only thing we care about is the number of objects in each bin.



# Stars and Bars: More Formally

- Suppose there are  $n$  objects and  $k$  bins. Bins are distinguishable, but objects are not. The only thing we care about is the number of objects in each bin.
- If each bin has to have at least one object in it:
  - Total number of ways =  $\binom{n-1}{k-1}$  (think of filling in gaps between objects)

# Stars and Bars: More Formally

- Suppose there are  $n$  objects and  $k$  bins. Bins are distinguishable, but objects are not. The only thing we care about is the number of objects in each bin.
- If each bin has to have at least one object in it:
  - Total number of ways =  $\binom{n-1}{k-1}$  (think of filling in gaps between objects)
- For nonnegative (not positive) constraints:
  - Total number of ways =  $\binom{n+k-1}{k-1}$  (think of arranging  $n$  objects and  $k-1$  dividers)

# Stars and Bars: Example

# Stars and Bars: Example

- Setup: Six children are choosing ice cream flavors from {vanilla, strawberry, chocolate, caramel}. Each child picks exactly one flavor. Requests are placed in the form: {# vanilla, # strawberry, # chocolate, # caramel}.

# Stars and Bars: Example

- Setup: Six children are choosing ice cream flavors from {vanilla, strawberry, chocolate, caramel}. Each child picks exactly one flavor. Requests are placed in the form: {# vanilla, # strawberry, # chocolate, # caramel}.
- Q1: How many different requests are possible if at least one child must choose each flavor?

# Stars and Bars: Example

- Setup: Six children are choosing ice cream flavors from {vanilla, strawberry, chocolate, caramel}. Each child picks exactly one flavor. Requests are placed in the form: {# vanilla, # strawberry, # chocolate, # caramel}.
- Q1: How many different requests are possible if at least one child must choose each flavor?
- Q2: How many different requests are possible without this restriction?