

# Chapter 5: Distributions

DSCC 462  
Computational Introduction to Statistics

Anson Kahng  
Fall 2022

# Random Variables

# Random Variables

- **Random variable:** A variable that can take a number of different values and whose outcome is determined by chance

# Random Variables

- **Random variable:** A variable that can take a number of different values and whose outcome is determined by chance
- **Discrete random variable:** A random variable whose possible outcomes are a list of discrete values (finite or countably infinite sample space)

# Random Variables

- **Random variable:** A variable that can take a number of different values and whose outcome is determined by chance
- **Discrete random variable:** A random variable whose possible outcomes are a list of discrete values (finite or countably infinite sample space)
  - Example: Coin flip (heads/tails)

# Random Variables

- **Random variable:** A variable that can take a number of different values and whose outcome is determined by chance
- **Discrete random variable:** A random variable whose possible outcomes are a list of discrete values (finite or countably infinite sample space)
  - Example: Coin flip (heads/tails)
- **Continuous random variable:** A random variable whose possible outcomes are any value in an interval (uncountable sample space)

# Random Variables

- **Random variable:** A variable that can take a number of different values and whose outcome is determined by chance
- **Discrete random variable:** A random variable whose possible outcomes are a list of discrete values (finite or countably infinite sample space)
  - Example: Coin flip (heads/tails)
- **Continuous random variable:** A random variable whose possible outcomes are any value in an interval (uncountable sample space)
  - Examples: Time required to run a mile

# Notation

# Notation

- Random variable: Uppercase letters (e.g.,  $X$ ,  $Y$ )

# Notation

- Random variable: Uppercase letters (e.g.,  $X, Y$ )
- Outcome of a random variable: Lowercase letters (e.g.,  $x, y$ )

# Notation

- Random variable: Uppercase letters (e.g.,  $X, Y$ )
- Outcome of a random variable: Lowercase letters (e.g.,  $x, y$ )
- Example: Let  $X =$  the number of surgeries a person has had

# Notation

- Random variable: Uppercase letters (e.g.,  $X, Y$ )
- Outcome of a random variable: Lowercase letters (e.g.,  $x, y$ )
- Example: Let  $X$  = the number of surgeries a person has had
  - $\Pr(X = 1)$ : Probability of having 1 surgery

# Notation

- Random variable: Uppercase letters (e.g.,  $X, Y$ )
- Outcome of a random variable: Lowercase letters (e.g.,  $x, y$ )
- Example: Let  $X$  = the number of surgeries a person has had
  - $\Pr(X = 1)$ : Probability of having 1 surgery
  - $\Pr(X = x)$ : Probability of having  $x$  surgeries

# Probability Distribution

# Probability Distribution

- **Probability distribution:** List of all possible values that a random variable can take, along with their corresponding probabilities

# Probability Distribution

- **Probability distribution:** List of all possible values that a random variable can take, along with their corresponding probabilities
  - Discrete: Probability mass function (PMF)

# Probability Distribution

- **Probability distribution:** List of all possible values that a random variable can take, along with their corresponding probabilities
  - Discrete: Probability mass function (PMF)
  - Continuous: Probability density function (PDF)

# Probability Distribution

- **Probability distribution:** List of all possible values that a random variable can take, along with their corresponding probabilities
  - Discrete: Probability mass function (PMF)
  - Continuous: Probability density function (PDF)
- Let  $X$  be a random variable defined over sample space  $S_X$

# Probability Distribution

- **Probability distribution:** List of all possible values that a random variable can take, along with their corresponding probabilities
  - Discrete: Probability mass function (PMF)
  - Continuous: Probability density function (PDF)
- Let  $X$  be a random variable defined over sample space  $S_X$
- For any  $E \subseteq S_X$ , we can define  $p_X(E) = \Pr(X \in E)$

$$p_X(E)$$

# Discrete Probability Distribution

# Discrete Probability Distribution

- For a discrete random variable  $X$  with sample space  $S_X$ , a probability mass function (PMF)  $p_X$  maps  $x \in S_X$  to a number in  $[0,1]$  such that:

# Discrete Probability Distribution

- For a discrete random variable  $X$  with sample space  $S_X$ , a probability mass function (PMF)  $p_X$  maps  $x \in S_X$  to a number in  $[0,1]$  such that:

$$0 \leq p_X(s) = \Pr(X = x) \leq 1$$

# Discrete Probability Distribution

- For a discrete random variable  $X$  with sample space  $S_X$ , a probability mass function (PMF)  $p_X$  maps  $x \in S_X$  to a number in  $[0,1]$  such that:

$$0 \leq p_X(s) = \Pr(X = x) \leq 1$$

$$\sum_{x \in S_X} p_X(x) = \sum_{x \in S_X} \Pr(X = x) = 1$$

# Discrete Probability Distribution

- For a discrete random variable  $X$  with sample space  $S_X$ , a probability mass function (PMF)  $p_X$  maps  $x \in S_X$  to a number in  $[0,1]$  such that:

$$0 \leq p_X(s) = \Pr(X = x) \leq 1$$

$$\sum_{x \in S_X} p_X(x) = \sum_{x \in S_X} \Pr(X = x) = 1$$

- The support  $S_X$  consists of all  $x$  for which  $p_X(x) > 0$  (all achievable outcomes)

# Discrete Probability Distribution: Example

# Discrete Probability Distribution: Example

- Setup: A fair coin is flipped 3 times. Let  $X$  be a random variable that counts the number of heads observed

# Discrete Probability Distribution: Example

- Setup: A fair coin is flipped 3 times. Let  $X$  be a random variable that counts the number of heads observed
- Fill in the following table:

# Discrete Probability Distribution: Example

- Setup: A fair coin is flipped 3 times. Let  $X$  be a random variable that counts the number of heads observed
- Fill in the following table:

$$3 \times \frac{1}{8} \left[ \begin{array}{l} \text{HTT} \\ \text{THT} \\ \text{TTT} \end{array} \right]$$

PMF

x	Pr(X = x)
0	1/8      12.5%
1	3/8      37.5%
2	3/8
3	1/8      12.5%

# Discrete Probability Distribution: Example

- Setup: A fair coin is flipped 3 times. Let  $X$  be a random variable that counts the number of heads observed
- Fill in the following table:
- Probability distribution tables resemble relative frequency distribution tables: probability of each outcome is the relative frequency distribution of each outcome in a large number of trials

$x$	$\Pr(X = x)$
0	
1	
2	
3	

# Continuous Probability Distribution

# Continuous Probability Distribution

- Specify continuous probability distributions through a *density function*,  $f(x)$

# Continuous Probability Distribution

- Specify continuous probability distributions through a *density function*,  $f(x)$
- Properties:

# Continuous Probability Distribution

- Specify continuous probability distributions through a *density function*,  $f(x)$
- Properties:

$$f(x) \geq 0, \text{ for all } x \in S_X \text{ (nonnegative density)}$$

# Continuous Probability Distribution

- Specify continuous probability distributions through a *density function*,  $f(x)$
- Properties:

$f(x) \geq 0$ , for all  $x \in S_X$  (nonnegative density)

$$\int_S f(x) dx = 1 \text{ (total probability is 1)}$$

# Continuous Probability Distribution

- Specify continuous probability distributions through a *density function*,  $f(x)$
- Properties:

$f(x) \geq 0$ , for all  $x \in S_X$  (nonnegative density)

$$\int f(x) dx = 1 \text{ (total probability is 1)}$$

- $X$  is continuous iff there is a density function  $f_X$  such that the following holds:

# Continuous Probability Distribution



- Specify continuous probability distributions through a *density function*,  $f(x)$
- Properties:

$f(x) \geq 0$ , for all  $x \in S_X$  (nonnegative density)

$$\int f(x) dx = 1 \text{ (total probability is 1)}$$

- $X$  is continuous iff there is a density function  $f_X$  such that the following holds:

$$\begin{aligned}\Pr(a \leq X \leq b) &= \int_a^b f_X(x) dx \\ &= \text{Area under } f \text{ between } a \text{ and } b\end{aligned}$$



# Continuous Probability Distribution

- Specify continuous probability distributions through a *density function*,  $f(x)$
- Properties:

$f(x) \geq 0$ , for all  $x \in S_X$  (nonnegative density)

$$\int f(x) dx = 1 \text{ (total probability is 1)}$$

- $X$  is continuous iff there is a density function  $f_X$  such that the following holds:

$$\begin{aligned} \Pr(a \leq X \leq b) &= \int_a^b f_X(x) dx \\ &= \text{Area under } f \text{ between } a \text{ and } b \end{aligned}$$

- The support  $S_X$  consists of all  $x$  for which  $f_X(x) > 0$

# Normalization

# Normalization

- We must ensure that probability distributions sum / integrate to 1 (i.e., total probability must equal 1)

# Normalization

- We must ensure that probability distributions sum / integrate to 1 (i.e., total probability must equal 1)
- **Normalization:** Scalar adjustment in order to ensure that  $\Pr(S_X) = 1$

# Normalization

- We must ensure that probability distributions sum / integrate to 1 (i.e., total probability must equal 1)
- **Normalization:** Scalar adjustment in order to ensure that  $\Pr(S_X) = 1$
- If  $g(x) > 0$  for all  $x \in S_X$ , then

# Normalization

- We must ensure that probability distributions sum / integrate to 1 (i.e., total probability must equal 1)
- **Normalization:** Scalar adjustment in order to ensure that  $\Pr(S_X) = 1$
- If  $g(x) > 0$  for all  $x \in S_X$ , then

Discrete:  $p(x) = \frac{g(x)}{\sum_{x^* \in S_X} g(x^*)}$

# Normalization

- We must ensure that probability distributions sum / integrate to 1 (i.e., total probability must equal 1)
- **Normalization:** Scalar adjustment in order to ensure that  $\Pr(S_X) = 1$
- If  $g(x) > 0$  for all  $x \in S_X$ , then

$$\text{Discrete: } p(x) = \frac{g(x)}{\sum_{x^* \in S_X} g(x^*)}$$

$$\text{Continuous: } f(x) = \frac{g(x)}{\int_{x^* \in S_X} g(x^*) dx^*}$$

$$\begin{aligned} & \int_{S_X} f(x) dx \\ &= \int_{S_X} \frac{g(x)}{\text{NL}} dx \\ &= \frac{1}{\text{NL}} \int_{S_X} g(x) dx \\ &= 1. \end{aligned}$$

# Normalization


- We must ensure that probability distributions sum / integrate to 1 (i.e., total probability must equal 1)
- **Normalization:** Scalar adjustment in order to ensure that  $\Pr(S_X) = 1$
- If  $g(x) > 0$  for all  $x \in S_X$ , then

$$\text{Discrete: } p(x) = \frac{g(x)}{\sum_{x^* \in S_X} g(x^*)}$$

$$\text{Continuous: } f(x) = \frac{g(x)}{\int_{x^* \in S_X} g(x^*) dx^*}$$


- *Normalization constant:* 1/denominator

# Normalization: Example



# Normalization: Example

- Suppose that we generally know that probability is distributed according to the following curve:



# Normalization: Example

# Normalization: Example

- We can generally define the shape of this curve as

# Normalization: Example

- We can generally define the shape of this curve as

$$g(x) = 1 - x^2, \quad -1 \leq x \leq 1$$

# Normalization: Example

- We can generally define the shape of this curve as

$$g(x) = 1 - x^2, \quad -1 \leq x \leq 1$$

- Is this a proper density?

$$\begin{aligned} \int_{-1}^1 g(x) dx &= \int_{-1}^1 (1 - x^2) dx = \left[ x - \frac{x^3}{3} \right]_{-1}^1 \\ &= \left( 1 - \frac{1}{3} \right) - \left( -1 + \frac{1}{3} \right) \\ &= \frac{2}{3} - \frac{2}{3} = \frac{4}{3} \end{aligned}$$

# Normalization: Example

- We can generally define the shape of this curve as

$$g(x) = 1 - x^2, \quad -1 \leq x \leq 1$$

- Is this a proper density? **No**

# Normalization: Example

- We can generally define the shape of this curve as

$$g(x) = 1 - x^2, \quad -1 \leq x \leq 1$$

- Is this a proper density?
- What's the normalization constant? 3/4

# Normalization: Example

- We can generally define the shape of this curve as

$$g(x) = 1 - x^2, \quad -1 \leq x \leq 1$$

- Is this a proper density?
- What's the normalization constant?

# Normalization: Example

- We can generally define the shape of this curve as

$$g(x) = 1 - x^2, \quad ? \geq 0 \quad -1 \leq x \leq 1$$

- Is this a proper density?
- What's the normalization constant?       $1/\sqrt{2}$

- What is  $f(x)$ ?

$$f(x) = \frac{1}{\sqrt{2}} (1 - x^2)^{-1/2}$$

# Normalization: Example

PDFs

- We can generally define the shape of this curve as

$$g(x) = 1 - x^2, \quad -1 \leq x \leq 1$$

- Is this a proper density?
- What's the normalization constant?
- What is  $f(x)$ ?

# Cumulative Distribution Functions (CDFs)

# Cumulative Distribution Functions (CDFs)

- The **cumulative distribution function (CDF)** of random variable  $X$  is

# Cumulative Distribution Functions (CDFs)

- The **cumulative distribution function (CDF)** of random variable  $X$  is

$$F_X(x) = \Pr(X \leq x) \text{ for all } x \in (-\infty, \infty)$$

# Cumulative Distribution Functions (CDFs)

- The **cumulative distribution function (CDF)** of random variable  $X$  is

$$F_X(x) = \Pr(X \leq x) \text{ for all } x \in (-\infty, \infty)$$

- If  $X$  is discrete with support  $S_X$ , then the CDF is defined as

# Cumulative Distribution Functions (CDFs)

- The **cumulative distribution function (CDF)** of random variable  $X$  is

$$F_X(x) = \Pr(X \leq x) \text{ for all } x \in (-\infty, \infty)$$

- If  $X$  is discrete with support  $S_X$ , then the CDF is defined as

$$F_X(x) = \Pr(X \leq x) = \sum_{u \in S_X : u \leq x} \Pr(X = u)$$

# Cumulative Distribution Functions (CDFs)

- The **cumulative distribution function (CDF)** of random variable  $X$  is

$$F_X(x) = \Pr(X \leq x) \text{ for all } x \in (-\infty, \infty)$$

- If  $X$  is discrete with support  $S_X$ , then the CDF is defined as

$$F_X(x) = \Pr(X \leq x) = \sum_{u \in S_X : u \leq x} \Pr(X = u)$$

- If  $X$  is continuous, then the CDF is defined as

# Cumulative Distribution Functions (CDFs)

- The **cumulative distribution function (CDF)** of random variable  $X$  is

$$F_X(x) = \Pr(X \leq x) \text{ for all } x \in (-\infty, \infty)$$

- If  $X$  is discrete with support  $S_X$ , then the CDF is defined as

$$F_X(x) = \Pr(X \leq x) = \sum_{u \in S_X : u \leq x} \Pr(X = u)$$

- If  $X$  is continuous, then the CDF is defined as

$$F_X(x) = \Pr(X \leq x) = \int_{-\infty}^x f_X(u) du$$


# Cumulative Distribution Functions (CDFs)

# Cumulative Distribution Functions (CDFs)

- Consider the parabolic density

# Cumulative Distribution Functions (CDFs)


- Consider the parabolic density



# Cumulative Distribution Functions (CDFs)

- Consider the parabolic density

$$f(x) = \begin{cases} 0 & x \in (-\infty, -1) \\ \frac{3}{4}(1 - x^2) & x \in [-1, 1] \\ 0 & x \in (1, \infty) \end{cases}$$




# Cumulative Distribution Functions (CDFs)

- Consider the parabolic density

$$f(x) = \begin{cases} 0 & x \in (-\infty, -1) \\ \frac{3}{4}(1 - x^2) & x \in [-1, 1] \\ 0 & x \in (1, \infty) \end{cases}$$

- Over the range  $x \in (-\infty, -1)$ , we have  $F(x) = 0$




# Cumulative Distribution Functions (CDFs)

- Consider the parabolic density

$$f(x) = \begin{cases} 0 & x \in (-\infty, -1) \\ \frac{3}{4}(1 - x^2) & x \in [-1, 1] \\ 0 & x \in (1, \infty) \end{cases}$$

- Over the range  $x \in (-\infty, -1)$ , we have  $F(x) = 0$
- Over the range  $x \in [-1, 1]$ , we have



$x \in (1, \infty)$

$$F(x) = \int_{-1}^x f(u) du = \int_{-1}^x \frac{3}{4}(1 - u^2) du$$


# Cumulative Distribution Functions (CDFs)

- Consider the parabolic density

$$f(x) = \begin{cases} 0 & x \in (-\infty, -1) \\ \frac{3}{4}(1 - x^2) & x \in [-1, 1] \\ 0 & x \in (1, \infty) \end{cases}$$

- Over the range  $x \in (-\infty, -1)$ , we have  $F(x) = 0$
- Over the range  $x \in [-1, 1]$ , we have

$$F(x) = \int_{-1}^x \frac{3}{4}(1 - u^2) du = -x^3/4 + 3x/4 + 1/2$$



# Cumulative Distribution Functions (CDFs)


- Consider the parabolic density

$$f(x) = \begin{cases} 0 & x \in (-\infty, -1) \\ \frac{3}{4}(1 - x^2) & x \in [-1, 1] \\ 0 & x \in (1, \infty) \end{cases}$$


- Over the range  $x \in (-\infty, -1)$ , we have  $F(x) = 0$
- Over the range  $x \in [-1, 1]$ , we have

$$F(x) = \int_{-1}^x \frac{3}{4}(1 - u^2) du = -x^3/4 + 3x/4 + 1/2$$

- Over the range  $x \in (1, \infty)$ , we have  $F(x) = 1$



# Cumulative Distribution Functions (CDFs)



- Consider the parabolic density

$$f(x) = \begin{cases} 0 & x \in (-\infty, -1) \\ \frac{3}{4}(1 - x^2) & x \in [-1, 1] \\ 0 & x \in (1, \infty) \end{cases}$$



- Over the range  $x \in (-\infty, -1)$ , we have  $F(x) = 0$

- Over the range  $x \in [-1, 1]$ , we have

$$F(x) = \int_{-1}^x \frac{3}{4}(1 - u^2) du = -x^3/4 + 3x/4 + 1/2$$

- Over the range  $x \in (1, \infty)$ , we have  $F(x) = 1$

$F(x) = \Pr(X \leq x)$



# Quantiles and Percentiles

# Quantiles and Percentiles

- Suppose that a student with an 85 on an exam scored higher than 72% of their classmates

# Quantiles and Percentiles

- Suppose that a student with an 85 on an exam scored higher than 72% of their classmates
- Then,  $\Pr(X \leq 85) = 0.72$

$$F(85)$$

# Quantiles and Percentiles

- Suppose that a student with an 85 on an exam scored higher than 72% of their classmates
- Then,  $\Pr(X \leq 85) = 0.72$
- We say that  $q = 85$  is the  $p = 0.72$  quantile of this distribution (also called the 72<sup>nd</sup> percentile)

# Quantiles and Percentiles


# Quantiles and Percentiles

- More generally: For a random variable  $X$ ,  $q$  is the  $p$ -quantile of  $X$  if

# Quantiles and Percentiles

- More generally: For a random variable  $X$ ,  $q$  is the  $p$ -quantile of  $X$  if

$$\Pr(X < q) \leq p \text{ and } \Pr(X > q) \leq 1 - p$$



# Quantiles and Percentiles

- More generally: For a random variable  $X$ ,  $q$  is the  $p$ -quantile of  $X$  if

$$\Pr(X < q) \leq p \text{ and } \Pr(X > q) \leq 1 - p$$

- The quantile function of  $X$  is then defined as

# Quantiles and Percentiles

- More generally: For a random variable  $X$ ,  $q$  is the  $p$ -quantile of  $X$  if

$$\Pr(X < q) \leq p \text{ and } \Pr(X > q) \leq 1 - p$$

- The quantile function of  $X$  is then defined as

$$Q(p) = \min\{x \in S_X : \Pr(X \leq x) \geq p\}$$

# Quantiles and Percentiles

- More generally: For a random variable  $X$ ,  $q$  is the  $p$ -quantile of  $X$  if

$$\Pr(X < q) \leq p \text{ and } \Pr(X > q) \leq 1 - p$$

- The quantile function of  $X$  is then defined as

$$Q(p) = \min\{x \in S_X : \Pr(X \leq x) \geq p\}$$

- If the CDF  $F_X(x)$  is continuous and strictly increasing on  $S_X$ , then

# Quantiles and Percentiles

- More generally: For a random variable  $X$ ,  $q$  is the  $p$ -quantile of  $X$  if

$$\Pr(X < q) \leq p \text{ and } \Pr(X > q) \leq 1 - p$$

- The quantile function of  $X$  is then defined as

$$Q(p) = \min\{x \in S_X : \Pr(X \leq x) \geq p\}$$

- If the CDF  $F_X(x)$  is continuous and strictly increasing on  $S_X$ , then

$$Q(p) = F_x^{-1}(p)$$

# Quantiles and Percentiles

- More generally: For a random variable  $X$ ,  $q$  is the  $p$ -quantile of  $X$  if

$$\Pr(X < q) \leq p \text{ and } \Pr(X > q) \leq 1 - p$$

- The quantile function of  $X$  is then defined as

$$Q(p) = \min\{x \in S_X : \Pr(X \leq x) \geq p\}$$

- If the CDF  $F_X(x)$  is continuous and strictly increasing on  $S_X$ , then

$$Q(p) = F_x^{-1}(p)$$

q<sup>norm</sup>

- Although  $Q(p)$  is uniquely defined, the  $p$ -quantile may not be unique

# Quantiles and Percentiles: Example

# Quantiles and Percentiles: Example

- Consider the parabolic density,  $f(x) = \frac{3}{4}(1 - x^2)$

# Quantiles and Percentiles: Example

- Consider the parabolic density,  $f(x) = \frac{3}{4}(1 - x^2)$   $S_x \in [-1, 1]$
- What is the 0.25-quantile?  $F(0.25)$  find  $x$  s.t.  $F(x) = 0.25$


$$Q(p) = \left\{ \min_{x \in S_x} : \underbrace{\Pr(X \leq x)}_{\text{find } x} \geq p \right\}$$

$$F(x) = \boxed{\int_{-1}^x f(x) dx} = \underbrace{\frac{3}{4} \left( x - \frac{x^3}{3} \right) + \frac{1}{2}}_{F(x)}$$

$$\frac{3}{4} \left( x - \frac{x^3}{3} \right) + \frac{1}{2} = \frac{1}{4} \rightarrow x = \{-1.53, \boxed{-0.35}, 1.879\}$$

# Quantiles and Percentiles: Example

- Consider the parabolic density,  $f(x) = \frac{3}{4}(1 - x^2)$
- What is the 0.25-quantile?



# Quantiles and Percentiles: Example

- Consider the parabolic density,  $f(x) = \frac{3}{4}(1-x^2)$
- What is the 0.25-quantile?


$$g(x) \text{ over } [-1, 1]$$

$\uparrow$

$s_x$

$$g(x) = \frac{1}{[-10, \cdot]} \quad \text{over} \quad [-10, 10] .$$

$[3, 10]$



# Summarizing Probability Distributions

# Summarizing Probability Distributions

- Many random variables can take a large number of values, so an explicit probability distribution may be quite complicated

# Summarizing Probability Distributions

- Many random variables can take a large number of values, so an explicit probability distribution may be quite complicated
- We can describe a probability distribution with measures of central tendency and dispersion

central tendency

dispersion

# Summarizing Probability Distributions

- Many random variables can take a large number of values, so an explicit probability distribution may be quite complicated
- We can describe a probability distribution with measures of central tendency and dispersion
- *Population mean:* Average value that a random variable takes

# Summarizing Probability Distributions

- Many random variables can take a large number of values, so an explicit probability distribution may be quite complicated
- We can describe a probability distribution with measures of central tendency and dispersion
- *Population mean*: Average value that a random variable takes
- *Population variance*: Dispersion of the values relative to the population mean

# Summarizing Probability Distributions

- Many random variables can take a large number of values, so an explicit probability distribution may be quite complicated
- We can describe a probability distribution with measures of central tendency and dispersion
- *Population mean*: Average value that a random variable takes
- *Population variance*: Dispersion of the values relative to the population mean
- *Population standard deviation*: The square root of the population variance

# Expected Value

# Expected Value

- **Expected value** of  $X$ , denoted  $E(X)$ , represents a theoretical average of an infinitely large sample

# Expected Value

- **Expected value** of  $X$ , denoted  $E(X)$ , represents a theoretical average of an infinitely large sample
- $E(X)$  is what we “expect”  $X$  to equal; the population mean of  $X$

# Expected Value

- **Expected value** of  $X$ , denoted  $E(X)$ , represents a theoretical average of an infinitely large sample
  - $E(X)$  is what we “expect”  $X$  to equal; the population mean of  $X$
- We use the notation  $\underline{\mu} = \mu_X = E(X)$

# Expected Value

# Expected Value

- If  $X$  is a discrete random variable:

# Expected Value

3 coins     $\frac{1}{2} H, \frac{1}{2} T$

$X = \# Hs$

- If  $X$  is a discrete random variable:

$$\mu_X = E(X) = \sum_{x \in S_X} x \cdot \Pr(X = x)$$

$$\Pr(X=0) = \frac{1}{8}$$

$$E[X] = \sum_{x \in \{0,1,2,3\}} x \cdot \Pr(X=x)$$

$$\Pr(X=1) = \frac{3}{8}$$

$$= 0 \times \frac{1}{8} + 1 \times \frac{3}{8} + 2 \times \frac{3}{8} + 3 \times \frac{1}{8}$$

$$\frac{2 \times 6 + 3}{8} = \boxed{1.5}$$

# Expected Value

- If  $X$  is a discrete random variable:

$$\mu_X = E(X) = \sum_{x \in S_X} x \cdot \Pr(X = x)$$

- If  $X$  is a continuous random variable:

# Expected Value

- If  $X$  is a discrete random variable:

$$\mu_X = E(X) = \sum_{x \in S_X} x \cdot \Pr(X = x)$$

- If  $X$  is a continuous random variable:

$$\mu_X = E(X) = \int_{-\infty}^{\infty} x \cdot f_X(x) dx$$

# Expected Value

- If  $X$  is a discrete random variable:

$$\mu_X = E(X) = \sum_{x \in S_X} x \cdot \Pr(X = x)$$

- If  $X$  is a continuous random variable:

$$\mu_X = E(X) = \int_{-\infty}^{\infty} x \cdot f_X(x) dx$$

- If  $c$  is a constant, then

# Expected Value

- If  $X$  is a discrete random variable:

$$\mu_X = E(X) = \sum_{x \in S_X} x \cdot \Pr(X = x)$$

- If  $X$  is a continuous random variable:

$$\mu_X = E(X) = \int_{-\infty}^{\infty} x \cdot f_X(x) dx$$

- If  $c$  is a constant, then

$$E(c) = c$$

# Linearity of Expectation

# Linearity of Expectation

- For any random variables  $X$  and  $Y$ :

# Linearity of Expectation

- For any random variables  $X$  and  $Y$ :

$$E(X + Y) = E(X) + E(Y)$$

# Linearity of Expectation

$$\sum_x \sum_y (x, y) \Pr(X=x, Y=y)$$

- For any random variables  $X$  and  $Y$ :

$$E(X + Y) = E(X) + E(Y)$$

- This holds even if  $X$  and  $Y$  are *not* independent

# Linearity of Expectation

- For any random variables  $X$  and  $Y$ :

$$E(X + Y) = E(X) + E(Y)$$


- This holds even if  $X$  and  $Y$  are *not* independent
- In general, for random variables  $X_1, \dots, X_n$ :

# Linearity of Expectation

- For any random variables  $X$  and  $Y$ :

$$E(X + Y) = E(X) + E(Y)$$

- This holds even if  $X$  and  $Y$  are *not* independent
- In general, for random variables  $X_1, \dots, X_n$ :

$$E\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n E(X_i)$$


# Mean of a Random Variable: Example

# Mean of a Random Variable: Example

- Q1: What is the expected number of heads when flipping a fair coin ( $1/2$  H,  $1/2$  T)?

# Mean of a Random Variable: Example

- Q1: What is the expected number of heads when flipping a fair coin ( $1/2$  H,  $1/2$  T)?
- Q2: What is the expected number of heads when flipping an unfair coin ( $2/3$  H,  $1/3$  T)?

# Mean of a Random Variable: Example

- Q1: What is the expected number of heads when flipping a fair coin (1/2 H, 1/2 T)?

$$0 \times \frac{1}{2} + 1 \times \frac{1}{2} = \frac{1}{2}$$

- Q2: What is the expected number of heads when flipping an unfair coin (2/3 H, 1/3 T)?

$$0 \times \frac{1}{3} + 1 \times \frac{2}{3} = \frac{2}{3}$$

- Q3: What is the expected number of heads when flipping three fair coins (1/2 H, 1/2 T) and two unfair coins (2/3 H, 1/3 T)?

$$3 \times \frac{1}{2} + 2 \times \frac{2}{3} = \frac{17}{6}$$


# Mean of a Random Variable: Example

# Mean of a Random Variable: Example

- Consider the parabolic density,  $f(x) = \frac{3}{4}(1 - x^2)$


# Mean of a Random Variable: Example

- Consider the parabolic density,  $f(x) = \frac{3}{4}(1 - x^2)$




# Mean of a Random Variable: Example

- Consider the parabolic density,  $f(x) = \frac{3}{4}(1 - x^2)$
- What is  $E(X)$ ?



# Mean of a Random Variable: Example

- Consider the parabolic density,  $f(x) = \frac{3}{4}(1 - x^2)$
- What is  $E(X)$ ?



- Intuition: By symmetry,  $E(X) = 0$

# Variance

# Variance

- The variance of  $X$ , denoted  $\text{var}(X)$ , measures the tendency of  $X$  to deviate from  $E(X)$  and is defined as follows

# Variance

$$E(c) = c$$

- The variance of  $X$ , denoted  $\text{var}(X)$ , measures the tendency of  $X$  to deviate from  $E(X)$  and is defined as follows

$$\begin{aligned} \text{var}(X) &= E((X - E(X))^2) \\ &= E(X^2) - E(X)^2 \end{aligned}$$

$E(X^2) \geq E(X)^2$

$$\begin{aligned} &= E(X^2) - 2XE(X) + E(X)^2 \\ &= E(X^2) - \cancel{2E(X)E(X)} + \underline{E(X)^2} \\ &= E(X^2) - E(X)^2 \end{aligned}$$

# Variance

- The variance of  $X$ , denoted  $\text{var}(X)$ , measures the tendency of  $X$  to deviate from  $E(X)$  and is defined as follows

$$\begin{aligned}\text{var}(X) &= E\left(\left(X - E(X)\right)^2\right) \\ &= E(X^2) - E(X)^2\end{aligned}$$

- We use the notation  $\sigma^2 = \sigma_X^2 = \text{var}(X)$

# Variance

- The variance of  $X$ , denoted  $\text{var}(X)$ , measures the tendency of  $X$  to deviate from  $E(X)$  and is defined as follows

$$\begin{aligned}\text{var}(X) &= E\left(\left(X - E(X)\right)^2\right) \\ &= E(X^2) - E(X)^2\end{aligned}$$

- We use the notation  $\sigma^2 = \sigma_X^2 = \text{var}(X)$
- The standard deviation is the square root of the variance:  $\sigma = \sigma_X = \sqrt{\text{var}(X)}$

# Variance

# Variance

- Recall:  $\text{var}(X) = E\left(\left(X - E(X)\right)^2\right)$

# Variance

- Recall:  $\text{var}(X) = E\left(\left(X - E(X)\right)^2\right)$
- Let  $X$  be a discrete random variable with mean  $\mu_X$ :

# Variance

- Recall:  $\text{var}(X) = E\left(\left(X - E(X)\right)^2\right)$
- Let  $X$  be a discrete random variable with mean  $\mu_X$ :

$$\sigma_X^2 = \sum_{S_X} (x - \mu_X)^2 \Pr(X = x)$$

# Variance

- Recall:  $\text{var}(X) = E\left(\left(X - E(X)\right)^2\right)$
- Let  $X$  be a discrete random variable with mean  $\mu_X$ :

$$\sigma_X^2 = \sum_{S_X} (x - \mu_X)^2 \Pr(X = x)$$

- Let  $X$  be a continuous random variable with mean  $\mu_X$

# Variance

$$E(X) = \sum_{x \in X} x \cdot \Pr(X=x)$$

- Recall:  $\text{var}(X) = E\left(\underline{(X - E(X))^2}\right)$
- Let  $X$  be a discrete random variable with mean  $\mu_X$ :

$$\sigma_X^2 = \sum_{S_X} (x - \mu_X)^2 \Pr(X=x)$$

- Let  $X$  be a continuous random variable with mean  $\mu_X$

$$\sigma_X^2 = \int_{-\infty}^{\infty} (x - \mu_X)^2 f_X(x) dx$$

# Variance: Example

# Variance: Example

- Setup: Flip two fair coins; let  $X$  be the number of heads

# Variance: Example

- Setup: Flip two fair coins; let  $X$  be the number of heads
- Q: What is  $\text{var}(X)$ ?

$$\begin{aligned}\sigma^2 &= \sum_{x \in S_X} (x - \mu_x)^2 \Pr(X = x) \\ &= (0-1)^2 \cdot \frac{1}{4} + (1-1)^2 \cdot \frac{1}{2} \\ &\quad + (2-1)^2 \cdot \frac{1}{4} = \boxed{\frac{1}{2}}\end{aligned}$$

$$\left[ \begin{array}{ll} 0 & : 1/4 \\ 1 & : 1/2 \\ 2 & : 1/4 \end{array} \right] \quad E(X) = 1$$

# Variance: Example

- Setup: Flip two fair coins; let  $X$  be the number of heads
- Q: What is  $\text{var}(X)$ ?
- Q: What is the standard deviation of  $X$ ?

$$\sqrt{1/2} = \sqrt{2}/2$$

# Variance: Example

- Setup: Flip two fair coins; let  $X$  be the number of heads
- Q: What is  $\text{var}(X)$ ?
- Q: What is the standard deviation of  $X$ ?

# Functions of Random Variables

# Functions of Random Variables

- Take random variable  $X$  and function  $g( \cdot )$

# Functions of Random Variables


- Take random variable  $X$  and function  $g( \cdot )$
- We can obtain a new random variable:  $Y = g(X)$

# Functions of Random Variables

- Take random variable  $X$  and function  $g( \cdot )$
- We can obtain a new random variable:  $Y = g(X)$
- This is what is called a *transformation of variables*

# Functions of Random Variables

- Take random variable  $X$  and function  $g( \cdot )$
- We can obtain a new random variable:  $Y = g(X)$
- This is what is called a *transformation of variables*
- In general, to get the distribution of  $Y$ , we have that for any event  $E \subseteq S_Y$ , we have  $p_Y(E) = p_X(g^{-1}(E))$



# Linear Transformations: Mean and Variance

# Linear Transformations: Mean and Variance

- Let  $g$  be a linear function of the form  $g(X) = aX + b$

# Linear Transformations: Mean and Variance

- Let  $g$  be a linear function of the form  $g(X) = aX + b$
- Let  $X$  be a random variable with mean  $\mu_X$  and variance  $\sigma_X^2$

# Linear Transformations: Mean and Variance

- Let  $g$  be a linear function of the form  $g(X) = aX + b$
- Let  $X$  be a random variable with mean  $\mu_X$  and variance  $\sigma_X^2$
- Define a new random variable  $Y = g(X) = aX + b$

# Linear Transformations: Mean and Variance

- Let  $g$  be a linear function of the form  $\underline{g(X) = aX + b}$
- Let  $X$  be a random variable with mean  $\underline{\mu_X}$  and variance  $\sigma_X^2$
- Define a new random variable  $Y = g(X) = aX + b$
- Finding the mean of  $Y$ :

$$\underline{a\mu_X + b}$$

# Linear Transformations: Mean and Variance

- Let  $g$  be a linear function of the form  $g(X) = aX + b$
- Let  $X$  be a random variable with mean  $\mu_X$  and variance  $\sigma_X^2$
- Define a new random variable  $Y = g(X) = aX + b$
- Finding the mean of  $Y$ :

$$\mu_Y = E(Y) = E(aX + b) = aE(X) + b = a\mu_X + b$$

# Linear Transformations: Mean and Variance

- Let  $g$  be a linear function of the form  $g(X) = aX + b$
- Let  $X$  be a random variable with mean  $\mu_X$  and variance  $\sigma_X^2$
- Define a new random variable  $Y = g(X) = aX + b$
- Finding the mean of  $Y$ :

$$\mu_Y = E(Y) = E(aX + b) = aE(X) + b = a\mu_X + b$$

- Finding the variance of  $Y$ :

# Linear Transformations: Mean and Variance

- Let  $g$  be a linear function of the form  $g(X) = aX + b$
- Let  $X$  be a random variable with mean  $\mu_X$  and variance  $\sigma_X^2$
- Define a new random variable  $Y = g(X) = aX + b$
- Finding the mean of  $Y$ :

$$\mu_Y = E(Y) = E(aX + b) = aE(X) + b = a\mu_X + b$$

- Finding the variance of  $Y$ :

$$\begin{aligned}\sigma_Y^2 &= \text{var}(Y) = E((aX + b - E(aX + b))^2) \\ &= E((aX + b - aE(X) - b)^2) = E((aX - aE(X))^2) \\ &= E(a^2(X - E(X))^2) = a^2 E((X - E(X))^2) \\ &= a^2 \cdot \text{var}(X) = \boxed{a^2 \cdot \sigma_X^2}\end{aligned}$$

# General Transformations: Mean

# General Transformations: Mean

- If we have  $Y = g(X)$  for general  $g(X)$ , then we have:

# General Transformations: Mean

- If we have  $Y = g(X)$  for general  $g(X)$ , then we have:

$$\mu_Y = E(\underline{Y}) = E(\underline{g(X)})$$

# General Transformations: Mean

- If we have  $Y = g(X)$  for general  $g(X)$ , then we have:

$$\mu_Y = E(Y) = E(g(X))$$

- We do **not** necessarily have that:

# General Transformations: Mean

- If we have  $Y = g(X)$  for general  $g(X)$ , then we have:

$$\mu_Y = E(Y) = E(g(X))$$

- We do **not** necessarily have that:

$$E(g(X)) = g(E(X))$$

# General Transformations: Mean

- If we have  $Y = g(X)$  for general  $g(X)$ , then we have:

$$\mu_Y = E(Y) = E(g(X))$$

- We do **not** necessarily have that:

$$E(g(X)) = g(E(X))$$

- Example: Consider  $X$  = the outcome of rolling a fair six-sided die, and let  $g(X) = X^2$

# General Transformations: Mean

- If we have  $Y = g(X)$  for general  $g(X)$ , then we have:

$$\mu_Y = E(Y) = E(g(X))$$

- We do **not** necessarily have that:

$$E(g(X)) = g(E(X))$$

- Example: Consider  $X$  = the outcome of rolling a fair six-sided die, and let  $g(X) = X^2$

$$E(g(X)) = \frac{1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2}{6} = \frac{91}{6} \approx 15.17$$

$\sum x \cdot \Pr(X=x)$

# General Transformations: Mean

- If we have  $Y = \underline{g(X)}$  for general  $g(X)$ , then we have:


$$\mu_Y = E(Y) = E(g(X))$$

$$\Pr(X=1) = \Pr(Y=1^2)$$

$$\Pr(X=2) = \Pr(Y=2^2)$$


- We do **not** necessarily have that:

$$E(g(X)) = g(E(X))$$



- Example: Consider  $X$  = the outcome of rolling a fair six-sided die, and let  $\underline{g(X)} = X^2$

$$E(g(X)) = \frac{1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2}{6} = \frac{91}{6} \approx 15.17$$



$$\text{Pr } g(E(X)) = \left( \frac{1+2+3+4+5+6}{6} \right)^2 = (3.5)^2 = 12.25$$

let cost =  $g(x)$

\$ - \$

# Independence

$$P(A, B) = P(A) \times P(B)$$

# Independence

- Two random variables  $X_1$  and  $X_2$  are independent if the following holds, for any two events  $E_1$  and  $E_2$ :

# Independence

- Two random variables  $X_1$  and  $X_2$  are independent if the following holds, for any two events  $E_1$  and  $E_2$ :

$$P(X_1 \in E_1 \cap X_2 \in E_2) = P(X_1 \in E_1) \cdot P(X_2 \in E_2)$$

# Independence

- Two random variables  $X_1$  and  $X_2$  are independent if the following holds, for any two events  $E_1$  and  $E_2$ :

$$P(X_1 \in E_1 \cap X_2 \in E_2) = P(X_1 \in E_1) \cdot P(X_2 \in E_2)$$

- Notation:

# Independence

- Two random variables  $X_1$  and  $X_2$  are independent if the following holds, for any two events  $E_1$  and  $E_2$ :

$$P(X_1 \in E_1 \cap X_2 \in E_2) = P(X_1 \in E_1) \cdot P(X_2 \in E_2)$$

- Notation:

$X_1 \perp X_2$  means  $X_1$  and  $X_2$  are independent

# Independence

- Two random variables  $X_1$  and  $X_2$  are independent if the following holds, for any two events  $E_1$  and  $E_2$ :

$$P(X_1 \in E_1 \cap X_2 \in E_2) = P(X_1 \in E_1) \cdot P(X_2 \in E_2)$$

- Notation:

$X_1 \perp X_2$  means  $X_1$  and  $X_2$  are independent

- If a collection of random variables  $X_1, X_2, \dots, X_n$  are all independent and have the same distribution, we say that they are i.i.d. (independent and identically distributed)

# Independence

- Two random variables  $X_1$  and  $X_2$  are independent if the following holds, for any two events  $E_1$  and  $E_2$ :

$$P(X_1 \in E_1 \cap X_2 \in E_2) = P(X_1 \in E_1) \cdot P(X_2 \in E_2)$$

- Notation:

$X_1 \perp X_2$  means  $X_1$  and  $X_2$  are independent

- If a collection of random variables  $X_1, X_2, \dots, X_n$  are all independent and have the same distribution, we say that they are i.i.d. (independent and identically distributed)
  - Example: Roll two dice, or flip three fair coins

# Covariance

# Covariance

- If two variables are not independent, we measure their dependency through their **covariance**

# Covariance

- If two variables are not independent, we measure their dependency through their **covariance**
- Let  $X$  and  $Y$  be two random variables with means  $\mu_X$  and  $\mu_Y$ , respectively

die :  $\{1, \dots, 6\}$

$X$  takes values in  $S_X$

$X$

$E \subseteq S_X$

$E = \{ \text{either a } 1 \text{ or } 6 \}$

$S_X = \{ \underline{1}, \dots, \underline{6} \}$

A diagram showing a set  $S_X$  containing elements 1 through 6. The element 6 is circled and labeled '1/6'. The element 1 is circled and labeled '1/6'.

# Covariance

- If two variables are not independent, we measure their dependency through their **covariance**
- Let  $X$  and  $Y$  be two random variables with means  $\mu_X$  and  $\mu_Y$ , respectively
- The covariance of  $X$  and  $Y$  is defined as follows:

# Covariance

- If two variables are not independent, we measure their dependency through their **covariance**
- Let  $X$  and  $Y$  be two random variables with means  $\mu_X$  and  $\mu_Y$ , respectively
- The covariance of  $X$  and  $Y$  is defined as follows:

$$\text{cov}(X, Y) = \sigma_{XY} = E((\underbrace{X - \mu_X}_{}) (\underbrace{Y - \mu_Y}_{})) = \underbrace{E(XY)}_{\text{ }} - E(X)E(Y)$$

$$\text{Var}(X) = E((\underbrace{(X - \mu_X)^2}_{}) ) = E(X^2) - E(X)^2$$

# Covariance

- If two variables are not independent, we measure their dependency through their **covariance**
- Let  $X$  and  $Y$  be two random variables with means  $\mu_X$  and  $\mu_Y$ , respectively
- The covariance of  $X$  and  $Y$  is defined as follows:


$$\text{cov}(X, Y) = \sigma_{XY} = E((X - \mu_X)(Y - \mu_Y)) = E(XY) - E(X)E(Y)$$

- Correlation (essentially normalized covariance):

# Covariance

$$\mathbf{x}' = (x_1 - \bar{x}, x_2 - \bar{x}, \dots, x_n - \bar{x})$$

$$\mathbf{y}' = (y_1 - \bar{y}, \dots, y_n - \bar{y})$$



- If two variables are not independent, we measure their dependency through their **covariance**

$$X = \{x : S_x\}$$

- Let  $X$  and  $Y$  be two random variables with means  $\mu_X$  and  $\mu_Y$ , respectively
- The covariance of  $X$  and  $Y$  is defined as follows:

$$\text{cov}(X, Y) = \sigma_{XY} = E((\underline{X - \mu_X})(\underline{Y - \mu_Y})) = E(XY) - E(X)E(Y)$$

- Correlation (essentially normalized covariance):

$$\text{corr}(X, Y) = \rho = \rho_{XY} = \frac{\text{cov}(X, Y)}{\sigma_X \sigma_Y}$$

**SAMPLE**

$$\frac{\sum_i (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\left(\sum_i (x_i - \bar{x})^2\right)\left(\sum_i (y_i - \bar{y})^2\right)}} \approx \cos \theta$$

$\|x'\| \quad \|y'\|$

# Properties of Covariance

# Properties of Covariance

- Given random variables  $X$  and  $Y$ , the following hold:

# Properties of Covariance

- Given random variables  $X$  and  $Y$ , the following hold:
  - If either  $X$  or  $Y$  is a constant, then  $\text{cov}(X, Y) = 0$  and  $\text{corr}(X, Y)$  is undefined

# Properties of Covariance

- Given random variables  $X$  and  $Y$ , the following hold:
  - If either  $X$  or  $Y$  is a constant, then  $cov(X, Y) = 0$  and  $corr(X, Y)$  is undefined
  - If  $X \perp Y$ , then  $cov(X, Y) = corr(X, Y) = 0$

# Properties of Covariance

- Given random variables  $X$  and  $Y$ , the following hold:
  - If either  $X$  or  $Y$  is a constant, then  $cov(X, Y) = 0$  and  $corr(X, Y)$  is undefined
  - If  $X \perp Y$ , then  $cov(X, Y) = corr(X, Y) = 0$
  - $cov(X, X) = var(X)$

# Properties of Covariance

- Given random variables  $X$  and  $Y$ , the following hold:
  - If either  $X$  or  $Y$  is a constant, then  $cov(X, Y) = 0$  and  $corr(X, Y)$  is undefined
  - If  $X \perp Y$ , then  $cov(X, Y) = corr(X, Y) = 0$
  - $cov(X, X) = var(X)$
  - $cov(X, Y) = cov(Y, X)$

# Linear Combinations

$$\rightarrow Y = ax + b$$

$$Z = \underline{ax} + \underline{by}$$

# Linear Combinations

- Suppose you have random variables  $X$  and  $Y$  with means  $\mu_X, \mu_Y$  and variances  $\sigma_X^2, \sigma_Y^2$

# Linear Combinations

- Suppose you have random variables  $X$  and  $Y$  with means  $\mu_X, \mu_Y$  and variances  $\sigma_X^2, \sigma_Y^2$
- Let  $Z = aX + bY$

# Linear Combinations

- Suppose you have random variables  $X$  and  $Y$  with means  $\mu_X, \mu_Y$  and variances  $\sigma_X^2, \sigma_Y^2$
- Let  $Z = aX + bY$
- The mean of  $Z$  is

$$E[Z] = E[aX + bY] = E[aX] + E[bY] = aE[X] + bE[Y]$$

$a\mu_X + b\mu_Y$        $\underbrace{\varepsilon}$

# Linear Combinations

- Suppose you have random variables  $X$  and  $Y$  with means  $\mu_X, \mu_Y$  and variances  $\sigma_X^2, \sigma_Y^2$
- Let  $Z = aX + bY$
- The mean of  $Z$  is

$$\mu_Z = E(Z) = E(aX + bY) = E(aX) + E(bY) = a\mu_X + b\mu_Y$$

# Linear Combinations

- Suppose you have random variables  $X$  and  $Y$  with means  $\mu_X, \mu_Y$  and variances  $\sigma_X^2, \sigma_Y^2$
- Let  $Z = aX + bY$
- The mean of  $Z$  is

$$\begin{aligned} Z &= aX \\ &+ bY \end{aligned}$$

$$\mu_Z = E(Z) = E(aX + bY) = E(aX) + E(bY) = a\mu_X + b\mu_Y$$

- The variance of  $Z$  is

$$\text{var}(Z) = \text{cov}(Z, Z) = \text{cov}(a\underline{X} + b\underline{Y}, a\underline{X} + b\underline{Y})$$

$$\sigma_Z^2 = \underbrace{a^2 \sigma_X^2}_{\geq 0} + \underbrace{b^2 \sigma_Y^2}_{\geq 0} + 2ab \underbrace{\text{cov}(X, Y)}_{\leq 0}$$

# Linear Combinations

- Suppose you have random variables  $X$  and  $Y$  with means  $\mu_X, \mu_Y$  and variances  $\sigma_X^2, \sigma_Y^2$
- Let  $Z = aX + bY$
- The mean of  $Z$  is

$$\mu_Z = E(Z) = E(aX + bY) = E(aX) + E(bY) = a\mu_X + b\mu_Y$$

- The variance of  $Z$  is

$$\sigma_Z^2 = a^2\sigma_X^2 + b^2\sigma_Y^2 + 2ab\sigma_{XY}$$

# Linear Combinations

- Suppose you have random variables  $X$  and  $Y$  with means  $\mu_X, \mu_Y$  and variances  $\sigma_X^2, \sigma_Y^2$
- Let  $Z = aX + bY$
- The mean of  $Z$  is

$$\mu_Z = E(Z) = E(aX + bY) = E(aX) + E(bY) = a\mu_X + b\mu_Y$$

- The variance of  $Z$  is
$$\sigma_Z^2 = a^2\sigma_X^2 + b^2\sigma_Y^2 + 2ab\sigma_{XY}$$
- The standard deviation of  $Z$  is

# Linear Combinations

- Suppose you have random variables  $X$  and  $Y$  with means  $\mu_X, \mu_Y$  and variances  $\sigma_X^2, \sigma_Y^2$
- Let  $Z = aX + bY$
- The mean of  $Z$  is

$$\mu_Z = E(Z) = E(aX + bY) = E(aX) + E(bY) = a\mu_X + b\mu_Y$$

- The variance of  $Z$  is
$$\sigma_Z^2 = a^2\sigma_X^2 + b^2\sigma_Y^2 + 2ab\sigma_{XY}$$
- The standard deviation of  $Z$  is

$$\sigma_Z = \sqrt{a^2\sigma_X^2 + b^2\sigma_Y^2 + 2ab\sigma_{XY}}$$

# Theoretical Distributions

# Theoretical Distributions

- Theoretical probability distributions describe what we expect to happen based on populations on a theoretical level

# Theoretical Distributions

- Theoretical probability distributions describe what we expect to happen based on populations on a theoretical level
- We will consider the following theoretical distributions (**D** = discrete, **C** = continuous):

# Theoretical Distributions

- Theoretical probability distributions describe what we expect to happen based on populations on a theoretical level
- We will consider the following theoretical distributions (**D** = discrete, **C** = continuous):
  - Bernoulli distribution (D)

# Theoretical Distributions

- Theoretical probability distributions describe what we expect to happen based on populations on a theoretical level
- We will consider the following theoretical distributions (**D** = discrete, **C** = continuous):
  - Bernoulli distribution (D)
  - Binomial distribution (D)

# Theoretical Distributions

- Theoretical probability distributions describe what we expect to happen based on populations on a theoretical level
- We will consider the following theoretical distributions (**D** = discrete, **C** = continuous):
  - Bernoulli distribution (D)
  - Binomial distribution (D)
  - Poisson distribution (D)

# Theoretical Distributions

- Theoretical probability distributions describe what we expect to happen based on populations on a theoretical level
- We will consider the following theoretical distributions (**D** = discrete, **C** = continuous):
  - Bernoulli distribution (D)
  - Binomial distribution (D)
  - Poisson distribution (D)
  - Geometric distribution (D)

# Theoretical Distributions

- Theoretical probability distributions describe what we expect to happen based on populations on a theoretical level
- We will consider the following theoretical distributions (**D** = discrete, **C** = continuous):
  - Bernoulli distribution (D)
  - Binomial distribution (D)
  - Poisson distribution (D)
  - Geometric distribution (D)
  - Uniform distribution (C)

# Theoretical Distributions

- Theoretical probability distributions describe what we expect to happen based on populations on a theoretical level
- We will consider the following theoretical distributions (**D** = discrete, **C** = continuous):
  - Bernoulli distribution (D)
  - Binomial distribution (D)
  - Poisson distribution (D)
  - Geometric distribution (D)
  - Uniform distribution (C)
  - Exponential distribution (C)

# Theoretical Distributions

- Theoretical probability distributions describe what we expect to happen based on populations on a theoretical level
- We will consider the following theoretical distributions (**D** = discrete, **C** = continuous):
  - Bernoulli distribution (D)
  - Binomial distribution (D)
  - Poisson distribution (D)
  - Geometric distribution (D)
  - Uniform distribution (C)
  - Exponential distribution (C)
  - Normal distribution (C)

# Bernoulli Distribution

# Bernoulli Distribution

- Let  $Y$  be a dichotomous random variable (takes one of two mutually exclusive values)

# Bernoulli Distribution

- Let  $Y$  be a dichotomous random variable (takes one of two mutually exclusive values)
  - Classic example: Coin flip

# Bernoulli Distribution

- Let  $Y$  be a dichotomous random variable (takes one of two mutually exclusive values)
  - Classic example: Coin flip
- Successes ( $= 1$ ) occur with probability  $p$  and failures ( $= 0$ ) occur with probability  $1 - p$ , for constant  $p \in [0,1]$

# Bernoulli Distribution

- Let  $Y$  be a dichotomous random variable (takes one of two mutually exclusive values)
  - Classic example: Coin flip
- Successes ( $= 1$ ) occur with probability  $p$  and failures ( $= 0$ ) occur with probability  $1 - p$ , for constant  $p \in [0,1]$
- Notation:  $\underbrace{Y \sim \text{Bern}(p)}$   
*"follows"*

# Bernoulli Distribution: Example

# Bernoulli Distribution: Example

- Let  $Y$  be a dichotomous random variable representing a coin flip

# Bernoulli Distribution: Example

- Let  $Y$  be a dichotomous random variable representing a coin flip
  - $Y = 1$ : heads

# Bernoulli Distribution: Example

- Let  $Y$  be a dichotomous random variable representing a coin flip
  - $Y = 1$ : heads
  - $Y = 0$ : tails

# Bernoulli Distribution: Example

- Let  $Y$  be a dichotomous random variable representing a coin flip
  - $Y = 1$ : heads
  - $Y = 0$ : tails
- If the coin is fair, then  $p = \frac{1}{2}$  and  $1 - p = \frac{1}{2}$

# Bernoulli Distribution: Example

- Let  $Y$  be a dichotomous random variable representing a coin flip
  - $Y = 1$ : heads
  - $Y = 0$ : tails
- If the coin is fair, then  $p =$       and  $1 - p =$
- If the coin has a 60% chance of landing heads, then  $p = .60$  and  $1 - p = .40$

# Bernoulli Distribution: Example

$$\sum x \cdot P(Y=x)$$

- Let  $Y$  be a dichotomous random variable representing a coin flip

- $Y = 1$ : heads

$$E(Y) = 1 \cdot p + 0 \cdot (1-p) = p$$

- $Y = 0$ : tails

- If the coin is fair, then  $p =$  and  $1 - p =$

- If the coin has a 60% chance of landing heads, then  $p =$  and  $1 - p =$

- What is  $E(Y)$  in terms of  $p$ ?

p

# Bernoulli Distribution: Example

- Let  $Y$  be a dichotomous random variable representing a coin flip

- $Y = 1$ : heads

$$var(Y) = E((Y - E(Y))^2) = \underbrace{E(Y^2)}_{E(Y) = p} - \underbrace{E(Y)^2}_p$$

- $Y = 0$ : tails

$$Y^2 : \begin{cases} 0^2 & (1-p) \\ 1^2 & p \end{cases} = p - p^2 = p(1-p)$$

- If the coin is fair, then  $p =$

$$\text{and } 1 - p =$$

- If the coin has a 60% chance of landing heads, then  $p =$

$$\text{and } 1 - p =$$

- What is  $E(Y)$  in terms of  $p$ ?

$$P$$

- What is  $var(Y)$  in terms of  $p$ ?

$$Pq$$

# Binomial Distribution

# Binomial Distribution

- Suppose we flip  $n$  i.i.d. coins instead of just one coin

# Binomial Distribution

- Suppose we flip  $n$  i.i.d. coins instead of just one coin
- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe

# Binomial Distribution

- Suppose we flip  $n$  i.i.d. coins instead of just one coin
- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe
- $X$  is binomially distributed

# Binomial Distribution

- Suppose we flip  $n$  i.i.d. coins instead of just one coin
- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe
- $X$  is binomially distributed
- **Binomial distribution:** If we have a sequence of  $n$  Bernoulli random variables, each with a probability of success  $p$ , then the total number of successes is a binomial random variable


# Binomial Distribution

- Suppose we flip  $n$  i.i.d. coins instead of just one coin
- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe
- $X$  is binomially distributed
- **Binomial distribution:** If we have a sequence of  $n$  Bernoulli random variables, each with a probability of success  $p$ , then the total number of successes is a binomial random variable
  - Assumptions: fixed number of trials, independent, constant  $p$

# Binomial Distribution

- Suppose we flip  $n$  i.i.d. coins instead of just one coin
- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe
- $X$  is binomially distributed
- **Binomial distribution:** If we have a sequence of  $n$  Bernoulli random variables, each with a probability of success  $p$ , then the total number of successes is a binomial random variable
  - Assumptions: fixed number of trials, independent, constant  $p$
  - Notation:  $X \sim Bin(n, p)$

# Binomial Distribution



# Binomial Coefficients

# Binomial Coefficients

- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe when we flip  $n$  i.i.d. coins

# Binomial Coefficients

- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe when we flip  $n$  i.i.d. coins
- Each coin has probability of heads  $p$ , and flips are independent

# Binomial Coefficients

- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe when we flip  $n$  i.i.d. coins
- Each coin has probability of heads  $p$ , and flips are independent
- Q: What is the probability of getting exactly  $x$  out of  $n$  successes?

# Binomial Coefficients

- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe when we flip  $n$  i.i.d. coins
- Each coin has probability of heads  $p$ , and flips are independent
- Q: What is the probability of getting exactly  $x$  out of  $n$  successes?
  - Choose which  $x$  trials succeed:  $\binom{n}{x}$

# Binomial Coefficients

- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe when we flip  $n$  i.i.d. coins
  - Each coin has probability of heads  $p$ , and flips are independent
  - Q: What is the probability of getting exactly  $x$  out of  $n$  successes?
    - Choose which  $x$  trials succeed:  $\binom{n}{x}$
    - Probability that these  $x$  trials succeed:

# Binomial Coefficients

- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe when we flip  $n$  i.i.d. coins
  - Each coin has probability of heads  $p$ , and flips are independent
  - Q: What is the probability of getting exactly  $x$  out of  $n$  successes?
    - Choose which  $x$  trials succeed:  $\binom{n}{x}$
    - Probability that these  $x$  trials succeed:  $p^x$
    - Probability that the other  $n - x$  trials fail:  $(1-p)^{n-x}$

# Binomial Coefficients

$$\begin{aligned} n &= 3 \\ p &= \frac{1}{3} \end{aligned}$$

- Let  $X = \sum_{i=1}^n X_i$  be the number of heads we observe when we flip  $n$  i.i.d. coins

- Each coin has probability of heads  $p$ , and flips are independent

- Q: What is the probability of getting exactly  $x$  out of  $n$  successes?

$$\Pr(X=2)$$

- Choose which  $x$  trials succeed:

$$\binom{3}{2}$$

$$\text{HHT} = \frac{1}{3} \cdot \frac{1}{3} \cdot \frac{2}{3}$$

$$\text{HTH} = \frac{1}{3} \cdot \frac{2}{3} \cdot \frac{1}{3}$$

$$\text{THT} = \frac{2}{3} \cdot \frac{1}{3} \cdot \frac{1}{3}$$

- Probability that these  $x$  trials succeed:

- Probability that the other  $n - x$  trials fail:

- In general,

$$\binom{n}{x}$$

$$p^x (1-p)^{n-x}$$

# Binomial Probabilities in R

# Binomial Probabilities in R

- Calculate probabilities in R:


# Binomial Probabilities in R

- Calculate probabilities in R:
  - Calculate  $\Pr(X = x)$  using `dbinom(x, n, p)`

# Binomial Probabilities in R

- Calculate probabilities in R:
  - Calculate  $\Pr(X = x)$  using `dbinom(x, n, p)`
  - Calculate  $\Pr(X \leq x)$  using `pbinom(x, n, p)`

# Binomial Probabilities in R



- Calculate probabilities in R:
  - Calculate  $\Pr(X = x)$  using `dbinom(x, n, p)`
  - Calculate  $\Pr(X \leq x)$  using `pbinom(x, n, p)`
  - Calculate  $\Pr(X \geq x)$  using `1-pbinom(x-1, n, p)`

# Binomial Distribution: Summary Measures

# Binomial Distribution: Summary Measures

- Note that a binomial distribution with parameters  $n$  and  $p$  is the sum of  $n$  independent Bernoulli distributions with parameter  $p$

# Binomial Distribution: Summary Measures

- Note that a binomial distribution with parameters  $n$  and  $p$  is the sum of  $n$  independent Bernoulli distributions with parameter  $p$

$$E(X) = \mu_X = np$$

$$E(X) = E(\sum X_i)$$

$$= \sum E(X_i)$$

$$= \sum_{i=1}^n p$$

# Binomial Distribution: Summary Measures

- Note that a binomial distribution with parameters  $n$  and  $p$  is the sum of  $n$  independent Bernoulli distributions with parameter  $p$

$$E(X) = \mu_X = np$$

$$\text{var}(X) = \sigma_X^2 = np(1 - p)$$

# Binomial Distribution: Summary Measures

- Note that a binomial distribution with parameters  $n$  and  $p$  is the sum of  $n$  independent Bernoulli distributions with parameter  $p$

$$E(X) = \mu_X = np$$

$$\text{var}(X) = \sigma_X^2 = np(1 - p)$$

$$\text{stdev}(X) = \sigma_X = \sqrt{np(1 - p)}$$

# Binomial Distribution: Summary Measures


- Note that a binomial distribution with parameters  $n$  and  $p$  is the sum of  $n$  independent Bernoulli distributions with parameter  $p$

$$E(X) = \mu_X = np$$

$$\text{var}(X) = \sigma_X^2 = np(1 - p)$$

$$\text{stdev}(X) = \sigma_X = \sqrt{np(1 - p)}$$

- Q: How does  $\text{var}(X)$  change with  $p \in [0,1]$ ?



# Binomial Distribution: Summary

# Binomial Distribution: Summary

- Main take-away points from the binomial distribution:

# Binomial Distribution: Summary

- Main take-away points from the binomial distribution:
  - Fixed number of independent Bernoulli trials,  $n$

# Binomial Distribution: Summary

- Main take-away points from the binomial distribution:
  - Fixed number of independent Bernoulli trials,  $n$
  - Constant probability of success,  $p$  (Bernoulli parameter)

# Binomial Distribution: Summary

- Main take-away points from the binomial distribution:
  - Fixed number of independent Bernoulli trials,  $n$
  - Constant probability of success,  $p$  (Bernoulli parameter)
  - Interested in the total number of successes in  $n$  trials (not order)

# Binomial Distribution: Summary

- Main take-away points from the binomial distribution:
  - Fixed number of independent Bernoulli trials,  $n$
  - Constant probability of success,  $p$  (Bernoulli parameter)
  - Interested in the total number of successes in  $n$  trials (not order)
  - Mean:  $\mu_X = np$

# Binomial Distribution: Summary

- Main take-away points from the binomial distribution:
  - Fixed number of independent Bernoulli trials,  $n$
  - Constant probability of success,  $p$  (Bernoulli parameter)
  - Interested in the total number of successes in  $n$  trials (not order)
  - Mean:  $\mu_X = np$
  - Variance:  $\sigma_X^2 = np(1 - p)$

# Binomial Distribution: Summary

- Main take-away points from the binomial distribution:
  - Fixed number of independent Bernoulli trials,  $n$
  - Constant probability of success,  $p$  (Bernoulli parameter)
  - Interested in the total number of successes in  $n$  trials (not order)
  - Mean:  $\mu_X = np$
  - Variance:  $\sigma_X^2 = np(1 - p)$
- Examples:

# Binomial Distribution: Summary

- Main take-away points from the binomial distribution:
  - Fixed number of independent Bernoulli trials,  $n$
  - Constant probability of success,  $p$  (Bernoulli parameter)
  - Interested in the total number of successes in  $n$  trials (not order)
  - Mean:  $\mu_X = np$
  - Variance:  $\sigma_X^2 = np(1 - p)$
- Examples:
  - Number of heads in 15 flips of a fair coin

# Poisson Distribution

# Poisson Distribution

- **Poisson distribution:** Probability of observing a certain number of discrete events within a known interval

# Poisson Distribution

- **Poisson distribution:** Probability of observing a certain number of discrete events within a known interval
- Models discrete events that occur infrequently in time or space

# Poisson Distribution

- **Poisson distribution:** Probability of observing a certain number of discrete events within a known interval
  - Models discrete events that occur infrequently in time or space
- Example:

# Poisson Distribution

- **Poisson distribution:** Probability of observing a certain number of discrete events within a known interval
  - Models discrete events that occur infrequently in time or space
- Example:
  - The number of babies born at Strong Memorial Hospital between 10 am and 2 pm today

# Poisson Distribution

- **Poisson distribution:** Probability of observing a certain number of discrete events within a known interval
  - Models discrete events that occur infrequently in time or space
- Example:
  - The number of babies born at Strong Memorial Hospital between 10 am and 2 pm today
  - The number of students who enter River Campus today

# Poisson Distribution

# Poisson Distribution

- Let  $X \in [0, \infty)$  be the number of occurrences of some event over a given interval

# Poisson Distribution

- Let  $X \in [0, \infty)$  be the number of occurrences of some event over a given interval
- Let  $\lambda > 0$  be the average number of occurrences of the event over the specified interval

# Poisson Distribution

- Let  $X \in [0, \infty)$  be the number of occurrences of some event over a given interval
- Let  $\lambda > 0$  be the average number of occurrences of the event over the specified interval
- In this case, we say that  $\overset{\sim}{X} \sim Pois(\lambda)$

# Poisson Distribution

- Let  $X \in [0, \infty)$  be the number of occurrences of some event over a given interval
- Let  $\lambda > 0$  be the average number of occurrences of the event over the specified interval
- In this case, we say that  $X \sim Pois(\lambda)$
- The probability function is given by  $P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$


$$\frac{\lambda^{\#}}{\#!}$$

# Poisson Distribution

- Let  $X \in [0, \infty)$  be the number of occurrences of some event over a given interval
- Let  $\lambda > 0$  be the average number of occurrences of the event over the specified interval
- In this case, we say that  $X \sim Pois(\lambda)$
- The probability function is given by  $P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$
- If  $X \sim Pois(\lambda)$ , then  $\mu_X = \sigma_X^2 = \lambda$

# Poisson Distribution

- Let  $X \in [0, \infty)$  be the number of occurrences of some event over a given interval
- Let  $\lambda > 0$  be the average number of occurrences of the event over the specified interval
- In this case, we say that  $X \sim Pois(\lambda)$
- The probability function is given by  $P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$
- If  $X \sim Pois(\lambda)$ , then  $\mu_X = \sigma_X^2 = \lambda$ 
  - For a Poisson distribution, both the mean and the variance are equal to  $\lambda$



# Poisson Distribution

# Poisson Distribution

- Poisson distribution assumptions:

# Poisson Distribution

- Poisson distribution assumptions:
  - The probability of an event occurring is proportional to the length of the interval

# Poisson Distribution

- Poisson distribution assumptions:
  - The probability of an event occurring is proportional to the length of the interval
  - Within an interval, an infinite number of events is theoretically possible

# Poisson Distribution

- Poisson distribution assumptions:
  - The probability of an event occurring is proportional to the length of the interval
  - Within an interval, an infinite number of events is theoretically possible
  - Events occur independently

# Poisson Distribution

- Poisson distribution assumptions:
  - The probability of an event occurring is proportional to the length of the interval
  - Within an interval, an infinite number of events is theoretically possible
  - Events occur independently
  - The number of events that occur must be non-negative

# Poisson Distribution: Example

# Poisson Distribution: Example

- Setup: We want to examine the probability of certain numbers of people developing a rare disease in the next year. On average, 1.95 people develop the disease per year

$$\lambda$$

# Poisson Distribution: Example


- Setup: We want to examine the probability of certain numbers of people developing a rare disease in the next year. On average, 1.95 people develop the disease per year
- Q: What is the probability of no one developing the disease in the next year?

$$P(X=0) = \frac{e^{-1.95} \cdot 1.95^0}{0!} = e^{-1.95} \approx 0.142$$

# Poisson Distribution: Example

- Setup: We want to examine the probability of certain numbers of people developing a rare disease in the next year. On average, 1.95 people develop the disease per year
- Q: What is the probability of no one developing the disease in the next year?


$\approx 0.142$



- Q: What is the probability of one person developing the disease in the next year?

$$\Pr(X=1) = \frac{e^{-1.95} \cdot 1.95^1}{1!} = 1.95 \times \approx 0.277$$

# Poisson Distribution: Visualized



# Poisson Probabilities in R

# Poisson Probabilities in R

- Calculate probabilities in R:

# Poisson Probabilities in R

- Calculate probabilities in R:
  - Calculate  $\Pr(X = x)$  using `dpois(x, lambda)`

# Poisson Probabilities in R

- Calculate probabilities in R:
  - Calculate  $\Pr(X = x)$  using `dpois(x, lambda)`
  - Calculate  $\Pr(X \leq x)$  using `ppois(x, lambda)`

# Poisson Probabilities in R

- Calculate probabilities in R:
  - Calculate  $\Pr(X = x)$  using dpois(x, lambda)
  - Calculate  $\Pr(X \leq x)$  using ppois(x, lambda)
  - Calculate  $\Pr(X \geq x)$  using 1 - ppois(x-1, lamda)

# Poisson Distribution: Summary

# Poisson Distribution: Summary

- Main take-away points from the Poisson distribution:

# Poisson Distribution: Summary

- Main take-away points from the Poisson distribution:
  - Fixed interval, independent events, interested in number of events in interval

# Poisson Distribution: Summary

- Main take-away points from the Poisson distribution:
  - Fixed interval, independent events, interested in number of events in interval
  - Unlimited number of events is theoretically possible

# Poisson Distribution: Summary

- Main take-away points from the Poisson distribution:
  - Fixed interval, independent events, interested in number of events in interval
  - Unlimited number of events is theoretically possible
  - Mean:  $\mu_X = \lambda$

# Poisson Distribution: Summary

- Main take-away points from the Poisson distribution:
  - Fixed interval, independent events, interested in number of events in interval
  - Unlimited number of events is theoretically possible
  - Mean:  $\mu_X = \lambda$
  - Variance:  $\sigma_X^2 = \lambda$

# Poisson Distribution: Summary

- Main take-away points from the Poisson distribution:
  - Fixed interval, independent events, interested in number of events in interval
  - Unlimited number of events is theoretically possible
  - Mean:  $\mu_X = \lambda$
  - Variance:  $\sigma_X^2 = \lambda$
- Examples:

# Poisson Distribution: Summary

- Main take-away points from the Poisson distribution:
  - Fixed interval, independent events, interested in number of events in interval
  - Unlimited number of events is theoretically possible
  - Mean:  $\mu_X = \lambda$
  - Variance:  $\sigma_X^2 = \lambda$
- Examples:
  - Number of calculators the book store sells this year

# Poisson Distribution: Summary

- Main take-away points from the Poisson distribution:
  - Fixed interval, independent events, interested in number of events in interval
  - Unlimited number of events is theoretically possible
  - Mean:  $\mu_X = \lambda$
  - Variance:  $\sigma_X^2 = \lambda$
- Examples:
  - Number of calculators the book store sells this year
  - Number of babies born today

# Geometric Distribution

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$
- Let  $X$  be the first index  $i$  for which  $Y_i = 1$  (location of first success)

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$
- Let  $X$  be the first index  $i$  for which  $Y_i = 1$  (location of first success)
- The probability mass function (PMF) is given by

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$
- Let  $X$  be the first index  $i$  for which  $Y_i = 1$  (location of first success)
- The probability mass function (PMF) is given by

$$P(X = x) = p(1 - p)^{x-1}$$

$\overbrace{\quad\quad\quad}^{\Pr(\text{failing first } x-1)}$

$\Pr(\text{succeeding on } x^+)$

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$
- Let  $X$  be the first index  $i$  for which  $Y_i = 1$  (location of first success)
- The probability mass function (PMF) is given by

$$P(X = x) = p(1 - p)^{x-1}$$

- Notation:  $X \sim \underline{\text{Geom}}(p)$

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$
- Let  $X$  be the first index  $i$  for which  $Y_i = 1$  (location of first success)
- The probability mass function (PMF) is given by

$$P(X = x) = p(1 - p)^{x-1}$$

- Notation:  $X \sim Geom(p)$
- Mean and variance:

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$
- Let  $X$  be the first index  $i$  for which  $Y_i = 1$  (location of first success)
- The probability mass function (PMF) is given by

$$P(X = x) = p(1 - p)^{x-1}$$

- Notation:  $X \sim Geom(p)$
- Mean and variance:

$$E(X) = \mu_X = \frac{1}{p}$$

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$
- Let  $X$  be the first index  $i$  for which  $Y_i = 1$  (location of first success)
- The probability mass function (PMF) is given by

$$P(X = x) = p(1 - p)^{x-1}$$

- Notation:  $X \sim Geom(p)$
- Mean and variance:

$$E(X) = \mu_X = \frac{1}{p}$$

$$var(X) = \sigma_X^2 = \frac{1-p}{p^2}$$

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$
- Let  $X$  be the first index  $i$  for which  $Y_i = 1$  (location of first success)
- The probability mass function (PMF) is given by

$$P(X = x) = \cancel{x} (1 - p)^{x-1} (1-p)$$

- Notation:  $X \sim Geom(p)$
- Mean and variance:

$$E(X) = \mu_X = \frac{1}{p}$$

$$var(X) = \sigma_X^2 = \frac{1-p}{p^2}$$

- CDF:  $\underline{P(X \leq x)} = 1 - \boxed{(1 - p)^x}$

$$\Pr(X \leq x) = 1 - \Pr(\text{didn't see } H \text{ in first } x \text{ flips})$$

# Geometric Distribution

- Suppose  $Y_1, Y_2, \dots$  is an *infinite* sequence of independent Bernoulli random variables with parameter  $p$
- Let  $X$  be the first index  $i$  for which  $Y_i = 1$  (location of first success)
- The probability mass function (PMF) is given by


$$P(X = x) = p(1 - p)^{x-1}$$

- Notation:  $X \sim Geom(p)$
- Mean and variance:

$$E(X) = \mu_X = \frac{1}{p}$$

$$var(X) = \sigma_X^2 = \frac{1-p}{p^2}$$

- CDF:  $P(X \leq x) = 1 - (1 - p)^x$



# Continuous Distributions

# Continuous Distributions

- Continuous random variables: Intuitively, discrete random variables that are infinitesimally close together

# Continuous Distributions

- Continuous random variables: Intuitively, discrete random variables that are infinitesimally close together
- Instead of having discrete bars for the density at each discrete value, we now have a continuous density curve

# Continuous Distributions

- Continuous random variables: Intuitively, discrete random variables that are infinitesimally close together
- Instead of having discrete bars for the density at each discrete value, we now have a continuous density curve
- The area under the curve equals 1 (law of total probability)

# Continuous Distributions

- Continuous random variables: Intuitively, discrete random variables that are infinitesimally close together
- Instead of having discrete bars for the density at each discrete value, we now have a continuous density curve
- The area under the curve equals 1 (law of total probability)
- Since the random variable  $X$  can take on an infinite number of values, the probability associated with any single outcome equals 0

# Continuous Distributions

- Continuous random variables: Intuitively, discrete random variables that are infinitesimally close together
- Instead of having discrete bars for the density at each discrete value, we now have a continuous density curve
- The area under the curve equals 1 (law of total probability)
- Since the random variable  $X$  can take on an infinite number of values, the probability associated with any single outcome equals 0
- The probability that  $X \in (x_1, x_2)$  is equal to the area under the curve that lies between these two values

# Uniform Distribution

# Uniform Distribution

- Let  $X$  be a continuous random variable which can take on any value between  $a$  and  $b$  with equal probability

# Uniform Distribution

- Let  $X$  be a continuous random variable which can take on any value between  $a$  and  $b$  with equal probability
- Any value outside the range  $[a, b]$  cannot occur

# Uniform Distribution

- Let  $X$  be a continuous random variable which can take on any value between  $a$  and  $b$  with equal probability
- Any value outside the range  $[a, b]$  cannot occur
- The uniform probability density is

# Uniform Distribution

- Let  $X$  be a continuous random variable which can take on any value between  $a$  and  $b$  with equal probability
- Any value outside the range  $[a, b]$  cannot occur
- The uniform probability density is

$$f_X(x) = \begin{cases} \frac{1}{b-a} & x \in [a, b] \\ 0 & \text{otherwise} \end{cases}$$

# Uniform Distribution

- Let  $X$  be a continuous random variable which can take on any value between  $a$  and  $b$  with equal probability
- Any value outside the range  $[a, b]$  cannot occur
- The uniform probability density is

$$f_X(x) = \begin{cases} \frac{1}{b-a} & x \in [a, b] \\ 0 & \text{otherwise} \end{cases}$$

- Notation:  $X \sim Unif(a, b)$

# Uniform Distribution

- Let  $X$  be a continuous random variable which can take on any value between  $a$  and  $b$  with equal probability
- Any value outside the range  $[a, b]$  cannot occur
- The uniform probability density is

$$f_X(x) = \begin{cases} \frac{1}{b-a} & x \in [a, b] \\ 0 & \text{otherwise} \end{cases}$$

- Notation:  $X \sim Unif(a, b)$
- $\mu_X = \frac{a+b}{2}$  and  $\sigma_X^2 = \frac{(b-a)^2}{12}$


# Uniform Distribution

- Let  $X$  be a continuous random variable which can take on any value between  $a$  and  $b$  with equal probability
- Any value outside the range  $[a, b]$  cannot occur
- The uniform probability density is

$$f_X(x) = \begin{cases} \frac{1}{b-a} & x \in [a, b] \\ 0 & \text{otherwise} \end{cases}$$

- Notation:  $X \sim \text{Unif}(a, b)$

$$\bullet \mu_X = \frac{a+b}{2} \text{ and } \sigma_X^2 = \frac{(b-a)^2}{12}$$



# Exponential Distribution

# Exponential Distribution

- A continuous random variable  $X$  is exponentially distributed if it follows the following density:

# Exponential Distribution

- A continuous random variable  $X$  is exponentially distributed if it follows the following density:

$$f_X(x) = \lambda e^{-\lambda x}, \lambda > 0$$

# Exponential Distribution

- A continuous random variable  $X$  is exponentially distributed if it follows the following density:

$$f_X(x) = \lambda e^{-\lambda x}, \lambda > 0$$

- Notation:  $X \sim Exp(\lambda)$

# Exponential Distribution

- A continuous random variable  $X$  is exponentially distributed if it follows the following density:

$$f_X(x) = \lambda e^{-\lambda x}, \lambda > 0$$


- Notation:  $X \sim Exp(\lambda)$
- Generalizes the geometric distribution

# Exponential Distribution

- A continuous random variable  $X$  is exponentially distributed if it follows the following density:

$$f_X(x) = \lambda e^{-\lambda x}, \lambda > 0$$

- Notation:  $X \sim Exp(\lambda)$
- Generalizes the geometric distribution




# Exponential Distribution

- A continuous random variable  $X$  is exponentially distributed if it follows the following density:

$$f_X(x) = \lambda e^{-\lambda x}, \lambda > 0$$

- Notation:  $X \sim Exp(\lambda)$
- Generalizes the geometric distribution

- $\mu_X = \frac{1}{\lambda}$  and  $\sigma_X^2 = \frac{1}{\lambda^2}$




# Exponential Distribution

- A continuous random variable  $X$  is exponentially distributed if it follows the following density:

$$f_X(x) = \lambda e^{-\lambda x}, \lambda > 0$$

- Notation:  $X \sim Exp(\lambda)$
- Generalizes the geometric distribution

- $\mu_X = \frac{1}{\lambda}$  and  $\sigma_X^2 = \frac{1}{\lambda^2}$
- CDF:  $F_X(x) = 1 - e^{-\lambda x}$



# Normal Distribution

# Normal Distribution

- The most common continuous distribution is the normal distribution (also called a Gaussian distribution or bell-shaped curve)

# Normal Distribution

- The most common continuous distribution is the normal distribution (also called a Gaussian distribution or bell-shaped curve)
  - Shape of the binomial distribution when  $p$  is constant but  $n \rightarrow \infty$

# Normal Distribution

- The most common continuous distribution is the normal distribution (also called a Gaussian distribution or bell-shaped curve)
  - Shape of the binomial distribution when  $p$  is constant but  $n \rightarrow \infty$
  - Shape of the Poisson distribution when  $\lambda \rightarrow \infty$

# Normal Distribution

- The most common continuous distribution is the normal distribution (also called a Gaussian distribution or bell-shaped curve)
  - Shape of the binomial distribution when  $p$  is constant but  $n \rightarrow \infty$
  - Shape of the Poisson distribution when  $\lambda \rightarrow \infty$
- Given  $\mu, \sigma$ , the density function is

# Normal Distribution

- The most common continuous distribution is the normal distribution (also called a Gaussian distribution or bell-shaped curve)
  - Shape of the binomial distribution when  $p$  is constant but  $n \rightarrow \infty$
  - Shape of the Poisson distribution when  $\lambda \rightarrow \infty$
- Given  $\mu, \sigma$ , the density function is

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2}$$

# Normal Distribution

- The most common continuous distribution is the normal distribution (also called a Gaussian distribution or bell-shaped curve)
  - Shape of the binomial distribution when  $p$  is constant but  $n \rightarrow \infty$
  - Shape of the Poisson distribution when  $\lambda \rightarrow \infty$
- Given  $\mu, \sigma$ , the density function is

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2}$$

- Notation:  $X \sim N(\mu, \sigma^2)$  – but in R, use standard deviation instead of variance


# Normal Distribution

- The most common continuous distribution is the normal distribution (also called a Gaussian distribution or bell-shaped curve)
  - Shape of the binomial distribution when  $p$  is constant but  $n \rightarrow \infty$
  - Shape of the Poisson distribution when  $\lambda \rightarrow \infty$
- Given  $\mu, \sigma$ , the density function is

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2}$$


- Notation:  $X \sim N(\mu, \sigma^2)$  – but in R, use standard deviation instead of variance
- Mean = median = mode =  $\mu$ , variance =  $\sigma^2$ , standard deviation =  $\sigma$

# Normal Distribution: Visualization




# Normal Distribution: Visualization

- $\mu$  (center) and  $\sigma^2$  (spread) fully define the normal distribution




# Normal Distribution: Visualization

- $\mu$  (center) and  $\sigma^2$  (spread) fully define the normal distribution
- Always symmetric



# Normal Distribution: Visualization

- $\mu$  (center) and  $\sigma^2$  (spread) fully define the normal distribution
- Always symmetric
- When  $\mu = 0$  and  $\sigma^2 = 1$ , we have the *standard normal distribution*



# Normal Distribution: z-scores

# Normal Distribution: z-scores

- Recall from Chapter 2 that a z-score tells us how many standard deviations an observation is from its mean:

# Normal Distribution: z-scores

- Recall from Chapter 2 that a z-score tells us how many standard deviations an observation is from its mean:

$$z = \frac{x - \mu}{\sigma}$$

# Normal Distribution: z-scores

- Recall from Chapter 2 that a z-score tells us how many standard deviations an observation is from its mean:

$$z = \frac{x - \mu}{\sigma}$$

- Z has the nice property that it will always be  $N(0,1)$

# Normal Distribution: z-scores

- Recall from Chapter 2 that a z-score tells us how many standard deviations an observation is from its mean:

$$z = \frac{x - \mu}{\sigma}$$

- $Z$  has the nice property that it will always be  $N(0,1)$
- Given  $X \sim N(\mu, \sigma)$ , we can calculate a z-score, which will be  $Z \sim N(0,1)$

# Normal Distribution: z-scores

- Recall from Chapter 2 that a z-score tells us how many standard deviations an observation is from its mean:

$$z = \frac{x - \mu}{\sigma}$$

- $Z$  has the nice property that it will always be  $N(0,1)$
- Given  $X \sim N(\mu, \sigma)$ , we can calculate a z-score, which will be  $Z \sim N(0,1)$
- Standardizes the procedures for all normal distribution problems

# Normal Distribution

# Normal Distribution

- Recall that for continuous distributions, we are interested in determining the probability of being in an interval:

# Normal Distribution

- Recall that for continuous distributions, we are interested in determining the probability of being in an interval:

$$\Pr(X \leq a), \Pr(X \geq b), \text{ or } \Pr(a \leq X \leq b)$$

# Normal Distribution

- Recall that for continuous distributions, we are interested in determining the probability of being in an interval:

$$\Pr(X \leq a), \Pr(X \geq b), \text{ or } \Pr(a \leq X \leq b)$$

- We can look at the plot of the normal distribution and determine the probability (= area under the curve between endpoints)

# Normal Distribution

- Recall that for continuous distributions, we are interested in determining the probability of being in an interval:

$$\Pr(X \leq a), \Pr(X \geq b), \text{ or } \Pr(a \leq X \leq b)$$

- We can look at the plot of the normal distribution and determine the probability (= area under the curve between endpoints)
- In general, we will use R to calculate area under the curve (i.e., probabilities)

# Normal Distribution

- Recall that for continuous distributions, we are interested in determining the probability of being in an interval:

$$\Pr(X \leq a), \Pr(X \geq b), \text{ or } \Pr(a \leq X \leq b)$$

- We can look at the plot of the normal distribution and determine the probability (= area under the curve between endpoints)
- In general, we will use R to calculate area under the curve (i.e., probabilities)
- By default, R works in terms of z-scores:

# Normal Distribution

- Recall that for continuous distributions, we are interested in determining the probability of being in an interval:

$$\Pr(X \leq a), \Pr(X \geq b), \text{ or } \Pr(a \leq X \leq b)$$

- We can look at the plot of the normal distribution and determine the probability (= area under the curve between endpoints)
- In general, we will use R to calculate area under the curve (i.e., probabilities)
- By default, R works in terms of z-scores:

$$\Pr(Z \leq z) : \text{pnorm}(z)$$

# Normal Distribution

- Recall that for continuous distributions, we are interested in determining the probability of being in an interval:

$$\Pr(X \leq a), \Pr(X \geq b), \text{ or } \Pr(a \leq X \leq b)$$

- We can look at the plot of the normal distribution and determine the probability (= area under the curve between endpoints)
- In general, we will use R to calculate area under the curve (i.e., probabilities)
- By default, R works in terms of z-scores:

$$\Pr(Z \leq z) : \text{pnorm}(z)$$

$$\Pr(Z \geq z) : 1 - \text{pnorm}(z)$$

# Normal Distribution

- Recall that for continuous distributions, we are interested in determining the probability of being in an interval:

$$\Pr(X \leq a), \Pr(X \geq b), \text{ or } \Pr(a \leq X \leq b)$$

- We can look at the plot of the normal distribution and determine the probability (= area under the curve between endpoints)
- In general, we will use R to calculate area under the curve (i.e., probabilities)
- By default, R works in terms of z-scores:

$$\Pr(Z \leq z) : \text{pnorm}(z)$$

$$\Pr(Z \geq z) : 1 - \text{pnorm}(z)$$

$$\Pr(z_1 \leq Z \leq z_2) : \text{pnorm}(z_2) - \text{pnorm}(z_1)$$

# Normal Distribution

# Normal Distribution

- The general process for calculating probabilities based on a normal distribution is as follows:

# Normal Distribution

- The general process for calculating probabilities based on a normal distribution is as follows:

- Calculate appropriate z-scores: 
$$z = \frac{X - \mu}{\sigma}$$

# Normal Distribution

- The general process for calculating probabilities based on a normal distribution is as follows:
  - Calculate appropriate z-scores: 
$$z = \frac{X - \mu}{\sigma}$$
  - Use R to calculate the probability based on this z-score (`pnorm(z)`)

# Normal Distribution: Example

# Normal Distribution: Example

- Suppose that test scores are normally distributed with mean 78 and standard deviation 9

# Normal Distribution: Example

- Suppose that test scores are normally distributed with mean 78 and standard deviation 9
- Q: What is the probability that a person scored below 60?

# Normal Distribution: Example

- Suppose that test scores are normally distributed with mean 78 and standard deviation 9
- Q: What is the probability that a person scored below 60?
- Q: What is the probability that a person scored between 80 and 90?

# Normal Probabilities in R (Shortcut)

# Normal Probabilities in R (Shortcut)

- We can let R do the entire process of calculating a z-score and probability for us

# Normal Probabilities in R (Shortcut)

- We can let R do the entire process of calculating a z-score and probability for us
- Let  $X \sim N(\mu, \sigma)$

# Normal Probabilities in R (Shortcut)

- We can let R do the entire process of calculating a z-score and probability for us
- Let  $X \sim N(\mu, \sigma)$

$\Pr(X \leq x)$  : `pnorm(x, mean, sd)`

# Normal Probabilities in R (Shortcut)

- We can let R do the entire process of calculating a z-score and probability for us
- Let  $X \sim N(\mu, \sigma)$

$\Pr(X \leq x)$  : `pnorm(x, mean, sd)`

$\Pr(X \geq x)$  : `1-pnorm(x, mean, sd)`

# Normal Probabilities in R (Shortcut)

- We can let R do the entire process of calculating a z-score and probability for us
- Let  $X \sim N(\mu, \sigma)$

$\Pr(X \leq x) : \text{pnorm}(x, \text{mean}, \text{sd})$

$\Pr(X \geq x) : 1 - \text{pnorm}(x, \text{mean}, \text{sd})$

$\Pr(x_1 \leq X \leq x_2) : \text{pnorm}(x_2, \text{mean}, \text{sd}) - \text{pnorm}(x_1, \text{mean}, \text{sd})$

# Normal Distribution

# Normal Distribution

- Suppose that BMI is normally distributed with mean 26.6 and standard deviation 3.2

# Normal Distribution

- Suppose that BMI is normally distributed with mean 26.6 and standard deviation 3.2
- What is the probability that a person has a BMI in the range (18.5, 24.9)?

# Normal Distribution

- Suppose that BMI is normally distributed with mean 26.6 and standard deviation 3.2
- What is the probability that a person has a BMI in the range (18.5, 24.9)?
- Find  $\Pr(18.5 \leq X \leq 24.9)$  :

# Normal Distribution

- Suppose that BMI is normally distributed with mean 26.6 and standard deviation 3.2
- What is the probability that a person has a BMI in the range (18.5, 24.9)?
- Find  $\Pr(18.5 \leq X \leq 24.9)$  :

$$z_1 = \frac{18.5 - 26.6}{3.2} = -2.53$$

# Normal Distribution

- Suppose that BMI is normally distributed with mean 26.6 and standard deviation 3.2
- What is the probability that a person has a BMI in the range (18.5, 24.9)?
- Find  $\Pr(18.5 \leq X \leq 24.9)$  :

$$z_1 = \frac{18.5 - 26.6}{3.2} = -2.53$$

$$z_2 = \frac{24.9 - 26.6}{3.2} = -0.53$$

# Normal Distribution

- Suppose that BMI is normally distributed with mean 26.6 and standard deviation 3.2
- What is the probability that a person has a BMI in the range (18.5, 24.9)?
- Find  $\Pr(18.5 \leq X \leq 24.9)$  :

$$z_1 = \frac{18.5 - 26.6}{3.2} = -2.53$$

$$z_2 = \frac{24.9 - 26.6}{3.2} = -0.53$$

$$\Pr(-2.53 \leq Z \leq -0.53) = 0.2924$$

# Normal Distribution

- Suppose that BMI is normally distributed with mean 26.6 and standard deviation 3.2
- What is the probability that a person has a BMI in the range (18.5, 24.9)?
- Find  $\Pr(18.5 \leq X \leq 24.9)$  :

$$z_1 = \frac{18.5 - 26.6}{3.2} = -2.53$$

$$z_2 = \frac{24.9 - 26.6}{3.2} = -0.53$$

$$\Pr(-2.53 \leq Z \leq -0.53) = 0.2924$$

- $\text{pnorm}(-0.53) - \text{pnorm}(-2.53) = 0.2924$

# Normal Distribution

- Suppose that BMI is normally distributed with mean 26.6 and standard deviation 3.2
- What is the probability that a person has a BMI in the range (18.5, 24.9)?
- Find  $\Pr(18.5 \leq X \leq 24.9)$  :

$$z_1 = \frac{18.5 - 26.6}{3.2} = -2.53$$

$$z_2 = \frac{24.9 - 26.6}{3.2} = -0.53$$

$$\Pr(-2.53 \leq Z \leq -0.53) = 0.2924$$

- $\text{pnorm}(-0.53) - \text{pnorm}(-2.53) = 0.2924$
- $\text{pnorm}(24.9, 26.6, 3.2) - \text{pnorm}(18.5, 26.6, 3.2) = 0.2919$

# Revisiting the Empirical Rule

# Revisiting the Empirical Rule

- How well does the empirical rule approximate the normal distribution?

# Revisiting the Empirical Rule

- How well does the empirical rule approximate the normal distribution?
  - $\Pr(-1 \leq Z \leq 1) = 0.683$

# Revisiting the Empirical Rule

- How well does the empirical rule approximate the normal distribution?
  - $\Pr(-1 \leq Z \leq 1) = 0.683$
  - $\Pr(-2 \leq Z \leq 2) = 0.954$

# Revisiting the Empirical Rule

- How well does the empirical rule approximate the normal distribution?
  - $\Pr(-1 \leq Z \leq 1) = 0.683$
  - $\Pr(-2 \leq Z \leq 2) = 0.954$
  - $\Pr(-3 \leq Z \leq 3) = 0.997$

# Revisiting the Empirical Rule

- How well does the empirical rule approximate the normal distribution?
  - $\Pr(-1 \leq Z \leq 1) = 0.683$
  - $\Pr(-2 \leq Z \leq 2) = 0.954$
  - $\Pr(-3 \leq Z \leq 3) = 0.997$
- Empirical rule (68%, 95%, 99.7%) appears to be quite good

# Normal Distribution: Percentiles

# Normal Distribution: Percentiles

- Given data  $x_1, \dots, x_n$ , what value of  $x$  corresponds to a probability of the  $p^{th}$  percentile?

# Normal Distribution: Percentiles

- Given data  $x_1, \dots, x_n$ , what value of  $x$  corresponds to a probability of the  $p^{th}$  percentile?
- Strategy:

# Normal Distribution: Percentiles

- Given data  $x_1, \dots, x_n$ , what value of  $x$  corresponds to a probability of the  $p^{th}$  percentile?
- Strategy:
  - Find  $z$  value such that  $\Pr(Z \leq z) = p$  (lower tail probability is  $p$ )

# Normal Distribution: Percentiles

- Given data  $x_1, \dots, x_n$ , what value of  $x$  corresponds to a probability of the  $p^{th}$  percentile?
- Strategy:
  - Find  $z$  value such that  $\Pr(Z \leq z) = p$  (lower tail probability is  $p$ )
  - Solve for  $x$  by inverting z-score:  $x = z \cdot \sigma + \mu$

# Normal Distribution: Percentiles

- Given data  $x_1, \dots, x_n$ , what value of  $x$  corresponds to a probability of the  $p^{th}$  percentile?
- Strategy:
  - Find  $z$  value such that  $\Pr(Z \leq z) = p$  (lower tail probability is  $p$ )
  - Solve for  $x$  by inverting z-score:  $x = z \cdot \sigma + \mu$
- Directly in R: `qnorm(p, mean, sd)` ; `qnorm(p)` for z value

# Normal Distribution: Example

# Normal Distribution: Example

- Setup: Let  $X$  be a random variable that represents weights of patients in American hospital EDs;  $X$  is normally distributed with  $\mu = 160$  and  $\sigma = 15$

# Normal Distribution: Example

- Setup: Let  $X$  be a random variable that represents weights of patients in American hospital EDs;  $X$  is normally distributed with  $\mu = 160$  and  $\sigma = 15$
- Q1: Find the probability that a randomly selected patient in the ED weighs between 140 pounds and 210 pounds

# Normal Distribution: Example

- Setup: Let  $X$  be a random variable that represents weights of patients in American hospital EDs;  $X$  is normally distributed with  $\mu = 160$  and  $\sigma = 15$
- Q1: Find the probability that a randomly selected patient in the ED weighs between 140 pounds and 210 pounds
- Q2: Find the value that cuts off the upper 10% of the curve in American ED patient weights

# Sampling Distributions

# Sampling Distributions

- Suppose we want to estimate the mean value of some continuous random variable of interest

# Sampling Distributions

- Suppose we want to estimate the mean value of some continuous random variable of interest
- We can take a sample from the population and use the sample mean as an estimate of the population mean:  $\bar{x}$  is an estimate for  $\mu$

# Sampling Distributions

- Suppose we want to estimate the mean value of some continuous random variable of interest
- We can take a sample from the population and use the sample mean as an estimate of the population mean:  $\bar{x}$  is an estimate for  $\mu$
- For a normally distributed population,  $\bar{x}$  is the *maximum likelihood estimator* for  $\mu$

# Sampling Distributions

- Suppose we want to estimate the mean value of some continuous random variable of interest
- We can take a sample from the population and use the sample mean as an estimate of the population mean:  $\bar{x}$  is an estimate for  $\mu$
- For a normally distributed population,  $\bar{x}$  is the *maximum likelihood estimator* for  $\mu$ 
  - Value of the parameter that is most likely to have produced the observed sample data

# Sampling Distributions

- Suppose we want to estimate the mean value of some continuous random variable of interest
- We can take a sample from the population and use the sample mean as an estimate of the population mean:  $\bar{x}$  is an estimate for  $\mu$
- For a normally distributed population,  $\bar{x}$  is the *maximum likelihood estimator* for  $\mu$ 
  - Value of the parameter that is most likely to have produced the observed sample data
- Different samples will have different means

# Sampling Distributions

# Sampling Distributions

- What if you continued sampling  $m$  times?

# Sampling Distributions

- What if you continued sampling  $m$  times?
  - You take one random sample and get mean  $\bar{x}_1$ , take another random sample and get mean  $\bar{x}_2$ , and repeat until you have  $\bar{x}_1, \bar{x}_2, \dots, \bar{x}_m$

# Sampling Distributions

- What if you continued sampling  $m$  times?
  - You take one random sample and get mean  $\bar{x}_1$ , take another random sample and get mean  $\bar{x}_2$ , and repeat until you have  $\bar{x}_1, \bar{x}_2, \dots, \bar{x}_m$
  - Take  $m$  random samples for a total of  $m$  sample means

# Sampling Distributions

- What if you continued sampling  $m$  times?
  - You take one random sample and get mean  $\bar{x}_1$ , take another random sample and get mean  $\bar{x}_2$ , and repeat until you have  $\bar{x}_1, \bar{x}_2, \dots, \bar{x}_m$
  - Take  $m$  random samples for a total of  $m$  sample means
- These  $m$  means form a distribution with mean  $\mu$  and variance  $\frac{\sigma^2}{n}$  where  $n$  is the sample size

# Sampling Distributions

- What if you continued sampling  $m$  times?
  - You take one random sample and get mean  $\bar{x}_1$ , take another random sample and get mean  $\bar{x}_2$ , and repeat until you have  $\bar{x}_1, \bar{x}_2, \dots, \bar{x}_m$
  - Take  $m$  random samples for a total of  $m$  sample means
- These  $m$  means form a distribution with mean  $\mu$  and variance  $\frac{\sigma^2}{n}$  where  $n$  is the sample size
- Key idea:  $\bar{X}$  has its own distribution

# Sampling Distributions

- What if you continued sampling  $m$  times?
  - You take one random sample and get mean  $\bar{x}_1$ , take another random sample and get mean  $\bar{x}_2$ , and repeat until you have  $\bar{x}_1, \bar{x}_2, \dots, \bar{x}_m$
  - Take  $m$  random samples for a total of  $m$  sample means
- These  $m$  means form a distribution with mean  $\mu$  and variance  $\frac{\sigma^2}{n}$  where  $n$  is the sample size
- Key idea:  $\bar{X}$  has its own distribution
  - Standard deviation of  $\bar{X}$  is  $\frac{\sigma}{\sqrt{n}}$ ; this is known as the **standard error**

# Sampling Normal Distributions

# Sampling Normal Distributions

- If the population we are sampling from is normal, then the distribution of  $\bar{X}$  will also be normal

# Sampling Normal Distributions

- If the population we are sampling from is normal, then the distribution of  $\bar{X}$  will also be normal
- If  $X \sim N(\mu, \sigma)$ , then  $\bar{X} \sim N\left(\mu, \frac{\sigma}{\sqrt{n}}\right)$

# Central Limit Theorem (CLT)

# Central Limit Theorem (CLT)

- If the population we are sampling from is not normal, then we can use the *Central Limit Theorem (CLT)* to get the distribution of  $\bar{X}$

# Central Limit Theorem (CLT)

- If the population we are sampling from is not normal, then we can use the *Central Limit Theorem (CLT)* to get the distribution of  $\bar{X}$
- **Central Limit Theorem:** If the population we are sampling from is not normal, then the shape of the distribution of  $\bar{X}$  will be normal as long as  $n$  is sufficiently large (typically  $n \geq 30$  suffices)

# Central Limit Theorem (CLT)

# Central Limit Theorem (CLT)

- In particular, given that the distribution of an underlying population has mean  $\mu$  and standard deviation  $\sigma$ , the distribution of the sample means computed for samples of size  $n$  has three important properties:

# Central Limit Theorem (CLT)

- In particular, given that the distribution of an underlying population has mean  $\mu$  and standard deviation  $\sigma$ , the distribution of the sample means computed for samples of size  $n$  has three important properties:
  - The mean of the sampling distribution equals the population mean  $\mu$

# Central Limit Theorem (CLT)

- In particular, given that the distribution of an underlying population has mean  $\mu$  and standard deviation  $\sigma$ , the distribution of the sample means computed for samples of size  $n$  has three important properties:
  - The mean of the sampling distribution equals the population mean  $\mu$
  - The standard deviation of the distribution of sample means is equal to  $\frac{\sigma}{\sqrt{n}}$ , which is the standard error of the mean

# Central Limit Theorem (CLT)

- In particular, given that the distribution of an underlying population has mean  $\mu$  and standard deviation  $\sigma$ , the distribution of the sample means computed for samples of size  $n$  has three important properties:
  - The mean of the sampling distribution equals the population mean  $\mu$
  - The standard deviation of the distribution of sample means is equal to  $\frac{\sigma}{\sqrt{n}}$ , which is the standard error of the mean
  - Given that  $n$  is sufficiently large, the shape of the sampling distribution is approximabely normal

# Central Limit Theorem (CLT)

- In particular, given that the distribution of an underlying population has mean  $\mu$  and standard deviation  $\sigma$ , the distribution of the sample means computed for samples of size  $n$  has three important properties:
  - The mean of the sampling distribution equals the population mean  $\mu$
  - The standard deviation of the distribution of sample means is equal to  $\frac{\sigma}{\sqrt{n}}$ , which is the standard error of the mean
  - Given that  $n$  is sufficiently large, the shape of the sampling distribution is approximabely normal
- In notation:  $\bar{X} \sim N\left(\mu, \frac{\sigma}{\sqrt{n}}\right)$

# Central Limit Theorem (CLT)

# Central Limit Theorem (CLT)

- The further the underlying population is from normal, the larger the sample size you need to ensure normality of the sampling distribution

# Central Limit Theorem (CLT)

- The further the underlying population is from normal, the larger the sample size you need to ensure normality of the sampling distribution
- However, if the underlying population is normal, you do not need the central limit theorem to ensure normality of the sampling distribution – normality will hold regardless of the sample size if the underlying population is normal

# Central Limit Theorem (CLT)

- The further the underlying population is from normal, the larger the sample size you need to ensure normality of the sampling distribution
- However, if the underlying population is normal, you do not need the central limit theorem to ensure normality of the sampling distribution – normality will hold regardless of the sample size if the underlying population is normal
- Since  $\bar{X} \sim N(\mu, \sigma/\sqrt{n})$ , we can standardize  $\bar{X}$  to a standard normal distribution as follows:

# Central Limit Theorem (CLT)

- The further the underlying population is from normal, the larger the sample size you need to ensure normality of the sampling distribution
- However, if the underlying population is normal, you do not need the central limit theorem to ensure normality of the sampling distribution – normality will hold regardless of the sample size if the underlying population is normal
- Since  $\bar{X} \sim N(\mu, \sigma/\sqrt{n})$ , we can standardize  $\bar{X}$  to a standard normal distribution as follows:

$$Z = \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

# Sampling Distributions Example

# Sampling Distributions Example

- Setup: Suppose house prices have a distribution with a mean of  $\mu = \$450,000$  and standard deviation  $\sigma = \$100,000$ . You draw a random sample of  $n = 64$  houses and determine their prices

# Sampling Distributions Example

- Setup: Suppose house prices have a distribution with a mean of  $\mu = \$450,000$  and standard deviation  $\sigma = \$100,000$ . You draw a random sample of  $n = 64$  houses and determine their prices
- Q1: What is the mean of the distribution of sample means?

# Sampling Distributions Example

- Setup: Suppose house prices have a distribution with a mean of  $\mu = \$450,000$  and standard deviation  $\sigma = \$100,000$ . You draw a random sample of  $n = 64$  houses and determine their prices
- Q1: What is the mean of the distribution of sample means?
- Q2: What is the standard error of the sample mean?

# Sampling Distributions Example

- Setup: Suppose house prices have a distribution with a mean of  $\mu = \$450,000$  and standard deviation  $\sigma = \$100,000$ . You draw a random sample of  $n = 64$  houses and determine their prices
- Q1: What is the mean of the distribution of sample means?
- Q2: What is the standard error of the sample mean?
- Q3: What distribution does the sample mean follow?

# Sampling Distributions Example

- Setup: Suppose house prices have a distribution with a mean of  $\mu = \$450,000$  and standard deviation  $\sigma = \$100,000$ . You draw a random sample of  $n = 64$  houses and determine their prices
- Q1: What is the mean of the distribution of sample means?
- Q2: What is the standard error of the sample mean?
- Q3: What distribution does the sample mean follow?
- Q4: What is the probability that the sample mean of  $n = 64$  house prices is greater than \$500,000?

# Sampling Distribution of a Proportion

# Sampling Distribution of a Proportion

- Suppose we are interested in the proportion of the time that an event occurs

# Sampling Distribution of a Proportion

- Suppose we are interested in the proportion of the time that an event occurs
- If we take a sample of size  $n$  and observe  $x$  successes, then we could estimate the population proportion  $p$  by  $\hat{p} = x/n$

# Sampling Distribution of a Proportion

- Suppose we are interested in the proportion of the time that an event occurs
- If we take a sample of size  $n$  and observe  $x$  successes, then we could estimate the population proportion  $p$  by  $\hat{p} = x/n$
- We can do this sampling process  $m$  times for a total of  $m$  different values of  $\hat{p}_i$  for  $i \in \{1, 2, \dots, m\}$

# Sampling Distribution of a Proportion

- Suppose we are interested in the proportion of the time that an event occurs
- If we take a sample of size  $n$  and observe  $x$  successes, then we could estimate the population proportion  $p$  by  $\hat{p} = x/n$
- We can do this sampling process  $m$  times for a total of  $m$  different values of  $\hat{p}_i$  for  $i \in \{1, 2, \dots, m\}$
- These  $m$  proportions form a distribution with mean  $p$

# Sampling Distribution of a Proportion

- Suppose we are interested in the proportion of the time that an event occurs
- If we take a sample of size  $n$  and observe  $x$  successes, then we could estimate the population proportion  $p$  by  $\hat{p} = x/n$
- We can do this sampling process  $m$  times for a total of  $m$  different values of  $\hat{p}_i$  for  $i \in \{1, 2, \dots, m\}$
- These  $m$  proportions form a distribution with mean  $p$
- Standard deviation of  $\hat{p}$ , known as standard error, is  $\sqrt{\frac{p(1-p)}{n}}$

# Sampling Distribution of a Proportion

# Sampling Distribution of a Proportion

- The shape of the distribution of  $\hat{p}$  will be approximately normal as long as two conditions are met:

# Sampling Distribution of a Proportion

- The shape of the distribution of  $\hat{p}$  will be approximately normal as long as two conditions are met:
  - $np \geq 5$

# Sampling Distribution of a Proportion

- The shape of the distribution of  $\hat{p}$  will be approximately normal as long as two conditions are met:
  - $np \geq 5$
  - $n(1 - p) \geq 5$

# Sampling Distribution of a Proportion

- The shape of the distribution of  $\hat{p}$  will be approximately normal as long as two conditions are met:
  - $np \geq 5$
  - $n(1 - p) \geq 5$
- If both of these conditions are met, then  $\hat{p} \sim N\left(p, \sqrt{\frac{p(1 - p)}{n}}\right)$