

# Chapter 12: Nonparametric Inference

DSCC 462

Computational Introduction to Statistics

Anson Kahng

Fall 2022

# Plan for Today

# Plan for Today

- Introduce nonparametric analogues to hypothesis tests

# Plan for Today

- Introduce nonparametric analogues to hypothesis tests
- *Wilcoxon Signed-Rank Test*

# Plan for Today

- Introduce nonparametric analogues to hypothesis tests
- *Wilcoxon Signed-Rank Test*
  - Nonparametric analog to the one-sample or paired t-test


# Plan for Today

- Introduce nonparametric analogues to hypothesis tests
- *Wilcoxon Signed-Rank Test*
  - Nonparametric analog to the one-sample or paired t-test
- *Wilcoxon Rank-Sum Test (Mann-Whitney U Test)*

# Plan for Today

- Introduce nonparametric analogues to hypothesis tests
- *Wilcoxon Signed-Rank Test*
  - Nonparametric analog to the one-sample or paired t-test
- *Wilcoxon Rank-Sum Test (Mann-Whitney U Test)*
  - Nonparametric analog to the two-sample t-test

# Plan for Today, Visualized





# Nonparametric Methods

# Nonparametric Methods

- Think about the statistical tests we've done so far (z-test, t-test,  $\chi^2$ -test, F-test, etc.)

# Nonparametric Methods

- Think about the statistical tests we've done so far (z-test, t-test,  $\chi^2$ -test, F-test, etc.)
- In all of these, we knew the distribution of the population and we only needed to perform inference on the unknown parameters

# Nonparametric Methods

- Think about the statistical tests we've done so far (z-test, t-test,  $\chi^2$ -test, F-test, etc.)
- In all of these, we knew the distribution of the population and we only needed to perform inference on the unknown parameters
  - *Parametric methods*

# Nonparametric Methods

- Think about the statistical tests we've done so far (z-test, t-test,  $\chi^2$ -test, F-test, etc.)
- In all of these, we knew the distribution of the population and we only needed to perform inference on the unknown parameters
  - *Parametric methods*
  - We knew what distribution the population followed

# Nonparametric Methods

- Think about the statistical tests we've done so far (z-test, t-test,  $\chi^2$ -test, F-test, etc.)
- In all of these, we knew the distribution of the population and we only needed to perform inference on the unknown parameters
  - *Parametric methods*
  - We knew what distribution the population followed
- What if we don't know the distribution of the population?

# Nonparametric Methods

- Think about the statistical tests we've done so far (z-test, t-test,  $\chi^2$ -test, F-test, etc.)
- In all of these, we knew the distribution of the population and we only needed to perform inference on the unknown parameters
  - *Parametric methods*
  - We knew what distribution the population followed
- What if we don't know the distribution of the population?
  - *Nonparametric methods*

# Nonparametric Methods



# Nonparametric Methods

- When do we use nonparametric methods?

# Nonparametric Methods

- When do we use nonparametric methods?
  - When we don't know the underlying population distribution

# Nonparametric Methods

- When do we use nonparametric methods?
  - When we don't know the underlying population distribution
  - Or the data do not meet the assumptions needed for particular parametric techniques (e.g., CLT doesn't hold, normal approximation for proportions doesn't hold, etc.)

# Nonparametric Methods

- When do we use nonparametric methods?
  - When we don't know the underlying population distribution
  - Or the data do not meet the assumptions needed for particular parametric techniques (e.g., CLT doesn't hold, normal approximation for proportions doesn't hold, etc.)
- In this case, we use *nonparametric methods*, which make fewer assumptions regarding the underlying distribution

# Nonparametric Methods

- When do we use nonparametric methods?
  - When we don't know the underlying population distribution
  - Or the data do not meet the assumptions needed for particular parametric techniques (e.g., CLT doesn't hold, normal approximation for proportions doesn't hold, etc.)
- In this case, we use *nonparametric methods*, which make fewer assumptions regarding the underlying distribution
  - Also known as *distribution-free methods*

# Nonparametric Methods

# Nonparametric Methods

- Although nonparametric testing procedures make different assumptions, they still follow the same general setup as all hypothesis tests we have discussed so far

# Nonparametric Methods

- Although nonparametric testing procedures make different assumptions, they still follow the same general setup as all hypothesis tests we have discussed so far
  - Make a claim, develop hypotheses, state significance level



# Nonparametric Methods

- Although nonparametric testing procedures make different assumptions, they still follow the same general setup as all hypothesis tests we have discussed so far
  - Make a claim, develop hypotheses, state significance level
  - Calculate a test statistic based on a random sample of data

# Nonparametric Methods

- Although nonparametric testing procedures make different assumptions, they still follow the same general setup as all hypothesis tests we have discussed so far
  - Make a claim, develop hypotheses, state significance level
  - Calculate a test statistic based on a random sample of data
  - Determine whether to reject or fail to reject the null hypothesis based on the test statistic and significance level

# Motivating Example #1

# Motivating Example #1

- Suppose we want to determine whether a new drug changes tumor size

# Motivating Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed

# Motivating Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed
- Let's say that we have a sample of  $n$  pairs of observations (tumor size before drug vs. tumor size after drug), where  $n = 13$

# Motivating Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed
- Let's say that we have a sample of  $n$  pairs of observations (tumor size before drug vs. tumor size after drug), where  $n = 13$
- Can we apply the CLT here?

# Motivating Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed
- Let's say that we have a sample of  $n$  pairs of observations (tumor size before drug vs. tumor size after drug), where  $n = 13$
- Can we apply the CLT here?
- This means that we need to use a nonparametric method



# Motivating Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed
- Let's say that we have a sample of  $n$  pairs of observations (tumor size before drug vs. tumor size after drug), where  $n = 13$
- Can we apply the CLT here?
- This means that we need to use a nonparametric method
  - *Wilcoxon Signed-Rank Test*

# Wilcoxon Signed-Rank Test

# Wilcoxon Signed-Rank Test

- Used to compare two samples from populations that are not independent

# Wilcoxon Signed-Rank Test

- Used to compare two samples from populations that are not independent
  - Nonparametric analog to the paired t-test

# Wilcoxon Signed-Rank Test

- Used to compare two samples from populations that are not independent
  - Nonparametric analog to the paired t-test
- Because we are considering paired data, we may look at the difference in values for each pair of observations

# Wilcoxon Signed-Rank Test

- Used to compare two samples from populations that are not independent
  - Nonparametric analog to the paired t-test
- Because we are considering paired data, we may look at the difference in values for each pair of observations
- Does not require populations to be normally distributed

# Wilcoxon Signed-Rank Test

- Used to compare two samples from populations that are not independent
  - Nonparametric analog to the paired t-test
- Because we are considering paired data, we may look at the difference in values for each pair of observations
- Does not require populations to be normally distributed
- Takes into account both the magnitudes of the differences and their signs

# Wilcoxon Signed-Rank Test

- Used to compare two samples from populations that are not independent
  - Nonparametric analog to the paired t-test
- Because we are considering paired data, we may look at the difference in values for each pair of observations
- Does not require populations to be normally distributed
- Takes into account both the magnitudes of the differences and their signs
- Null hypothesis: In the underlying population differences among pairs, the median difference is equal to 0



# Wilcoxon Signed-Rank Test

- Used to compare two samples from populations that are not independent
  - Nonparametric analog to the paired t-test
- Because we are considering paired data, we may look at the difference in values for each pair of observations
- Does not require populations to be normally distributed
- Takes into account both the magnitudes of the differences and their signs
- Null hypothesis: In the underlying population differences among pairs, the median difference is equal to 0
  - Note that we consider medians for nonparametric tests as opposed to means

# Wilcoxon Signed-Rank Test: Back to Example #1

# Wilcoxon Signed-Rank Test: Back to Example #1

- Suppose we want to determine whether a new drug changes tumor size

# Wilcoxon Signed-Rank Test: Back to Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed

# Wilcoxon Signed-Rank Test: Back to Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed
- Let's say that we have a sample of  $n$  pairs of observations (tumor size before drug vs. tumor size after drug), where  $n = 13$

# Wilcoxon Signed-Rank Test: Back to Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed
- Let's say that we have a sample of  $n$  pairs of observations (tumor size before drug vs. tumor size after drug), where  $n = 13$
- $H_0$  : The median difference in tumor size equals 0

# Wilcoxon Signed-Rank Test: Back to Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed
- Let's say that we have a sample of  $n$  pairs of observations (tumor size before drug vs. tumor size after drug), where  $n = 13$
- $H_0$  : The median difference in tumor size equals 0
- $H_1$  : The median difference in tumor size is different from 0

# Wilcoxon Signed-Rank Test: Back to Example #1

- Suppose we want to determine whether a new drug changes tumor size
- We cannot assume that tumor sizes are normally distributed
- Let's say that we have a sample of  $n$  pairs of observations (tumor size before drug vs. tumor size after drug), where  $n = 13$
- $H_0$  : The median difference in tumor size equals 0
- $H_1$  : The median difference in tumor size is different from 0
- Test at the  $\alpha = 0.05$  significance level



# Wilcoxon Signed-Rank Test: Steps

# Wilcoxon Signed-Rank Test: Steps

- Next, take the difference for each pair of observations

# Wilcoxon Signed-Rank Test: Steps

- Next, take the difference for each pair of observations
- Ignoring the sign of these observations, rank their absolute values from smallest to largest

# Wilcoxon Signed-Rank Test: Steps

- Next, take the difference for each pair of observations
- Ignoring the sign of these observations, rank their absolute values from smallest to largest
  - A difference of 0 is not ranked

# Wilcoxon Signed-Rank Test: Steps

- Next, take the difference for each pair of observations
- Ignoring the sign of these observations, rank their absolute values from smallest to largest
  - A difference of 0 is not ranked
  - Remove pair from data set and reduce number of pairs by 1

# Wilcoxon Signed-Rank Test: Steps

- Next, take the difference for each pair of observations
- Ignoring the sign of these observations, rank their absolute values from smallest to largest
  - A difference of 0 is not ranked
  - Remove pair from data set and reduce number of pairs by 1
- Tied observations are assigned an average rank

$d_i$	$\rightarrow$	<u>0.5</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>
Ranks	$\rightarrow$	1	3	-3	3	3	1	5

# Wilcoxon Signed-Rank Test: Steps

- Next, take the difference for each pair of observations
- Ignoring the sign of these observations, rank their absolute values from smallest to largest
  - A difference of 0 is not ranked
  - Remove pair from data set and reduce number of pairs by 1
- Tied observations are assigned an average rank
- Finally separate the ranks by sign to either + or –

# Wilcoxon Signed-Rank Test: Data Table

↓

Subject	Tumor Size (mm)		Difference	Rank	Signed Rank	
	Before	After			+	-
1	36.3	27.1	9.2			
2	21.7	17.4	4.3	4.5		
3	45.1	33.1	12.0			
4	27.8	32.1	-4.3	4.5		
5	5.1	8.3	→ -2.2	2		
6	23.4	22.1	→ 1.3	1		
7	25.0	31.2	-6.2			
8	12.6	16.4	→ -3.8	3		
9	19.9	12.5	7.4			
10	22.1	22.1	→ 0	—	—	—
11	18.6	4.8	13.8			
12	8.9	22.6	-13.7			
13	12.7	6.4	6.3			
14	29.3	18.3	9.0			
15	26.4	21.8	4.6			



# Wilcoxon Signed-Rank Test: Data Table

Subject	Tumor Size (mm)		Difference	Rank	Signed Rank	
	Before	After			+	-
1	36.3	27.1	9.2	11	11	
2	21.7	17.4	4.3	4.5		
3	45.1	33.1	12.0	12		
4	27.8	32.1	-4.3	4.5		4.5
5	5.1	8.3	-2.2	2		
6	23.4	22.1	1.3	1		
7	25.0	31.2	-6.2	7		
8	12.6	16.4	-3.8	3		
9	19.9	12.5	7.4	9		
10	22.1	22.1	0	-		
11	18.6	4.8	13.8	14		
12	8.9	22.6	-13.7	13		
13	12.7	6.4	6.3	8		
14	29.3	18.3	9.0	10		
15	26.4	21.8	4.6	6		

# Wilcoxon Signed-Rank Test: Data Table

Subject	Tumor Size (mm)		Difference	Rank	Signed Rank	
	Before	After			+	-
1	36.3	27.1	9.2	11	11	
2	21.7	17.4	4.3	4.5	4.5	
3	45.1	33.1	12.0	12	12	
4	27.8	32.1	-4.3	4.5		4.5
5	5.1	8.3	2.2	2		2
6	23.4	22.1	1.3	1	1	
7	25.0	31.2	-6.2	7		7
8	12.6	16.4	-3.8	3		3
9	19.9	12.5	7.4	9	9	
10	22.1	22.1	0	-		
11	18.6	4.8	13.8	14	14	
12	8.9	22.6	-13.7	13		13
13	12.7	6.4	6.3	8	8	
14	29.3	18.3	9.0	10	10	
15	26.4	21.8	4.6	6	6	

$\Sigma \rightarrow T^+$

$\bar{\Sigma} \rightarrow T^-$

# Wilcoxon Signed-Rank Test

# Wilcoxon Signed-Rank Test

- Calculate the sum of the positive ranks,  $T^+$ , and the sum of the negative ranks,  $T^-$

# Wilcoxon Signed-Rank Test

- Calculate the sum of the positive ranks,  $T^+$ , and the sum of the negative ranks,  $T^-$
- Calculate  $T = T^+ - T^-$

# Wilcoxon Signed-Rank Test

- Calculate the sum of the positive ranks,  $T^+$ , and the sum of the negative ranks,  $T^-$
- Calculate  $T = T^+ - T^-$
- Under the null hypothesis, the median of the underlying population differences is equal to 0

# Wilcoxon Signed-Rank Test

$$T^+ = \sum \text{ranks all } +$$

- Calculate the sum of the positive ranks,  $T^+$ , and the sum of the negative ranks,  $T^-$
- Calculate  $T = T^+ - T^-$
- Under the null hypothesis, the median of the underlying population differences is equal to 0
- Thus, we expect approximately equal numbers of positive and negative ranks

$$- \frac{n(n+1)}{2} \quad / \quad \frac{n(n+1)}{2}$$

# Wilcoxon Signed-Rank Test

- Calculate the sum of the positive ranks,  $T^+$ , and the sum of the negative ranks,  $T^-$
- Calculate  $T = T^+ - T^-$ 

$-3$

$-2$

$-1$

$0$

$1$

$2$

$3$
- Under the null hypothesis, the median of the underlying population differences is equal to 0
- Thus, we expect approximately equal numbers of positive and negative ranks
- Additionally, the sum of the positive ranks should be approximately equal to the sum of the negative ranks, so  $T$  should be approximately 0



# Wilcoxon Signed-Rank Test

# Wilcoxon Signed-Rank Test

- Evaluate the null hypothesis using the test statistic:

# Wilcoxon Signed-Rank Test

- Evaluate the null hypothesis using the test statistic:

$$z_T = \frac{T - \mu_T}{\sigma_T}$$

# Wilcoxon Signed-Rank Test

- Evaluate the null hypothesis using the test statistic:

$$z_T = \frac{T - \mu_T}{\sigma_T}$$

- Note that

# Wilcoxon Signed-Rank Test

- Evaluate the null hypothesis using the test statistic:

$$z_T = \frac{T - \mu_T}{\sigma_T}$$

- Note that

$$\mu_T = 0$$

# Wilcoxon Signed-Rank Test


- Evaluate the null hypothesis using the test statistic:

$$z_T = \frac{T - \mu_T}{\sigma_T}$$

- Note that

$$\mu_T = 0$$


$$\sigma_T = \sqrt{\frac{n(n+1)(2n+1)}{6}}$$



A handwritten blue formula for the standard deviation  $\sigma_T$ , which is  $\sqrt{\sum_{k=1}^n k^2}$ . A blue arrow points from this handwritten formula towards the printed formula for  $\sigma_T$  in the previous block.

# Wilcoxon Signed-Rank Test

- Evaluate the null hypothesis using the test statistic:

$$z_T = \frac{T - \mu_T}{\sigma_T}$$



- Note that

$$\mu_T = 0$$

$$\sigma_T = \sqrt{\frac{n(n+1)(2n+1)}{6}}$$

- $Z_T \sim N(0,1)$  given that  $n$  is large enough (typically  $n > 12$ )

# Histogram of Differences





# Wilcoxon Signed-Rank Test

Subject	Signed Rank	
	+	-
1	11	
2	4.5	
3	12	
4		4.5
5		2
6	1	
7		7
8		3
9	9	
10		
11	14	
12		13
13	8	
14	10	
15	6	

# Wilcoxon Signed-Rank Test

- Based on the histogram, the difference in tumor size does not appear to be normally distributed, so we want to use the Wilcoxon signed-rank test

Subject	Signed Rank	
	+	-
1	11	
2	4.5	
3	12	
4		4.5
5		2
6	1	
7		7
8		3
9	9	
10		
11	14	
12		13
13	8	
14	10	
15	6	

# Wilcoxon Signed-Rank Test

- Based on the histogram, the difference in tumor size does not appear to be normally distributed, so we want to use the Wilcoxon signed-rank test
- $T^+ =$

Subject	Signed Rank	
	+	-
1	11	
2	4.5	
3	12	
4		4.5
5		2
6	1	
7		7
8		3
9	9	
10		
11	14	
12		13
13	8	
14	10	
15	6	

# Wilcoxon Signed-Rank Test

- Based on the histogram, the difference in tumor size does not appear to be normally distributed, so we want to use the Wilcoxon signed-rank test
- $T^+ =$
- $T^- =$

Subject	Signed Rank	
	+	-
1	11	
2	4.5	
3	12	
4		4.5
5		2
6	1	
7		7
8		3
9	9	
10		
11	14	
12		13
13	8	
14	10	
15	6	

# Wilcoxon Signed-Rank Test

- Based on the histogram, the difference in tumor size does not appear to be normally distributed, so we want to use the Wilcoxon signed-rank test
- $T^+ =$
- $T^- =$
- $T =$

Subject	Signed Rank	
	+	-
1	11	
2	4.5	
3	12	
4		4.5
5		2
6	1	
7		7
8		3
9	9	
10		
11	14	
12		13
13	8	
14	10	
15	6	

# Wilcoxon Signed-Rank Test

- Based on the histogram, the difference in tumor size does not appear to be normally distributed, so we want to use the Wilcoxon signed-rank test
- $T^+ = 75.5$
- $T^- = 29.5$
- $T = 46$
- $n = 14 > 12$

Subject	Signed Rank	
	+	-
1	11	
2	4.5	
3	12	
4		4.5
5		2
6	1	
7		7
8		3
9	9	
10		
11	14	
12		13
13	8	
14	10	
15	6	

75.5 29.5

# Wilcoxon Signed-Rank Test

# Wilcoxon Signed-Rank Test

- Given  $T = 46$ , we then have the following:



# Wilcoxon Signed-Rank Test

- Given  $T = 46$ , we then have the following:

$$\mu_T = 0$$

# Wilcoxon Signed-Rank Test

$$n = 14$$

- Given  $T = 46$ , we then have the following:

$$\mu_T = 0$$

$$\sigma_T = \sqrt{\frac{n(n+1)(2n+1)}{6}} = 31.86$$

# Wilcoxon Signed-Rank Test

- Given  $T = 46$ , we then have the following:

$$\mu_T = 0$$

$$\sigma_T = \sqrt{\frac{n(n+1)(2n+1)}{6}} =$$

- Thus,

# Wilcoxon Signed-Rank Test

- Given  $T = 46$ , we then have the following:

$$\mu_T = 0$$

$$\sigma_T = \sqrt{\frac{n(n+1)(2n+1)}{6}} = 31.86$$

- Thus,

$$z_T = \frac{T - \mu_T}{\sigma_T} = \frac{46 - 0}{31.86} = \underline{1.44}$$

# Wilcoxon Signed-Rank Test

# Wilcoxon Signed-Rank Test

- Calculating the p-value, we have

# Wilcoxon Signed-Rank Test

$$Z_T = \frac{T - \mu_T}{\sigma_T} \sim N(0,1)$$

- Calculating the p-value, we have

$$p = 2 \cdot \Pr(Z > 1.44) = 2 \cdot (1 - \text{pnorm}(1.44)) = 0.149$$

- Conclusion:  $p = 0.149 > \alpha = 0.05$

# Wilcoxon Signed-Rank Test



# Wilcoxon Signed-Rank Test

- If the sample size is  $n \leq 12$ , we cannot use the normal approximation

# Wilcoxon Signed-Rank Test

- If the sample size is  $n \leq 12$ , we cannot use the normal approximation
- In that case, we can use `psignrank(T, n)` in R to calculate the exact p-value

# Wilcoxon Signed-Rank Test

- If the sample size is  $n \leq 12$ , we cannot use the normal approximation
- In that case, we can use `psignrank(T, n)` in R to calculate the exact p-value
  - $2 * \text{psignrank}(\underset{24.5}{\cancel{74}}, n=14) = \underset{0.135}{\cancel{0.173}}$

**\*** R requires  $T = \min(T^+, T^-)$  for this to work correctly!

# Wilcoxon Signed-Rank Test: R Code

$$d_i = x_i - y_i$$

$$z \sim N(0,1)$$

```
> wilcox.test(before, after, paired=T, exact=F, correct=F)
```

Wilcoxon signed rank test

data: before and after

V = 64, p-value = 0.1961

alternative hypothesis: true location shift is not equal to 0

$\uparrow$   $pnorm(T, n)$

```
> wilcox.test(before, after, paired=T, exact=T, correct=F)
```

Wilcoxon signed rank test

data: before and after

V = 64, p-value = 0.2163

alternative hypothesis: true location shift is not equal to 0

$\uparrow$   $p_{signed\ rank}(T, n)$

# Motivating Example #2

# Motivating Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature

# Motivating Example #2


- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed

# Motivating Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed
- Take a sample of  $n_1 = 12$  people who do not have the disease



# Motivating Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed
- Take a sample of  $n_1 = 12$  people who do not have the disease
- Take a sample of  $n_2 = 15$  people who do have the disease

# Motivating Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed
- Take a sample of  $n_1 = 12$  people who do not have the disease
- Take a sample of  $n_2 = 15$  people who do have the disease
- How can we compare the median body temperature for these two populations?

# Motivating Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed
- Take a sample of  $n_1 = 12$  people who do not have the disease
- Take a sample of  $n_2 = 15$  people who do have the disease
- How can we compare the median body temperature for these two populations?
  - *Wilcoxon Rank-Sum Test (Mann-Whitney U Test)*

# Wilcoxon Rank-Sum Test

# Wilcoxon Rank-Sum Test

- Used to compare samples from independent populations


# Wilcoxon Rank-Sum Test

- Used to compare samples from independent populations
  - Nonparametric analog to the two-sample t-test

# Wilcoxon Rank-Sum Test

- Used to compare samples from independent populations
  - Nonparametric analog to the two-sample t-test
- Does not require populations to be normally distributed

# Wilcoxon Rank-Sum Test



- Used to compare samples from independent populations
  - Nonparametric analog to the two-sample t-test
- Does not require populations to be normally distributed
- Requires the two populations to have the same general shape



# Wilcoxon Rank-Sum Test

- Used to compare samples from independent populations
  - Nonparametric analog to the two-sample t-test
- Does not require populations to be normally distributed
- Requires the two populations to have the same general shape
- $H_0$  : The medians of the two populations are identical

# Wilcoxon Rank-Sum Test: Back to Example #2

# Wilcoxon Rank-Sum Test: Back to Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature

# Wilcoxon Rank-Sum Test: Back to Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed

# Wilcoxon Rank-Sum Test: Back to Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed
- Take a sample of  $n_1 = 12$  people who do not have the disease

# Wilcoxon Rank-Sum Test: Back to Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed
- Take a sample of  $n_1 = 12$  people who do not have the disease
- Take a sample of  $n_2 = 15$  people who do have the disease

# Wilcoxon Rank-Sum Test: Back to Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed
- Take a sample of  $n_1 = 12$  people who do not have the disease
- Take a sample of  $n_2 = 15$  people who do have the disease
- $H_0$  : The median body temperature for those without the disease is greater than or equal to those with the disease

# Wilcoxon Rank-Sum Test: Back to Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed
- Take a sample of  $n_1 = 12$  people who do not have the disease
- Take a sample of  $n_2 = 15$  people who do have the disease
- $H_0$  : The median body temperature for those without the disease is greater than or equal to those with the disease
- $H_1$  : The median body temperature for those without the disease is less than those with the disease



# Wilcoxon Rank-Sum Test: Back to Example #2

- Suppose that we want to determine whether having a certain disease is associated with a raised body temperature
- Cannot assume that body temperature is normally distributed
- Take a sample of  $n_1 = 12$  people who do not have the disease
- Take a sample of  $n_2 = 15$  people who do have the disease
- $H_0$  : The median body temperature for those without the disease is greater than or equal to those with the disease
- $H_1$  : The median body temperature for those without the disease is less than those with the disease
- Test at the  $\alpha = 0.05$  significance level

# Wilcoxon Rank-Sum Test: Steps

# Wilcoxon Rank-Sum Test: Steps

- Combine all data from the two samples and rank the observations from smallest to largest

# Wilcoxon Rank-Sum Test: Steps

- Combine all data from the two samples and rank the observations from smallest to largest
- If ranks are tied, we assign the average rank to those values

# Wilcoxon Rank-Sum Test: Steps

- Combine all data from the two samples and rank the observations from smallest to largest
- If ranks are tied, we assign the average rank to those values
- We then find the sum of ranks for each of the two original samples, denoted  $W_1$  and  $W_2$ , and then let  $W = \min(W_1, W_2)$

# Wilcoxon Rank-Sum Test: Steps

- Combine all data from the two samples and rank the observations from smallest to largest
- If ranks are tied, we assign the average rank to those values
- We then find the sum of ranks for each of the two original samples, denoted  $W_1$  and  $W_2$ , and then let  $W = \min(W_1, W_2)$
- Under  $H_0$ , the underlying populations have the same median, so we would expect ranks to be randomly distributed between the two groups

# Wilcoxon Rank-Sum Test: Steps

- Combine all data from the two samples and rank the observations from smallest to largest
- If ranks are tied, we assign the average rank to those values
- We then find the sum of ranks for each of the two original samples, denoted  $W_1$  and  $W_2$ , and then let  $W = \min(W_1, W_2)$
- Under  $H_0$ , the underlying populations have the same median, so we would expect ranks to be randomly distributed between the two groups
- Thus, the average ranks for the two samples (i.e.,  $W_1/n_1$  and  $W_2/n_2$ ) should be approximately equal

# Data Table

No Disease		Disease	
Temp	Rank	Temp	Rank
98.1	1	99.3	8
98.5	2	99.4	[ 4.5 4.5
98.6	3	99.4	
98.8	4	99.5	
98.9	5	99.5	
99.0	6	99.6	
99.2	7	99.7	
99.5		99.7	
99.6		100.0	
99.7		100.0	
100.5		100.1	
101.0		100.1	
		100.1	
		101.1	
		101.9	



# Data Table

No Disease		Disease	
Temp	Rank	Temp	Rank
98.1	1	99.3	8
98.5	2	99.4	9.5
98.6	3	99.4	9.5
98.8	4	99.5	12
98.9	5	99.5	12
99.0	6	99.6	14.5
99.2	7	99.7	17
99.5	12	99.7	17
99.6	14.5	100.0	19.5
99.7	17	100.0	19.5
100.5	24	100.1	22
101.0	25	100.1	22
		100.1	22
		101.1	26
		101.9	27

$w_1 =$

$w_2 =$

# Wilcoxon Rank-Sum Test: Steps

# Wilcoxon Rank-Sum Test: Steps

- Evaluate the null hypothesis using the test statistic  $z_W = \frac{W - \mu_W}{\sigma_W}$

# Wilcoxon Rank-Sum Test: Steps

- Evaluate the null hypothesis using the test statistic  $z_W = \frac{W - \mu_W}{\sigma_W}$
- Let  $n_1$  be the number of observations in the sample with the smaller sum of ranks

# Wilcoxon Rank-Sum Test: Steps

- Evaluate the null hypothesis using the test statistic  $z_W = \frac{W - \mu_W}{\sigma_W}$
- Let  $n_1$  be the number of observations in the sample with the smaller sum of ranks
- Let  $n_2$  be the number of observations in the sample with the larger sum of ranks

# Wilcoxon Rank-Sum Test: Steps

- Evaluate the null hypothesis using the test statistic  $z_W = \frac{W - \mu_W}{\sigma_W}$
- Let  $n_1$  be the number of observations in the sample with the smaller sum of ranks
- Let  $n_2$  be the number of observations in the sample with the larger sum of ranks
- Then,

# Wilcoxon Rank-Sum Test: Steps

$$W - \binom{n_1}{2}$$

- Evaluate the null hypothesis using the test statistic  $z_W = \frac{W - \mu_W}{\sigma_W}$
- Let  $n_1$  be the number of observations in the sample with the smaller sum of ranks
- Let  $n_2$  be the number of observations in the sample with the larger sum of ranks

• Then,

$$\mu_W = \frac{n_1(n_1 + n_2 + 1)}{2} \text{ and } \sigma_W = \sqrt{\frac{n_1 n_2 (n_1 + n_2 + 1)}{12}}$$

$$\mu = n_1 \left( \frac{n_1 + n_2}{2} \right)$$

avg rank

# Wilcoxon Rank-Sum Test: Steps



- Evaluate the null hypothesis using the test statistic  $z_W = \frac{W - \mu_W}{\sigma_W}$
- Let  $n_1$  be the number of observations in the sample with the smaller sum of ranks
- Let  $n_2$  be the number of observations in the sample with the larger sum of ranks
- Then,

- $\mu_W = \frac{n_1(n_1 + n_2 + 1)}{2}$  and  $\sigma_W = \sqrt{\frac{n_1 n_2 (n_1 + n_2 + 1)}{12}}$

- $z_W \sim N(0,1)$  when  $n_1$  and  $n_2$  are large enough ( $n_1, n_2 > 10$ )



# Histograms of Samples



# Wilcoxon Rank-Sum Test

# Wilcoxon Rank-Sum Test

- Based on the histograms, the two samples do not appear to be coming from normally distributed populations, so we want to use the Wilcoxon rank sum test

# Wilcoxon Rank-Sum Test

- Based on the histograms, the two samples do not appear to be coming from normally distributed populations, so we want to use the Wilcoxon rank sum test
- Wilcoxon rank sum test is appropriate since both populations have similar shapes

# Wilcoxon Rank-Sum Test

- Based on the histograms, the two samples do not appear to be coming from normally distributed populations, so we want to use the Wilcoxon rank sum test
- Wilcoxon rank sum test is appropriate since both populations have similar shapes
- The sum of ranks for sample 1 is 120.5

# Wilcoxon Rank-Sum Test

- Based on the histograms, the two samples do not appear to be coming from normally distributed populations, so we want to use the Wilcoxon rank sum test
- Wilcoxon rank sum test is appropriate since both populations have similar shapes
- The sum of ranks for sample 1 is 120.5  $= W_1$
- The sum of ranks for sample 2 is 257.5  $= W_2$

$$W = \min(W_1, W_2) = \underline{120.5}$$

# Wilcoxon Rank-Sum Test

- Based on the histograms, the two samples do not appear to be coming from normally distributed populations, so we want to use the Wilcoxon rank sum test
- Wilcoxon rank sum test is appropriate since both populations have similar shapes
- The sum of ranks for sample 1 is 120.5
- The sum of ranks for sample 2 is 257.5
- Thus,  $W = 120.5$ ,  $n_1 = 12$ , and  $n_2 = 15$

# Wilcoxon Rank-Sum Test



# Wilcoxon Rank-Sum Test

- Given  $W = 120.5$ ,  $n_1 = 12$ , and  $n_2 = 15$ , we then have

# Wilcoxon Rank-Sum Test

- Given  $W = 120.5$ ,  $n_1 = 12$ , and  $n_2 = 15$ , we then have
- $\mu_W = \frac{n_1(n_1 + n_2 + 1)}{2} =$

# Wilcoxon Rank-Sum Test

$$\frac{1}{n_1 + n_2} \cdot 1 + \frac{1}{n_1 + n_2} \cdot 2 + \dots + \frac{1}{n_1 + n_2} \cdot (n_1 + n_2)$$

- Given  $W = 120.5$ ,  $n_1 = 12$ , and  $n_2 = 15$ , we then have

- $\mu_W = \frac{n_1(n_1 + n_2 + 1)}{2} =$

- $\sigma_W = \sqrt{\frac{n_1 n_2 (n_1 + n_2 + 1)}{12}} =$

	A	B
$x_1$	21	$y_1$
$\vdots$		$\vdots$
$x_{n_1}$		$y_{n_2}$

Ranks

1  
 $\vdots$   
 $n_1 + n_2$

$$\frac{(n_1 + n_2 + 1)}{2}$$

# Wilcoxon Rank-Sum Test

- Given  $W = 120.5$ ,  $n_1 = 12$ , and  $n_2 = 15$ , we then have

- $$\mu_W = \frac{n_1(n_1 + n_2 + 1)}{2} = \frac{12(12+15+1)}{2} = 168$$

- $$\sigma_W = \sqrt{\frac{n_1 n_2 (n_1 + n_2 + 1)}{12}} = 20.49$$

- Thus, we have 
$$z_W = \frac{W - \mu_W}{\sigma_W} = \frac{120.5 - 168}{20.49} = -2.318$$

# Wilcoxon Rank-Sum Test

# Wilcoxon Rank-Sum Test

- Calculating the p-value:

# Wilcoxon Rank-Sum Test

- Calculating the p-value:

$$p = \Pr(Z < -2.318) = \text{pnorm}(-2.318) = 0.01$$

- Conclusion:

$$p < \alpha \rightarrow \text{Reject } H_0.$$

# Wilcoxon Rank-Sum Test



# Wilcoxon Rank-Sum Test

- If  $n_1$  and  $n_2$  are very small (i.e., either is less than or equal to 10), we cannot use the normal approximation

# Wilcoxon Rank-Sum Test

- If  $n_1$  and  $n_2$  are very small (i.e., either is less than or equal to 10), we cannot use the normal approximation
- When sample sizes are small, we can use the exact distribution to calculate p-values

# Wilcoxon Rank-Sum Test

- If  $n_1$  and  $n_2$  are very small (i.e., either is less than or equal to 10), we cannot use the normal approximation
- When sample sizes are small, we can use the exact distribution to calculate p-values
- In R, we use `pwilcox( $W_{\text{obs}}$ ,  $n_1$ ,  $n_2$ )`


# Wilcoxon Rank-Sum Test

- If  $n_1$  and  $n_2$  are very small (i.e., either is less than or equal to 10), we cannot use the normal approximation
- When sample sizes are small, we can use the exact distribution to calculate p-values
- In R, we use `pwilcox( $W_{\text{obs}}$ ,  $n_1$ ,  $n_2$ )`
  - In this case,  $W_{\text{obs}} = \underline{W} - \frac{n_1(n_1 + 1)}{2}$

# Wilcoxon Rank-Sum Test

$$W = \min(W_1, W_2)$$

$$W^* = W - \frac{n_1(n_1+1)}{2}$$

- If  $n_1$  and  $n_2$  are very small (i.e., either is less than or equal to 10), we cannot use the normal approximation
- When sample sizes are small, we can use the exact distribution to calculate p-values
- In R, we use `pwilcox( $W_{\text{obs}}$ ,  $n_1$ ,  $n_2$ )`
  - In this case,  $W_{\text{obs}} = W - \frac{n_1(n_1+1)}{2}$  
  - `pwilcox(120.5-78, 12, 15) = 0.0093`

# Wilcoxon Rank-Sum Test: R Code

$F \rightarrow N(0,1)$   
 $T \rightarrow \text{Wilcox distr.}$

```
> wilcox.test(sample1, sample2, exact=F, correct=F, alt="less")
```

Wilcoxon rank sum test

data: sample1 and sample2

W = 42.5, p-value = 0.01009

alternative hypothesis: true location shift is less than 0

# Nonparametric Methods: Pros and Cons

# Nonparametric Methods: Pros and Cons

- Advantages:



# Nonparametric Methods: Pros and Cons

- Advantages:
  - Do not impose restrictive assumptions

# Nonparametric Methods: Pros and Cons

- Advantages:
  - Do not impose restrictive assumptions
  - Do not require normally distributed populations

# Nonparametric Methods: Pros and Cons

- Advantages:
  - Do not impose restrictive assumptions
  - Do not require normally distributed populations
  - Are sometimes easier to compute by hand

# Nonparametric Methods: Pros and Cons

- Advantages:
  - Do not impose restrictive assumptions
  - Do not require normally distributed populations
  - Are sometimes easier to compute by hand
  - Ranks are less sensitive to measurement error

# Nonparametric Methods: Pros and Cons

- Advantages:
  - Do not impose restrictive assumptions
  - Do not require normally distributed populations
  - Are sometimes easier to compute by hand
  - Ranks are less sensitive to measurement error
  - Permits the use of ordinal data

# Nonparametric Methods: Pros and Cons

- Advantages:

- Do not impose restrictive assumptions
- Do not require normally distributed populations
- Are sometimes easier to compute by hand
- Ranks are less sensitive to measurement error
- Permits the use of ordinal data

- Disadvantages:

# Nonparametric Methods: Pros and Cons

- Advantages:
  - Do not impose restrictive assumptions
  - Do not require normally distributed populations
  - Are sometimes easier to compute by hand
  - Ranks are less sensitive to measurement error
  - Permits the use of ordinal data
- Disadvantages:
  - If a parametric test can be used, it is more powerful than its nonparametric counterpart

# Nonparametric Methods: Pros and Cons

- Advantages:
  - Do not impose restrictive assumptions
  - Do not require normally distributed populations
  - Are sometimes easier to compute by hand
  - Ranks are less sensitive to measurement error
  - Permits the use of ordinal data
- Disadvantages:
  - If a parametric test can be used, it is more powerful than its nonparametric counterpart
  - Hypotheses tend to be less specific for nonparametric tests



# Nonparametric Methods: Pros and Cons

- Advantages:
  - Do not impose restrictive assumptions
  - Do not require normally distributed populations
  - Are sometimes easier to compute by hand
  - Ranks are less sensitive to measurement error
  - Permits the use of ordinal data
- Disadvantages:
  - If a parametric test can be used, it is more powerful than its nonparametric counterpart
  - Hypotheses tend to be less specific for nonparametric tests
  - Variances are typically overestimated

# Nonparametric Methods: Summary

# Nonparametric Methods: Summary

- Sometimes we want to run tests on variables but do not know their distributions

# Nonparametric Methods: Summary

- Sometimes we want to run tests on variables but do not know their distributions
- Nonparametric tests are a flexible but sometimes underpowered way of doing so

# Nonparametric Methods: Summary

- Sometimes we want to run tests on variables but do not know their distributions
- Nonparametric tests are a flexible but sometimes underpowered way of doing so
- Nonparametric analog to the one-sample or paired t-test: Wilcoxon Signed-Rank Test

# Nonparametric Methods: Summary

- Sometimes we want to run tests on variables but do not know their distributions
- Nonparametric tests are a flexible but sometimes underpowered way of doing so
- Nonparametric analog to the one-sample or paired t-test: Wilcoxon Signed-Rank Test
- Nonparametric analog to the two-sample t-test: Wilcoxon Rank-Sum Test (Mann-Whitney U Test)

# Review Sessions Next Week

# Review Sessions Next Week

- TA review sessions (Wegmans 1201):



# Review Sessions Next Week

- TA review sessions (Wegmans 1201):
  - Tuesday, November 1, from 2 - 3 pm (Lucinda's normal OH)

# Review Sessions Next Week

- TA review sessions (Wegmans 1201):
  - Tuesday, November 1, from 2 - 3 pm (Lucinda's normal OH)
  - Wednesday, November 2, from 6 - 8 pm

# Review Sessions Next Week

- TA review sessions (Wegmans 1201):
  - Tuesday, November 1, from 2 - 3 pm (Lucinda's normal OH)
  - Wednesday, November 2, from 6 - 8 pm
- Instructor: Tuesday and Thursday during class

# Review Sessions Next Week

- TA review sessions (Wegmans 1201):
  - Tuesday, November 1, from 2 - 3 pm (Lucinda's normal OH)
  - Wednesday, November 2, from 6 - 8 pm
- Instructor: Tuesday and Thursday during class
  - Submit specific requests via Google Form