#### Chapter 1: Introduction to Statistical Data

DSCC 462
Computational Introduction to Statistics

Anson Kahng Fall 2022

# Plan for Today

## Plan for Today

Cover the basics of statistics

## Plan for Today

- Cover the basics of statistics
- Introduce types of data

• Statistics: Collection, organization, analysis, and interpretation of data

- Statistics: Collection, organization, analysis, and interpretation of data
- Descriptive statistics: methods for organizing and summarizing data

- Statistics: Collection, organization, analysis, and interpretation of data
- Descriptive statistics: methods for organizing and summarizing data
- Statistical inference: methods for inferring properties of a *population* based on a *sample*

• Sample: Subset of a group of interest we have data for

- Sample: Subset of a group of interest we have data for
- Population: The entire group of interest

- Sample: Subset of a group of interest we have data for
- Population: The entire group of interest
- Using the sample, we can:

- Sample: Subset of a group of interest we have data for
- Population: The entire group of interest
- Using the sample, we can:
  - Create estimates and plots, perform inference, summarize results

- Sample: Subset of a group of interest we have data for
- Population: The entire group of interest
- Using the sample, we can:
  - Create estimates and plots, perform inference, summarize results
- Goal: Have sample be representative of population so that we can generalize results

- Sample: Subset of a group of interest we have data for
- Population: The entire group of interest
- Using the sample, we can:
  - Create estimates and plots, perform inference, summarize results
- Goal: Have sample be representative of population so that we can generalize results
- Notation:

- Sample: Subset of a group of interest we have data for
- Population: The entire group of interest
- Using the sample, we can:
  - Create estimates and plots, perform inference, summarize results
- Goal: Have sample be representative of population so that we can generalize results
- Notation:
  - Population: parameter ( $\mu$ ,  $\sigma^2$ , etc.)

- Sample: Subset of a group of interest we have data for
- Population: The entire group of interest
- Using the sample, we can:
  - Create estimates and plots, perform inference, summarize results
- Goal: Have sample be representative of population so that we can generalize results
- Notation:
  - Population: parameter ( $\mu$ ,  $\sigma^2$ , etc.)
  - Sample: estimate ( $\bar{x}$ ,  $s^2$ , etc.)

A study was conducted on 423 male children between the age of 5 and 15.
 From this study, it was concluded that advanced maternal age is associated with higher risk of a male child having autism

ALL

- A study was conducted on 423 male children between the age of 5 and 15. From this study, it was concluded that advanced maternal age is associated with higher risk of a male child having autism
- Population:

A study was conducted on 423 male children between the age of 5 and 15.
 From this study, it was concluded that advanced maternal age is associated with higher risk of a male child having autism

• Population: male children aged 5-15

• Sample: 423 children in the study

• Data are pieces of information about **subjects** that are organized into **variables** 

- Data are pieces of information about **subjects** that are organized into **variables**
- **Subjects** are the particular people or objects we are interested in studying (i.e., the people in our sample)

- Data are pieces of information about subjects that are organized into variables
- **Subjects** are the particular people or objects we are interested in studying (i.e., the people in our sample)
- **Variables** are the characteristics we are interested in measuring for each subject (e.g., weight, height, eye color)

- Data are pieces of information about subjects that are organized into variables
- **Subjects** are the particular people or objects we are interested in studying (i.e., the people in our sample)
- **Variables** are the characteristics we are interested in measuring for each subject (e.g., weight, height, eye color)
- Different types of summaries and analyses are appropriate for different types of data

### Example Dataset: Emergency Room Patients

	Age	Weight (kg)	Eye Color	Smoker?	Pain Rating
Patient 1	25	82	Brown	Yes	Medium
Patient 2	42	60	Green	No	High
Patient 3	31	105	Blue	No	Low

### Example Dataset: Emergency Room Patients

	Age	Weight (kg)	Eye Color	Smoker?	Pain Rating
Patient 1	25	82	Brown	Yes	Medium
Patient 2	42	60	Green	No	High
Patient 3	31	105	Blue	No	Low

• Subjects: Patient 1, Patient 2, Patient 3

#### Example Dataset: Emergency Room Patients



- Subjects: Patient 1, Patient 2, Patient 3
- Variables: Age, Weight, Eye Color, Smoker, Pain Rating

• Categorical (qualitative) data: Data that are measured on a scale consisting of sets of groups or categories

- Categorical (qualitative) data: Data that are measured on a scale consisting of sets of groups or categories
  - Place subjects in one of the categories

- Categorical (qualitative) data: Data that are measured on a scale consisting of sets of groups or categories
  - Place subjects in one of the categories
  - Usually, care about the count / proportion in each category

- Categorical (qualitative) data: Data that are measured on a scale consisting of sets of groups or categories
  - Place subjects in one of the categories
  - Usually, care about the count / proportion in each category
- Examples:

- Categorical (qualitative) data: Data that are measured on a scale consisting of sets of groups or categories
  - Place subjects in one of the categories
  - Usually, care about the count / proportion in each category
- Examples:
  - Nominal variables, ordinal variables, discrete interval variables with few values, continuous variables that have been grouped into a small number of categories

# Types of Data

• Numerical (quantitative) data: Data that are counts or measured on a numeric scale

# Types of Data

- Numerical (quantitative) data: Data that are counts or measured on a numeric scale
  - Discrete if measurements are integers

# Types of Data

- **Numerical (quantitative) data**: Data that are counts or measured on a numeric scale
  - Discrete if measurements are integers
  - Continuous if measurements can take any value within a range

Categorical:

- Categorical:
  - Smoking status (yes/no)

#### Categorical:

- Smoking status (yes/no)
- Class year (first-year/sophomore/junior/senior/graduate student)

#### Categorical:

- Smoking status (yes/no)
- Class year (first-year/sophomore/junior/senior/graduate student)
- Hair color (black, brown, blonde, red, other)

#### Categorical:

- Smoking status (yes/no)
- Class year (first-year/sophomore/junior/senior/graduate student)
- Hair color (black, brown, blonde, red, other)

#### • Quantitative:

#### Categorical:

- Smoking status (yes/no)
- Class year (first-year/sophomore/junior/senior/graduate student)
- Hair color (black, brown, blonde, red, other)

#### • Quantitative:

Age

#### Categorical:

- Smoking status (yes/no)
- Class year (first-year/sophomore/junior/senior/graduate student)
- Hair color (black, brown, blonde, red, other)

#### Quantitative:

- Age
- Steps per day

#### • Categorical:

- Smoking status (yes/no)
- Class year (first-year/sophomore/junior/senior/graduate student)
- Hair color (black, brown, blonde, red, other)

#### • Quantitative:

- Age
- Steps per day
- GPA

#### Categorical:

- Smoking status (yes/no)
- Class year (first-year/sophomore/junior/senior/graduate student)
- Hair color (black, brown, blonde, red, other)

#### • Quantitative:

- Age
- Steps per day
- GPA
- Number of siblings

#### Nominal Data:

• Unordered categories or classes

- Unordered categories or classes
- Order of the categories is irrelevant

- Unordered categories or classes
- Order of the categories is irrelevant
- Examples:

- Unordered categories or classes
- Order of the categories is irrelevant
- Examples:
  - Department: philosophy, data science, statistics, linguistics, art history

- Unordered categories or classes
- Order of the categories is irrelevant
- Examples:
  - Department: philosophy, data science, statistics, linguistics, art history
  - Hair color: black, brown, blonde, red, other

#### Ordinal Data:

• Ordered categories or classes ("natural ordering")

- Ordered categories or classes ("natural ordering")
- Distances between categories are unknown

- Ordered categories or classes ("natural ordering")
- Distances between categories are unknown
- Care about the ordering itself, not the magnitude

- Ordered categories or classes ("natural ordering")
- Distances between categories are unknown
- Care about the ordering itself, not the magnitude
- Examples:

- Ordered categories or classes ("natural ordering")
- Distances between categories are unknown
- Care about the ordering itself, not the magnitude
- Examples:
  - Pain scale: low, medium, high

# Categorical Data strong disgree • Ordinal Data: -3, 2, -1, 2, 3

- Ordered categories or classes ("natural ordering")
- Distances between categories are unknown
- Care about the ordering itself, not the magnitude
- Examples:
  - Pain scale: low, medium, high
  - Course evaluations: unsatisfactory, neutral, satisfactory, excellent

#### Ranked Data:

 Arrange a group of observations from highest to lowest (or reversed) according to their magnitude

- Arrange a group of observations from highest to lowest (or reversed) according to their magnitude
- Assign ranks corresponding to each observation's place in the sequence

- Arrange a group of observations from highest to lowest (or reversed) according to their magnitude
- Assign ranks corresponding to each observation's place in the sequence
- Example:

- Arrange a group of observations from highest to lowest (or reversed) according to their magnitude
- Assign ranks corresponding to each observation's place in the sequence
- Example:
  - GPAs:  $(93.1, 86.2, 98.5, 89.8) \rightarrow (2, 4, 5)$

## Quantitative Data

### Quantitative Data

Discrete Data:

#### Discrete Data:

• Both order and magnitude are important

- Both order and magnitude are important
- Numbers represent actual values instead of labels

- Both order and magnitude are important
- Numbers represent actual values instead of labels
- Often integers or counts (isolated points on a number line)

- Both order and magnitude are important
- Numbers represent actual values instead of labels
- Often integers or counts (isolated points on a number line)
- Examples:

- Both order and magnitude are important
- Numbers represent actual values instead of labels
- Often integers or counts (isolated points on a number line)
- Examples:
  - Number of steps walked in a day

- Both order and magnitude are important
- Numbers represent actual values instead of labels
- Often integers or counts (isolated points on a number line)
- Examples:
  - Number of steps walked in a day
  - Number of prospective students who come to Admissions this week

#### Continuous Data:

• Data can take any value within a given interval (entire interval on a number line)

- Data can take any value within a given interval (entire interval on a number line)
- Distance between measurements is meaningful (both order and magnitude matter)

- Data can take any value within a given interval (entire interval on a number line)
- Distance between measurements is meaningful (both order and magnitude matter)
- Examples:

- Data can take any value within a given interval (entire interval on a number line)
- Distance between measurements is meaningful (both order and magnitude matter)
- Examples:
  - Height in cm

- Data can take any value within a given interval (entire interval on a number line)
- Distance between measurements is meaningful (both order and magnitude matter)
- Examples:
  - Height in cm
  - Time in minutes spent on an assignment

• Interval level of measurement:

- Interval level of measurement:
  - Data can be arranged in some order, and the difference between any two data values is meaningful

- Interval level of measurement:
  - Data can be arranged in some order, and the difference between any two data values is meaningful
  - There is no natural zero or starting point

- Interval level of measurement:
  - Data can be arranged in some order, and the difference between any two data values is meaningful
  - There is no natural zero or starting point
  - Ex: Years (1000, 1359, 2009, 2017)

- Interval level of measurement:
  - Data can be arranged in some order, and the difference between any two data values is meaningful
  - There is no natural zero or starting point
  - Ex: Years (1000, 1359, 2009, 2017)
- Ratio level of measurement:


- Interval level of measurement:
  - Data can be arranged in some order, and the difference between any two data values is meaningful
  - There is no natural zero or starting point
  - Ex: Years (1000, 1359, 2009, 2017)
- Ratio level of measurement:
  - Data can be arranged in some order, and the difference or ratio between any two data values is meaningful

- Interval level of measurement:
  - Data can be arranged in some order, and the difference between any two data values is meaningful
  - There is no natural zero or starting point
  - Ex: Years (1000, 1359, 2009, 2017)
- Ratio level of measurement:
  - Data can be arranged in some order, and the difference or ratio between any two data values is meaningful
  - There is a natural zero or starting point

- Interval level of measurement:
  - Data can be arranged in some order, and the difference between any two data values is meaningful
  - There is no natural zero or starting point
  - Ex: Years (1000, 1359, 2009, 2017)
- Ratio level of measurement:
  - Data can be arranged in some order, and the difference or ratio between any two data values is meaningful
  - There is a natural zero or starting point
  - Ex: Price of textbooks (\$0 represents no cost; \$100 costs twice as much as \$50)

## Four Levels of Measurement





Disc Cont

Categorical

Quantitative