

# Chapter 10: Inference on Proportions

DSCC 462  
Computational Introduction to Statistics

Anson Kahng  
Fall 2022

# Inference on Proportions

# Inference on Proportions

- So far, we have considered inference for when we have continuous data

# Inference on Proportions

- So far, we have considered inference for when we have continuous data
- We can also extend inferential methods to cover count data

# Inference on Proportions

- So far, we have considered inference for when we have continuous data
- We can also extend inferential methods to cover count data
- In particular, we are often interested in the proportion of times a dichotomous (i.e., yes/no) event occurs

# Sampling Distribution of a Proportion

# Sampling Distribution of a Proportion

- Recall that the sample mean is distributed like  $\hat{p} \sim N\left(p, \sqrt{\frac{(p(1-p)}{n}}\right)$ , given that  $\underline{np \geq 5}$  and  $\underline{n(1-p) \geq 5}$

$$\begin{aligned} p &= \frac{\text{Binom}(n, p)}{n} && \xrightarrow{\text{CLT}} && N\left(\frac{np}{n}, \sqrt{\frac{np(1-p)}{n}}\right) \\ &&& \rightarrow && N\left(p, \sqrt{\frac{p(1-p)}{n}}\right) \end{aligned}$$

# Sampling Distribution of a Proportion

- Recall that the sample mean is distributed like  $\hat{p} \sim N\left(p, \sqrt{\frac{(p(1-p)}{n}}\right)$ , given that  $np \geq 5$  and  $n(1-p) \geq 5$
- Thus,  $Z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$  has a standard normal distribution

# Confidence Intervals for Proportions

# Confidence Intervals for Proportions

- Confidence intervals for population proportions follows the same procedure as what we used for population means

# Confidence Intervals for Proportions

- Confidence intervals for population proportions follows the same procedure as what we used for population means
- Draw a sample of size  $n$  and compute  $\hat{p} = \frac{x}{n}$

# Confidence Intervals for Proportions

- Confidence intervals for population proportions follows the same procedure as what we used for population means
- Draw a sample of size  $n$  and compute  $\hat{p} = \frac{x}{n}$
- $\hat{p}$  is a point estimate of population proportion  $p$

$$\hat{p} \sim N(p, \sigma^2)$$

# Confidence Intervals for Proportions

- Confidence intervals for population proportions follows the same procedure as what we used for population means
- Draw a sample of size  $n$  and compute  $\hat{p} = \frac{x}{n}$
- $\hat{p}$  is a point estimate of population proportion  $p$
- We know from above that  $Z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$  is a standard normal random variable, given that  $n$  is sufficiently large (i.e.,  $np \geq 5$  and  $n(1 - p) \geq 5$ )

# Confidence Intervals for Proportions

# Confidence Intervals for Proportions

- For a standard normal distribution, 95% of possible outcomes lie between  $\text{qnorm}(0.025) = -1.96$  and  $\text{qnorm}(0.975) = 1.96$

# Confidence Intervals for Proportions

- For a standard normal distribution, 95% of possible outcomes lie between  $\text{qnorm}(0.025) = -1.96$  and  $\text{qnorm}(0.975) = 1.96$

- Thus,  $\Pr\left(-1.96 \leq \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}} \leq 1.96\right) = 0.95$

$$\Pr\left(-1.96 \sqrt{\frac{p(1-p)}{n}} + \hat{p} \leq p \leq 1.96 \sqrt{\frac{p(1-p)}{n}} + \hat{p}\right) = 0.95$$

# Confidence Intervals for Proportions

- For a standard normal distribution, 95% of possible outcomes lie between  $\text{qnorm}(0.025) = -1.96$  and  $\text{qnorm}(0.975) = 1.96$

- Thus,  $\Pr\left(-1.96 \leq \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}} \leq 1.96\right) = 0.95$
- This can be rearranged to give  
$$\Pr\left(\hat{p} - 1.96\sqrt{\frac{p(1-p)}{n}} \leq p \leq \hat{p} + 1.96\sqrt{\frac{p(1-p)}{n}}\right) = 0.95$$

# Confidence Intervals for Proportions

- For a standard normal distribution, 95% of possible outcomes lie between  $\text{qnorm}(0.025) = -1.96$  and  $\text{qnorm}(0.975) = 1.96$

- Thus,  $\Pr\left(-1.96 \leq \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}} \leq 1.96\right) = 0.95$

- This can be rearranged to give

$$\Pr\left(\hat{p} - 1.96\sqrt{\frac{p(1-p)}{n}} \leq p \leq \hat{p} + 1.96\sqrt{\frac{p(1-p)}{n}}\right) = 0.95$$

- Note that this confidence interval depends on the (unknown) value of  $p$ !

# Confidence Intervals for Proportions

# Confidence Intervals for Proportions

- So how do we estimate  $p$ ? Use  $\hat{p}$ , our sample estimate (Wald)

# Confidence Intervals for Proportions

- So how do we estimate  $p$ ? Use  $\hat{p}$ , our sample estimate (Wald)
- Therefore, our confidence interval calculation becomes

$$\Pr \left( \hat{p} - 1.96 \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \leq p \leq \hat{p} + 1.96 \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \right) = 0.95$$

# Confidence Intervals for Proportions

- So how do we estimate  $p$ ? Use  $\hat{p}$ , our sample estimate (Wald)

- Therefore, our confidence interval calculation becomes

$$\Pr \left( \hat{p} - 1.96 \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \leq p \leq \hat{p} + 1.96 \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \right) = 0.95$$

- In other words, we are 95% confident that the interval

$$\left( \hat{p} - 1.96 \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}, \hat{p} + 1.96 \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \right)$$

contains the true population proportion  $p$

given Wald estimate  $P = \hat{P}$ .

# Confidence Intervals for Proportions

# Confidence Intervals for Proportions

- In general, an approximate two-sided  $(1 - \alpha) \cdot 100\%$  confidence interval for  $p$  is given by  $\left( \hat{p} - z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}, \hat{p} + z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \right)$

# Confidence Intervals for Proportions

- In general, an approximate two-sided  $(1 - \alpha) \cdot 100\%$  confidence interval for  $p$  is given by  $\left( \hat{p} - z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}, \hat{p} + z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \right)$
- A one-sided lower  $(1 - \alpha) \cdot 100\%$  confidence interval for  $p$  is given by  $\left( \hat{p} - z_{\alpha} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}, 1 \right)$

# Confidence Intervals for Proportions

$$\hat{P} = \hat{p}$$

- In general, an approximate two-sided  $(1 - \alpha) \cdot 100\%$  confidence interval for  $p$  is given by  $\left( \hat{p} - z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}, \hat{p} + z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \right)$
- A one-sided lower  $(1 - \alpha) \cdot 100\%$  confidence interval for  $p$  is given by  $\left( \hat{p} - z_{\alpha} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}, 1 \right)$
- A one-sided upper  $(1 - \alpha) \cdot 100\%$  confidence interval for  $p$  is given by  $\left( 0, \hat{p} + z_{\alpha} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \right)$

# Wald vs. Wilson Intervals

# Wald vs. Wilson Intervals

- Recall that we estimate  $p$  using  $\hat{p}$ , our sample estimate (Wald)


# Wald vs. Wilson Intervals

- Recall that we estimate  $p$  using  $\hat{p}$ , our sample estimate (Wald)
- In general, this provides poor coverage when  $\hat{p}$  is close to extremes (0 or 1)

# Wald vs. Wilson Intervals

- Recall that we estimate  $p$  using  $\hat{p}$ , our sample estimate (Wald)
- In general, this provides poor coverage when  $\hat{p}$  is close to extremes (0 or 1)
  - Less than  $(1 - \alpha) \cdot 100\%$  confidence interval

$$\sqrt{\frac{p(1-p)}{n}}$$



# Wald vs. Wilson Intervals

- Recall that we estimate  $p$  using  $\hat{p}$ , our sample estimate (Wald)
- In general, this provides poor coverage when  $\hat{p}$  is close to extremes (0 or 1)
  - Less than  $(1 - \alpha) \cdot 100\%$  confidence interval
- (One) alternative method: Wilson (what `prop.test()` uses in R)

# Wald vs. Wilson Intervals

- Recall that we estimate  $p$  using  $\hat{p}$ , our sample estimate (Wald)
- In general, this provides poor coverage when  $\hat{p}$  is close to extremes (0 or 1)
  - Less than  $(1 - \alpha) \cdot 100\%$  confidence interval
- (One) alternative method: Wilson (what `prop.test()` uses in R)
  - Solve for  $p$  in terms of  $Z, \hat{p}$  from the approximation  $Z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$

# Wald vs. Wilson Intervals

- Recall that we estimate  $p$  using  $\hat{p}$ , our sample estimate (Wald)
- In general, this provides poor coverage when  $\hat{p}$  is close to extremes (0 or 1)
  - Less than  $(1 - \alpha) \cdot 100\%$  confidence interval
- (One) alternative method: Wilson (what `prop.test()` uses in R)
  - Solve for  $p$  in terms of  $Z, \hat{p}$  from the approximation  $Z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$
  - Get an estimate of  $p$  that is a weighted average of  $\hat{p}$  and  $\frac{1}{2}$ , where the weight on  $\hat{p}$  increases with  $n$

# ~~Wald~~ vs. Wilson Intervals R

- Recall that we estimate  $p$  using  $\hat{p}$ , our sample estimate (Wald)
- In general, this provides poor coverage when  $\hat{p}$  is close to extremes (0 or 1)
  - Less than  $(1 - \alpha) \cdot 100\%$  confidence interval
- (One) alternative method: Wilson (what `prop.test()` uses in R)
  - Solve for  $p$  in terms of  $Z, \hat{p}$  from the approximation  $Z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$
  - Get an estimate of  $p$  that is a weighted average of  $\hat{p}$  and  $\frac{1}{2}$ , where the weight on  $\hat{p}$  increases with  $n$
  - Better coverage!

MN

# Confidence Intervals for Proportions: Example

# Confidence Intervals for Proportions: Example

- Setup: We are interested in determining what proportion of a population is right-handed. Suppose we have a sample of  $n = 62$  subjects and 53 of these subjects are right-handed. Find a 95% confidence interval for the population proportion of right-handed people in the population

# Confidence Intervals for Proportions: Example

- Setup: We are interested in determining what proportion of a population is right-handed. Suppose we have a sample of  $n = 62$  subjects and 53 of these subjects are right-handed. Find a 95% confidence interval for the population proportion of right-handed people in the population
- Find  $\hat{p}$  (our estimate of  $p$ ):

# Confidence Intervals for Proportions: Example

- Setup: We are interested in determining what proportion of a population is right-handed. Suppose we have a sample of  $n = 62$  subjects and 53 of these subjects are right-handed. Find a 95% confidence interval for the population proportion of right-handed people in the population
- Find  $\hat{p}$  (our estimate of  $p$ ):
- Check normality assumptions:

# Confidence Intervals for Proportions: Example

- Setup: We are interested in determining what proportion of a population is right-handed.  
Suppose we have a sample of  $n = 62$  subjects and 53 of these subjects are right-handed. Find a 95% confidence interval for the population proportion of right-handed people in the population

- Find  $\hat{p}$  (our estimate of  $p$ ):

$$\hat{p} = \frac{53}{62} = 0.855$$

- Check normality assumptions:

$$n\hat{p} = 53$$

$$n(1-\hat{p}) = 9$$

$\alpha = 0.05$

- Apply a two-sided 95% confidence interval:

$$\hat{p} \pm z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

"by hand"  $(\underline{0.717}, \underline{0.943})$

prop. test  $\rightarrow (0.75, 0.95)$

# Normal Approximations of Binomial Distributions

# Normal Approximations of Binomial Distributions

- Note that the true distribution for proportions is a binomial distribution (number of “successes” out of a certain number of trials)

# Normal Approximations of Binomial Distributions

- Note that the true distribution for proportions is a binomial distribution (number of “successes” out of a certain number of trials)
- However, confidence intervals are based on the normal distribution

# Normal Approximations of Binomial Distributions

- Note that the true distribution for proportions is a binomial distribution (number of “successes” out of a certain number of trials)
- However, confidence intervals are based on the normal distribution
- We are using the normal distribution as an approximation for a binomial distribution

# Normal Approximations of Binomial Distributions

- Note that the true distribution for proportions is a binomial distribution (number of “successes” out of a certain number of trials)
- However, confidence intervals are based on the normal distribution
- We are using the normal distribution as an approximation for a binomial distribution
- Normal approximation (Wilson) confidence intervals can be calculated in R using `prop.test(x, n)`

# Normal Approximations of Binomial Distributions

- Note that the true distribution for proportions is a binomial distribution (number of “successes” out of a certain number of trials)
- However, confidence intervals are based on the normal distribution
- We are using the normal distribution as an approximation for a binomial distribution
- Normal approximation (Wilson) confidence intervals can be calculated in R using `prop.test(x, n)`
- Exact binomial (Clopper-Pearson) confidence intervals can be calculated in R using `binom.test(x, n)`

# Sample Size Estimation

# Sample Size Estimation

- For confidence intervals on proportions, we have that the margin of error is

$$m = z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \text{ (half the length of the confidence interval)}$$

# Sample Size Estimation

- For confidence intervals on proportions, we have that the margin of error is
$$m = z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \quad (\text{half the length of the confidence interval})$$
- Just like before, if we want a certain margin of error at the same confidence level, we can determine the number of subjects ( $n$ ) needed to get the desired results

# Sample Size Estimation

- For confidence intervals on proportions, we have that the margin of error is

$$m = z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \quad (\text{half the length of the confidence interval})$$

- Just like before, if we want a certain margin of error at the same confidence level, we can determine the number of subjects ( $n$ ) needed to get the desired results

- Thus,  $n = \left\lceil \frac{z_{\alpha/2}^2 p(1 - p)}{m^2} \right\rceil$

# Sample Size Estimation

- For confidence intervals on proportions, we have that the margin of error is
$$m = z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \quad (\text{half the length of the confidence interval})$$
- Just like before, if we want a certain margin of error at the same confidence level, we can determine the number of subjects ( $n$ ) needed to get the desired results
- Thus, 
$$n = \left\lceil \frac{z_{\alpha/2}^2 p(1 - p)}{m^2} \right\rceil$$
- If we can estimate  $p$  based on previous studies or information, use that


# Sample Size Estimation

- For confidence intervals on proportions, we have that the margin of error is

$$m = z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \quad (\text{half the length of the confidence interval})$$

- Just like before, if we want a certain margin of error at the same confidence level, we can determine the number of subjects ( $n$ ) needed to get the desired results

- Thus,  $n = \left\lceil \frac{z_{\alpha/2}^2 p(1 - p)}{m^2} \right\rceil$



- If we can estimate  $p$  based on previous studies or information, use that
- Otherwise, use  $p = 0.5$  to get the most conservative estimate of the standard error (overestimate of the number of subjects needed)

# Sample Size Estimation: Example

# Sample Size Estimation: Example

- Setup: We want to determine what proportion of college students have an iPhone within a margin of error of 8 percentage points with 95% confidence

$$m = 0.08$$

# Sample Size Estimation: Example

- Setup: We want to determine what proportion of college students have an iPhone within a margin of error of 8 percentage points with 95% confidence
- Q1: How large of a sample should you take?

$$n = \left\lceil \frac{z_{\alpha/2}^2 \cdot p(1-p)}{m^2} \right\rceil$$

Annotations:

- $m = 0.08$
- $p = 0.5$
- $z_{\alpha/2} = 1.96$
- $q_{90\%}(0.95) = 0.5$
- $n = 151$

# Sample Size Estimation: Example

- Setup: We want to determine what proportion of college students have an iPhone within a margin of error of 8 percentage points with 95% confidence
- Q1: How large of a sample should you take?
- Q2: A national study determined that 38% of all Americans own iPhones. Now, how large of a sample should you take?



$$\begin{aligned}m &= 0.08 \\P &= 0.38 \\1 - P &= 0.62\end{aligned}$$

$$n = \left\lceil \frac{1.96^2 \cdot 0.38 \cdot 0.62}{0.08^2} \right\rceil = 142$$

40

# Hypothesis Testing for Proportions

# Hypothesis Testing for Proportions

- Just as we used hypothesis tests to see if a population mean was equal to some hypothesized value, we can also test whether a population proportion is equal to some value

# Hypothesis Testing for Proportions

- Just as we used hypothesis tests to see if a population mean was equal to some hypothesized value, we can also test whether a population proportion is equal to some value
- Consider a two-tailed test at the  $\alpha = 0.05$  significance level

# Hypothesis Testing for Proportions

- Just as we used hypothesis tests to see if a population mean was equal to some hypothesized value, we can also test whether a population proportion is equal to some value
- Consider a two-tailed test at the  $\alpha = 0.05$  significance level
- $H_0 : p = p_0$  vs.  $H_1 : p \neq p_0$

# Hypothesis Testing for Proportions

- Just as we used hypothesis tests to see if a population mean was equal to some hypothesized value, we can also test whether a population proportion is equal to some value
- Consider a two-tailed test at the  $\alpha = 0.05$  significance level
- $H_0 : p = p_0$  vs.  $H_1 : p \neq p_0$
- Draw a random sample of size  $n$  observations from the underlying population (each observation is a dichotomous yes/no)

$$\hat{P} = \frac{x}{n}$$

# Hypothesis Testing for Proportions

- Just as we used hypothesis tests to see if a population mean was equal to some hypothesized value, we can also test whether a population proportion is equal to some value
- Consider a two-tailed test at the  $\alpha = 0.05$  significance level
- $H_0 : p = p_0$  vs.  $H_1 : p \neq p_0$
- Draw a random sample of size  $n$  observations from the underlying population (each observation is a dichotomous yes/no)

Calculate a z-statistic: 
$$z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1 - p_0)}{n}}}$$

CI : 
$$z = \frac{\hat{p} - p}{\sqrt{\frac{p(1 - p)}{n}}}$$

# Hypothesis Testing for Proportions

# Hypothesis Testing for Proportions

- For sufficiently large  $n$  and when  $H_0$  is true, we can compare  $z$  to a standard normal distribution to calculate the probability of obtaining a proportion as extreme or more extreme than  $\hat{p}$

# Hypothesis Testing for Proportions

- For sufficiently large  $n$  and when  $H_0$  is true, we can compare  $z$  to a standard normal distribution to calculate the probability of obtaining a proportion as extreme or more extreme than  $\hat{p}$
- Calculate p-value in R by `p=2*pnorm(-abs(z))`

# Hypothesis Testing for Proportions

- For sufficiently large  $n$  and when  $H_0$  is true, we can compare  $z$  to a standard normal distribution to calculate the probability of obtaining a proportion as extreme or more extreme than  $\hat{p}$
- Calculate p-value in R by `p=2*pnorm(-abs(z))`
- If  $p \leq 0.05$ , we reject the null hypothesis and conclude that  $p \neq p_0$

# Hypothesis Testing for Proportions

- For sufficiently large  $n$  and when  $H_0$  is true, we can compare  $z$  to a standard normal distribution to calculate the probability of obtaining a proportion as extreme or more extreme than  $\hat{p}$
- Calculate p-value in R by `p=2*pnorm (-abs (z) )`
- If  $p \leq 0.05$ , we reject the null hypothesis and conclude that  $p \neq p_0$
- If  $p > 0.05$ , we fail to reject the null hypothesis and conclude that there is not significant evidence to say that  $p \neq p_0$

# Hypothesis Testing for Proportions: Example

# Hypothesis Testing for Proportions: Example

- Consider the dominant hand example ( $n = 62$ ,  $\hat{p} = 53/62 = 0.855$ )

# Hypothesis Testing for Proportions: Example

- Consider the dominant hand example ( $n = 62$ ,  $\hat{p} = 53/62 = 0.855$ )
- Test at the  $\alpha = 0.05$  level whether the true population proportion of right-handed people is equal to 0.9

# Hypothesis Testing for Proportions: Example

- Consider the dominant hand example ( $n = 62$ ,  $\hat{p} = 53/62 = 0.855$ )
- Test at the  $\alpha = 0.05$  level whether the true population proportion of right-handed people is equal to 0.9
- $H_0 : p = 0.9$  vs.  $H_1 : p \neq 0.9$

# Hypothesis Testing for Proportions: Example

- Consider the dominant hand example ( $n = 62$ ,  $\hat{p} = 53/62 = 0.855$ )
- Test at the  $\alpha = 0.05$  level whether the true population proportion of right-handed people is equal to 0.9
- $H_0 : p = 0.9$  vs.  $H_1 : p \neq 0.9$
- Check normality assumptions based on  $p$ :

# Hypothesis Testing for Proportions: Example

- Consider the dominant hand example ( $n = 62$ ,  $\hat{p} = 53/62 = 0.855$ )
- Test at the  $\alpha = 0.05$  level whether the true population proportion of right-handed people is equal to 0.9
- $H_0 : p = 0.9$  vs.  $H_1 : p \neq 0.9$
- Check normality assumptions based on  $p$ :
- Calculate z-score:

# Hypothesis Testing for Proportions: Example

- Consider the dominant hand example ( $n = 62$ ,  $\hat{p} = 53/62 = \underline{\underline{0.855}}$ )
- Test at the  $\alpha = 0.05$  level whether the true population proportion of right-handed people is equal to 0.9
- $H_0 : p = \underline{\underline{0.9}}$  vs.  $H_1 : p \neq 0.9$
- Check normality assumptions based on  $p$ :  $np = 62 \cdot 0.9 > 5$        $n(1-p) = 62 \cdot 0.1 = 6.2$

$$\mu = p_0, \sigma = \sqrt{\frac{p_0(1-p_0)}{n}}$$

Calculate z-score:

$$z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}} = \frac{0.855 - 0.9}{\sqrt{\frac{0.9 \times 0.1}{62}}} = -1.18$$

- Calculate p-value:



$$2 \cdot p_{norm}(-1.18) = \underline{\underline{0.238}}$$

# Confidence Intervals vs. Hypothesis Tests for Proportions

# Confidence Intervals vs. Hypothesis Tests for Proportions

- When looking at sample means, confidence intervals and hypothesis tests are essentially equivalent

# Confidence Intervals vs. Hypothesis Tests for Proportions

- When looking at sample means, confidence intervals and hypothesis tests are essentially equivalent
- This is no longer the case for proportions!

# Confidence Intervals vs. Hypothesis Tests for Proportions

- When looking at sample means, confidence intervals and hypothesis tests are essentially equivalent
- This is no longer the case for proportions!
  - Intuition: For sample means, there are two parameters of interest ( $\mu$ ,  $\sigma$ ), whereas for proportions,  $p$  determines both the mean and variance

# Confidence Intervals vs. Hypothesis Tests for Proportions

- When looking at sample means, confidence intervals and hypothesis tests are essentially equivalent
- This is no longer the case for proportions!
  - Intuition: For sample means, there are two parameters of interest ( $\mu$ ,  $\sigma$ ), whereas for proportions,  $p$  determines both the mean and variance
- For proportion hypothesis tests, we calculate the standard error based on  $p_0$  as  $\sqrt{\frac{p_0(1 - p_0)}{n}}$  (i.e., our frame of reference is centered at the null hypothesis)

# Confidence Intervals vs. Hypothesis Tests for Proportions

- When looking at sample means, confidence intervals and hypothesis tests are essentially equivalent
- This is no longer the case for proportions!
  - Intuition: For sample means, there are two parameters of interest ( $\mu$ ,  $\sigma$ ), whereas for proportions,  $p$  determines both the mean and variance
- For proportion hypothesis tests, we calculate the standard error based on  $p_0$  as  $\sqrt{\frac{p_0(1 - p_0)}{n}}$  (i.e., our frame of reference is centered at the null hypothesis)
- For proportion (Wald) confidence intervals, we calculate the standard error based on  $\hat{p}$  as  $\sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$  (i.e., our frame of reference is centered at our observed sample proportion)

$$H_0 : p = 0.3 \quad H_1 : p \neq 0.3$$

# Confidence Intervals vs. Hypothesis Tests for Proportions

- When looking at sample means, confidence intervals and hypothesis tests are essentially equivalent
- This is no longer the case for proportions!
  - Intuition: For sample means, there are two parameters of interest ( $\mu, \sigma$ ), whereas for proportions,  $p$  determines both the mean and variance
- For proportion hypothesis tests, we calculate the standard error based on  $p_0$  as  $\sqrt{\frac{p_0(1 - p_0)}{n}}$  (i.e., our frame of reference is centered at the null hypothesis)
- For proportion (Wald) confidence intervals, we calculate the standard error based on  $\hat{p}$  as  $\sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$  (i.e., our frame of reference is centered at our observed sample proportion)
- As with confidence intervals, we can perform an exact test based on the binomial distribution instead of using the normal approximation: `binom.test(x, n, p=p0)`

$$\Pr(\hat{x} | p_0, n)$$

# One-Sided Hypothesis Tests

# One-Sided Hypothesis Tests

- With two-sided hypothesis tests, we were only concerned with whether or not there was a difference from the postulated population proportion

# One-Sided Hypothesis Tests

- With two-sided hypothesis tests, we were only concerned with whether or not there was a difference from the postulated population proportion
  - $H_1 : p \neq p_0$

# One-Sided Hypothesis Tests

- With two-sided hypothesis tests, we were only concerned with whether or not there was a difference from the postulated population proportion
  - $H_1 : p \neq p_0$
- However, we are sometimes interested in deviations only in one direction

# One-Sided Hypothesis Tests

- With two-sided hypothesis tests, we were only concerned with whether or not there was a difference from the postulated population proportion
  - $H_1 : p \neq p_0$
- However, we are sometimes interested in deviations only in one direction
  - $H_1 : p > p_0$

# One-Sided Hypothesis Tests

- With two-sided hypothesis tests, we were only concerned with whether or not there was a difference from the postulated population proportion
  - $H_1 : p \neq p_0$
  - However, we are sometimes interested in deviations only in one direction
    - $H_1 : p > p_0$
    - $H_1 : p < p_0$

# One-Sided Hypothesis Tests

- With two-sided hypothesis tests, we were only concerned with whether or not there was a difference from the postulated population proportion
  - $H_1 : p \neq p_0$
- However, we are sometimes interested in deviations only in one direction
  - $H_1 : p > p_0$
  - $H_1 : p < p_0$
- For two-sided tests, we are concerned with the area in both tails of the distribution

# One-Sided Hypothesis Tests

- With two-sided hypothesis tests, we were only concerned with whether or not there was a difference from the postulated population proportion
  - $H_1 : p \neq p_0$
- However, we are sometimes interested in deviations only in one direction
  - $H_1 : p > p_0$
  - $H_1 : p < p_0$
- For two-sided tests, we are concerned with the area in both tails of the distribution
- For one-sided tests, we are concerned with the area in only one tail of the distribution

# One-Sided Hypothesis Tests

- With two-sided hypothesis tests, we were only concerned with whether or not there was a difference from the postulated population proportion
  - $H_1 : p \neq p_0$
- However, we are sometimes interested in deviations only in one direction
  - $H_1 : p > p_0$
  - $H_1 : p < p_0$
- For two-sided tests, we are concerned with the area in both tails of the distribution
- For one-sided tests, we are concerned with the area in only one tail of the distribution
- Analyses follow directly as they did for one-sided tests for sample means

# Comparison of Two Proportions

# Comparison of Two Proportions

- We can extend hypothesis tests to situations where we compare proportions for two groups

# Comparison of Two Proportions

- We can extend hypothesis tests to situations where we compare proportions for two groups
- Interested in testing whether the proportions from two independent populations are the same

# Comparison of Two Proportions

- We can extend hypothesis tests to situations where we compare proportions for two groups
- Interested in testing whether the proportions from two independent populations are the same
  - $H_0 : p_1 = p_2$  or  $H_0 : p_1 - p_2 = 0$

# Comparison of Two Proportions

- We can extend hypothesis tests to situations where we compare proportions for two groups
- Interested in testing whether the proportions from two independent populations are the same
  - $H_0 : p_1 = p_2$  or  $H_0 : p_1 - p_2 = 0$
  - Our alternative hypothesis is that there is a difference between these groups

# Comparison of Two Proportions

- We can extend hypothesis tests to situations where we compare proportions for two groups
- Interested in testing whether the proportions from two independent populations are the same
  - $H_0 : p_1 = p_2$  or  $H_0 : p_1 - p_2 = 0$
  - Our alternative hypothesis is that there is a difference between these groups
    - $H_1 : p_1 \neq p_2$  or  $H_1 : p_1 - p_2 \neq 0$

# Comparison of Two Proportions

# Comparison of Two Proportions

- We draw a sample of size  $n_1$  from the first population and a sample of size  $n_2$  from the second population

# Comparison of Two Proportions

- We draw a sample of size  $n_1$  from the first population and a sample of size  $n_2$  from the second population
- There are  $x_1$  successes in the first sample and  $x_2$  successes in the second sample

# Comparison of Two Proportions

- We draw a sample of size  $n_1$  from the first population and a sample of size  $n_2$  from the second population
- There are  $x_1$  successes in the first sample and  $x_2$  successes in the second sample
- Sample proportion for each group:

# Comparison of Two Proportions

- We draw a sample of size  $n_1$  from the first population and a sample of size  $n_2$  from the second population
- There are  $x_1$  successes in the first sample and  $x_2$  successes in the second sample
- Sample proportion for each group:
  - $\hat{p}_1 = \frac{x_1}{n_1}$

# Comparison of Two Proportions

- We draw a sample of size  $n_1$  from the first population and a sample of size  $n_2$  from the second population
- There are  $x_1$  successes in the first sample and  $x_2$  successes in the second sample
- Sample proportion for each group:
  - $\hat{p}_1 = \frac{x_1}{n_1}$
  - $\hat{p}_2 = \frac{x_2}{n_2}$

# Comparison of Two Proportions

$$H_0 : p_1 = p_2$$

# Comparison of Two Proportions

- Under the null hypothesis,  $p_1 = p_2 = p$

# Comparison of Two Proportions

- Under the null hypothesis,  $p_1 = p_2 = p$
- Thus, the data from both samples can be combined to estimate this common parameter

# Comparison of Two Proportions

- Under the null hypothesis,  $p_1 = p_2 = p$
- Thus, the data from both samples can be combined to estimate this common parameter

$$\bullet \hat{p} = \frac{x_1 + x_2}{n_1 + n_2} = \frac{n_1 \hat{p}_1 + n_2 \hat{p}_2}{n_1 + n_2}$$

$$\hat{p}_1 = \frac{x_1}{n_1}$$

# Comparison of Two Proportions

- Under the null hypothesis,  $p_1 = p_2 = p$
- Thus, the data from both samples can be combined to estimate this common parameter

$$\bullet \hat{p} = \frac{x_1 + x_2}{n_1 + n_2} = \frac{n_1\hat{p}_1 + n_2\hat{p}_2}{n_1 + n_2}$$

- $\hat{p}$  is the weighted average of the two sample proportions (or total successes over total trials)

# Comparison of Two Proportions

- Under the null hypothesis,  $p_1 = p_2 = p$
- Thus, the data from both samples can be combined to estimate this common parameter
  - $\hat{p} = \frac{x_1 + x_2}{n_1 + n_2} = \frac{n_1\hat{p}_1 + n_2\hat{p}_2}{n_1 + n_2}$
  - $\hat{p}$  is the weighted average of the two sample proportions (or total successes over total trials)
  - The estimator of the standard error of  $\hat{p}_1 - \hat{p}_2$  can now be based on this common  $\hat{p}$

# Comparison of Two Proportions

- Under the null hypothesis,  $p_1 = p_2 = p$
- Thus, the data from both samples can be combined to estimate this common parameter

$$\hat{p} = \frac{x_1 + x_2}{n_1 + n_2} = \frac{n_1 \hat{p}_1 + n_2 \hat{p}_2}{n_1 + n_2}$$

$$\hat{p}_1 - \hat{p}_2 \sim N(p_1 - p_2, \text{ wr: } \left( \frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2} \right))$$

- $\hat{p}$  is the weighted average of the two sample proportions (or total successes over total trials)
- The estimator of the standard error of  $\hat{p}_1 - \hat{p}_2$  can now be based on this common  $\hat{p}$

$$\text{Standard error: } \sqrt{\hat{p}(1-\hat{p}) \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}$$

$$P_1 = P_2 = P$$

$$\hat{p}_1 \sim N(p_1, \sqrt{\frac{p_1(1-p_1)}{n_1}})$$

$$\hat{p}_2 \sim N(p_2, \sqrt{\frac{p_2(1-p_2)}{n_2}})$$

$$\sigma = \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{(p_2(1-p_2))}{n_2}}$$

# Comparison of Two Proportions

- Under the null hypothesis,  $p_1 = p_2 = p$
- Thus, the data from both samples can be combined to estimate this common parameter

- $\hat{p} = \frac{x_1 + x_2}{n_1 + n_2} = \frac{n_1\hat{p}_1 + n_2\hat{p}_2}{n_1 + n_2}$

- $\hat{p}$  is the weighted average of the two sample proportions (or total successes over total trials)
- The estimator of the standard error of  $\hat{p}_1 - \hat{p}_2$  can now be based on this common  $\hat{p}$

- Standard error:  $\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}$

- Similar to the “pooled” estimate for sample means

# Comparison of Two Proportions

$$\hat{p}_1 - \hat{p}_2 \sim N\left(p_1 - p_2, \sqrt{p(1-p)\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}\right)$$

# Comparison of Two Proportions

- Putting these pieces together, we get our z-statistic:

# Comparison of Two Proportions

- Putting these pieces together, we get our z-statistic:

$$z = \frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$$

Wald

$$H_0 : p_1 = p_2$$
$$p_1 \neq p_2$$

$$p(1-p)$$

# Comparison of Two Proportions

- Putting these pieces together, we get our z-statistic:

$$z = \frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$$

- If  $n_1$  and  $n_2$  are sufficiently large, this  $z$  statistic is approximately a standard normal (mean 0, standard deviation 1)

# Comparison of Two Proportions

- Putting these pieces together, we get our z-statistic:

$$z = \frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$$

- If  $n_1$  and  $n_2$  are sufficiently large, this  $z$  statistic is approximately a standard normal (mean 0, standard deviation 1)
- Typically, we want  $n_1\hat{p}_1$ ,  $n_1(1 - \hat{p}_1)$ ,  $n_2\hat{p}_2$ , and  $n_2(1 - \hat{p}_2)$  to all be greater than 5 (this is a conservative standard)

# Comparison of Two Proportions

# Comparison of Two Proportions

- With these conditions satisfied, we compare the value of the  $z$  statistic with the critical value to find a p-value,  $p$

# Comparison of Two Proportions

- With these conditions satisfied, we compare the value of the  $z$  statistic with the critical value to find a p-value,  $p$
- If  $p \leq \alpha$ , we reject the null hypothesis

# Comparison of Two Proportions

- With these conditions satisfied, we compare the value of the  $z$  statistic with the critical value to find a p-value,  $p$
- If  $p \leq \alpha$ , we reject the null hypothesis
- If  $p > \alpha$ , we fail to reject the null hypothesis

# Comparison of Two Proportions

# Comparison of Two Proportions

- Consider our dominant hand example. Suppose we are interested in knowing whether the right-handedness rate is different for Group A and Group B

# Comparison of Two Proportions

- Consider our dominant hand example. Suppose we are interested in knowing whether the right-handedness rate is different for Group A and Group B
- At the  $\alpha = 0.01$  significance level, we will test the following hypotheses:

# Comparison of Two Proportions

- Consider our dominant hand example. Suppose we are interested in knowing whether the right-handedness rate is different for Group A and Group B
- At the  $\alpha = 0.01$  significance level, we will test the following hypotheses:
  - $H_0 : p_A = p_B$  vs.  $H_1 : p_A \neq p_B$

# Comparison of Two Proportions

- Consider our dominant hand example. Suppose we are interested in knowing whether the right-handedness rate is different for Group A and Group B
- At the  $\alpha = 0.01$  significance level, we will test the following hypotheses:
  - $H_0 : p_A = p_B$  vs.  $H_1 : p_A \neq p_B$
  - We take samples of  $n_A = 54$  and  $n_B = 62$

# Comparison of Two Proportions

- Consider our dominant hand example. Suppose we are interested in knowing whether the right-handedness rate is different for Group A and Group B
- At the  $\alpha = 0.01$  significance level, we will test the following hypotheses:
  - $H_0 : p_A = p_B$  vs.  $H_1 : p_A \neq p_B$
  - We take samples of  $n_A = \underline{54}$  and  $n_B = \underline{62}$
  - We observe  $x_A = \underline{48}$  and  $x_B = \underline{60}$  subjects being right handed

# Comparison of Two Proportions

# Comparison of Two Proportions

- Calculate proportions:

$$\hat{P}_A = \frac{48}{54} = 0.889$$

$$\hat{P}_B = \frac{62}{62} = 0.968$$

# Comparison of Two Proportions

- Calculate proportions:

0.889

0.968

- Is this difference too large to be attributed to chance?

# Comparison of Two Proportions

- Calculate proportions:

$$\hat{p}_A = .889$$

$$\hat{p}_B = 0.968$$

- Is this difference too large to be attributed to chance?
- Under  $H_0, p_A = p_B$ , so we can estimate their common value  $p$

$$p = \frac{x_A + x_B}{n_A + n_B} = \frac{48 + 60}{54 + 62} = \frac{108}{116} = 0.931$$

# Comparison of Two Proportions

# Comparison of Two Proportions

- Checking normality assumptions:

# Comparison of Two Proportions

- Checking normality assumptions:
  - $n_A p_A = 48 > 5$

# Comparison of Two Proportions

- Checking normality assumptions:

- $n_A p_A = 48 > 5$

- $n_A(1 - p_A) = 6 > 5$

# Comparison of Two Proportions

- Checking normality assumptions:

- $n_A p_A = 48 > 5$

- $n_A(1 - p_A) = 6 > 5$

- $n_B p_B = 60 > 5$

# Comparison of Two Proportions

- Checking normality assumptions:
  - $n_A p_A = 48 > 5$
  - $n_A(1 - p_A) = 6 > 5$
  - $n_B p_B = 60 > 5$
  - $n_B(1 - p_B) = 2 < 5 \implies$  proceed with caution

# Comparison of Two Proportions

# Comparison of Two Proportions

- Calculate z-statistic:

# Comparison of Two Proportions

$$H_0 : p_A - p_B = 0$$

- Calculate z-statistic:

$$z = \frac{(\hat{p}_A - \hat{p}_B) - (p_A - p_B)}{\sqrt{\hat{p}(1-\hat{p})\left(\frac{1}{n_A} + \frac{1}{n_B}\right)}} = \frac{0.889 - 0.963}{\sqrt{0.931(0.069)\left(\frac{1}{51} + \frac{1}{62}\right)}} = -1.675$$
$$P = 2 \cdot \Pr(Z < -1.675) = 0.09$$

- Conclusion:

$$\alpha = 0.01$$

# Comparison of Two Proportions

# Comparison of Two Proportions

- We can also calculate a confidence interval for the difference of two proportions

# Comparison of Two Proportions

- We can also calculate a confidence interval for the difference of two proportions
- As in the one-sample case, the standard error is not the same for the confidence interval and hypothesis test

# Comparison of Two Proportions

- We can also calculate a confidence interval for the difference of two proportions
- As in the one-sample case, the standard error is not the same for the confidence interval and hypothesis test
- For a two-sided confidence interval, we are  $(1 - \alpha) \cdot 100\%$  confident that the interval  $\hat{p}_1 - \hat{p}_2 \pm z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}}$  contains the true population difference,  $p_1 - p_2$

$$\sqrt{\hat{p}_1(1 - \hat{p}_1) \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}$$

(Wald)

# Comparison of Two Proportions

# Comparison of Two Proportions

- Continuing with our dominant hand by group example, we can construct a two-sided 95% confidence interval for  $p_A - p_B$  as follows:

# Comparison of Two Proportions

- Continuing with our dominant hand by group example, we can construct a two-sided 95% confidence interval for  $p_A - p_B$  as follows:

$$\begin{aligned} & \hat{p}_A - \hat{p}_B \\ & (0.889 - 0.968) \pm 1.96 \sqrt{\frac{0.889 \cdot 0.111}{54} + \frac{0.968 \cdot 0.032}{62}} \\ & = (-0.173, 0.016) \end{aligned}$$

The handwritten annotations show the formula for the standard error of the difference in proportions. A red bracket groups the two terms:  $\frac{p_A(1-p_A)}{n_1}$  and  $\frac{p_B(1-p_B)}{n_2}$ . A red '+' sign is placed between the two terms.

# Comparison of Two Proportions

- Continuing with our dominant hand by group example, we can construct a two-sided 95% confidence interval for  $p_A - p_B$  as follows:

$$(0.889 - 0.968) \pm 1.96 \sqrt{\frac{0.889 \cdot 0.111}{54} + \frac{0.968 \cdot 0.032}{62}}$$
$$= (-0.173, 0.016)$$

- We are 95% confident that the interval  $(-0.173, 0.016)$  contains the true difference in the proportion of members of Group A and Group B who are right-handed

# Comparison of Two Proportions

# Comparison of Two Proportions

- One tail, lower bound:

# Comparison of Two Proportions

- One tail, lower bound:

$$\left( \hat{p}_1 - \hat{p}_2 - z_{\alpha} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}}, 1 \right)$$

# Comparison of Two Proportions

- One tail, lower bound:

$$\left( \hat{p}_1 - \hat{p}_2 - z_{\alpha} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}}, 1 \right)$$

- One tail, upper bound:

# Comparison of Two Proportions

- One tail, lower bound:

$$\left( \hat{p}_1 - \hat{p}_2 - z_\alpha \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}}, 1 \right)$$

- One tail, upper bound:

$$\left( -1, \hat{p}_1 - \hat{p}_2 + z_\alpha \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} \right)$$

# DSCC 462 Midpoint Survey


$$z_B \cdot \sqrt{\frac{\sigma}{n}} = \mu_0 + z_{1-\alpha} \cdot \sqrt{\frac{\sigma}{n}} - \mu_1$$

$$(z_B - z_{1-\alpha}) \sqrt{\frac{\sigma}{n}} = \mu_0 - \mu_1$$

- <https://forms.gle/Zt3Qzrb7S7UXFXY28>

- If you fill it out: +2.5% on midterm
- If at least 90% of the class fills it out: +2.5% on each person's midterm
  - Tell your friends!

$$(z_B - z_\alpha) \sqrt{\frac{\sigma}{n}} = \mu_0 - \mu_1$$
$$n = \left( \frac{z_\alpha + z_B}{\mu_0 - \mu_1} \right)^2 \cdot \sigma^2$$



$$z_{1-\alpha} = \frac{\bar{x} - \mu}{\sqrt{\frac{\sigma}{n}}}$$

$$z_{1-\alpha} \cdot \sqrt{\frac{\sigma}{n}} + \mu_0 = \bar{x}$$


$$z_B = \frac{\bar{x} - \mu_1}{\sqrt{\frac{\sigma}{n}}}$$

# DSCC 462 Midpoint Survey

$$\bar{x} \leftarrow x$$

plot (mu, some function of  
mu, -)

- <https://forms.gle/Zt3Qzrb7S7UXFXY28>
- If you fill it out: +2.5% on midterm
- If at least 90% of the class fills it out: +2.5% on each person's midterm
  - Tell your friends!



# DSCC 462 Midpoint Survey

- <https://forms.gle/Zt3Qzrb7S7UXFXY28>
- If you fill it out: +2.5% on midterm
- If at least 90% of the class fills it out: +2.5% on each person's midterm
  - Tell your friends!