

Midterm Cheat Sheet


Daxiang Na (那达翔)

2022-11-01

Contents

| | | |
|----------|--|----------|
| 1 | Type I and Type II Error, calculating power: | 2 |
| 2 | Skewness | 2 |
| 3 | Quantiles | 2 |
| 3.1 | Percentiles and quartiles | 2 |
| 3.2 | Modified Boxplot | 2 |
| 4 | Variance | 3 |
| 5 | Relationships Between Variables | 3 |
| 6 | Three Variables | 3 |
| 7 | Empirical Rule | 3 |
| 8 | Transformation: | 3 |
| 8.1 | Box-Cox Power Transformation | 3 |
| 8.2 | For right skewed data, use a function that tends to reduce larger values in proportion to smaller ones (i.e., an increasing function whose slope is decreasing): | 4 |
| 9 | Check normality: qqplot | 4 |

1 Type I and Type II Error, calculating power:



2 Skewness

Left (negative) skew: The left tail extends farther out than the right tail
Right (positive skew): The right tail extends farther out than the left tail

3 Quantiles

3.1 Percentiles and quartiles

The p sample quantile is the value below which a proportion p of the data are located.

E.g., if your birth weight is at the 95th percentile, then you weighed more than 0.95 of all newborn babies.

IQR: Interquartile Range = $Q3 - Q1$

1st quartile ($Q1$): 25th percentile, 2nd quartile ($Q2$): 50th percentile (median), 3rd quartile ($Q3$): 75th percentile.

3.2 Modified Boxplot

An outlier is a data point that is either:

- Less than: $Q1 - 1.5 \times (Q3 - Q1)$ = lower fence of box
- Greater than: $Q3 + 1.5 \times (Q3 - Q1)$ = upper fence of boxplot

- Standard span: $1.5 \times (Q3 - Q1) = 1.5 \times IQR$

4 Variance

5 Relationships Between Variables

Case CQ: Categorical and Quantitative

Case CC: Categorical and Categorical

Case QQ: Quantitative and Quantitative

6 Three Variables

add color as the third dimension

7 Empirical Rule

If a distribution is symmetric, unimodal, and bell-shaped (i.e., normally distributed), then the following hold:

- Approximately 68% of observations fall within one SD of the mean: $x \pm s$, or $(x - s, x + s)$
- Approximately 95% of observations fall within two SDs of the mean: $x \pm 2s$, or $(x - 2s, x + 2s)$
- Approximately 99.7% of observations fall within three SDs of the mean: $x \pm 3s$, or $(x - 3s, x + 3s)$

8 Transformation:

8.1 Box-Cox Power Transformation

$$y_{\lambda} = \begin{cases} \frac{x^{\lambda}-1}{\lambda}, & \lambda \neq 0 \\ \log(x), & \lambda = 0 \end{cases}$$

R code:

```
```r
library(MASS)
bc1 <- boxcox(x ~ 1)
bc1$x[bc1$y == max(bc1$y)]
```
```

```

```r
Example code from assignment
library(MASS)
bc1 <- boxcox(df$price ~ 1)
lambda <- bc1$x[bc1$y == max(bc1$y)]
trans <- (df$price^lambda - 1)/lambda
summary(trans)
```

```

8.2 For right skewed data, use a function that tends to reduce larger values in proportion to smaller ones (i.e., an increasing function whose slope is decreasing):

8.2.1 Log transformation:

in R: `log()`

8.2.2 Square-root transformation:

in R: `sqrt`

9 Check normality: qqplot

in R: `qqnorm(data); qqline(data)`