

Assignment t-test

Date: 05/05/2020 Name: D.Saravanan

t-test test the null hypothesis H_0 against the alternative hypothesis H_1 .

For univariate samples, t-test performs a Student t test. The test statistic is assumed to follow a Student t-Distribution $[df]$.

For multivariate samples, t-test performs Hotelling's t^2 test. The test statistic is assumed to follow a Hotelling t-square distribution $[p, df]$ where p is the dimension of data.

The degrees of freedom df , used to specify the distribution of the test statistic, depend on the sample size, number of samples, and in the case of two univariate samples, the results of a test for equal variances.

For the t-test, a cutoff α is chosen such that H_0 is rejected only if $p < \alpha$. The value of α used for the "Test Conclusion" and "Short Test Conclusion" properties is controlled by the Significance Level option. This value α is also used in diagnostic tests of assumptions, including tests for normality, equal variance, and symmetry. By default, α is set to 0.05.

1. Two sets of ten students selected at random from a college were taken. One set was given memory test as they were and the other was given the memory test after two weeks of training and the scores are given below.

Set A: 10 8 7 9 8 10 9 6 7 8

Set B: 12 8 8 10 8 11 9 8 9 9

Do you think there is a significant effect due to training?

Solution:

Null hypothesis: $H_0 : \mu_1 = \mu_2$, No significant effect due to training

Alternative hypothesis: $H_1 : \mu_1 \neq \mu_2$, Significant effect due to training

Set A:

Mean:

$$\bar{x}_1 = \frac{\sum_{i=1}^{n_1} x_{1i}}{n_1} = \frac{10 + 8 + 7 + 9 + 8 + 10 + 9 + 6 + 7 + 8}{10} = 8.2$$

Variance:

$$s_1^2 = \frac{\sum_{i=1}^{n_1} (x_{1i} - \bar{x}_1)^2}{n_1 - 1} = \frac{\sum_{i=1}^{10} (x_{1i} - 8.2)^2}{10 - 1} = 1.73333$$

Standard Deviation:

$$s_1 = \sqrt{\frac{\sum_{i=1}^{n_1} (x_{1i} - \bar{x}_1)^2}{n_1 - 1}} = \sqrt{\frac{\sum_{i=1}^{10} (x_{1i} - 8.2)^2}{10 - 1}} = \sqrt{1.73333} = 1.31656$$

Set B:

Mean:

$$\bar{x}_2 = \frac{\sum_{i=1}^{n_2} x_{2i}}{n_2} = \frac{12 + 8 + 8 + 10 + 8 + 11 + 9 + 8 + 9 + 9}{10} = 9.2$$

Variance:

$$s_2^2 = \frac{\sum_{i=1}^{n_2} (x_{2i} - \bar{x}_2)^2}{n_2 - 1} = \frac{\sum_{i=1}^{10} (x_{2i} - 9.2)^2}{10 - 1} = 1.95556$$

Standard Deviation:

$$s_2 = \sqrt{\frac{\sum_{i=1}^{n_2} (x_{2i} - \bar{x}_2)^2}{n_2 - 1}} = \sqrt{\frac{\sum_{i=1}^{10} (x_{2i} - 9.2)^2}{10 - 1}} = \sqrt{1.95556} = 1.39841$$

Calculation of s_1/s_2 :

$$\frac{s_1}{s_2} = \frac{1.31656}{1.39841} = 0.94147$$

Test statistic:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{s_1^2/n_1 + s_2^2/n_2}}$$

where n_1 and n_2 are the sample sizes, \bar{x}_1 and \bar{x}_2 are the sample means, and s_1^2 and s_2^2 are the sample variances.

If equal variances are assumed ($0.5 < s_1/s_2 < 2$), then the formula reduces to:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{s_p \sqrt{1/n_1 + 1/n_2}}$$

where

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$$

Calculation of s_p :

$$s_p = \sqrt{\frac{(10 - 1) \times 1.31656^2 + (10 - 1) \times 1.39841^2}{10 + 10 - 2}} = 1.35810$$

t-statistic:

$$t = \frac{8.2 - 9.2}{1.35810 \times \sqrt{1/10 + 1/10}} = -1.64647$$

degrees of freedom:

$$d.o.f = n_1 + n_2 - 2 = 10 + 10 - 2 = 18$$

The critical value for t (from t -distribution table) with degrees of freedom = 18 and $\alpha = 0.05$ is 2.101

Conclusion: $|t| < t_{0.05/2}$, we fail to reject the Null hypothesis. That is there is no significant effect in the scores of the memory test due to training.

Standard Error:

$$S.E = s_p \sqrt{1/n_1 + 1/n_2} = 1.35810 \times \sqrt{1/10 + 1/10} = 0.60736$$

Confidence Interval: 95% fiducial limits

The confidence interval is given by

$$\begin{aligned} & (\bar{x}_1 - \bar{x}_2) \pm t_{0.05/2} \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \\ & (8.2 - 9.2) \pm 2.101 \sqrt{\frac{(10 - 1)(1.31656)^2 + (10 - 1)(1.39841)^2}{10 + 10 - 2}} \sqrt{\frac{1}{10} + \frac{1}{10}} \\ & (-1.0) \pm 2.101 \sqrt{\frac{33.19993}{18}} 0.44721 \\ & - 2.27606 < \mu < 0.27606 \end{aligned}$$

Program:

```
#!/usr/bin/env python3
import numpy as np
from scipy.stats import t
import matplotlib.pyplot as plt
plt.style.use('seaborn-darkgrid')

def similar_variance(n1, s1_mean, n2, s2_mean):
    dof = n1+n2-2
    sp = np.sqrt(((n1-1)*s1_stdv**2+(n2-1)*s2_stdv**2)/(dof))
    return (s1_mean-s2_mean)/(sp * np.sqrt(1/n1 + 1/n2)), dof, sp

def non_similar_variance(n1, s1_mean, n2, s2_mean):
    sd = np.sqrt(s1_stdv**2/n1 + s2_stdv**2/n2)
    dof = (sd**4)/(((s1_stdv**2/n1)**2/(n1-1)) + ((s2_stdv**2/n2)**2/(n2-1)))
    return (s1_mean-s2_mean)/sd, dof, sd

def ttest_and_variance(s1_stdv, s2_stdv):
    if 0.5 < s1_stdv/s2_stdv < 2:
        return similar_variance(n1, s1_mean, n2, s2_mean)
    else:
        return non_similar_variance(n1, s1_mean, n2, s2_mean)

print("_____Two-sample t-test for unpaired data_____\\n")

sample1 = [float(value) for value in input("Sample_1 values: ").split()]
sample2 = [float(value) for value in input("Sample_2 values: ").split()]

# Number of Observations
n1 = len(sample1); n2 = len(sample2)

# Mean and Standard Deviation
s1_mean = np.mean(sample1); s1_stdv = np.std(sample1, ddof=1)
s2_mean = np.mean(sample2); s2_stdv = np.std(sample2, ddof=1)

# level of significance, confidence level
alpha = 0.05; clevel = 1 - alpha

# standard error (SE) of mean of sample1
std_err1 = s1_stdv/np.sqrt(n1)

print("\\nSample 1: \\n\\t Number of Observations = {} \\n\\t Mean = {:.5f}".format(n1, s1_mean))
print("\\t Standard Deviation = {:.5f}".format(s1_stdv))
print("\\t Standard Error of the Mean = {:.5f}".format(std_err1))

# standard error (SE) of mean of sample2
std_err2 = s2_stdv/np.sqrt(n2)
```

```

print("\nSample 2: \n\t Number of Observations = {} \n\t Mean = {:.5f}".format(n2,s2_mean))
print("\t Standard Deviation = {:.5f}".format(s2_stdv))
print("\t Standard Error of the Mean = {:.5f}".format(std_err2))

# calculation of t-statistic and degrees of freedom
tstatistic, dof, sp = ttest_and_variance(s1_stdv, s2_stdv)
print("\nt-statistic: {:.5f}".format(tstatistic))

# calculation of critical values
tcritical_l = t.ppf(q = alpha/2, df = dof)
tcritical_u = -tcritical_l
print("\nCritical values are {:.5f}, {:.5f}".format(tcritical_l, tcritical_u))

# decision making: t-statistic and critical values
if tstatistic < tcritical_l or tstatistic > tcritical_u:
    print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

# calculation of p-value
if tstatistic < 0: pvalue = 2*t.cdf(tstatistic, df = dof)
else: pvalue = 2*(1 - t.cdf(tstatistic, df = dof))
print("\np-value: {:.5f}".format(pvalue))

# decision making: p-value and level of significance
if pvalue < alpha: print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

# standard error
std_error = sp * np.sqrt(1/n1 + 1/n2)
print("\nStandard Error: {:.5f}".format(std_error))

# confidence interval
cnf_int = (s1_mean - s2_mean) + std_error * np.array([tcritical_l, tcritical_u])
print("Confidence Interval: {}".format(cnf_int))

x1=np.linspace(-10,tcritical_l,1000); y1=t.pdf(x1,df=dof)
x2=np.linspace(tcritical_u,10,1000); y2=t.pdf(x2,df=dof)
x3=np.linspace(-4,4,1000); y3=t.pdf(x3,df=dof)
ax=plt.figure().add_subplot(111); ax.plot(x3,y3,color='brown')
ax.fill_between(x1,y1,0,alpha=0.5,color='brown',label='Rejection Region')
ax.fill_between(x2,y2,0,alpha=0.5,color='brown'); ax.set(xlabel='X',ylabel='P(X)')
ax.axvline(x=tstatistic,ls='--',c='b',label='t-statistic = {:.5f}'.format(tstatistic))
ax.set(xlim=[-4,4],title="t-distribution (degrees of freedom = {:.0f)".format(dof))
plt.legend(); plt.savefig('tplot3a.pdf',dpi=72,bbox_inches='tight'); plt.show()

```

Output:

_____Two-sample t-test for unpaired data_____

Sample_1 values: 10 8 7 9 8 10 9 6 7 8
Sample_2 values: 12 8 8 10 8 11 9 8 9 9

Sample 1:
Number of Observations = 10
Mean = 8.20000
Standard Deviation = 1.31656
Standard Error of the Mean = 0.41633


Sample 2:
Number of Observations = 10
Mean = 9.20000
Standard Deviation = 1.39841
Standard Error of the Mean = 0.44222

t-statistic: -1.64646

Critical values are -2.10092, 2.10092
Fail to reject the Null hypothesis.

p-value: 0.11702
Fail to reject the Null hypothesis.

Standard Error: 0.60736
Confidence Interval: [-2.27602071 0.27602071]



Program: t-test with in-built scipy.stats.ttest

```
#!/usr/bin/env python3
import numpy as np
from scipy.stats import t, ttest_ind
import matplotlib.pyplot as plt
plt.style.use('seaborn-darkgrid')

print("_____Two-sample t-test for unpaired data_____\\n")

sample1 = [float(value) for value in input("Sample_1 values: ").split()]
sample2 = [float(value) for value in input("Sample_2 values: ").split()]

# Number of Observations
n1 = len(sample1); n2 = len(sample2)

# Mean and Standard Deviation
s1_mean = np.mean(sample1); s1_stdv = np.std(sample1, ddof=1)
s2_mean = np.mean(sample2); s2_stdv = np.std(sample2, ddof=1)

# calculation of degrees of freedom and pooled variance
if 0.5 < s1_stdv/s2_stdv < 2:
    dof = n1+n2-2
    sp = np.sqrt(((n1-1)*s1_stdv**2+(n2-1)*s2_stdv**2)/(dof))
else:
    sp = np.sqrt(s1_stdv**2/n1 + s2_stdv**2/n2)
    dof = (sp**4)/(((s1_stdv**2/n1)**2/(n1-1)) + ((s2_stdv**2/n2)**2/(n2-1)))
```

```

# level of significance, confidence level
alpha = 0.05; clevel = 1 - alpha

# standard error (SE) of mean of sample1
std_err1 = s1_stdv/np.sqrt(n1)

print("\nSample 1: \n\t Number of Observations = {} \n\t Mean = {:.5f}".format(n1,s1_mean))
print("\t Standard Deviation = {:.5f}".format(s1_stdv))
print("\t Standard Error of the Mean = {:.5f}".format(std_err1))

# standard error (SE) of mean of sample2
std_err2 = s2_stdv/np.sqrt(n2)

print("\nSample 2: \n\t Number of Observations = {} \n\t Mean = {:.5f}".format(n2,s2_mean))
print("\t Standard Deviation = {:.5f}".format(s2_stdv))
print("\t Standard Error of the Mean = {:.5f}".format(std_err2))

# calculation of t-statistic and p-value
tstatistic, pvalue = ttest_ind(sample1, sample2)
print("\nt-statistic: {:.5f}".format(tstatistic))

# calculation of critical values
tcritical_l = t.ppf(q = alpha/2, df = dof)
tcritical_u = -tcritical_l
print("\nCritical values are {:.5f}, {:.5f}".format(tcritical_l, tcritical_u))

# decision making: t-statistic and critical values
if tstatistic < tcritical_l or tstatistic > tcritical_u:
    print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

print("\np-value: {:.5f}".format(pvalue))

# decision making: p-value and level of significance
if pvalue < alpha: print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

# standard error
std_error = sp * np.sqrt(1/n1 + 1/n2)
print("\nStandard Error: {:.5f}".format(std_error))

# confidence interval
cnf_int = (s1_mean - s2_mean) + std_error * np.array([tcritical_l, tcritical_u])
print("Confidence Interval: {}".format(cnf_int))

x1=np.linspace(-10,tcritical_l,1000); y1=t.pdf(x1,df=dof)
x2=np.linspace(tcritical_u,10,1000); y2=t.pdf(x2,df=dof)
x3=np.linspace(-4,4,1000); y3=t.pdf(x3,df=dof)
ax=plt.figure().add_subplot(111); ax.plot(x3,y3,color='brown')
ax.fill_between(x1,y1,0,alpha=0.5,color='brown',label='Rejection Region')
ax.fill_between(x2,y2,0,alpha=0.5,color='brown'); ax.set(xlabel='X',ylabel='P(X)')
ax.axvline(x=tstatistic,ls='--',c='b',label='t-statistic = {:.5f}'.format(tstatistic))
ax.set(xlim=[-4,4],title="t-distribution (degrees of freedom = {:.0f})".format(dof))
plt.legend(); plt.savefig('ttest2.pdf',dpi=72,bbox_inches='tight'); plt.show()

```

Output:

_____Two-sample t-test for unpaired data_____

Sample_1 values: 10 8 7 9 8 10 9 6 7 8
Sample_2 values: 12 8 8 10 8 11 9 8 9 9

Sample 1:
Number of Observations = 10

Mean = 8.20000
 Standard Deviation = 1.31656
 Standard Error of the Mean = 0.41633

Sample 2:


Number of Observations = 10
 Mean = 9.20000
 Standard Deviation = 1.39841
 Standard Error of the Mean = 0.44222

t-statistic: -1.64646

Critical values are -2.10092, 2.10092
 Fail to reject the Null hypothesis.

p-value: 0.11702
 Fail to reject the Null hypothesis.

Standard Error: 0.60736
 Confidence Interval: [-2.27602071 0.27602071]



2. A group of 5 patients treated with medicine A weighs 42, 29, 48, 60 and 41 kg. A second group of 7 patients from the same hospital treated with medicine B weighs 38, 42, 56, 64, 68, 69 and 62 kg. Do you agree with the claim that medicine B increases weight significantly.

Solution:

Null hypothesis: $H_0 : \mu_1 = \mu_2$, No significant increase in weight

Alternative hypothesis: $H_1 : \mu_1 \neq \mu_2$, Significant increase in weight

Medicine A:

Mean:

$$\bar{x}_1 = \frac{\sum_{i=1}^{n_1} x_{1i}}{n_1} = \frac{42 + 29 + 48 + 60 + 41}{5} = 44.0$$

Variance:

$$s_1^2 = \frac{\sum_{i=1}^{n_1} (x_{1i} - \bar{x}_1)^2}{n_1 - 1} = \frac{\sum_{i=1}^5 (x_{1i} - 44.0)^2}{5 - 1} = 127.5$$

Standard Deviation:

$$s_1 = \sqrt{\frac{\sum_{i=1}^{n_1} (x_{1i} - \bar{x}_1)^2}{n_1 - 1}} = \sqrt{\frac{\sum_{i=1}^5 (x_{1i} - 44.0)^2}{5 - 1}} = \sqrt{127.5} = 11.29159$$

Medicine B:

Mean:

$$\bar{x}_1 = \frac{\sum_{i=1}^{n_1} x_{1i}}{n_1} = \frac{38 + 42 + 56 + 64 + 68 + 69 + 62}{7} = 57.0$$

Variance:

$$s_1^2 = \frac{\sum_{i=1}^{n_1} (x_{1i} - \bar{x}_1)^2}{n_1 - 1} = \frac{\sum_{i=1}^7 (x_{1i} - 57.0)^2}{7 - 1} = 154.3$$

Standard Deviation:

$$s_1 = \sqrt{\frac{\sum_{i=1}^{n_1} (x_{1i} - \bar{x}_1)^2}{n_1 - 1}} = \sqrt{\frac{\sum_{i=1}^7 (x_{1i} - 57.0)^2}{7 - 1}} = \sqrt{154.3} = 12.42310$$

Calculation of s_1/s_2 :

$$\frac{s_1}{s_2} = \frac{11.29159}{12.42310} = 0.90892$$

Test statistic:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{s_1^2/n_1 + s_2^2/n_2}}$$

If equal variances are assumed ($0.5 < s_1/s_2 < 2$), then the formula reduces to:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{s_p \sqrt{1/n_1 + 1/n_2}}$$

where

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$$

Calculation of s_p :

$$s_p = \sqrt{\frac{(5 - 1) \times 11.29159^2 + (7 - 1) \times 12.42310^2}{5 + 7 - 2}} = 11.98332$$

t-statistic:

$$t = \frac{44.0 - 57.0}{11.98332 \times \sqrt{1/5 + 1/7}} = -1.85272$$

degrees of freedom:

$$d.o.f = n_1 + n_2 - 2 = 5 + 7 - 2 = 10$$

The critical value for t (from t -distribution table) with degrees of freedom = 10 and $\alpha = 0.05$ is 2.228

Conclusion: $|t| < t_{0.05/2}$, we fail to reject the Null hypothesis. That is there is no significant increase in weight due to medicine B when compared with medicine A.

Standard Error:

$$S.E = s_p \sqrt{1/n_1 + 1/n_2} = 11.98332 \times \sqrt{1/5 + 1/7} = 7.01671$$

Confidence Interval: 95% fiducial limits

The confidence interval is given by

$$\begin{aligned} & (\bar{x}_1 - \bar{x}_2) \pm t_{0.05/2} \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \\ & (44.0 - 57.0) \pm 2.228 \sqrt{\frac{(5 - 1)(11.29159)^2 + (7 - 1)(12.42310)^2}{5 + 7 - 2}} \sqrt{\frac{1}{5} + \frac{1}{7}} \\ & (-13.0) \pm 2.228 \sqrt{\frac{1436.00050}{10}} 0.58554 \\ & -28.63324 < \mu < 2.63324 \end{aligned}$$

Program:

```
#!/usr/bin/env python3
import numpy as np
from scipy.stats import t
import matplotlib.pyplot as plt
plt.style.use('seaborn-darkgrid')

def similar_variance(n1, s1_mean, n2, s2_mean):
    dof = n1+n2-2
    sp = np.sqrt(((n1-1)*s1_stdv**2+(n2-1)*s2_stdv**2)/(dof))
    return (s1_mean-s2_mean)/(sp * np.sqrt(1/n1 + 1/n2)), dof, sp

def non_similar_variance(n1, s1_mean, n2, s2_mean):
    sd = np.sqrt(s1_stdv**2/n1 + s2_stdv**2/n2)
    dof = (sd**4)/(((s1_stdv**2/n1)**2/(n1-1)) + ((s2_stdv**2/n2)**2/(n2-1)))
    return (s1_mean-s2_mean)/sd, dof, sd

def ttest_and_variance(s1_stdv, s2_stdv):
    if 0.5 < s1_stdv/s2_stdv < 2:
        return similar_variance(n1, s1_mean, n2, s2_mean)
    else:
        return non_similar_variance(n1, s1_mean, n2, s2_mean)

print("_____Two-sample t-test for unpaired data_____\\n")

sample1 = [float(value) for value in input("Sample_1 values: ").split()]
sample2 = [float(value) for value in input("Sample_2 values: ").split()]

# Number of Observations
n1 = len(sample1); n2 = len(sample2)

# Mean and Standard Deviation
s1_mean = np.mean(sample1); s1_stdv = np.std(sample1, ddof=1)
s2_mean = np.mean(sample2); s2_stdv = np.std(sample2, ddof=1)

# level of significance, confidence level
alpha = 0.05; clevel = 1 - alpha
```

```

# standard error (SE) of mean of sample1
std_err1 = s1_stdv/np.sqrt(n1)

print("\nSample 1: \n\t Number of Observations = {} \n\t Mean = {:.5f}".format(n1,s1_mean))
print("\t Standard Deviation = {:.5f}".format(s1_stdv))
print("\t Standard Error of the Mean = {:.5f}".format(std_err1))

# standard error (SE) of mean of sample2
std_err2 = s2_stdv/np.sqrt(n2)

print("\nSample 2: \n\t Number of Observations = {} \n\t Mean = {:.5f}".format(n2,s2_mean))
print("\t Standard Deviation = {:.5f}".format(s2_stdv))
print("\t Standard Error of the Mean = {:.5f}".format(std_err2))

# calculation of t-statistic and degrees of freedom
tstatistic, dof, sp = ttest_and_variance(s1_stdv, s2_stdv)
print("\nt-statistic: {:.5f}".format(tstatistic))

# calculation of critical values
tcritical_l = t.ppf(q = alpha/2, df = dof)
tcritical_u = -tcritical_l
print("\nCritical values are {:.5f}, {:.5f}".format(tcritical_l, tcritical_u))

# decision making: t-statistic and critical values
if tstatistic < tcritical_l or tstatistic > tcritical_u:
    print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

# calculation of p-value
if tstatistic < 0: pvalue = 2*t.cdf(tstatistic, df = dof)
else: pvalue = 2*(1 - t.cdf(tstatistic, df = dof))
print("\np-value: {:.5f}".format(pvalue))

# decision making: p-value and level of significance
if pvalue < alpha: print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

# standard error
std_error = sp * np.sqrt(1/n1 + 1/n2)
print("\nStandard Error: {:.5f}".format(std_error))

# confidence interval
cnf_int = (s1_mean - s2_mean) + std_error * np.array([tcritical_l, tcritical_u])
print("Confidence Interval: {}".format(cnf_int))

x1=np.linspace(-10,tcritical_l,1000); y1=t.pdf(x1,df=dof)
x2=np.linspace(tcritical_u,10,1000); y2=t.pdf(x2,df=dof)
x3=np.linspace(-4,4,1000); y3=t.pdf(x3,df=dof)
ax=plt.figure().add_subplot(111); ax.plot(x3,y3,color='brown')
ax.fill_between(x1,y1,0,alpha=0.5,color='brown',label='Rejection Region')
ax.fill_between(x2,y2,0,alpha=0.5,color='brown'); ax.set(xlabel='X',ylabel='P(X)')
ax.axvline(x=tstatistic,ls='--',c='b',label='t-statistic = {:.5f}'.format(tstatistic))
ax.set(xlim=[-4,4],title="t-distribution (degrees of freedom = {:.0f})".format(dof))
plt.legend(); plt.savefig('tscript41.pdf',dpi=72,bbox_inches='tight'); plt.show()

```

Output:

_____Two-sample t-test for unpaired data_____

Sample_1 values: 42 29 48 60 41

Sample_2 values: 38 42 56 64 68 69 62

Sample 1:

Number of Observations = 5
Mean = 44.00000
Standard Deviation = 11.29159
Standard Error of the Mean = 5.04975

Sample 2:


Number of Observations = 7
Mean = 57.00000
Standard Deviation = 12.42310
Standard Error of the Mean = 4.69549

t-statistic: -1.85272

Critical values are -2.22814, 2.22814
Fail to reject the Null hypothesis.

p-value: 0.09363
Fail to reject the Null hypothesis.

Standard Error: 7.01671
Confidence Interval: [-28.63421472 2.63421472]



Program: t-test with in-built scipy.stats.ttest

```
#!/usr/bin/env python3
import numpy as np
from scipy.stats import t, ttest_ind
import matplotlib.pyplot as plt
plt.style.use('seaborn-darkgrid')

print("_____Two-sample t-test for unpaired data_____\\n")

sample1 = [float(value) for value in input("Sample_1 values: ").split()]
sample2 = [float(value) for value in input("Sample_2 values: ").split()]

# Number of Observations
n1 = len(sample1); n2 = len(sample2)
```

```

# Mean and Standard Deviation
s1_mean = np.mean(sample1); s1_stdv = np.std(sample1, ddof=1)
s2_mean = np.mean(sample2); s2_stdv = np.std(sample2, ddof=1)

# calculation of degrees of freedom and pooled variance
if 0.5 < s1_stdv/s2_stdv < 2:
    dof = n1+n2-2
    sp = np.sqrt(((n1-1)*s1_stdv**2+(n2-1)*s2_stdv**2)/(dof))
else:
    sp = np.sqrt(s1_stdv**2/n1 + s2_stdv**2/n2)
    dof = (sp**4)/(((s1_stdv**2/n1)**2/(n1-1)) + ((s2_stdv**2/n2)**2/(n2-1)))

# level of significance, confidence level
alpha = 0.05; clevel = 1 - alpha

# standard error (SE) of mean of sample1
std_err1 = s1_stdv/np.sqrt(n1)

print("\nSample 1: \n\t Number of Observations = {} \n\t Mean = {:.5f}".format(n1,s1_mean))
print("\t Standard Deviation = {:.5f}".format(s1_stdv))
print("\t Standard Error of the Mean = {:.5f}".format(std_err1))

# standard error (SE) of mean of sample2
std_err2 = s2_stdv/np.sqrt(n2)

print("\nSample 2: \n\t Number of Observations = {} \n\t Mean = {:.5f}".format(n2,s2_mean))
print("\t Standard Deviation = {:.5f}".format(s2_stdv))
print("\t Standard Error of the Mean = {:.5f}".format(std_err2))

# calculation of t-statistic and p-value
tstatistic, pvalue = ttest_ind(sample1, sample2)
print("\nt-statistic: {:.5f}".format(tstatistic))

# calculation of critical values
tcritical_l = t.ppf(q = alpha/2, df = dof)
tcritical_u = -tcritical_l
print("\nCritical values are {:.5f}, {:.5f}".format(tcritical_l, tcritical_u))

# decision making: t-statistic and critical values
if tstatistic < tcritical_l or tstatistic > tcritical_u:
    print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

print("\np-value: {:.5f}".format(pvalue))

# decision making: p-value and level of significance
if pvalue < alpha: print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

# standard error
std_error = sp * np.sqrt(1/n1 + 1/n2)
print("\nStandard Error: {:.5f}".format(std_error))

# confidence interval
cnf_int = (s1_mean - s2_mean) + std_error * np.array([tcritical_l, tcritical_u])
print("Confidence Interval: {}".format(cnf_int))

x1=np.linspace(-10,tcritical_l,1000); y1=t.pdf(x1,df=dof)
x2=np.linspace(tcritical_u,10,1000); y2=t.pdf(x2,df=dof)
x3=np.linspace(-4,4,1000); y3=t.pdf(x3,df=dof)
ax=plt.figure().add_subplot(111); ax.plot(x3,y3,color='brown')
ax.fill_between(x1,y1,0,alpha=0.5,color='brown',label='Rejection Region')
ax.fill_between(x2,y2,0,alpha=0.5,color='brown'); ax.set(xlabel='X',ylabel='P(X)')
ax.axvline(x=tstatistic,ls='--',c='b',label='t-statistic = {:.5f}'.format(tstatistic))
ax.set(xlim=[-4,4],title="t-distribution (degrees of freedom = {:.0f})".format(dof))

```

```
plt.legend(); plt.savefig('tscript42.pdf',dpi=72,bbox_inches='tight'); plt.show()
```

Output:

_____Two-sample T-test for unpaired data_____

Sample 1:

Number of Observations = 5
Mean = 44.00000
Standard Deviation = 10.09950
Standard Error of the Mean = 4.51664

Sample 2:


Number of Observations = 7
Mean = 57.00000
Standard Deviation = 11.50155
Standard Error of the Mean = 4.34718

T statistics: -1.85272

Critical values are -2.22814, 2.22814
Fail to reject the Null hypothesis.

p-value: 0.09363
Fail to reject the Null hypothesis.

Standard Error: 6.41885
Confidence Interval: [-27.30208861 1.30208861]



3. Samples of two types of electric bulbs were tested for length of life and the following data were obtained

	Type I	Type II
No. of Samples:	8	7
Mean (hours):	1134	1024
SD (hours):	35	40

Test at 5 percent level, whether the difference in sample mean is significant.

Solution:**Null hypothesis:** $H_0 : \mu_1 = \mu_2$, No significant difference in sample mean**Alternative hypothesis:** $H_1 : \mu_1 > \mu_2$, (Right tailed test)**Type I:**

Mean = 1134

Variance = 1225

Standard Deviation = 35

Type II:

Mean = 1024

Variance = 1600

Standard Deviation = 40

Calculation of s_1/s_2 :

$$\frac{s_1}{s_2} = \frac{35}{40} = 0.875$$

Test statistic:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{s_1^2/n_1 + s_2^2/n_2}}$$

where n_1 and n_2 are the sample sizes, \bar{x}_1 and \bar{x}_2 are the sample means, and s_1^2 and s_2^2 are the sample variances.If equal variances are assumed ($0.5 < s_1/s_2 < 2$), then the formula reduces to:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{s_p \sqrt{1/n_1 + 1/n_2}}$$

where

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$$

Calculation of s_p :

$$s_p = \sqrt{\frac{(8 - 1) \times 35^2 + (7 - 1) \times 40^2}{8 + 7 - 2}} = 37.39087$$

t-statistic:

$$t = \frac{1134 - 1024}{37.39087 \times \sqrt{1/8 + 1/7}} = 5.68428$$

degrees of freedom:

$$d.o.f = n_1 + n_2 - 2 = 8 + 7 - 2 = 13$$

The critical value for t (from t -distribution table) with degrees of freedom = 13 and $\alpha = 0.05$ is 1.771**Conclusion:** $|t| > t_{0.05}$, we reject the Null hypothesis. That is there is significant difference in sample mean.

Standard Error:

$$S.E = s_p \sqrt{1/n_1 + 1/n_2} = 37.39087 \times \sqrt{1/8 + 1/7} = 19.35161$$

Confidence Interval: 95% fiducial limits

The confidence interval for the small sample mean is given by

$$(\bar{x}_1 - \bar{x}_2) \pm t_{0.05} \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$$
$$(1134 - 1024) \pm 1.771 \sqrt{\frac{(8 - 1)(35)^2 + (7 - 1)(40)^2}{8 + 7 - 2}} \sqrt{\frac{1}{8} + \frac{1}{7}}$$
$$(110) \pm 1.771 \sqrt{\frac{18175}{13}} 0.51755$$
$$75.72824 < \mu < 144.27176$$

Program:

```
#!/usr/bin/env python3
import numpy as np
from scipy.stats import t
import matplotlib.pyplot as plt
plt.style.use('seaborn-darkgrid')

def similar_variance(n1,s1_mean,n2,s2_mean):
    dof = n1+n2-2
    sp = np.sqrt(((n1-1)*s1_stdv**2+(n2-1)*s2_stdv**2)/(dof))
    return (s1_mean-s2_mean)/(sp * np.sqrt(1/n1 + 1/n2)), dof, sp

def non_similar_variance(n1,s1_mean,n2,s2_mean):
    sd = np.sqrt(s1_stdv**2/n1 + s2_stdv**2/n2)
    dof = (sd**4)/(((s1_stdv**2/n1)**2/(n1-1)) + ((s2_stdv**2/n2)**2/(n2-1)))
    return (s1_mean-s2_mean)/sd, dof, sd

def ttest_and_variance(s1_stdv, s2_stdv):
    if 0.5 < s1_stdv/s2_stdv < 2:
        return similar_variance(n1,s1_mean,n2,s2_mean)
    else:
        return non_similar_variance(n1,s1_mean,n2,s2_mean)

print("_____Two-sample t-test for unpaired data_____\\n")

# Number of Observations
n1 = 8; n2 = 7

# Mean and Standard Deviation
s1_mean = 1134; s1_stdv = 35
s2_mean = 1024; s2_stdv = 40

# level of significance, confidence level
alpha = 0.05; clevel = 1 - alpha

# standard error (SE) of mean of sample1
std_err1 = s1_stdv/np.sqrt(n1)

print("\\nSample 1: \\n\\t Number of Observations = {} \\n\\t Mean = {:.5f}".format(n1,s1_mean))
print("\\t Standard Deviation = {:.5f}".format(s1_stdv))
print("\\t Standard Error of the Mean = {:.5f}".format(std_err1))

# standard error (SE) of mean of sample2
std_err2 = s2_stdv/np.sqrt(n2)

print("\\nSample 2: \\n\\t Number of Observations = {} \\n\\t Mean = {:.5f}".format(n2,s2_mean))
```

```

print("\\t Standard Deviation = {:.5f}".format(s2_stdv))
print("\\t Standard Error of the Mean = {:.5f}".format(std_err2))

# calculation of t-statistic and degrees of freedom
tstatistic, dof, sp = ttest_and_variance(s1_stdv, s2_stdv)
print("\\nT statistics: {:.5f}".format(tstatistic))

# calculation of critical values
tcritical_l = t.ppf(q = alpha, df = dof)
tcritical_u = -tcritical_l
print("\\nCritical values are {:.5f}, {:.5f)".format(tcritical_l, tcritical_u))

# decision making: t-statistic and critical values
if tstatistic < tcritical_l or tstatistic > tcritical_u:
    print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

# calculation of p-value
if tstatistic < 0: pvalue = 2*t.cdf(tstatistic, df = dof)
else: pvalue = 2*(1 - t.cdf(tstatistic, df = dof))
print("\\np-value: {:.5f)".format(pvalue))

# decision making: p-value and level of significance
if pvalue < alpha: print("Reject the Null hypothesis.")
else: print("Fail to reject the Null hypothesis.")

# standard error
std_error = sp * np.sqrt(1/n1 + 1/n2)
print("\\nStandard Error: {:.5f)".format(std_error))

# confidence interval
cnf_int = (s1_mean - s2_mean) + std_error * np.array([tcritical_l, tcritical_u])
print("Confidence Interval: {}".format(cnf_int))

#x1=np.linspace(-10,tcritical_l,1000); y1=t.pdf(x1,df=dof)
#x2=np.linspace(tcritical_u,10,1000); y2=t.pdf(x2,df=dof)
#x3=np.linspace(-6,6,1000); y3=t.pdf(x3,df=dof)
ax=plt.figure().add_subplot(111); ax.plot(x3,y3,color='brown')
ax.fill_between(x2,y2,0,alpha=0.5,color='brown',label='Rejection Region')
ax.fill_between(x2,y2,0,alpha=0.5,color='brown'); ax.set(xlabel='X',ylabel='P(X)')
ax.axvline(x=tstatistic,ls='--',c='b',label='t-statistic = {:.5f}'.format(tstatistic))
ax.set(xlim=[-6,6],title="t-distribution (degrees of freedom = {:.0f})".format(dof))
plt.legend(); plt.savefig('t2script3.pdf',dpi=72,bbox_inches='tight'); plt.show()

```

Output:

_____Two-sample t-test for unpaired data_____

Sample 1:

```

Number of Observations = 8
Mean = 1134.00000
Standard Deviation = 35.00000
Standard Error of the Mean = 12.37437

```

Sample 2:

```

Number of Observations = 7
Mean = 1024.00000
Standard Deviation = 40.00000
Standard Error of the Mean = 15.11858

```

T statistics: 5.68428

Critical values are -1.77093, 1.77093
Reject the Null hypothesis.

p-value: 0.00007
Reject the Null hypothesis.

Standard Error: 19.35161
Confidence Interval: [75.7295839 144.2704161]

