

Lab06

1. Apply Hadamard gate to $|\psi\rangle=\alpha|0\rangle+\beta|1\rangle$ and report the result as both Dirac notation and Vector notation.

$$\text{a) } |\psi\rangle=\alpha|0\rangle+\beta|1\rangle = \frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle = \frac{1}{\sqrt{2}}\begin{bmatrix} 1 \\ 0 \end{bmatrix} + \frac{1}{\sqrt{2}}\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$\text{b) } |\psi\rangle=\alpha|0\rangle+\beta|1\rangle = \frac{1}{\sqrt{2}}|0\rangle - \frac{1}{\sqrt{2}}|1\rangle = \frac{1}{\sqrt{2}}\begin{bmatrix} 1 \\ 0 \end{bmatrix} - \frac{1}{\sqrt{2}}\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} \end{bmatrix}$$

2. Mathematically show that the resulted qubit from the previous question follows the probability requirement (i.e., the summation of all probabilities of all measurement results should be 1.),

$$\text{a) } \alpha^2 + \beta^2 = \frac{1}{\sqrt{2}}^2 + \frac{1}{\sqrt{2}}^2 = \frac{1}{2} + \frac{1}{2} = 1$$

$$\text{b) } \alpha^2 + \beta^2 = \frac{1}{\sqrt{2}}^2 + \left(-\frac{1}{\sqrt{2}}\right)^2 = \frac{1}{2} + \frac{1}{2} = 1$$

3. Mathematically show that the reverse of Hadamard gate is itself. Note that a matrix A is its own reverse if $AA = I$, where I is the identity matrix.

$$\text{a) } \frac{1}{\sqrt{2}}\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} = \frac{1}{\sqrt{2}}\begin{bmatrix} \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \end{bmatrix} = \frac{1}{\sqrt{2}}\begin{bmatrix} \frac{2}{\sqrt{2}} \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\text{b) } \frac{1}{\sqrt{2}}\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} \end{bmatrix} = \frac{1}{\sqrt{2}}\begin{bmatrix} \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \end{bmatrix} = \frac{1}{\sqrt{2}}\begin{bmatrix} 0 \\ \frac{2}{\sqrt{2}} \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$