Relativistic Stark resonances in Debye plasmas: screening undoes the relativistic suppression of ionization widths

C. D. Reinoso[1, *]

[1] *Facultad de Ciencias, Escuela Superior Politécnica de Chimborazo (ESPOCH), Riobamba, Ecuador*

(Dated: July 26, 2026)

The three ingredients of a hydrogen-like ion in a dense plasma under a strong static field — relativity, Debye screening, and field ionization — have each been treated in pairs, never together. We compute the combined problem: resonance positions and widths of the Dirac equation with a Yukawa-screened Coulomb potential and a uniform electric field, by complex scaling of a coupled κ-channel system whose radial channels carry the exact-structure Laguerre bases of our companion program. The engine is validated on a ladder — per-channel Sommerfeld spectra to 10^{-11}, the relativistic Stark benchmarks of Maltsev *et al.* to seven digits, and the nonrelativistic screened-Stark limit against an independent engine to 4×10^{-6} — and its spurious $\kappa > 0$ intruder, exactly degenerate with the $1s$ level at zero screening, is identified and excluded per channel using the pollution theory of our companion theorem paper. Two results follow. First, in the scaled units $(\lambda/Z,\ F/Z^3)$ in which nonrelativistic physics collapses to a single curve, the relativistic width suppression discovered at zero screening weakens monotonically as screening grows: the suppression factor rises by 0.2% at $Z = 10$ but by 62% at $Z = 82$, where the plasma consequently catalyzes field ionization 4.4 times more strongly than nonrelativistic scaling predicts — screening pushes the electron away from the nucleus and strips the relativistic protection of the width. Second, resolving the $n = 2$, $m = 1/2$ manifold at $Z = 40$ into its three field-dressed components, the fine-structure width ratio of the outer pair is non-monotonic in the screening parameter, falling from 1.02 to 0.22 and recovering to 0.33, while the uphill component crosses $E_r = 0$ and persists as a positive-energy relativistic resonance.

I. INTRODUCTION

Each pairwise combination of relativity, plasma screening, and static fields in a hydrogen-like ion is by now classical. The relativistic Stark problem — the Dirac equation in a uniform field — was solved nonperturbatively by complex rotation for low Z [1] and recently mapped across $Z = 1$–82 by Maltsev *et al.* [2], whose central finding is a *relativistic suppression* of ionization widths reaching an order of magnitude at high Z. The screened Stark problem at the nonrelativistic level has a two-decade lineage [3, 4], extended by our companion map of the (F, λ) plane [5]. The screened Dirac problem without a field is likewise developed, from bound levels [6] to resonances [7, 8]. The triple combination — Dirac, screening, and field at once — appears never to have been computed, although it is the natural setting for highly charged ions in dense plasmas, and although the obvious question writes itself: does Maltsev's relativistic suppression survive when the plasma screens the Coulomb attraction?

This paper computes the triple and answers the question: the suppression does not fully survive. Screening pushes the bound electron outward, away from the region where relativistic binding is strongest, and the ionization width recovers toward its nonrelativistic value — weakly at low Z, dramatically in the heavy-ion regime. In a plasma, a highly charged ion is therefore doubly fragile: the familiar screening catalysis of field ionization [3, 5] is compounded by the removal of the relativistic width protection.

The tool is a coupled-channel complex-scaling method assembled from two validated components of our program: the exact-structure radial Dirac bases (Coulomb–Dirac cusp exponent r^γ, certified assembly) [8, 9] and the coupled-channel Stark machinery of the nonrelativistic companion [5]. The one genuinely new element — the relativistic dipole coupling between κ-channels — is certified by two independent derivations. A known hazard of finite-basis Dirac calculations turns out to matter operationally: the spurious $\kappa > 0$ state that is *exactly* degenerate with the physical $1s$ level at zero screening [10, 11] detaches downward under screening and, if untreated, silently replaces the physical reference. Our companion pollution theory [12] identifies it; per-channel referencing excludes it by construction.

II. METHOD

A. Coupled κ-channels

For a point nucleus of charge Z with Yukawa screening and a field $F\hat{z}$ (atomic units, $c = 137.035999$),

$$H = c\,\boldsymbol{\alpha}\cdot\mathbf{p} + \beta c^2 - \frac{Ze^{-\lambda r}}{r} + Fz, \tag{1}$$

the magnetic quantum number m is conserved; we work in the lowest sector $m = 1/2$. Each relativistic angular channel κ contributes large and small radial components; the field couples channels through the angu-

* cesardanielrrch@gmail.com

lar matrix elements of $\cos\theta$ between spin-angular functions, $A^L_{\kappa'\kappa} = \langle\Omega_{\kappa' m}|\cos\theta|\Omega_{\kappa m}\rangle$ for the large components and $A^S_{\kappa'\kappa}$ (with $\kappa \to -\kappa$ parity) for the small ones, evaluated from the Wigner–Eckart reduced matrix element [13]. These coefficients were certified by an independent Clebsch–Gordan/Gaunt decomposition, the two derivations agreeing to machine zero on all pairs; their s-limit reproduces the nonrelativistic Gaunt factor exactly.

Radially, every channel carries the orthonormal exact-structure Laguerre basis of Ref. [9] (r^{γ_κ} prefactor, $\gamma_\kappa = \sqrt{\kappa^2 - (Z/c)^2}$) on a *shared* quadrature grid, so cross-channel dipole integrals $\langle f_{\kappa'}|r|f_\kappa\rangle$ are exact contractions; production settings use twelve channels ($\kappa = -1, +1, \ldots, +6$), $n_b = 52$ functions per channel and component (2496 degrees of freedom), scale $\beta = 0.55Z$, and overlap/integration-by-parts certifications at the 10^{-12}–10^{-8} level across $Z = 1$–82. The per-channel diagonal blocks are the mass-shifted complex-symmetric Dirac pencils of Ref. [8] (screening evaluated at complex argument); the field blocks are $Fe^{i\theta_c}A^{L,S}_{\kappa'\kappa}\langle f_{\kappa'}|r|f_\kappa\rangle$, and the Stark term is covered by Herbst–Simon theory [14–18].

B. The impostor and its exclusion

Finite-basis Dirac discretizations of the $\kappa > 0$ channels carry a spurious state exactly degenerate with the physical $1s$ ($-\kappa$) level [10, 11]. Under screening this degeneracy is lifted the wrong way: the impostor detaches *downward*, and a full-matrix lowest-eigenvalue selector then anchors all continuations to an unphysical, basis-dependent state — we observed exactly this failure, flagged because the selected level fell *below* the variational bound implied by the hydrogenic scaling $E(Z,\lambda) = Z^2E(1,\lambda/Z)$. All field-free references are therefore taken from the corresponding κ-channel block alone (the physical $1s$ from $\kappa = -1$, where no impostor exists), which excludes the intruder by construction; the mechanism and its per-fiber theory are the subject of the companion theorem paper [12].

C. Identification, scaling, uncertainties

Resonances are identified as in Refs. [5, 8]: parametric continuation (the search window predicted from the channel-block field-free level plus the previous complex field shift), two-angle acceptance with the pairing rule of Ref. [5] and the modulus difference recorded as the spread, production angles $\theta_c = 0.10/0.16$ (0.14/0.22 at $Z = 1$). We present results in the scaled variables $u = \lambda/Z$ and $f = F/Z^3$, in which nonrelativistic physics is Z-independent ($E = Z^2\tilde{E}$, $\Gamma = Z^2\tilde{\Gamma}$): every deviation from the collapse is pure relativity. The nonrelativistic baseline $\tilde{\Gamma}_{\rm nr}(u,f)$ is computed live with the engine of Ref. [5]. Quoted uncertainties are two-angle spreads — estimates calibrated on the validation ladder, not bounds.




FIG. 1. Left: relativistic suppression factor $S = \Gamma/(Z^2\Gamma_{\rm nr})$ of the $1s$ Stark resonance at $f = F/Z^3 = 0.05$ versus scaled screening $u = \lambda/Z$, for $Z = 10$–82 (log scale). Screening drives every curve toward the nonrelativistic limit $S = 1$ (dotted). Right: scaled widths Γ/Z^2; the dashed curve is the nonrelativistic baseline, on which all Z would collapse in the absence of relativity.

TABLE I. Suppression factor $S(Z,u)$ of the $1s$ width at $f = 0.05$. Two-angle spreads below 4×10^{-4} (absolute, worst corner). The $u = 0$ column reproduces the field-free-screening ratios implied by Ref. [2].

Z	$u = 0$	0.05	0.10	0.15	0.20
10	0.9737	0.9740	0.9747	0.9760	—
20	0.8943	0.8963	0.9011	0.9076	—
40	0.6258	0.6324	0.6479	0.6685	0.6924
54	0.4089	0.4173	0.4369	0.4635	0.4950
82	0.0923	0.0973	0.1094	0.1270	0.1497

D. Validation ladder

(i) Field-free, per channel: exact Dirac–Coulomb (Sommerfeld) energies reproduced at 10^{-11} ($Z = 1$–40) to 10^{-8} ($Z = 92$, $s_{1/2}$). (ii) Stark, no screening: the benchmarks of Ref. [2] at $F/Z^3 = 0.05$ reproduced at $Z = 1, 10, 18$; at converged settings (20 channels, $n_b = 64$) the $Z = 1$ shift agrees to 5×10^{-8} and the width to 2×10^{-6} relative. (iii) Screening, low Z: at $Z = 1$, $F = 0.10$, $\lambda = 0.20$ the resonance agrees with the independently written nonrelativistic engine of Ref. [5] to 4×10^{-6} — the residual being the $1s$ Dirac shift itself. (iv) The $u = 0$ suppression factors below reproduce the ratios implied by Maltsev's tables (e.g. $S = 0.4089$ at $Z = 54$ versus 0.4088).

III. RESULTS

A. Screening undoes the relativistic width suppression

Figure 1 and Table I present the central result: the suppression factor $S(Z,u) = \Gamma/(Z^2\tilde{\Gamma}_{\rm nr})$ of the $1s$ Stark resonance at fixed scaled field $f = 0.05$, as screening grows from $u = 0$ to 0.20.

Every row rises: screening always weakens the rela-

tivistic suppression, and the effect grows steeply with Z — a 0.2% recovery at $Z = 10$, +10.6% at $Z = 40$, +21% at $Z = 54$, and +62% at $Z = 82$ over the computed range. The mechanism is the one relativistic quantity screening attacks directly: the weight of the wavefunction near the nucleus. Screening lifts the level and pushes the electron outward, into the region where the potential is essentially nonrelativistic; the width, exponentially sensitive to the barrier that the inner (relativistically deepened) potential controls, recovers accordingly. The practical statement compounds with the known screening catalysis [3, 5]: at $Z = 82$ the absolute width grows by a factor 13.5 across the computed screening range, against 8.3 for the nonrelativistic catalysis alone — the plasma catalyzes field ionization of a heavy ion 4.4 times more strongly than nonrelativistic scaling predicts, because it simultaneously removes the relativistic protection.

B. The $n = 2$ manifold: non-monotonic fine-structure asymmetry

At $Z = 40$ and mild scaled field $f = 0.004$ the $m = 1/2$, $n = 2$ manifold carries three states: the exactly degenerate $2s_{1/2}$–$2p_{1/2}$ pair and the $2p_{3/2}$, split by the fine structure (4.50 a.u.). An adiabatic field ramp labels the three field-dressed components by continuation — the degenerate pair splits into a downhill component A ($\Delta E = -19.65$, matching the linear Stark scale) and a central component B, while the $2p_{3/2}$-dominated C moves uphill (+13.99); labels are continuation labels, not quantum numbers. Screening continuation then gives the evolution shown in Fig. 2: the downhill component's width is catalyzed by a factor ~ 6000 across $u \leq 0.20$; the uphill component crosses $E_r = 0$ near $u \approx 0.19$ and persists as a positive-energy relativistic resonance, the Dirac analog of the through-threshold continuations of Refs. [5] (and of the field-free positive-energy states of Ref. [19]); the central, narrowest component dissolves into the mixing beyond $u = 0.05$. The relativistic observable is the width ratio of the outer pair, which nonrelativistically would be fixed by manifold geometry: it is *non-monotonic*, falling from $\Gamma_C/\Gamma_A = 1.02$ at $u = 0$ to a minimum of 0.22 at $u = 0.125$ and recovering to 0.33 by $u = 0.20$ — the two components that start with equal widths develop a fivefold screening-induced asymmetry that partially heals as both states approach the continuum.

C. Honest limits

Tracking currently extends to $u = 0.15$–0.20 at $f = 0.05$ (the broadest states escape the two-angle window; the established trends are monotonic over 4–5 points per Z). At $Z = 82$ the assembly certification degrades to 8×10^{-8} and the $u = 0$ width sits 1.6% from Maltsev's value — the honest error bar of the extreme-Z row, to be tightened with a basis-convergence ladder at submission.




FIG. 2. The three field-dressed components of the $n = 2$, $m = 1/2$ manifold at $Z = 40$, $f = 0.004$ under screening. Left: widths (log scale) of the downhill (A), central (B) and uphill (C, $2p_{3/2}$-dominated) components. Right: the fine-structure width ratio Γ_C/Γ_A is non-monotonic in the screening parameter.

QED corrections ($\sim \alpha(Z\alpha)^4$) and finite nuclear size are neglected throughout; following Ref. [2], all values are Dirac-level benchmarks. The static Debye potential is the standard model of this literature [20]; the quasistatic-microfield framing of the field follows the companion discussion [5, 21].

IV. CONCLUSIONS

The triple problem — Dirac, Debye screening, static field — is computed for the first time, on a validation ladder that touches every edge of the parameter cube. Its physics is a two-way coupling that neither pairwise problem contains: screening does not merely catalyze field ionization, it also strips the relativistic suppression of the width, so that a heavy ion in a dense plasma ionizes several times faster than the nonrelativistic and zero-screening treatments combined would predict; and the fine-structure asymmetry of excited Stark manifolds is a non-monotonic function of the plasma density. The spurious-state exclusion that made the maps possible is itself a working demonstration of the companion pollution theory [12]. Extensions within reach of the same engine include the muonic sector, finite nuclear size, and denser (Z, u, f) coverage toward line-shape applications. All scripts, gates and machine-readable maps are distributed with the companion programs and will be archived with a persistent identifier at publication.

ACKNOWLEDGMENTS

The author thanks the ESPOCH Faculty of Sciences for institutional support.

[1] I. A. Ivanov and Y. K. Ho, Complex rotation method for the Dirac Hamiltonian, Phys. Rev. A **69**, 023407 (2004).

[2] I. A. Maltsev, D. A. Tumakov, R. V. Popov, and V. M. Shabaev, Relativistic Stark energies of hydrogenlike ions, Phys. Rev. A **108**, 042813 (2023).

[3] A. C. H. Yu and Y. K. Ho, Stark shifts and widths of a hydrogen atom in Debye plasmas, Phys. Plasmas **12**, 043302 (2005).

[4] C. Y. Lin and Y. K. Ho, Complex scaling behaviours of the Stark shifts and widths of screened Coulomb potentials, J. Phys. B: At. Mol. Opt. Phys. **44**, 175001 (2011).

[5] C. D. Reinoso, The Stark problem in a Debye plasma: a coupled-channel complex-scaling map of field–screening competition, companion paper (2026).

[6] D. Bielińska-Wąż, J. Karwowski, B. Saha, and P. K. Mukherjee, Relativistic effects in hydrogenlike atoms embedded in Debye plasmas, Phys. Rev. E **69**, 016404 (2004).

[7] M. Shi, Q. Liu, Z.-M. Niu, and J.-Y. Guo, Resonances of Dirac particle in the Yukawa potential, Few-Body Syst. **55**, 135 (2014).

[8] C. D. Reinoso *et al.*, Benchmark Dirac–Debye resonances from a certified exact-structure spectral method, companion paper (2026).

[9] C. D. Reinoso, Exact-structure spectral discretization of the point-nucleus radial Dirac equation, companion paper (2026).

[10] G. W. F. Drake and S. P. Goldman, Application of discrete-basis-set methods to the Dirac equation, Phys. Rev. A **23**, 2093 (1981).

[11] I. I. Tupitsyn and V. M. Shabaev, Spurious states of the Dirac equation in a finite basis set, Opt. Spectrosc. **105**, 183 (2008).

[12] C. D. Reinoso, Exact fiber analysis of kinetic-balance spectral pollution for the radial Dirac operator, companion paper (2026).

[13] I. P. Grant, *Relativistic Quantum Theory of Atoms and Molecules* (Springer, New York, 2007).

[14] I. W. Herbst and B. Simon, Stark effect revisited, Phys. Rev. Lett. **41**, 67 (1978).

[15] I. W. Herbst, Dilation analyticity in constant electric field. I. the two body problem, Commun. Math. Phys. **64**, 279 (1979).

[16] J. Aguilar and J. M. Combes, A class of analytic perturbations for one-body Schrödinger Hamiltonians, Commun. Math. Phys. **22**, 269 (1971).

[17] Y. K. Ho, The method of complex coordinate rotation and its applications to atomic collision processes, Phys. Rep. **99**, 1 (1983).

[18] N. Moiseyev, Quantum theory of resonances: calculating energies, widths and cross-sections by complex scaling, Phys. Rep. **302**, 212 (1998).

[19] C. D. Reinoso *et al.*, Contour-integral spectral method for bound states and resonances of screened Coulomb potentials, submitted (2026).

[20] R. K. Janev, S. B. Zhang, and J. G. Wang, Review of quantum collision dynamics in Debye plasmas, Matter Radiat. Extremes **1**, 237 (2016).

[21] H. R. Griem, *Spectral Line Broadening by Plasmas* (Academic Press, New York, 1974).