


Collisions in isogeny graphs, and the security of the SIDH-based identification protocol

Federico Pintore

Department of Mathematics, University of Bari (IT)

Joint work with W. Ghantous (University of Oxford, UK), S. Katsumata (AIST, JP) and M. Veroni (NTNU, NO)



Tor Vergata - Roma - 30/11/2021

PREVIEW

SIDH is the the best-established isogeny-based cryptosystem.

An identification protocol ID_{SIDH} deduced from SIDH was turned into a digital signature scheme DS_{SIDH} .

The security of DS_{SIDH} is deduced from two properties of ID_{SIDH} :

- honest-verifier zero-knowledge
- **special soundness**.

We **dispute** the **correctness of the proofs**
for the **special soundness** in the literature.

ROADMAP

1. Digital Signatures & Identification Protocols
2. Post-quantum Cryptography, SIDH and ID_{SIDH}
3. Counterexamples to the special soundness of ID_{SIDH}
4. Collisions in isogeny graphs

DIGITAL SIGNATURES

A digital signature is a triple $DS = (\text{KeyGen}, \text{Sign}, \text{Verify})$ of PPT algorithms:


- $(vk, sk) \leftarrow \text{KeyGen}(\lambda)$: vk is the **verification key**, sk the **secret key**;
- $\sigma \leftarrow \text{Sign}(sk, m)$: it outputs a **signature** on input sk and a **message** m ;
- $1/0 \leftarrow \text{Verify}(m, \sigma, vk)$: it deterministically verifies σ (on m) w.r.t. vk .

SECURITY OF DIGITAL SIGNATURES

The standard security notion for digital signatures is **existential unforgeability**.

Challenger \mathcal{C}

Adversary \mathcal{A}



\mathcal{A} wins the game if: a) $m^* \notin Q$, b) $1 \leftarrow \text{Verify}(m^*, \sigma^*, vk)$.


DS is **existential unforgeable** if the winning probability of any \mathcal{A} is negligible in λ .

IDENTIFICATION PROTOCOLS

Given $R \subset X \times W$, an identification protocol for R

$$\text{ID} = (P = (P_1, P_2), V = (V_1, V_2))$$


is a **three-move interactive protocol** between a **prover** (holding a verification-secret key pair $(vk, sk) \in R$) and a **verifier** (holding vk).



SPECIAL SOUNDNESS OF AN ID

Required properties:

- Correctness
- Honest-Verifier Zero-Knowledge
- **Special Soundness**
- ...



There exists an **extractor** Ex that, **on input two valid transcripts** $(vk, com, ch, resp), (vk, com, ch', resp')$, **outputs** sk s.t. $(vk, sk) \in R$.

FROM AN ID TO A DIGITAL SIGNATURE

When ch varies in an exponential-size set, ID can be turned into a digital signature DS.

- Fiat-Shamir Transform
- Unruh Transform
- Fischlin Transform



$V_1(\text{com})$ is replaced with $H(m, \text{com})$,
where H is a **hash function**.

If ID satisfies **HVZK** and **special soundness**, and R is a **hard relation**
the obtained DS is **existential unforgeable**.

Proof by reduction:

- the adversary \mathcal{A} against the unforgeability game is run twice;
- **thanks to special soundness sk is extracted.**

ROADMAP

1. Identification Protocols & Digital Signatures



2. Post-quantum Cryptography, SIDH and ID_{SIDH}

3. Counterexamples to the special soundness of ID_{SIDH}

4. Collisions in isogeny graphs

POST-QUANTUM CRYPTOGRAPHY

In modern cryptography, security of cryptosystems must be formally proven (**provable security**).

The security proof of a public-key cryptosystem is given under the assumption that a **mathematical problem is hard** (e.g. security of DS obtained from ID).


Hard mathematical problems: integer factorisation, **ECDLP**.

Shor (1994): **quantum algorithms** to solve both problems in **polynomial time**.

Post-quantum Cryptography: public-key cryptosystems from **mathematical problems** (supposed to be) **hard** even **for quantum computers**.

ISOGENY-BASED CRYPTOGRAPHY

$p=457, \ell = 3$



Let p be a prime.

Vertices: supersingular elliptic curves over \mathbb{F}_{p^2} (modulo isomorphism)

Edges: isogenies over \mathbb{F}_{p^2} between elliptic curves (modulo equivalence)

Isogeny problem:
given two vertices,
find a path between them.

ON ELLIPTIC CURVES AND ISOGENIES - 1

Let $E : y^2 = x^3 + Ax + B$ be an elliptic curve over \mathbb{F}_q (i.e. $A, B \in \mathbb{F}_q$). Then

$$E(\overline{\mathbb{F}}_q) = \{(x_0, y_0) \in \overline{\mathbb{F}}_q^2 \mid y_0^2 = x_0^3 + Ax_0 + B\} \cup \{\infty\}$$

is an abelian group.

An isogeny $\varphi : E_0 \rightarrow E_1$ is **non-constant morphism** which sends ∞ in ∞ .



$$\varphi(x, y) \mapsto (f_1(x, y)/f_2(x, y), g_1(x, y)/g_2(x, y))$$

with $f_1, f_2, g_1, g_2 \in \overline{\mathbb{F}}_q[x, y]$

$\deg(\varphi)$ is the degree of φ as a morphism. The degree is **multiplicative** w.r.t. \circ (comp.)

ON ELLIPTIC CURVES AND ISOGENIES - 2

Isomorphisms are isogenies of degree 1, which preserve j -invariants

$$j(E) = 1728 \frac{4A^3}{4A^3 + 27B^2}$$

An endomorphism of E is an isogeny $\varphi : E \rightarrow E$. $(\text{End}(E), +, \circ)$ is a **ring**.

$(\text{End}(E), +, \circ)$ is isomorphic to either an **order** in a quadratic field or a maximal order in a quaternion algebra. In the latter case, E is **supersingular**.

The number of points of E is predictable

ON ELLIPTIC CURVES AND ISOGENIES -

An isogeny $\varphi : E_0 \rightarrow E_1$ admits a dual $\hat{\varphi} : E_1 \rightarrow E_0$ s.t. $\varphi \circ \hat{\varphi} = \hat{\varphi} \circ \varphi = [\deg(\varphi)]$.

Given $G \leq E(\overline{\mathbb{F}}_q)$, there exists an isogeny $\varphi : E \rightarrow E'$ s.t. $\ker(\varphi) = G$.

- E' is denoted by E/G
- E/G and φ are **unique** modulo isomorphism and equivalence, resp.

If $(\ell, q) = 1$, $E[\ell] = \{P \in E(\overline{\mathbb{F}}_q) \mid [\ell]P = \infty\} \simeq \mathbb{Z}_\ell \times \mathbb{Z}_\ell$.

THE SIDH SETTING

- a prime $p = \ell_1^{e_1} \ell_2^{e_2} \pm 1$ (ℓ_1, ℓ_2 small primes)
- a supersingular elliptic curve E_0 over \mathbb{F}_{p^2}
- P_1, Q_1 s.t. $\langle P_1, Q_1 \rangle = E_0[\ell_1^{e_1}]$
- P_2, Q_2 s.t. $\langle P_2, Q_2 \rangle = E_0[\ell_2^{e_2}]$

$$\#E_0(\mathbb{F}_{p^2}) = (\ell_1^{e_1} \ell_2^{e_2})^2$$

Alice

Bob

Samples $m_1 \in \mathbb{Z}_{\ell_1^{e_1}}$

Computes $\varphi_A : E_0 \rightarrow E_A = E_0 / \langle P_1 + [m_1]Q_1 \rangle$

$$\xrightarrow{E_A, \varphi_A(P_2), \varphi_A(Q_2)}$$

Samples $m_2 \in \mathbb{Z}_{\ell_2^{e_2}}$

Computes $\varphi_B : E_0 \rightarrow E_B = E_0 / \langle P_2 + [m_2]Q_2 \rangle$

$$\xleftarrow{E_B, \varphi_B(P_1), \varphi_B(Q_1)}$$

$$E_B / \langle \varphi_B(P_1) + [m_1]\varphi_B(Q_1) \rangle \simeq E_A / \langle \varphi_A(P_2) + [m_2]\varphi_A(Q_2) \rangle$$

THE IDENTIFICATION PROTOCOL ID_{SIDH}

$$X = \{(E_1, P', Q') \mid E_1 \text{ sup.} \wedge \langle P', Q' \rangle = E_1[\ell_2^{e_2}]\}$$

$$Y = \{\varphi \mid \varphi \text{ cyclic isog.}\}$$

$$R = \{((E_1, P', Q'), \varphi) \mid \varphi : E_0 \rightarrow E_1 \wedge \deg(\varphi) = \ell_1^{e_1} \wedge \varphi(P_2) = P', \varphi(Q_2) = Q'\}$$

Prover

$$(vk = (E_1, P', Q'), sk = \varphi)$$

$$m_2 \in \mathbb{Z}_{\ell_2^{e_2}}$$


$$rsp = \begin{cases} m_2 & \text{if } ch = 0 \\ \phi(\ker(\varphi)) & \text{if } ch = 1 \end{cases}$$

Verifier


$$vk = (E_1, P', Q')$$

$$ch \leftarrow \{0, 1\}$$

$$1/0 \leftarrow V_2(com, ch, rsp)$$



(CLAIMED) SPECIAL SOUNDNESS OF ID_{SIDH}





Two valid transcripts give the isogeny $\hat{\phi}' \circ \psi \circ \phi$ between E_0 and E_1

In four papers, the special soundness of ID_{SIDH} is proven by means of the extractor

$$\begin{aligned} \hat{\phi}(T) &\leftarrow \text{EX}_{\text{SIDH}}(\phi, \psi, \phi') \\ &\parallel \\ \ker \left(\hat{\phi}' \circ \psi \circ \phi \right) \cap E_0[\ell_1^{e_1}] \end{aligned}$$

ROADMAP

1. Identification Protocols & Digital Signatures 
2. Post-quantum Cryptography, SIDH and ID_{SIDH} 
3. Counterexamples to the special soundness of ID_{SIDH}
4. Collisions in isogeny graphs


COUNTEREXAMPLES TO SPECIAL SOUNDNESS

Let

$$\psi : E_2 = E_0 / \langle P_2 + [m_2]Q_2 \rangle \rightarrow E_3 = E_1 / \langle P' + [m_2]Q' \rangle$$

be the isogeny with kernel $\phi(\ker(\varphi))$

Suppose there exists $\tilde{\psi} : E_2 \rightarrow E_3$ cyclic,
non equivalent to ψ , with $\ker(\tilde{\psi}) = \langle \tilde{T} \rangle$
and $\deg(\tilde{\psi}) = \ell_1^{e_1}$.



$((E_1, P', Q'), (E_2, E_3), 1, \tilde{T})$ is a **valid transcript!**

On input $(\phi, \tilde{\psi}, \phi')$, Ex_{SIDH} does not output
a **valid secret key** for (E_1, P', Q')

CONCRETE COUNTEREXAMPLES TO SPECIAL SOUNDNESS - 1

The scenario described in the previous slide is **not** only **theoretical**.

We obtained a **concrete instance** for the biggest set of parameters for SIDH, i.e. p_{751} .

The instance considers $E_2 = E_0$, with $j(E_0) = 0$, for which $\text{End}(E_0)$ is known.

The alternative isogeny $\tilde{\psi}$ is found by looking for
a **cyclic endomorphism of degree $\ell_1^{2e_1}$**




This corresponds to the resolution of a norm equation in the quaternion algebra.

COUNTEREXAMPLES TO SPECIAL SOUNDNESS

Theorem (Ghantous, Katsumata, _ , Veroni - 2021)

The inputs that make the extractor Ex_{SIDH} fail are precisely those that fall within the framework we described.

ROADMAP

1. Identification Protocols & Digital Signatures 
2. Post-quantum Cryptography, SIDH and ID_{SIDH} 
3. Counterexamples to the special soundness of ID_{SIDH} 
4. Collisions in isogeny graphs

MITIGATIONS FOR DS_{SIDH}

Replace special soundness with **relaxed special soundness**.

A bigger relation \tilde{R} , with $R \subseteq \tilde{R}$, is considered. The extractor Ex is only required to extract sk such that $(vk, sk) \in \tilde{R}$.

As long as \tilde{R} is a hard relation, the digital signature from ID is existential unforgeable.

$$\tilde{R} = \{((E_1, P', Q'), \varphi) \mid \varphi : E_0 \rightarrow E_1\}$$

The problem of computing **any** isogeny between E_0 and E_1 is supposed to be hard even for quantum computers

QUANTIFYING COLLISIONS IN ISOGENY GRAPHS

Given the distinct primes p, ℓ and $e \in \mathbb{N}$, we call **collision in $\mathcal{G}_{p^2}(\ell)$** any pair of non-equivalent cyclic isogenies $\psi, \tilde{\psi} : E \rightarrow E_1$ with $\deg(\psi) = \deg(\tilde{\psi}) = \ell^e$.

We denote by **$\text{Coll}_{\ell^e}(E)$** the number of such collisions originating from the curve E

Collisions in $\mathcal{G}_{p^2}(\ell)$ are related to endomorphisms of degree ℓ^{2e} , which are quantified by the **Brandt matrix of degree ℓ^{2e}** .

We denote by **$\mathcal{C}_E(\ell^{2e})$** the number of cyclic endomorphisms of E .

QUANTIFYING COLLISIONS IN ISOGENY GRAPHS

Lemma (Ghantous, Katsumata, _ , Veroni - 2021)

$$\mathcal{C}_E(\ell^{2e}) \leq \text{Coll}_{\ell^e}(E) \leq \mathcal{C}_E(\ell^{2e}) + \sum_{r=1}^{e-1} \mathcal{C}_E(\ell^{2r})(\ell - 1)\ell^{e-1-r}$$

Let n be the number of vertices of $\mathcal{G}_{p^2}(\ell)$, which is approximately $p/12$.

Lemma (Ghantous, Katsumata, _ , Veroni - 2021)

$$\sum_{i=1}^n \mathcal{C}_{E(i)}(\ell^{2e}) \leq \frac{\ell^{2e+1}}{\ell - 1}$$

QUANTIFYING COLLISIONS IN ISOGENY GRAPHS

Theorem (Ghantous, Katsumata, _ , Veroni - 2021)

$$\sum_{i=1}^n \text{Coll}_{\ell^e}(E_{(i)}) \leq \frac{\ell^{2e}(\ell + 1)}{\ell - 1}.$$

Corollary (Ghantous, Katsumata, _ , Veroni - 2021)

$$\mathbb{E}_E[\text{Coll}_{\ell^e}(E)] := \frac{1}{n} \sum_{i=1}^n \text{Coll}_{\ell^e}(E_{(i)}) \leq \frac{\ell^{2e}(\ell + 1)}{n(\ell - 1)}.$$

When $p \approx \ell^{2e}$ (SIDH setting), the upper bound of the above expectation is in $\mathcal{O}(1)$.

QUANTIFYING COLLISIONS IN ISOGENY GRAPHS

Obtaining lower bounds is trickier, as it involves incomplete character sums.

By considering a statistical model which makes use of Bernoulli random variables we obtained the following.

Theorem (Ghantous, Katsumata, _ , Veroni - 2021)

$$\frac{1}{4n} \ell^{e-1} (\ell + 1) (2\ell^e - 1) \leq \mathbb{E}_E(\text{Coll}_{\ell^e}(E))$$

When $p \approx \ell^{2e}$ (SIDH setting), the lower bound of the above expectation is in $\mathcal{O}(1)$.

Thanks for your attention

Federico Pintore

Department of Mathematics, University of Bari (IT)

federico.pintore@uniba.it