- nonpolyposis colorectal cancer syndrome. Cancer Res., 62,
- 14. Wu,Y., Berends,M.J., Post,J.G., Mensink,R.G., Verlind,E., van der Sluis, T., Kempinga, C., Sijmons, R.H., van der Zee, A.G., Hollema.H. et al. (2001) Germline mutations of EXO1 gene in patients with hereditary nonpolyposis colorectal cancer (HNPCC) and atypical HNPCC forms. Gastroenterology, 120, 1580-1587.
- 15. Schaetzlein, S., Kodandaramireddy, N.R., Ju, Z., Lechel, A., Stepczynska, A., Lilli, D.R., Clark, A.B., Rudolph, C., Kuhnel, F., Wei, K. et al. (2007) Exonuclease-1 deletion impairs DNA damage signaling and prolongs lifespan of telomere-dysfunctional mice. Cell, 130, 863-877.
- 16. Bolderson, E., Richard, D.J., Edelmann, W. and Khanna, K.K. (2009) Involvement of Exo1b in DNA damage-induced apoptosis. Nucleic Acids Res., 37, 3452-3463.
- 17. Tomimatsu, N., Mukherjee, B. and Burma, S. (2009) Distinct roles of ATR and DNA-PKcs in triggering DNA damage responses in ATM-deficient cells. EMBO Rep., 10, 629-635.
- 18. Richard, D.J., Bolderson, E., Cubeddu, L., Wadsworth, R.I., Savage, K., Sharma, G.G., Nicolette, M.L., Tsvetanov, S., McIlwraith, M.J., Pandita, R.K. et al. (2008) Single-stranded DNA-binding protein hSSB1 is critical for genomic stability. Nature, **453**, 677–681.
- 19. Uematsu, N., Weterings, E., Yano, K., Morotomi-Yano, K., Jakob, B., Taucher-Scholz, G., Mari, P.O., van Gent, D.C., Chen, B.P. and Chen, D.J. (2007) Autophosphorylation of DNA-PKCS regulates its dynamics at DNA double-strand breaks. J. Cell Biol., 177, 219-229.
- 20. Khanna, K.K., Keating, K.E., Kozlov, S., Scott, S., Gatei, M., Hobson, K., Tava, Y., Gabrielli, B., Chan, D., Lees-Miller, S.P. et al. (1998) ATM associates with and phosphorylates p53: mapping the region of interaction. Nat. Genet., 20, 398-400.
- 21. Young, D.B., Jonnalagadda, J., Gatei, M., Jans, D.A., Meyn, S. and Khanna, K.K. (2005) Identification of domains of ataxia-telangiectasia mutated required for nuclear localization and chromatin association. J. Biol. Chem., 280, 27587-27594.
- 22. Li,Y., Bolderson,E., Kumar,R., Muniandy,P.A., Xue,Y., Richard, D., Seidman, M., Pandita, T.K., Khanna, K.K. and Wang, W. (2009) hSSB1 and hSSB2 form similar multi-protein complexes that participate in DNA damage response. \hat{J} . Biol. Chem., 284, 23525-23531.
- 23. Gupta, A., Sharma, G.G., Young, C.S., Agarwal, M., Smith, E.R., Paull, T.T., Lucchesi, J.C., Khanna, K.K., Ludwig, T. and Pandita, T.K. (2005) Involvement of human MOF in ATM function. Mol. Cell Biol., 25, 5292-5305.
- 24. Gravel, S., Chapman, J.R., Magill, C. and Jackson, S.P. (2008) DNA helicases Sgs1 and BLM promote DNA double-strand break resection. Genes Dev., 22, 2767-2772.
- 25. Rogakou, E.P., Pilch, D.R., Orr, A.H., Ivanova, V.S. and Bonner, W.M. (1998) DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. J. Biol. Chem., 273, 5858-5868
- 26. Chen, H.T., Bhandoola, A., Difilippantonio, M.J., Zhu, J., Brown, M.J., Tai, X., Rogakou, E.P., Brotz, T.M., Bonner, W.M.,

- Ried, T. et al. (2000) Response to RAG-mediated VDJ cleavage by NBS1 and gamma-H2AX. Science, 290, 1962-1965.
- 27. Peterson, C.L. (2001) Chromatin: mysteries solved? Biochem. Cell Biol., 79, 219-225.
- 28. Litman, R., Peng, M., Jin, Z., Zhang, F., Zhang, J., Powell, S., Andreassen, P.R. and Cantor, S.B. (2005) BACH1 is critical for homologous recombination and appears to be the Fanconi anemia gene product FANCJ. Cancer Cell, 8, 255-265.
- 29. Pierce, A.J., Johnson, R.D., Thompson, L.H. and Jasin, M. (1999) XRCC3 promotes homology-directed repair of DNA damage in mammalian cells. Genes Dev., 13, 2633-2638.
- 30. Zhou, B.B. and Elledge, S.J. (2000) The DNA damage response: putting checkpoints in perspective. Nature, 408, 433-439.
- 31. Mimitou, E.P. and Symington, L.S. (2008) Sae2, Exo1 and Sgs1 collaborate in DNA double-strand break processing. Nature, 455, 770-774
- 32. Zhu, Z., Chung, W.H., Shim, E.Y., Lee, S.E. and Ira, G. (2008) Sgs1 helicase and two nucleases Dna2 and Exo1 resect DNA double-strand break ends. Cell, 134, 981-994.
- 33. Bakkenist, C.J. and Kastan, M.B. (2003) DNA damage activates ATM through intermolecular autophosphorylation and dimer dissociation. Nature, 421, 499-506.
- 34. Buis, J., Wu, Y., Deng, Y., Leddon, J., Westfield, G., Eckersdorff, M., Sekiguchi, J.M., Chang, S. and Ferguson, D.O. (2008) Mre11 nuclease activity has essential roles in DNA repair and genomic stability distinct from ATM activation. Cell, 135, 85-96.
- 35. Morin, I., Ngo, H.P., Greenall, A., Zubko, M.K., Morrice, N. and Lydall, D. (2008) Checkpoint-dependent phosphorylation of Exol modulates the DNA damage response. EMBO J., 27, 2400-2410.
- 36. Matsuoka, S., Ballif, B.A., Smogorzewska, A., McDonald, E.R. III, Hurov, K.E., Luo, J., Bakalarski, C.E., Zhao, Z., Solimini, N., Lerenthal, Y. et al. (2007) ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. Science, 316, 1160-1166.
- 37. van der Horst, A. and Khanna, K.K. (2009) The peptidyl-prolyl isomerase Pin1 regulates cytokinesis through Cep55. Cancer Res., **69**, 6651-6659.
- 38. Nimonkar, A.V., Ozsoy, A.Z., Genschel, J., Modrich, P. and Kowalczykowski, S.C. (2008) Human exonuclease 1 and BLM helicase interact to resect DNA and initiate DNA repair. Proc. Natl Acad. Sci. USA, 105, 16906-16911.
- 39. El-Shemerly, M., Hess, D., Pyakurel, A.K., Moselhy, S. and Ferrari,S. (2008) ATR-dependent pathways control hEXO1 stability in response to stalled forks. Nucleic Acids Res., 36,
- 40. Di Virgilio, M., Ying, C.Y. and Gautier, J. (2009) PIKK-dependent phosphorylation of Mrel1 induces MRN complex inactivation by disassembly from chromatin. DNA Repair, 8, 1311-1320.
- 41. Beucher, A., Birraux, J., Tchouandong, L., Barton, O., Shibata, A., Conrad,S., Goodarzi,A.A., Krempler,A., Jeggo,P.A. and Lobrich, M. (2009) ATM and Artemis promote homologous recombination of radiation-induced DNA double-strand breaks in G2. EMBO J., 28, 3413-3427.