

# Stereometry

Maxim Akhmedov  
Yandex

September 28th, 2018

This paper contains basic definitions, facts and formulas related to the 3d geometry useful for programming competitions. Almost no algorithms nor geometrical data structures included.

## 1 Basic definitions

A 3d *point* is defined by its coordinates  $(x, y, z)$ .

A 3d *vector* is also defined by its coordinates  $\vec{v} = (x, y, z)$ . We will identify points and vectors, that allows us to add vector to a point without writing extra classes and methods.

Let  $\vec{v}_1 = (x_1, y_1, z_1)$  and  $\vec{v}_2 = (x_2, y_2, z_2)$ . Then:

- $\vec{v}_1 + \vec{v}_2 = (x_1 + x_2, y_1 + y_2, z_1 + z_2)$  is the sum of vectors;
- $\vec{v}_1 - \vec{v}_2 = (x_1 - x_2, y_1 - y_2, z_1 - z_2)$  is the difference of vectors;
- $k\vec{v}_1 = \vec{v}_1 k = (kx_1, ky_1, kz_1)$  is the vector multiplied by the real value  $k$ ;
- $\vec{v}_1 \cdot \vec{v}_2 = (v_1, v_2) = x_1x_2 + y_1y_2 + z_1z_2$  is the *scalar* product of vectors;

As well as in the 2d case, the operations above have intuitive geometrical meaning. Sum of vectors may be find by triangle or parallelogram rule, the difference of vectors is a vector connecting endpoints of two vectors.

For vector  $\vec{v} = (x, y, z)$  its Euclidean norm is defined as  $\sqrt{x^2 + y^2 + z^2}$ . If  $\vec{v} \neq \vec{0}$ ,  $\vec{v}/|\vec{v}|$  always has a unit norm (i.e. 1).

Scalar product  $(\vec{u}, \vec{v})$  has the following geometrical meaning: if  $u \neq 0$ ,  $(\vec{u}, \vec{v}) = \text{proj}_{\vec{u}} \vec{v} \times |u|$  where  $\text{proj}_{\vec{u}} \vec{v}$  is the signed projection length of vector  $\vec{v}$  on the vector  $\vec{u}$ . In particular, when  $\vec{u}$  has unit norm,  $(\vec{u}, \vec{v})$  is a length of a component of vector  $\vec{v}$  that is parallel to  $\vec{u}$ .

The geometrical meaning of  $\text{sgn}(\vec{u}, \vec{v})$  for non-zero vectors: if it is zero,  $\vec{u}$  and  $\vec{v}$  are orthogonal, if it is positive, the angle between them is acute, otherwise it is obtuse. If one of vectors is a zero-vector, scalar product is always zero.

The following equation is held:

$$(\vec{u}, \vec{v}) = |\vec{v}||\vec{u}| \cos \angle(\vec{u}, \vec{v})$$

## 2 Vector and mixed products

A vector product of  $\vec{v}_1$  and  $\vec{v}_2$  is

$$[\vec{v}_1, \vec{v}_2] = \begin{vmatrix} \vec{e}_x & \vec{e}_y & \vec{e}_z \\ x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \end{vmatrix} = (y_1z_2 - y_2z_1, z_1x_2 - z_2x_1, x_1y_2 - x_2y_1)$$

Note that in 3d case the vector product is a vector!

$||\vec{v}_1, \vec{v}_2||$  is an area of a parallelogram built by vector  $\vec{v}_1$  and  $\vec{v}_2$ . The following equation is held:

$$||\vec{v}_1, \vec{v}_2|| = |v||u| \sin \angle(\vec{u}, \vec{v})$$

Vector product is always orthogonal to both  $\vec{v}_1$  and  $\vec{v}_2$  (it may be proven by taking scalar product with any of  $\vec{v}_1$  and  $\vec{v}_2$ ).

A mixed product  $(\vec{v}_1, \vec{v}_2, \vec{v}_3)$  is  $(\vec{v}_1, [\vec{v}_2, \vec{v}_3])$ . Also, by definition:

$$(\vec{v}_1, \vec{v}_2, \vec{v}_3) = (x_1, y_1, z_1) \cdot \begin{vmatrix} \vec{e}_x & \vec{e}_y & \vec{e}_z \\ x_2 & y_2 & z_2 \\ x_3 & y_3 & z_3 \end{vmatrix} = \begin{vmatrix} x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \\ x_3 & y_3 & z_3 \end{vmatrix}$$

Geometric meaning of a mixed product is a signed volume of a parallelepiped built by  $v_1, v_2, v_3$  as a edges. A tetrahedron volume is 6 times smaller than the parallelepiped volume.

The  $\text{sgn}(\vec{u}, \vec{v}, \vec{w})$  has several geometric meanings:

- if  $\vec{u}$  is a fixed vector, then  $\text{sgn}(\vec{u}, \vec{v}, \vec{w})$  is zero if  $\vec{v}, \vec{w}$  and  $\vec{u}$  lie in the same plane, is positive if  $\vec{w}$  is counter-clockwise from  $\vec{v}$  when watching straight at the end of  $\vec{u}$  (in direction of  $-\vec{u}$ ), and negative otherwise;
- $\text{sgn}(\vec{u}, \vec{v}, \vec{w})$  depends on which half-space  $w$  belongs in respect to the plane containing  $\vec{u}$  and  $\vec{v}$ .

Let  $\text{vol}(A, B, C, D) = \text{vol } ABCD = (\vec{AB}, \vec{AC}, \vec{AD})$ .

An angle between vectors  $\vec{u}$  and  $\vec{v}$  may be found using the formula:

$$\angle(\vec{u}, \vec{v}) = \text{atan2}(|[\vec{u}, \vec{v}]|, (\vec{u}, \vec{v}))$$

Note that, contrary to 2d case, this angle is always between 0 and  $\pi$  and may not be treated as a signed angle (in particular, there is no such notion as clockwise or counter-clockwise when we talk about arbitrary angle in the plane).

**Exercise:** given a line  $AB$  and  $n$  points  $p_1, \dots, p_n$  that belong to one half-space not containing the line  $AB$ , sort these points by their angle around the line  $AB$  (use the sign of mixed product as a comparison predicate, its similar to using vector product sign in 2d case).

### 3 Planes

The plane equation:  $Ax + By + Cz = D$  where at least one of  $A, B$  and  $C$  is not zero. This is equivalent to the following:  $(A, B, C) \cdot (x, y, z) = D$  that shows that  $(A, B, C)$  is a *normal* vector to the plane (i.e. vector orthogonal to the plane). Indeed, for any two points  $p_1$  and  $p_2$  belonging to the plane  $(A, B, C)(\vec{p}_1 - \vec{p}_2) = D - D = 0$ .

The *signed distance* from the point  $(x_0, y_0, z_0)$  to the plane  $Ax + By + Cz = D$  is  $\frac{Ax_0 + By_0 + Cz_0 - D}{\sqrt{A^2 + B^2 + C^2}}$ ; it is zero if  $(x_0, y_0, z_0)$  belongs to the plane, it is positive if  $(x_0, y_0, z_0)$  lies in the half-space in the direction of vector  $(A, B, C)$ , and it is negative otherwise.

Plane may also be defined by three non-collinear points on it. If  $A, B$  and  $C$  belong to the plane, then normal vector may be restored as  $\vec{n} = [\vec{AB}, \vec{AC}]$ .

## 4 Lines, line division rule

The easiest way to define the line in 3d space is the choose two points on it.

The barycentric line division rule: if  $X$  belongs to the line  $AB$ , then  $\vec{X} = \frac{\vec{AX}}{\vec{AB}} \vec{B} + \frac{\vec{XB}}{\vec{AB}} \vec{A}$ . Here  $\frac{\vec{u}}{\vec{v}}$  is the proportion between two vectors belonging to the same line: it is  $\frac{|u|}{|v|}$  if they have the same direction and  $-\frac{|u|}{|v|}$  otherwise. Note that  $\frac{\vec{AX}}{\vec{AB}} + \frac{\vec{XB}}{\vec{AB}} = \frac{\vec{AX} + \vec{XB}}{\vec{AB}} = \frac{\vec{AB}}{\vec{AB}} = 1$ .

Equivalent statement: if  $\lambda$  is a real number, then  $\lambda \vec{B} + (1 - \lambda) \vec{A} = \vec{A} + \lambda(\vec{B} - \vec{A})$  is always a point of the line  $AB$  that divides segment  $AB$  in proportion of  $\lambda : (1 - \lambda)$ .

Barycentric line division rule is a useful thing. We will illustrate how to apply it to intersect line with a plane.

Suppose you have a line  $AB$  and plane  $CDE$ . Then their intersection point  $O$  may be found using the following formula:

$$O = \frac{\text{vol } CDEB \cdot \vec{A} + \text{vol } ACDE \cdot \vec{B}}{\text{vol } CDEB + \text{vol } ACDE}$$

This is true because  $\text{vol } CDEB : \text{vol } ACDE$  is equal to the ratio of signed distances between  $B$  and  $CDE$  and between  $A$  and  $CDE$ , which is equal to the ratio in which plane  $CDE$  divides the segment (line)  $AB$ .

If the plane is defined by a normal vector  $\vec{n}$  (and the equation  $(\vec{n}, (x, y, z)) = D$ ), then the intersection of a line  $AB$  and plane may also be found by using a barycentric rule:

$$O = \frac{((\vec{n}, \vec{B}) - D) \vec{A} + (D - (\vec{n}, \vec{A})) \vec{B}}{(\vec{n}, \vec{AB})}$$

This is true because  $(\vec{n}, \vec{B}) - D$  is also proportional to the signed distance between  $B$  and  $CDE$  (with a coefficient of  $|\vec{n}|$ ) and similar for the second vector.

**Exercise:** write down similar formula for the 2d case and the line intersection.

Now we will discuss the following problem: you are given two lines in 3d space, find the distance between them. The full solution is now missing in this text, but it may eventually get here.

## 5 Polytopes

Polytope may be define by specifying its faces. Each face is a polygone located in some plane. Vertices are connected with edges, each edge is adjacent to exactly two faces.

**Euler Formula:** for a polytope that has at least three non-collinear points,  $V - E + F = 2$  where  $V$  is the number of vertices,  $E$  is the number of edges and  $F$  is the number of faces.

**Lemma:**  $E \leq 3V - 6$ . Proof is left as an exercise.

These facts are enough to show that for any two numbers  $x$  and  $y$  among  $V, E, F$  one may bound  $x = \mathcal{O}(y)$ , for example, number of edges and number of faces are both linear in terms of number of vertices.

In order to calculate the volume of a polytope, one should express it as a sum of several pyramids, some of which will be included with a negative sign (as well as in triangle formula for calculating polygon area in 2d case). Suppose that all faces are oriented counter-clockwise. Then  $\text{vol } P = \sum_{\text{face } f} \text{vol}(O, f)$  where  $\text{vol}(O, f)$  is the area of a pyramid with its vertex at  $O$  and

its base in  $f$ . Volume of pyramid may also be found as a sum of volumes of several tetrahedrons forming the pyramid. All volumes are considered to be signed.

If not all faces are oriented counter-clockwise, you may orient each face  $f$  such that the signed volume  $\text{vol}(p_i, f)$  is positive where  $p_i$  is any vertex of the polygon non-belonging to  $f$ .

Polytope may also be defined as a convex hull of a given set. A naive approach to find all faces of a polytope defined as a convex hull is to try each of the possible  $\Theta(n^3)$  triples of points, find a plane passing through them and then check that all other points lie by one side from it, resulting in a  $\Theta(n^4)$  algorithm.

More efficient approach is to perform a gift wrapping: suppose you have an edge  $v_i v_j$  of a convex hull. Sort all points around the line  $v_i v_j$ , then find two outermost points  $p_a$  and  $p_b$  in this order. They yield two faces  $v_i v_j v_a$  and  $v_i v_j v_b$  and four new edges. Continue considering edges in a DFS or BFS like procedure. Such an algorithm works in  $\mathcal{O}(n^2)$ .

There are much more complicated algorithms for finding convex hull in  $\mathcal{O}(n \log n)$  or even  $\mathcal{O}(n \log h)$  where  $h$  is the number of points in a convex hull, but we will not discuss them.

## 6 Geometry of masses

Center of mass of a set of a finite set of points  $p_i$  associated with the masses  $w_i$  may be found using the formula  $(\sum_i p_i w_i) / (\sum_i w_i)$ . Center of mass of a solid tetrahedron  $ABCD$  is located in a point  $(A + B + C + D)/4$  (as well as if we had only 4 points  $A, B, C, D$ ).

When dealing with points with masses, you may replace a system of points (or a solid body like tetrahedron) with their center of mass, associating a total mass of replaced points with it. For example, you may replace a tetrahedron  $ABCD$  with a point  $(A + B + C + D)/4$  of mass  $\text{vol } ABCD$ .

## 7 Spherical geometry

The main idea: keep sphere radius  $r = 1$  and use Euclidean coordinate system, that makes everything simpler.

Suppose  $\vec{v} = (x, y, z)$ ,  $|\vec{v}| = 1$ . The spherical coordinates are: latitude  $\psi$  and longitude  $\phi$ .

Spherical coordinates  $\rightarrow$  Euclidean coordinates:

$$z = \sin \psi$$

$$x = \cos \psi \cos \phi$$

$$y = \cos \psi \sin \phi$$

Euclidean coordinates  $\rightarrow$  spherical coordinates:

$$\psi = \text{atan2}(z, \sqrt{x^2 + y^2})$$

$$\phi = \text{atan2}(y, x)$$

Distance on sphere between points with radius-vectors  $\vec{u}$  and  $\vec{v}$  is  $d(\vec{u}, \vec{v}) = \angle(\vec{u}, \vec{v}) = \text{atan2}(|[\vec{u}, \vec{v}]|, \vec{u} \cdot \vec{v})$ .

Arc intersection on sphere: if  $A_1 B_1$  and  $A_2 B_2$  are two arcs on the sphere:

$$\vec{n}_1 = [\vec{A}_1, \vec{B}_1], \quad \vec{n}_2 = [\vec{A}_2, \vec{B}_2], \quad \vec{X} = [\vec{n}_1, \vec{n}_2] / |\vec{n}_1, \vec{n}_2|$$