



## DIGITAL SIGNAL PROCESSING LAB

Debagnik Kar

Roll: 1804373, ETC -06, EC - 3096

## Index Page

Experiment No.	Aim of the Experiment	Page No.	Date of Experiment	Date of Submission
01	To write a function for finding circular convolution and correlation. Prove that autocorrelation is an even function and $R_{xy}(n) = R_{yx}(-n)$	2-6	23/12/2020	25/12/2020
02	To solve a given differential equation to find its impulse and step responses. Plot the response and verify the solution	7-9	06/01/2021	13/01/2021
03	To find the DTFT of an arbitrary sequence and prove the convolution property	10-	13/01/2021	20/01/2021
04	To write a function to find DFT and IDFT and use it to find circular and linear convolution of given sequences.			
05	To perform sectional convolution using overlap add and overlap save method.			
06	To design FIR LPF, HPF and BPF using windowing techniques.			
07	To Study various parameters required for designing an IIR filter and design LPF, HPF and BPF Butterworth filters for specific cut-off frequencies.			
08	To study up sampling and down sampling of the sequences and study the frequency response			
09	Open Ended – I			
10	Open Ended - II			

<b>Experiment Number</b>	03
<b>Date of Experiment</b>	13/01/2021
<b>Date of Submission</b>	20/01/2021
<b>Name of the Student</b>	Debagnik Kar
<b>Roll Number</b>	1804373
<b>Section</b>	ETC - 06

### **Aim of The Experiment :-**

To find the DTFT of an arbitrary sequence and prove the convolution property


### **Software Required :-**

- MATLAB R2018a

### **Theory :-**

In mathematics, the discrete-time Fourier transform (DTFT) is a form of Fourier analysis that is applicable to a sequence of values.

The DTFT is often used to analyze samples of a continuous function. The term discrete-time refers to the fact that the transform operates on discrete data, often samples whose interval has units of time. From uniformly spaced samples it produces a function of frequency that is a periodic summation of the continuous Fourier transform of the original continuous function. Under certain theoretical conditions, described by the sampling theorem, the original continuous function can be recovered perfectly from the DTFT and thus from the original discrete samples. The DTFT itself is a continuous function of frequency, but discrete samples of it can be readily calculated via the discrete Fourier transform (DFT) (see Sampling the DTFT), which is by far the most common method of modern Fourier analysis.



**Code :-**

```
<<<File: expt3_1.m Comment: Main driver file for expt 3>>>
```

```
% Main driver program
% Written by Debagnik Kar 1804373
```

```
clear all
close all
clc
```

```
w = -2*pi:0.01:2*pi
n = 0:1:100
x = 0.5.^n
```

```
X = dtft(x, n, w)
```

```
subplot 221
plot(w/(pi),abs(X),'g')
ylabel('Magnitude')
xlabel('Frequency')
title('Magnitude plot')
subplot 223
plot(w/pi,angle(X),'r')
ylabel('Phase')
xlabel('Frequency')
title('Phase plot')
subplot 222
plot(w/pi,real(X),'g')
title('Real Plot')
xlabel('Frequency')
ylabel('Real part of f(x)')
subplot 224
plot(w/pi,imag(X),'r')
title('Imaginary Plot')
xlabel('Frequency')
ylabel('Imaginary part of f(x)')
```

```
<<<File: dtft.m Comment: Functional Dependencies for the previous programme>>>
```

```
% Function for performing DTFT
% Written by Debagnik Kar 1804373
```

```
function [X] = dtft(x,n,w)
```

```
    temp = n'*w;
    temp = -1i*temp;
    e = exp(temp);
    X = x*e;
```

```
end
```

<<<File: expt3\_2.m Comment: For analyzing DTFT properties>>>

```
% Program for analysing DTFT Functions  
% Writing by Debagnik Kar 1804373
```

```
clc  
clear all  
close all
```

```
n = 1  
w = linspace(-pi,pi,500)  
x1 = [1,3,5,7,9,11,13,15,17]  
x2 = [1,-2,3,-2,1]
```

```
y = conv(x1,x2)  
h1 = freqz(x1,n,w)  
h2 = freqz(x2,n,w)  
h12 = h1.*h2  
h3 = freqz(y',n,w)
```

```
subplot 321  
stem(x1,'k')  
title('Input Sequence one')
```

```
subplot 322  
stem(x2,'k')  
title('Input Sequence two')
```

```
subplot 323  
plot(w/pi,abs(h12),'g')  
title('Magnitude Plot of f(x) without using Convolution')
```

```
subplot 324  
plot(w/pi,abs(h3),'r')  
title('Magnitude Plot of f(x) with using Convolution')
```

```
subplot 325  
plot(w/pi,angle(h12),'g')  
title('Phase Plot of f(x) without using Convolution')
```

```
subplot 326  
plot(w/pi,angle(h3),'r')  
title('Phase Plot of f(x) with using Convolution')
```


**Graph/Output :-**

Fig 3.2: DTFT of the given arbitrary sequence




Fig 3.3: Analysis of DTFT properties

**Discussion/Inference of the experiment :-**

In the figure 3.2 we plotted the graph of a given arbitrary number.

From the figure 3.3 we observed that the DTFT magnitude and phase spectra obtained by performing pointwise multiplication of the two DTFT's of the original sequences are identical to those obtained by performing time domain convolution of the two original sequences; this verifies the convolution property of the DTFT.

**Conclusion :-**

In this experiment a function for computing DTFT of an arbitrary function is written in MATLAB and we successfully verified the convoluting property of DTFT programmatically. We will like to conclude by stating that the aim of the experiment is fulfilled.