
Maths Class 11 Chapter 3. Trigonometric Functions

Angle


When a ray OA starting from its initial position OA rotates about its end point 0 and takes the final position OB, we say that angle

AOB (written as \angle AOB) has been formed. The amount of rotation from the initial side to the terminal side is Called the measure of the angle.



Positive and Negative Angles

An angle formed by a rotating ray is said to be positive or negative depending on whether it moves in an anti-clockwise or a clockwise direction, respectively.



Measurement of Angles

There are three system for measuring angles,

1. Sexagesimal System/Degree Measure (English System)

In this system, a right angle is divided into 90 equal parts, called degrees. The symbol 1° is used to denote one degree. Each degree is divided into 60 equal parts, called minutes and one minute is divided into 60 equal parts, called seconds. Symbols 1' and 1" are used to denote one minute and one second, respectively.

2. Centesimal System (French System)

In this system, a right angle is divided into 100 equal parts, called 'grades'. Each grade is subdivided into 100 min and each minute is divided into 100 s.

```
i.e., 1 right angle = 100 \text{ grades} = 100^g

1^g = 100'

1' = 100''
```

3. Circular System (Radian System)

In this system, angle is measured in radian.

A radian is the angle subtended at the centre of a circle by an arc, whose length is equal to the radius of the circle.

The number of radians in an angle subtended by an arc of circle at the centre is equal to arc/radius.

Relationships

(i)
$$\pi$$
 radian = 180°
or 1 radian (180°/ π)= 57°16'22"
where, π = 22/7 = 3.14159

(ii)
$$1^{\circ} = (\pi/180)$$
 rad = 0.01746 rad

(iii) If D is the number of degrees, R is the number of radians and G is the number of grades in an angle θ , then

$$\frac{D}{90} = \frac{G}{100} = \frac{2R}{\pi}$$

(iv) $\theta = 1/r$ where $\theta =$ angle subtended by arc of length / at the centre of the circle, r = radius of the circle.

Trigonometric Ratios

Relation between different sides and angles of a right angled triangle are called trigonometric ratios or T-ratios

Trigonometric ratios can be represented as
$$\sin\theta = \frac{\text{Perpendicular}}{\text{Hypotenuse}} = \frac{BC}{AC}, \cos\theta = \frac{\text{Base}}{\text{Hypotenuse}} = \frac{AB}{AC},$$


$$\tan\theta = \frac{\text{Perpendicular}}{\text{Base}} = \frac{BC}{AB}, \csc\theta = \frac{1}{\sin\theta}$$

$$\sec\theta = \frac{1}{\cos\theta}, \cot\theta = \frac{\cos\theta}{\sin\theta} = \frac{1}{\tan\theta}$$

Trigonometric (or Circular) Functions

Let X'OX and YOY' be the coordinate axes. Taking 0 as the centre and a unit radius, draw a circle, cutting the coordinate axes at A,B, A' and B', as shown in the figure.

$$\left[\because \angle AOP = \frac{\operatorname{arc} AP}{\operatorname{radius} OP} = \frac{\theta}{1} = \theta^{\circ}, \text{ using } \theta = \frac{l}{r}\right]$$



Now, the six circular functions may be defined as under

- (i) $\cos \theta = x$
- (ii) $\sin \theta = y$
- (iii) $\sec \theta = 1/x, x \neq 0$
- (iv) cosec $\theta = 1/y$, $y \neq 0$
- (v) $\tan \theta = y/x, x \neq 0$
- (vi) cot $\theta = x/y$, $y \neq 0$

Domain and Range

Trigonometric Ratios	Domain	Range		
sinθ	R	[-1,1]		
cos θ	R	[-1,1]		
tanθ	$R - \left\{ (2n+1) \frac{\pi}{2} : n \in I \right\}$	R		
cosec 0	$R - \{n\pi : n \in I\}$	R - (-1,1)		
sec θ	$R - \left\{ (2n+1) \frac{\pi}{2} : n \in I \right\}$	R - (-1,1)		
cote	$R - \{n\pi : n \in I\}$	R		

Range of Modulus Functions

 $|\sin \theta| \le 1$, $|\cos \theta| \le 1$, $|\sec \theta| \ge 1$, $|\operatorname{Cosec} \theta| \ge 1$ for all values of 0, for which the functions are defined.

Trigonometric Identities

An equation involving trigonometric functions which is true for all those angles for which the functions are defined is called trigonometrical identity. Some identities are

(i)
$$\sin \theta = \frac{1}{\csc \theta}$$
 or $\csc \theta = \frac{1}{\sin \theta}$

(ii)
$$\cos \theta = \frac{1}{\sec \theta}$$
 or $\sec \theta = \frac{1}{\cos \theta}$

(iii)
$$\cot \theta = \frac{1}{\tan \theta} = \frac{\cos \theta}{\sin \theta} \text{ or } \tan \theta = \frac{1}{\cot \theta} = \frac{\sin \theta}{\cos \theta}$$

(iv)
$$\cos^2\theta + \sin^2\theta = 1$$
 or $1 - \cos^2\theta = \sin^2\theta$ or $1 - \sin^2\theta = \cos^2\theta$

(v)
$$1 + \tan^2 \theta = \sec^2 \theta$$

(vi)
$$1 + \cot^2 \theta = \csc^2 \theta$$

Sign of Trigonometric Ratios

Quadrant	sin θ	cos0	tan 0	cot0	sec θ	cosec θ
I (0, 90°)	+	+	+	+	+	+
II (90°, 180°)	+	-	_	_	_	+
(180°, 270°)	-	-	+	+	-	-
IV (270°, 360°)	_	+		-	+	_

Angle	0°	30°	45°	60°	90°	120°	135°	150°	180°
sin	0	1/2	$\frac{1}{\sqrt{2}}$	√3 2	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	1/2	0
cos	1	√3 2	$\frac{1}{\sqrt{2}}$	1/2	0	$-\frac{1}{2}$	$-\frac{1}{\sqrt{2}}$	$-\frac{\sqrt{3}}{2}$	-1
tan	0	$\frac{1}{\sqrt{3}}$	1 _	√3	00	- √3	-1	$-\frac{1}{\sqrt{3}}$	0
cot	00	√3	1	$\frac{1}{\sqrt{3}}$	0	$-\frac{1}{\sqrt{3}}$	-1	- √3	- 00
sec	1	$\frac{2}{\sqrt{3}}$	√2	2	00	- 2	- √2	$-\frac{2}{\sqrt{3}}$	-1
cosec	00	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1	$\frac{2}{\sqrt{3}}$	√2	2	90

Trigonometric Ratios of Some Special Angles

Angle	7 10	15°	22 1°	18°	36°
sinθ	$\frac{\sqrt{4-\sqrt{2}-\sqrt{6}}}{2\sqrt{2}}$	$\frac{\sqrt{3}-1}{2\sqrt{2}}$	$\frac{1}{2}\sqrt{2-\sqrt{2}}$	$\frac{\sqrt{5}-1}{4}$	$\frac{1}{4}\sqrt{10-2\sqrt{5}}$
cos θ	$\frac{\sqrt{4+\sqrt{2}+\sqrt{6}}}{2\sqrt{2}}$	$\frac{\sqrt{3}+1}{2\sqrt{2}}$	$\frac{1}{2}\sqrt{2+\sqrt{2}}$	$\frac{1}{4}\sqrt{10+2\sqrt{5}}$	$\frac{\sqrt{5}+1}{4}$
tanθ	$(\sqrt{3}-\sqrt{2})(\sqrt{2}-1)$	2 - √3	$\sqrt{2}-1$	√25-10√15 5	√5-2√5

Trigonometric Ratios of Allied Angles

Two angles are said to be allied when their sum or difference is either zero or a multiple of 90° . The angles — θ , $90^{\circ} \pm \theta$, $180^{\circ} \pm \theta$, $270^{\circ} + \theta$, 360° — θ etc., are angles allied to the angle θ , if θ is measured in degrees.

0	sin 0	cosec 0	cosθ	sec 0	tan 0	cote
- O	– sin θ	- cosec θ	cos θ	sec 0	– tan θ	- cot 0
90° − 0	cos θ	sec 0	sin 0	cosec 0	cat 0	tan 0
90° + 0	cos θ	sec θ	– sin θ	- cosec θ	- cot θ	- tan θ
180° – θ	sin θ	cosec θ	- cos θ	- sec θ	– tan θ	- cot θ
180° + θ	– sin θ	- cosec θ	- cos θ	- sec θ	tan 0	cot 0
270° - 0	- cos θ	- sec θ	– sin θ	- cosec θ	cat 0	tan θ
270° + 0	- cos θ	- sec θ	sin 0	cosec 0	- cot θ	– tan θ
360° - 0	- sin 0	- cosec θ	cos θ	sec 0	− tan θ	- cot 0

Trigonometric Periodic Functions

A function f(x) is said to be periodic, if there exists a real number T > 0 such that f(x + T) = f(x) for all x. T is called the period of the function, all trigonometric functions are periodic.

0	sin 0	cosec 0	cosθ	sec 0	tan 0	cote
- O	– sin θ	- cosec θ	cos θ	sec 0	– tan θ	- cot θ
90° − 0	cos θ	sec 0	sin 0	cosec 0	cot 0	tan θ
90° + 0	cos θ	sec θ	– sin θ	- cosec θ	- cot θ	- tan θ
180° – θ	sin θ	cosec θ	- cos θ	- sec θ	− tan θ	- cot θ
180° + θ	– sin θ	- cosec θ	- cos θ	– sec θ	tan 0	cot 0
270° − θ	- cos θ	- sec θ	- sin θ	- cosec θ	cat 0	tan θ
270° + 0	- cos θ	- sec θ	sin 0	cosec 0	- cot θ	- tan θ
360° - 0	– sin θ	- cosec θ	cos θ	sec 0	− tan θ	- cot θ

Maximum and Minimum Values of Trigonometric Expressions

- (i) Maximum value of $a \cos \theta \pm b \sin \theta = \sqrt{a^2 + b^2}$ Minimum value of $a \cos \theta \pm b \sin \theta = -\sqrt{a^2 + b^2}$
- (ii) Maximum value of $a \cos \theta \pm b \sin \theta + c = c + \sqrt{a^2 + b^2}$ Minimum value of $a \cos \theta \pm b \sin \theta + c = c - \sqrt{a^2 + b^2}$

Trigonometric Ratios of Compound Angles

The algebraic sum of two or more angles are generally called compound angles and the angles are known as the constituent angle. Some standard formulas of compound angles have been given below.

(i)
$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

(ii)
$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

(iii)
$$\cos(A+B) = \cos A \cos B - \sin A \sin B$$

(iv)
$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

(v)
$$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

(vi)
$$\tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

(vii)
$$\cot(A + B) = \frac{\cot A \cot B - 1}{\cot B + \cot A}$$

(viii)
$$\cot(A - B) = \frac{\cot A \cot B + 1}{\cot B - \cot A}$$

(ix)
$$\sin (A + B) \sin (A - B) = \sin^2 A - \sin^2 B = \cos^2 B - \cos^2 A$$

(x)
$$\cos(A + B)\cos(A - B) = \cos^2 A - \sin^2 B = \cos^2 B - \sin^2 A$$

(xi)
$$\sin(A + B + C) = \cos A \cos B \sin C + \cos A \sin B \cos C$$

$$+\sin A\cos B\cos C - \sin A\sin B\sin C$$

or
$$\sin(A+B+C) = \cos A \cos B \cos C(\tan A + \tan B + \tan C$$

 $-\tan A \tan B \tan C$

(xii)
$$\cos(A + B + C) = \cos A \cos B \cos C - \sin A \sin B \cos C$$

$$-\sin A\cos B\sin C - \cos A\sin B\sin C$$

or
$$\cos(A+B+C) = \cos A \cos B \cos C(1-\tan A \tan B - \tan B \tan C)$$

 $-\tan C \tan A$

(xiii)
$$\tan(A+B+C) = \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan B \tan C - \tan C \tan A}$$

If A+B+C=0, then $\tan A+\tan B+\tan C=\tan A$ $\tan B$ $\tan C$

(xiv) (a)
$$\sin(A_1 + A_2 + ... + A_n) = (\cos A_1 \cos A_2 \cos A_3 ... \cos A_n)$$

$$\times (S_1 - S_2 + S_5 - S_7 + ...)$$

(b)
$$\cos(A_1 + A_2 + ... + A_n) = (\cos A_1 \cos A_2 \cos A_3 ... \cos A_n)$$

$$\times (1 - S_2 + S_4 - S_6 + ...)$$

(c)
$$\tan(A_1 + A_2 + ... + A_n) = \frac{S_1 - S_3 + S_5 - S_7 + ...}{1 - S_2 + S_4 - S_6 + ...}$$

where.

$$S_1 = \tan A_1 + \tan A_2 + \dots + \tan A_n$$

$$S_2 = \tan A_1 \tan A_2 + \tan A_2 \tan A_3 + \dots$$

$$\tilde{S_3} = \tan A_1 \tan A_2 \tan A_3 + \tan A_2 \tan A_3 \tan A_4 + \dots$$

Transformation Formulae

(i)
$$2 \sin A \cos B = \sin (A + B) + \sin (A - B)$$

(ii)
$$2\cos A\cos B = \sin (A + B) - \sin (A - B)$$

(iii)
$$2\cos A\cos B = \cos(A+B) + \cos(A-B)$$

(iv)
$$2 \sin A \sin B = \cos (A - B) - \cos (A + B)$$

(v)
$$\sin C + \sin D = 2 \sin \left(\frac{C+D}{2}\right) \cos \left(\frac{C-D}{2}\right)$$

(vi)
$$\sin C - \sin D = 2\cos\left(\frac{C+D}{2}\right)\sin\left(\frac{C-D}{2}\right)$$

(vii)
$$\cos C + \cos D = 2\cos\left(\frac{C+D}{2}\right)\cos\left(\frac{C-D}{2}\right)$$

$$\begin{split} \text{(viii)} & \cos C - \cos D = -2 \sin \left(\frac{C+D}{2} \right) \sin \left(\frac{C-D}{2} \right) \\ & = \sin \left(\frac{C+D}{2} \right) \sin \left(\frac{D-C}{2} \right) \end{split}$$

Trigonometric Ratios of Multiple Angles

(i)
$$\sin 2A = 2 \sin A \cos A = \frac{2 \tan A}{1 + \tan^2 A}$$

(ii)
$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1$$

$$= 1 - 2\sin^2 A = \frac{1 - \tan^2 A}{1 + \tan^2 A}$$

(iii)
$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

(iv)
$$\sin 3A = 3\sin A - 4\sin^3 A$$

$$(v) \cos 3A = 4\cos^3 A - 3\cos A$$

(vi)
$$\tan 3A = \frac{3\tan A - \tan^3 A}{1 - 3\tan^2 A}$$

(vii)
$$\sin A = 2 \sin \frac{A}{2} \cos \frac{A}{2} = \frac{2 \tan \frac{A}{2}}{1 + \tan^2 \frac{A}{2}}$$

(viii)
$$\cos A = \frac{1 - \tan^2 \frac{A}{2}}{1 + \tan^2 \frac{A}{2}}$$

(ix)
$$1 - \cos A = 2\sin^2 \frac{A}{2}$$

(x)
$$1 + \cos A = 2\cos^2 \frac{A}{2}$$

(xi)
$$\frac{1-\cos A}{1+\cos A} = \tan^2 \frac{A}{2}$$

(xii)
$$\sin\left(\frac{A}{2}\right) + \cos\left(\frac{A}{2}\right) = \pm\sqrt{1 + \sin A}$$

(xiii)
$$\sin\left(\frac{A}{2}\right) - \cos\left(\frac{A}{2}\right) = \pm \sqrt{1 - \sin A}$$

Trigonometric Ratios of Some Useful Angles

(i)
$$\sin 75^\circ = \frac{\sqrt{3+1}}{2\sqrt{2}} = \cos 15^\circ$$

(ii)
$$\cos 75^\circ = \frac{\sqrt{3} - 1}{2\sqrt{2}} = \sin 15^\circ$$

(iii)
$$\tan 75^{\circ} = 2 + \sqrt{3} = \cot 15^{\circ}$$

(iv)
$$\cot 75^{\circ} = 2 - \sqrt{3} = \tan 15^{\circ}$$

(v)
$$\sin 9^\circ = \frac{\sqrt{3 + \sqrt{5}} - \sqrt{5 - \sqrt{5}}}{4} = \cos 81^\circ$$

(vi)
$$\cos 9^{\circ} = \frac{\sqrt{3 + \sqrt{5} + \sqrt{5 - \sqrt{5}}}}{4} = \sin 81^{\circ}$$

Important Results

1. Product of Trigonometric Ratio

(i)
$$\sin \theta \sin (60^{\circ} - \theta) \sin (60^{\circ} + \theta) = \frac{1}{4} \sin 3\theta$$

(ii)
$$\cos \theta \cos (60^{\circ} - \theta) \cos (60^{\circ} + \theta) = \frac{1}{4} \cos 3\theta$$

(iii)
$$\tan \theta \tan (60^\circ - \theta) \tan (60^\circ + \theta) = \tan 3\theta$$

(iv)
$$\cos 36^{\circ} \cos 72^{\circ} = \frac{1}{4}$$

(v)
$$\cos A \cos 2A \cos 4A ... \cos 2^{n-1}A = \frac{1}{2^n \sin A} \sin(2^n A)$$

2. Sum of Trigonometric Ratios

(i)
$$\sin A + \sin (A + B) + \sin (A + 2B) + ... + \sin (A + nB)$$

$$=\frac{\sin\left\{A+(n-1)\frac{B}{2}\right\}\sin\frac{nB}{2}}{\sin\frac{B}{2}}$$

(ii)
$$\cos A + \cos(A + B) + \cos(A + 2B) + ... + \cos(A + nB)$$

$$= \frac{\sin\frac{nB}{2}}{\sin\frac{B}{2}}\cos\left\{A + \frac{(n-1)B}{2}\right\}$$

3. A, B and C are Angles of a Triangle

(i) (a)
$$\sin (B + C) = \sin A$$

(b)
$$cos(B+C) = -cos A$$

(c)
$$\sin\left(\frac{B+C}{2}\right) = \cos\frac{A}{2}$$

(d)
$$\cos\left(\frac{B+C}{2}\right) = \sin\frac{A}{2}$$

(ii)
$$\sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C$$

(iii)
$$\cos 2A + \cos 2B + \cos 2C = -1 - 4\cos A\cos B\cos C$$

(iv)
$$\sin A + \sin B + \sin C = 4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}$$

(v)
$$\cos A + \cos B + \cos C = 1 + 4\sin \frac{A}{2}\sin \frac{B}{2}\sin \frac{C}{2}$$

(vi)
$$\tan A + \tan B + \tan C = \tan A \tan B \tan C$$

(vii)
$$\cot B \cot C + \cot C \cot A + \cot A \cot B = 1$$

(viii)
$$\cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} = \cot \frac{A}{2} \cot \frac{B}{2} \cot \frac{C}{2}$$

(ix)
$$\tan \frac{A}{2} \tan \frac{B}{2} + \tan \frac{B}{2} \tan \frac{C}{2} + \tan \frac{C}{2} \tan \frac{A}{2} = 1$$

4. Trigonometric Equations

(i)
$$\sin n\pi = 0$$
 and $\cos n\pi = (-1)^n$

(ii)
$$\sin \theta = \sin \alpha \Rightarrow \theta = n\pi + (-1)^n \alpha, n \in I$$

d many more)