

De Solutione Problematum per Motum

Author: Isaac Newton

Source: MS Add. 3958.3, ff. 68r-76v, Cambridge University Library, Cambridge, UK

Published online: October 2011

<68r>

De Solutione Problematum per Motum

Ne hujusmodi operationes obscuræ nimis evadant, Lemmata 6 sequentia breviter ideoque non demonstrata præmittam.

Lemma 1. Si corpus A in circumferentiâ circuli vel sphaeræ ADCE moveatur versus ejus centrum B: velocitas ejus ad unaquæque circumferentiæ puncta D, C, E, est ut cordæ AD, AC, AE, ductæ a Corpore A ad ista puncta D, C, E.

Lemma 2. Sit $\triangle ADC$ similis $\triangle AEC$, etsi non sunt in eodem plano. Inquam, Si tria Corpora a puncto A, primum ad D, secundum ad E, tertium ad C, uniformiter et in æqualibus temporibus moveantur: Motus tertij componetur ex motibus primi et secundi.

Notetur, quòd hic per corpus intelligitur ejus centrum gravitatis, vel aliquod ejus punctum.

Lemma 3. Omnia puncta corporis parallelismum servantis æqualiter moventur.

Lemma 4. Si corpus solo motu circulari circa axim quemvis rotetur; motus omnium ejus punctorum sunt ut distantiae ab isto axi. Et hi duo motus simplices vocentur.

Lemma 5. Si motus corporis consideretur ut mixtus e motibus simplicibus: motus omnium ejus punctorum componetur ex motibus eorum simplicibus, eo modo quo motus ab A ad C, in Lemmate 2^{do} componitur ex motibus ab A ad D et E.

Nota. quod motus quilibet ad unum horum 3^{um} casuum reduci poterit. Et in casu tertio, linea quævis pro axe, (vel si linea aut plana superficies moveatur in plano, quodvis punctum istius plani pro centro) motus assumi potest.

Lemma 6. Sint AE, AH, lineæ motæ et continuè secantes; Ducantur AB, AD, AC, CB, CD. Dico quòd, datis quæ requirantur ad proportionem et positionem harum quinque linearum AB, AD, AC, CB, DC, determinandas; illæ designent proportionem et positionem h_{orum} quinque motu_{um}, viz: puncti A in lineâ AE fixi et versus B moventis, puncti A in lineâ AH fixi et versus D moventis, Puncti intersectionis A in plano ABCD moventis versus C, (lineæ enim 5 istæ semper sunt in eodem plano etsi AE AH non sunt), Puncti intersectionis A in lineâ AE moventis secundum ordinem litterarum C, B, et Puncti intersectionis A in lineâ AH moventis secundum ordinem C. D.

Nota. Quòd linea recta tangens curvam motu descriptam (ut AB, AD, AC) vel istæ tangenti parallela (ut CB, CD,), dicitur designare positionem istius motûs in puncto contactûs.

Nota etiam quod lineâ AH quiescente (ut in Fig 1, et 4), punctum D et A coincident et punctum C in lineâ AH, modò sit recta (Fig 1,), aliter in ejus tangente AC (fig 4) reperietur.

Prop 1. Ducere Tangentem ad Ellipsin.

Sit ACB filum per quod ellipsis describi solet, et CE Tangens. Cùm filum AC augetur eadem velocitate quâ BC diminuitur, i.e: C habet eandem velocitatem versus D et B; erit $\angle DCE = \angle ECB$. Per Lem 1. Idem de reliquis Conicis intelligatur.

Prop 2. Ducere Tangentem ad Conchoiden

Sint GLC, ALF, GAE, regulæ quibus concha describi solet: fiat $GT \parallel AF \perp CB = \parallel MN$: et $NG = CL \perp TN \parallel RL$. Et, Cùm æqualitas proportionalitatis simplicior est, ponatur lineam $CB = NM$ esse æqualem velocitati puncti C versus B, vel puncti N versus M. Erit $NT =$ motui circulari puncti N circa G rotantis versus T, (Lem 1): et $LR =$ Motui circulari puncti L in lineâ GL fixi, circa G (Lem 4): et $LG =$ motui puncti intersectionis L (i.e: velocitati puncti C) a puncto G, sive versus punctum D in lineâ C G moventis, (Lem 6). Iam cum habeatur duplex velocitas puncti C viz CB versus B et LG versus D, fiat $FD \perp DC = LG$: Et motus puncti C erit in lineâ FC, Diametro nempe circuli transientis per puncta CDBF, (Lem 1): Quæ proinde Concham tangit in puncto C.

<68v>





Prop 3. Invenire punctum C distinguens concavam a convexâ Conchæ portione.

Iis in priori propositione suppositis: Fiat triangulus GFH similis triangulo GNT sive LBC: et $DF \perp FR \parallel HK = 2GL$; Iungantur F, K, et fiat $KP \parallel FD$. Si Linea DF solum motum Parallelum per CD vel FR directum haberet, (quia $CD = GL$) motus omnium ejus punctorum esset FR, (Lem 3): Et si solum motum circularem circa centrum G haberet, motus puncti F, in istâ lineâ DF fixi, esset FH, (Lem 4). At motus istius F ex istis duobus componitur, proinde erit FK, (Lem 5., 2.) et motus puncti intersectionis F per lineas AF, DF, facti, et in AF moventis, erit FP (Lem 6). Iam si linea CF Concham tangit in puncto quæsito C, facillè deprehendatur motum puncti intersectionis F esse nullum; proinde P et F coincidere; sive DF et FK in directum jacere; et $\triangle GDF$, $\triangle FKH$ esse similes.

Quæ ut calculo subijciantur, fiat $AG = b$. $CL = c$. $CB = y$. tum, $BL = \sqrt{cc - yy}$. $2GL = \frac{2bc}{y} = HK$. $LD = \frac{cb + cy}{y}$. $DF = \frac{cb + cy}{\sqrt{cc - yy}}$. Etc.

$\sqrt{cc - yy} : y :: BL : CB :: GF : FH :: DF : KH :: \frac{cb + cy}{\sqrt{cc - yy}} : \frac{2bc}{y} :: by + yy : 2b\sqrt{cc - yy}$. Quare, $2bcc - 2byy = byy + y^3$. sive $y^3 + 3byy - 2bcc = 0$. <

insertion from f 69r > Equatio $y^3 + 3byy - 2bcc = 0$, priùs inventa, ita resolve, fac $Ao = \frac{8c^4}{27b^3}$. af = c = AE duc {of} et cum diametro of describe circumulum fmao, in quo inscribe fm=b=AG. et cum radio om, fac circumulum mv, et a puncto intersectionis v duc vd \perp ad. erit 3b:2c::VD:GL. unde datur punctum c. < text from f 68v resumes >



Vsus autem hujus methodi (ut intelligo) præcipuus est in lineis Mechanicis, ubi deficit Algebraica calculatio. Exempli, Tangens Quadratricis ita ducetur.

Sint AG, MC regulæ quibus uniformitèr motis Quadratrix describi intelligitur. Et CB vocetur motus puncti C in lineâ CM fixi et versus B moventis; tum arcus GK erit motus puncti G circa A rotantis, (sup) et arcus CL erit motus puncti C in lineâ AG fixi et circa G rotantis, (lem 4). Quare (si fiat CL=CD⊥AG∥DE, et BF∥CM.) motus puncti intersectionis C in planò AEK erit CF (Lem 6), quæ proinde Quadratricem tangit in C.

<69r>
<70r>

If vce is a Conchoïdes, g its pole, &c: ga=b. ae=lc=vb=c. ma=y & c the point betwixt its convexity & concavity, then is $y^3 + 3byy - 2bcc = 0$. (see pag: 259, lin: 10.). Which Equation hath one affirmative roote (ma) referred to the point c. and two negative rootes whereof the greater is referred to the lower Conchoïdes, & the lesser (I thinke) uselesse to this question.

The rootes of this Equation may bee thus found by the helpe of the described Conchoïdes & a circle. viz: Suppose ga=b, ae=lc=vb=c. ma=y. (as before) a{o}=s. oc=r. And thereby may bee found this Equation $\frac{y^3 + rr - ss + bb - cc \times yy - 2bccy - bcc}{2s + 2b} = 0$, to bee compared with the former, (see pag 261 lin 20). But their rootes cannot become equal by reason of their third termes. Therefore I alter the rootes of the first equation, as, suppose I make $y = z - b$. Then is $z^3 - 3bbz + 2b^3 - 2bcc = 0$. To bee compared with the 2^d equation: which cannot yet bee done without a contradiction, there being but two unknowne quantitys, r & s to bee found by three Equations resulting from the comparison of their 2^d 3^d & 4th termes. But if I make $z = \frac{4ccx - 4bbx}{3bb}$, & substitute this valor into its place in the precedent Equation, the result is $x^3 * \frac{-27b^6x}{16c^4 - 32bbcc + 16b^4} - \frac{-27b^7}{32c^4 - 64bbcc + 32b^4} = 0$. The termes of which being compared with the termes of the 2^d Equation the 3^d or 4th give $\frac{2bbc}{2s + 2b} = \frac{27b^6}{16c^4 - 32bbcc + 16b^4}$. Or $16c^6 - 32bbcc^4 + 16b^4cc = 27b^6 + 27b^5s$. & $s = \frac{16c^6}{27b^5} - \frac{32c^4}{27b^3} + \frac{16cc}{27b} - b = ao$. Their second termes give $\frac{rr - ss + bb - cc}{2s + 2b} = 0$. Or $r = \sqrt{ss + cc - bb} = ov$. Therefore from the intersection point v (made by the Conchoïdes & a circle whose radius is ov & center o) let fall vd⊥ad; & ar=x=vd, is the roote of the 2^d & last equation. Which being found make $y = \frac{4ccx - 4bbx}{3bb} - b = am$.

Which was to bee done.

Prop 1. Suppose ab=x. bd=y⊥ab. And that the nature of the line addc is such that the valor of y is rationall & consists of no fractions in whose denominator x is, or else wholly of such fractions in whose denominators x is, but not of divers dimensions: If I then multiply the valor of y by x, & divide each terme of that valor by so many units as it hath dimensions in that terme; the result shall signifie the area abd of the afforesaide line addc.

As for example. If y=1, or y=x, or y=xx, or x³, or x⁴ &c then the area abd is $\frac{1xx}{1}$, or $\frac{xxx}{2}$, or $\frac{xxxx}{3}$, or $\frac{x^3xx}{4}$, or $\frac{x^4xx}{5}$ &c: so if $y = \frac{a}{b}$, or $y = \frac{ax}{b}$, or $\frac{aax}{b}$ &c: then the area abd is $\frac{ax}{b}$, $\frac{aax}{2b}$, $\frac{aax^2}{3b}$ &c. In like manner if $y = \frac{a}{xx}$, or $y = \frac{a}{x^3}$, or $y = \frac{a}{x^4}$ &c then $\frac{a}{\pm x} = \frac{-a}{x}$, or $\frac{-a}{-2xx} = \frac{a}{2x^2}$, or $\frac{-a}{-3x^3} = \frac{a}{3x^3}$, &c: is the area abd (as is by others demonstrated). [so if $\frac{a}{x} = y$. then $\frac{a}{0} = \frac{ax^0}{0}$ is the area abd; viz: tis infinite.]. Lastly if $y = 1 + x + aax - bx^3 - \frac{x^4}{d} + \frac{cx^{15}}{e+f}$ &c. the area abd is $x + \frac{1}{2}xx + \frac{aax}{3} - \frac{bx^4}{4} - \frac{x^5}{5d} + \frac{cx^6}{16e+16f}$ &c. For the area abd is compounded of those areas which are related to & generated by those quantitys of which the valor of y is compounded. & what those areas are appeare by the former example. (Note that Parabolicall & Hyperbolicall (i.e: in respect of bd,) affirmative & negative) areas (thus considered) cannot compound any 3^d area, because they are not on the same side of the line bd.)

Prop 2. $\frac{aa}{b+x} = \frac{aa}{b} - \frac{aax}{bb} + \frac{aaxx}{b^3} - \frac{aax^3}{b^4} + \frac{aax^4}{b^5} - \frac{aax^5}{b^6} + \frac{aax^6}{b^7}$ &c. For these termes $\frac{aa}{b+x}$. aa. aab+aax. aabb+2aabx+aaxx. aab³+3aabbx+3aabxx+aax³, & which termes may bee thus ordered This last appeares by multiplying both parts by b+x

<70v>

$$\begin{array}{cccccc} \frac{aa}{b+x} . & aa . & aab . & aabb . & aab^3 . & aab^4 . \\ & aax . & 2aabx . & 3aabbx . & 4aab^3x . & \&c: \\ & aaxx . & 3aabxx . & 6aabbxx . & & \\ & aax^3 . & 4aabx^3 . & & & \\ & aax^4 . & & & & \end{array}$$

Or by supplying the vacant places

$$\begin{array}{cccccc} \frac{aa}{b+x} . & 1 \times aa . & 1 \times aab . & 1 \times aabb . & 1 \times aab^3 . & 1 \times aab^4 . \\ & 0 \times \frac{aax}{b} . & 1 \times aax . & 2 \times aabx . & 3 \times aabbx . & 4 \times aab^3x . \\ & 0 \times \frac{aaxx}{bb} . & 0 \times \frac{aax^2}{b} . & 1 \times aaxx . & 3 \times aabbx . & 6 \times aabbxx . \\ & 0 \times \frac{aax^3}{b} . & 0 \times \frac{aax^3}{bb} . & 0 \times \frac{aax^3}{b} . & 1 \times aax^3 . & 4 \times aabx^3 . \\ & 0 \times \frac{aax^4}{b^4} . & 0 \times \frac{aax^4}{b^3} . & 0 \times \frac{aax^4}{bb} . & 0 \times \frac{aax^4}{b} . & 1 \times aax^4 . \end{array}$$

Now to reduce the first terme $\frac{aa}{b+x}$ to the same forme with the rest, I consider in what progressions the numbers prefixed to these termes proceede, & find them to bee such that any number added to the number above it is equall to the number following it. Whence any termes may bee found which are wanting, as in the annexed Table. Also any terme, to which these numbers are prefixed, being multiplied by b produceth the following litterall terme. Or the higher terme multiplyed by $\frac{x}{b}$, produceth the lower terme. As in the following table

1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
 -4. -3. -2. -1. 0. 1. 2. 3. 4. 5. 6. 7.
 10. 6. 3. 1. 0. 0. 1. 3. 6. 10. 15. 21.
 -20. -10. -4. -1. 0. 0. 0. 1. 4. 10. 20. 35. &c
 35. 15. 5. 1. 0. 0. 0. 0. 1. 5. 15. 35.
 -56. -21. -6. -1. 0. 0. 0. 0. 0. 1. 6. 21.
 84. 28. 7. 1. 0. 0. 0. 0. 0. 0. 1. 7.

| | | | | | | | |
|------------------------|-----------------------|----------------------|------------------------------|------------------------------|-------------------------------|-------------------------------|-------------------------------|
| $\frac{aa}{(b+x)^3}$ | $\frac{aa}{(b+x)^2}$ | $\frac{aa}{b+x}$ | aa | $aa \times b + x$ | $a^2 \times \overline{b+x^2}$ | $a^2 \times \overline{b+x^3}$ | $a^2 \times \overline{b+x^4}$ |
| = | = | = | = | = | = | = | = |
| $+\frac{aa}{b^3}$ | $+\frac{aa}{bb}$ | $+\frac{aa}{b}$ | $+aa$ | $+aab$ | $+aabb$ | $+aab^3$ | $+aab^4$ |
| $-\frac{3aax}{b^4}$ | $-\frac{2aax}{b^3}$ | $-\frac{aax}{bb}$ | $\frac{0 \times aax}{b}$ | $+aax$ | $+2aabb$ | $+3aabbx$ | $+4aab^3x$ |
| $+\frac{6aax^2}{b^5}$ | $+\frac{3aax^2}{b^4}$ | $+\frac{aaxx}{b^3}$ | $\frac{0 \times aax^2}{b^2}$ | $\frac{0 \times aaxx}{b}$ | $+aaxx$ | $+3aabbx$ | $+6aabbx^2$ |
| $-\frac{10aax^3}{b^6}$ | $-\frac{4aax^3}{b^5}$ | $-\frac{aax^3}{b^4}$ | $\frac{0 \times aax^3}{b^3}$ | $\frac{0 \times aax^3}{bb}$ | $\frac{0 \times aax^3}{b}$ | $+aax^3$ | $+4aabbx^3$ |
| $+\frac{15aax^4}{b^7}$ | $+\frac{5aax^4}{b^6}$ | $+\frac{aax^4}{b^5}$ | $\frac{0 \times aax^4}{b^4}$ | $\frac{0 \times aax^4}{b^3}$ | $\frac{0 \times aax^4}{bb}$ | $\frac{0 \times aax^4}{b}$ | $+aax^4$ |
| &c | &c | &c | &c | &c | &c | &c | &c |

Whence it appearesy^t

| | |
|--|----------------------|
| $\frac{aa}{b+x} = \frac{aa}{b} - \frac{aax}{bb} + \frac{aaxx}{b^3} - \frac{aax^3}{b^4}$ | &c. Ory ^t |
| $\frac{aa}{b-x} = \frac{aa}{b} + \frac{aax}{bb} + \frac{aaxx}{b^3} + \frac{aax^3}{b^4}$ | &c. Ory ^t |
| $\frac{aa}{b+x} = \frac{aa}{x} - \frac{aab}{xx} + \frac{aabb}{x^3} - \frac{aabb^3}{x^4}$ | &c. Ory ^t |
| $\frac{aa}{bb+2bx+xx} = \frac{aa}{bb} - \frac{2a2x}{b^3} + \frac{3aaxx}{b^4}$ | &c. |

Prop: 3^d. If ab=x. y=db⊥ab⊥ac. (fa=b) & $\frac{aa}{b+x} = y = \left(\text{prop2}\right) \frac{aa}{b} - \frac{aax}{bb} + \frac{aaxx}{b^3} - \frac{aax^3}{b^4} + \frac{aax^4}{b^5}$ &c. Then (by prop 1),

$\frac{aax}{b} - \frac{aaxx}{2bb} + \frac{aax^3}{3b^3} - \frac{aax^4}{4b^4} + \frac{aax^5}{5b^5} - \frac{aax^6}{6b^6}$ &c: is abde, the area of the Hyperbola. So if $\frac{aa}{b-x} = y$: &c. In like manner if


$\frac{aa}{bb+2bx+xx} = y = \left(\text{prop2}\right) \frac{aa}{bb} - \frac{2aax}{b^3} + \frac{3aaxx}{b^4} - \frac{4aax^3}{b^5}$ &c. Then (by prop 1)

$\frac{aax}{bb} - \frac{aaxx}{b^3} + \frac{aax^3}{b^5} - \frac{aax^4}{b^6}$ &c = $\left(\text{prop2}\right) \frac{aa}{bb} - \frac{aa}{b+x}$ is the area abde. (which may also thus appeare viz: if fb=b+x=z. then

$\frac{aa}{bb+2bx+xx} = \frac{aa}{zz} = y$. Therefore (prop 1) the area dbm = $\frac{aa}{z} = \frac{aa}{b+x}$. & eam = $\frac{aa}{b}$. so that eadb = $\frac{aa}{b} - \frac{aa}{b+x}$). And so of

the rest. As if $\frac{x^3}{aa+bx+x^2} = y = \left(\text{prop2}^d\right) \frac{x^3}{aa} - \frac{bx^4-x^5}{a^4} + \frac{bbx^5+2bx^6+x^7}{a^6} - \frac{bbbx^6-3bbx^7-3bx^8-x^9}{a^8}$ &c. The area abde is

$$\frac{4x^4}{aa} - \frac{5bx^5-6x^6}{a^4} + \frac{6bbx^6+14bx^7+8x^8}{a^6} - \frac{7b^3x^7-24bbx^8-27bx^9-10x^{10}}{a^8} \text{ &c.}$$



Prop 4th. To find two or three intermediate terms in the above mentioned table of numeral

progressions, I observe that those progressions are of this nature viz

| | | | | |
|----|------|---------|------------|---------------|
| a. | a. | a | a. | a. |
| b. | b+c. | b+2c. | b+3c. | b+4c. |
| d. | d+e. | d+2e+f. | d+3e+ef. | d+4e+6f. |
| g. | g+h. | g+2h+i. | g+3h+3i+k. | g+4h+6i+4k. |
| l. | l+m. | l+2m+n. | l+3m+3n+p. | l+4m+6n+4p+q. |
| r. | r+s. | r+2s+t. | r+3s+3t+v. | r+4s+6t+4v+w. |

<71r>

terms in the 3^d progression

3. *. 1. *. 0. *. 0. *. 1. *. 3.
 d-4e+10f. d-3e+6f. d-2e+3f. d-e+f d. d+e d+2e+f. d+3e+f d+4e+6f. d+5e+10f. d+6e+15f.

I compare the terms of that progression & of the correspondent litterall progression & find d=0=2e+f. 4e+6f=1. subduct 4e+2f=0, Or 12e+6f=0 from 4e+6f=1. & the rest is 4f=1. Or -8e=1. which terms being found viz d=0. e = $-\frac{1}{8}$. f = $\frac{1}{4}$. the progression must be $3 \cdot \frac{9}{8} \cdot 1 \cdot \frac{3}{8} \cdot 0 \cdot \frac{-1}{8} \cdot 0 \cdot \frac{3}{8} \cdot 1 \cdot \frac{15}{8} \cdot 3$ &c. Hence may

be deduced this table viz

| | | | | | | | | | | | | | | | | |
|------|------------------------|-----|-----------------------|-----|----------------------|-----|----------------------|-----|--------------------|-----|---------------------|-----|--------------------|-----|---------------------|-----|
| 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. |
| -3. | $-\frac{5}{2}$. | -2. | $-\frac{3}{2}$. | -1. | $-\frac{1}{2}$. | 0. | $\frac{1}{2}$. | 1. | $\frac{3}{2}$. | 2. | $\frac{5}{2}$. | 3. | $\frac{7}{2}$. | 4. | $\frac{9}{2}$. | 5. |
| 6. | $\frac{35}{8}$. | 3. | $\frac{15}{8}$. | 1. | $\frac{3}{8}$. | 0. | $-\frac{1}{8}$. | 0. | $\frac{3}{8}$. | 1. | $\frac{15}{8}$. | 3. | $\frac{35}{8}$. | 6. | $\frac{63}{8}$. | 10. |
| -10. | $-\frac{105}{16}$. | -4. | $-\frac{35}{16}$. | -1. | $-\frac{5}{16}$. | 0. | $\frac{1}{16}$. | 0. | $-\frac{1}{16}$. | 0. | $\frac{5}{16}$. | 1. | $\frac{35}{16}$. | 4. | $\frac{105}{16}$. | &c |
| 15. | $\frac{1155}{128}$. | 5. | $\frac{315}{128}$. | 1. | $\frac{35}{128}$. | 0. | $-\frac{5}{128}$. | 0. | $\frac{3}{128}$. | 0. | $-\frac{5}{128}$. | 0. | $\frac{35}{128}$. | 1. | $\frac{315}{128}$. | 5. |
| -21. | $-\frac{3003}{256}$. | -6. | $-\frac{693}{256}$. | -1. | $-\frac{63}{256}$. | 0. | $\frac{7}{256}$. | 0. | $-\frac{3}{256}$. | 0. | $\frac{7}{256}$. | 0. | $-\frac{7}{256}$. | 0. | $\frac{63}{256}$. | 1. |
| +28. | $\frac{15015}{1024}$. | 7. | $\frac{3003}{1024}$. | 1. | $\frac{231}{1024}$. | 0. | $-\frac{21}{1024}$. | 0. | $\frac{7}{1024}$. | 0. | $-\frac{5}{1024}$. | 0. | $\frac{7}{1024}$. | 0. | $\frac{21}{1024}$. | 0. |
| &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: | &c: |

Note that the progression $1 \cdot \frac{1}{2} \cdot \frac{-1}{8} \cdot \frac{1}{16} \cdot \frac{-5}{128} \cdot \frac{7}{256}$ &c: may bee deduced from hence

$$\frac{1 \times 1 \times -1 \times -3 \times -5 \times -7 \times -9 \times -11}{1 \times 2 \times 4 \times 6 \times 8 \times 10 \times 12 \times 14} \text{ &c \& one intermediate terme given the rest are easily deduced thence.}$$

In like manner if I would find two meanes twixt every terme of that numeral progression I compare the numeral & correspondent litterall progressions, suppose in the 3^d progression.

1. *. *. 0. *. *. 0. *. *. 1.

d-3e+6f. d-2e+3f. d-e+f. d. d+e d+2e+f. d+3e+3f. d+4e+6f. d+5e+10f. d+63+15f. &c.

And find that d=0=3e+3f. & 6f-3e=1. To which adding 3e+3f=0, or -6f-6e=0. the result is 9f=1, or -9e=1. So that the progression must bee

1. $\frac{5}{9}$. $\frac{2}{9}$. 0. $-\frac{1}{9}$. $-\frac{1}{9}$. 0. $\frac{2}{9}$. $\frac{5}{9}$. 1. $\frac{14}{9}$. $\frac{20}{9}$. 3. &c. Hence may bee composed this

Table

| | | | | | | | | | | | | |
|----|---------------------|--------------------|----|--------------------|--------------------|----|--------------------|---------------------|----|--------------------|---------------------|----|
| 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. |
| 0. | $\frac{1}{3}$. | $\frac{2}{3}$. | 1. | $\frac{4}{3}$. | $\frac{5}{3}$. | 2. | $\frac{7}{3}$. | $\frac{8}{3}$. | 3. | $\frac{10}{3}$. | $\frac{11}{3}$. | 4. |
| 0. | $-\frac{1}{9}$. | $-\frac{1}{9}$. | 0. | $\frac{2}{9}$. | $\frac{5}{9}$. | 1. | $\frac{14}{9}$. | $\frac{20}{9}$. | 3. | $\frac{35}{9}$. | $\frac{44}{9}$. | 6. |
| 0. | $\frac{5}{81}$. | $\frac{4}{81}$. | 0. | $-\frac{4}{81}$. | $-\frac{5}{81}$. | 0. | $\frac{14}{81}$. | $\frac{40}{81}$. | 1. | $\frac{140}{81}$. | $\frac{220}{81}$. | 4. |
| 0. | $-\frac{10}{243}$. | $-\frac{7}{243}$. | 0. | $\frac{+5}{243}$. | $\frac{+5}{243}$. | 0. | $-\frac{7}{243}$. | $-\frac{10}{243}$. | 0. | $\frac{25}{243}$. | $\frac{110}{243}$. | 1. |

Note that this progression viz

$1 \times 1 \times -2 \times -5 \times -8 \times -11 \times -14$ &c gives the second term $1 + \frac{1}{3} - \frac{1}{9} + \frac{5}{81} - \frac{10}{243} + \frac{22}{729}$ &c. & this

$1 \times 3 \times 6 \times 9 \times 12 \times 15 \times 18$
 $1 \times 2 \times -1 \times -4 \times -7 \times -10 \times -13$ &c gives the third terme. Also this progression
 $1 \times 3 \times 6 \times 9 \times 12 \times 15 \times 18$

$1 \times 4 \times 1 \times -2 \times -5 \times -8 \times -11 \times -14$ &c. gives the 4th.
 $1 \times 3 \times 6 \times 9 \times 12 \times 15 \times 18 \times 21$

And in Generall if the 2^d quantity of any terme is $\frac{x}{y}$, this progression gives all the rest viz

$\frac{1 \times x \times x-y \times x-2y \times x-3y \times x-4y \times x-5y \times x-6y}{1 \times y \times 2y \times 3y \times 4y \times 5y \times 6y \times 7y}$ &c. And $\frac{x}{y}$ is ever given by supposition for it signifieth the distance of

the terme from 1.0.0.0.


| | | | | | | | | | |
|--|----|----|----|----|----|-----|-----|------------------|---|
| Note alsoy ^t any
of these progressions
w th their intermediate
termes | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1 = y. |
| | 0. | 1. | 2. | 3. | 4. | 5. | | | x = y. |
| | 0. | 0. | 1. | 3. | 6. | 10. | &c. | may bee designed | xx - x = 2y. |
| | 0. | 0. | 0. | 1. | 4. | 10. | | by Geometricall | x ³ - 3xx + 2x = 6y. |
| | 0. | 0. | 0. | 0. | 1. | 5. | | lines | x ⁴ - 6x ³ + 11xx - 6x = 24y. |
| | 0. | 0. | 0. | 0. | 0. | 1. | | | x ⁵ - 10x ⁴ + 35x ³ - 50x ² + 24x = 120y. |
| | 0. | 0. | 0. | 0. | 0. | 0. | | | x ⁶ - 15x ⁵ + 85x ⁴ - 225x ³ + 274xx - 120x = 720y. |

In which x

signifieth the distance of any terme from the first 1.0.0.0.0.0. & y is the quantity of that terme.

Prop 5^t.

<72r>



If lab is an Hyperbola; cde, ck its Asymptotes, a its vertex, & cag its axis; if adef is a square & he||ad & cd=1, & de=x. then be = $\frac{1}{1+x}$. If also, ef=1. eg=1+x. eh=1+2x+x &c: (the progression continued is 1+3x+3xx+x³. 1+4x+6x²+4x³+x⁴.

1+5x+10x²+10x³+5x⁴+x⁵ &c). Then, shall the areas of those lines proceede in this progression. *=adeb. x=adeb.

$x + \frac{xx}{2} = adeg$. adeh = $x + \frac{2xx}{2} + \frac{x^3}{3}$. $x + \frac{3xx}{2} + \frac{3x^3}{3} + \frac{x^4}{4}$. $x + \frac{4xx}{2} + \frac{6x^3}{3} + \frac{4x^4}{4} + \frac{x^5}{5}$ &c. As in this table. In which the first area is also inserted. The composition of which table may be

| | | | | | | | | | | |
|----------------------------|---|-----|----|----|----|----|----|-----|-----|-----|
| x | × | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. |
| $\frac{xx}{2}$ | × | -1. | 0. | 1. | 2. | 3. | 4. | 5. | 6. | 7. |
| $\frac{x^3}{3}$ | × | 1. | 0. | 0. | 1. | 3. | 6. | 10. | 15. | 21. |
| $\frac{x^4}{4}$ | × | -1. | 0. | 0. | 0. | 1. | 4. | 10. | 20. | 35. |
| $\frac{x^5}{5}$ | × | 1. | 0. | 0. | 0. | 0. | 1. | 5. | 15. | 35. |
| $\frac{x^6}{6}$ | × | -1. | 0. | 0. | 0. | 0. | 0. | 1. | 6. | 21. |
| $\frac{1}{7}x^7$ | × | 1. | 0. | 0. | 0. | 0. | 0. | 0. | 1. | 7. |
| = = = = = | | | | | | | | | | |
| adeb. adeb. adeg. adeh. &c | | | | | | | | | | |

deduced from hence; viz: The sume of any figure &y^e figure above it is equall toy^e figure following & it. Byw^{ch}table it may appeary^ty^earea ofy^eHyperbola adeb is

$$x - \frac{xx}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \frac{x^5}{5} - \frac{x^6}{6} + \frac{x^7}{7} - \frac{x^8}{8} + \frac{x^9}{9} - \frac{x^{10}}{10}$$

Suppose that adck is a Square abc a circle agc a Parabola. &c. & that de=x. ad||fe=1=bd. & that the progression in which the lines fe, be, ge, he, ie, ne &c proceedes is $1.\sqrt{1-xx}. 1-xx. 1-xx\sqrt{1-xx}. 1-2xx+x^4. 1-2xx+x^4\sqrt{1-xx}. 1-3xx+3x^4-x^6.$ &c. Then will their areas fade, bade, gade, hade, iade, &c be in this progression.


x.*.x- $\frac{xxx}{3}$.*.x- $\frac{2}{3}x^3+\frac{1}{5}x^5$.*.x- $\frac{3x^3}{3}+\frac{3x^5}{5}-\frac{x^7}{7}$.*.x- $\frac{4x^3}{3}+\frac{6x^5}{5}-\frac{4x^7}{7}+\frac{x^9}{9}$. &c: as in this table following in which the intermediate termes are inserted. The property of which table is that the

| | | | | | | | | | | | | |
|------------------|---|-----|---------------------|----|---------------------|----|-------------------|----|--------------------|----|--------------------|----|
| x | × | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. | 1. |
| $-\frac{x^3}{3}$ | × | -1. | $-\frac{1}{2}$. | 0. | $\frac{1}{2}$. | 1. | $\frac{3}{2}$. | 2. | $\frac{5}{2}$. | 3. | $\frac{7}{2}$. | 4. |
| $\frac{x^5}{5}$ | × | 1. | $\frac{3}{8}$. | 0. | $-\frac{1}{8}$. | 0. | $\frac{3}{8}$. | 1. | $\frac{15}{8}$. | 3. | $\frac{35}{8}$. | 6. |
| $-\frac{x^7}{7}$ | × | -1. | $-\frac{5}{16}$. | 0. | $\frac{3}{48}$. | 0. | $-\frac{1}{16}$. | 0. | $\frac{5}{16}$. | 1. | $\frac{35}{16}$. | 4. |
| $\frac{x^9}{9}$ | × | 1. | $+\frac{35}{128}$. | 0. | $-\frac{15}{384}$. | 0. | $\frac{3}{128}$. | 0. | $-\frac{5}{128}$. | 0. | $\frac{35}{128}$. | 1. |
| | | | | | | | | | | | | |
| | | | | | | | | | | | | |

summe of any

figure & the figure above it is equall to the figure next after it save one. Also the numerall progressions are of these formes.

- a. a. a. a.
- a+b. 2a+b. 3a+b.



- c. b+c. a+2b+c. 3a+3b+c. &
d. c+d. b+2c+d. a+3b+3c+d.
e. d+e c+2d+e. b+3c+3d+e.

Where the calculation of the intermediate termes may bee easily performed. The area abed depends upon the 4th Collume 1. $\frac{1}{2} \cdot -\frac{1}{8} \cdot \frac{3}{48}$ &c: (which progression may bee continued at pleasure by the helpe of this rule $\frac{0 \times 1 \times -1 \times 3 \times -5 \times 7 \times -9 \times 11 \times -13 \times 15}{0 \times 2 \times -4 \times 6 \times -8 \times 10 \times -12 \times -14 \times 16 \times -18}$ &c.) Whereby it

may appeare that, what ever the sine de=x is, the area abed is $x - \frac{x^3}{6} - \frac{x^5}{40} - \frac{x^7}{112} - \frac{5x^9}{1152} - \frac{7x^{11}}{2816} - \frac{21x^{13}}{13312} - \frac{11x^{15}}{10240}$ &c. (& the area afb is $\frac{x^3}{6} + \frac{x^5}{40} + \frac{x^7}{112}$ &c.)

Whereby also the area & angle adb may bee found. The same may bee done thus. the areas afd, abd, agd, ahd &c are in this progression

$\frac{x}{2} \cdot \frac{x}{2} + \frac{x^3}{3} \cdot \frac{x}{2} + \frac{2x^5}{6} - \frac{3x^7}{10} \cdot \frac{x}{2} + \frac{3x^9}{6} - \frac{9x^{11}}{10} + \frac{5x^{13}}{14} \cdot \frac{x}{2} + \frac{4x^{15}}{6} - \frac{18x^{17}}{10} + \frac{40x^{19}}{14} - \frac{7x^{21}}{18}$. &c As in this following Table

| | | | | | | | |
|--------------------|----------|------------------|-----|-----------------|-----|------------------|---|
| $\frac{1}{2}x$ | \times | 1. | 1. | 1. | 1. | 1. | 1 |
| $\frac{1}{6}x^3$ | \times | $\frac{1}{2}$ | 1. | $\frac{3}{2}$ | 2. | $\frac{5}{2}$ | 3 |
| $-\frac{3}{10}x^5$ | \times | $-\frac{1}{8}$ | 0. | $\frac{3}{8}$ | 1. | $\frac{15}{8}$ | 3 |
| $\frac{5}{14}x^7$ | \times | $\frac{1}{16}$ | 0. | $-\frac{1}{16}$ | 0. | $\frac{5}{16}$ | 1 |
| $-\frac{7}{18}x^9$ | \times | $-\frac{5}{128}$ | 0. | $\frac{3}{128}$ | 0. | $-\frac{5}{128}$ | 0 |
| | | = | = | = | = | = | |
| | | abd | agd | ahd | aid | and | |

By which it may bee perceived that $abd = \frac{1}{2}x + \frac{1}{12}x^3 + \frac{3}{80}x^5 + \frac{5}{224}x^7 + \frac{35x^9}{2304} + \frac{63x^{11}}{5632}$. &c. And by this

meanes haveing the area abd, (which the angle adb gives) de the sine of the angle adb may bee found. Corol: If $de=x$. & $\sqrt{1+xx} = eb$. then abc is an Hyperbola. & its area dabe is $x + \frac{x^3}{6} - \frac{x^5}{40} + \frac{x^7}{112} - \frac{5x^9}{1152} + \frac{7x^{11}}{2816}$. &c.

<73r>




1. Fiat $bd \parallel ad$, et ducantur rectæ ab, gd, ag, bd: Conicarum portiones bas, gdt, erunt æquales, item et gasb, bdtg. (fig 1.2) Bisectis enim bg, ad, in punctis r, et e; et ducto cre; erunt brg aed ordinatim applicatæ ad diametrum cure: proinde portiones reasb=redtg, et trapezia reab=redg. Ergo eorum differentiæ bas=gdt. Deinde $\triangle abg=dgb$ (per 37.1. Elem), Ergo portiones gasbv=bdgtg.

Coroll. Hinc pateat modus ducendi tangentes ad Conicas, ignoratis eorum diametris: A dato enim (fig 3) puncto a, duc ac, ab, et ijs parallelas bd, cf: linea df erit parallela tangenti ah. Si non, fiat $ae \parallel df$, et duc de, ad, af, cg; et erit portio ecd=abf=bac=aecd, quod est impos:


Porro fiat $dn=nf$, et duc ank; fiat $am=mk$, et duc pmq. Erunt ak, pq diametri.

Diametro av describatur circulus abcdv. Et centro a, describantur circuli cr, dm, gn, hq. et a punctis c, d, g, h, duc ce, df, gk, hp, perpendiculares ad diametrum av; & duc abe, acf, adk, agp. In triangulis gqh, pqh &c: angulus hgp=hpg & $\angle gqh=\angle pqh=\text{recto}$. Ergo, si ponantur trianguli qgh, qph esse infinitè parvi, ut latera hg, hq sit rectæ; erunt similes et æquales, proinde $gq=pq$. Hinc omnes lineæ gp, dk, cf, be, sunt duplices linearum gq, dn cm, br, &c. hoc est longitudo Trochoid: an est duplex longit: rectæ ag.



<74r>

De Gravitate Conicarum.



Sit $\lambda v f$ Ellipsis, et $\lambda w f$ Parabola. Ita nempe, ut λd vocatâ x; ordinatim applicata sit $dr = \sqrt{rx - \frac{rx^2}{q}}$, & $dp = \frac{rx}{2} - \frac{rx^2}{2q}$. Fiat $\lambda c = cf = \frac{1}{2}q$. & $d\{\text{illeg}\}w \parallel hfz \parallel kb \parallel rp$, & $vh \parallel wz \parallel yg \parallel \beta t$, tangentes vel sec~~illeg~~ sese aut curvas in punctis v, l, k, h, r, y, s, g, c, d, e, f, β , q, t, p, w, a, b, z.

Dico quòd, posito λf axe gravitatis, pondus parallelogrammi vkec est ad pondus portionis vsec, sicut parallelogrammi vbec ad portionem wqec.

Ponatur enim pondus lineæ dr esse $\frac{1}{2}dr \times dr$ sive $\frac{rx}{2} - \frac{rx^2}{2q}$, hoc uest dp. Erit pondus omnium linearum dr in superficie vsec contentarum, hoc est pondus superficiei vsec, æquali superficiei wqec. Et eâdem ratione pondus parallelogrammi vkec erit parallelogrammum wbec. Q.E.D.

Coroll. Pondus portionis vks est wbq, est portionis vsy est wq β , et portionis esf est eqf, & portionis fsg est fqt &c. Quæ omnia dat quadratura Parabolæ, est enim $3\beta wq = 2\beta wbq$, &c.


Eâdem ratione gravitas cujuslibet portionis Hyperbolæ cognoscatur, modò axis gravitatis transeat per centrum Hyperbolæ. Et si quævis plana superficies conicis sectionibus itâ terminetur, ut omnia conicarum centra sint in eâdem rectâ lineâ: gravitas istius superficiei inveniri potest, positâ rectâ gravitatis axi. Denique, centrum gravitatis cujusvis planæ & finitæ superficiei conicis sectionibus ita terminatæ inveniri potest, datâ quantitate istius superficiei, & vice versâ, modò centrum gravitatis axi gravitatis non coïncidat.

Sint ac, af Asymptota Hyperbolæ gc, in infinitum versus c continuatæ. Duc $de=ab=af$. & $eb = ad = \sqrt{af \times fg}$. & $fg \parallel ac$. Dico quod. Parallelogrammum ae, & superficies afgc æquiponderant circa axim abc: Etsi superficies afgc versus c sit longitudine & quantitate infinita, & non habet centrum gravitatis.

fiat enim $ar=ap$. & duc $pq \parallel ab$. & $vr \parallel ad$. et sit $pq \times \frac{1}{2}ap$ gravitas lineæ pq, erit $vr \times \frac{1}{2}vr$ gravitas lineæ vr (nom $\angle vrb=\angle apq$). sed $pq \times \frac{1}{2}pq = \frac{af \times fg}{2} = \frac{ad \times ad}{2} = vr \times \frac{1}{2}vr$. æquiponderant ergo lineæ pq, vr. Sed numerus linearum pq in superficie acgf est æqualis numero linearum vr in parallelogrammo ae nam $af=ab$, ergo ae & ag æquiponderant.

Add to the end of the Paragraph. See the mystery. In the first instance from the series $1 + \frac{1}{2} + \frac{1}{3}$ &c subduct $\frac{1}{2} + \frac{1}{3} +$ &c. & then will remain $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5}$ &c $-\frac{1}{2} - \frac{1}{3} - \frac{1}{4} - \frac{1}{5}$ &c=1=
 $1 - \frac{1}{2} + \frac{1}{2} - \frac{1}{3} + \frac{1}{3} - \frac{1}{4} + \frac{1}{4} - \frac{1}{5}$ &c = $\frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \frac{1}{4 \times 5}$ &c. And so of the rest Or take all the terms but the three first & there will remain $\frac{11}{6} = \frac{3}{1 \times 4} + \frac{3}{2 \times 5} + \frac{3}{3 \times 6}$ &c Or from this series $\frac{1}{1} + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \frac{1}{9}$ &c take all the termes but the first & there will remain $1 = \frac{2}{1 \times 3} + \frac{2}{3 \times 5} + \frac{2}{5 \times 7} + \frac{2}{7 \times 9}$ &c.

| | | | | |
|-----------|--------|---------|-----------|--------|
| 23883.12 | 107474 | 19902.6 | 3654 × 76 | 21924 |
| 214948.08 | | 1791234 | | 25578 |
| | | 895617 | | 277704 |



<75r>


Problems of Curves.

Probl. 1.

To draw Tangents or Perpendiculars to any given point of a given Curve (V B) (Sup. VA=x, AB=y)

Put the equation of the Curve=0, multip. by this Progression. $+0, \frac{y}{x}, \frac{2y}{x}, \frac{3y}{x}$ &c. according to the dimensions of x for a numerator, change the signs and multip. by this Progression $0, \frac{1}{y}, \frac{2}{y}, \frac{3}{y}$ &c for a Denominator, which shall make the Fraction designing AD to be taken forwards from A, if affirmative, otherwise, backwards.

Examp. 1. If $ax+by-xy+y^3=0$, then $\frac{byy-2xyy}{bx-xx+3yy} = AD = v$ (the Subperpendicular.




Coroll.

Hence for AF (the Subtangent) multiply by the Dimensions of y for a Numerator & by this Progr. $0, \frac{1}{x}, \frac{2}{x}, \frac{3}{x}$ &c according to the Dimension of x for a Denominator; thus results $\frac{bx-xx+3yy}{-b+2x} = AF = t$.

Prob. 2^d.

From a given point H to draw a Tangent (HB) to any Curve (VB)

Sup. VG=p, GH=q and AF=t and it is $y-q:x+p::y:t$. Th. $\frac{pq^y+xy}{y-q} = t$, or $ty-tq=xy+py$. If into this you substitute the value of t (found per Prob. 1.) you will have the nature of a Line which described will cut the propounded Curve in the desired Tangent points.



Ex^a. Thus substituting $\frac{bx-ax+3yy}{-b+xx}$ for t, there comes $bxy-xyy+3y^3-bqx+qx^2-3qy^2=2x^2y+2pxy-bx^4+bpy=0$, the Curve to be described ** But note that this may be reduced to a simpler line by adding or subtracting the nature of the given line, viz. ordering this results, it is $3y^3-3x^2y+2bxy-2pay+qxx-3qyy-bqx+bpy=0$. from whence subtracting $y^3-xyy+bxy+athrice$ there rests $-bxy-2pxy+qx^2-3qyy-bqx+bqy-3a=0$, a Conic section which described will cut the Curve in the desired tangent points.

Thus if $ax + \frac{b}{a}xx - yy = 0$, then $\frac{2yy}{a + \frac{2b}{a}x} = t = \frac{py+xy}{y-q}$ <75v> or $2yy - 2qy = \frac{2b}{a}xx + ax + \frac{2b}{a}px + ap$ & subtracting $ax + \frac{b}{a}xx - yy = 0$ twice (viz so often as the Curve has dimensions) there comes $\frac{2bpx}{a} - ax + 2qy + ap = 0$, a strait line which drawn will cut the Curve in the points desired & so of the rest.

Where note that this Problem in any Curve is ever solvable by a line of an inferior degree. And also that a line drawn from a given point may touch a Curve of two Dimensions in 1×2 points, of three in 2×3 points of 4, in 3×4 points &c unless it be polar &c.

Probl. 3.

From a given point (C) to draw a Perpendicular (CB) to any Curve (VB)

Make VE=r, EC=s and AD=v so is $y-s:r-x::y:v$ or $\frac{ry-xy}{y-s} = v$ put this valor of v in the room of v found per Prob. 1. and you have a Curve which described will cut the propounded Curve in the points to which the perpendicular may be drawn [but ever try if by means of the nature of the given Curve you can reduce this resulting to any simpler form or degree.



Ex^a. If $ax + \frac{b}{a}xx - yy = 0$, then $\frac{1}{2}a + \frac{b}{a}x = v$, or $ry-xy = \frac{1}{2}y - \frac{1}{2}a + \frac{b}{a}xy - \frac{b}{a}sx$, which appears not reducible to a simpler form.

Yet it may be worth while to try if it may be solved by a Circle by assuming $d+ex+fy=xx+yy$ and comparing these 3 equations of the given Conick, found Hyperbola and assum'd Circle to find d, e & f which if they be found by plane Geometry that Circle described will cut the Curve in the defined points. This, I say, might be tried but it would be found impossible (unless the Conic be a Parabola) because there are 4 given points thro' which the Circle must pass, which make the Probl. contradicting, since 3 are {now} to determine a Circle.

This might be done by the Parabola, but the Hyperbola falling in so naturally, and being as easily described, 'tis not worth the while.

What is said of the Conics may be applied to any other Curves.

From hence it appears that this Probl. is ever solvable by a Curve of the same degree and sometimes phaps by a Curve of an inferior Degree.


<76r>

Also from a given point may be drawn so many perpendicular, as the square of the Curves Dimensions, unless some part extraordinary sum out to infinity, or two parts come together to make it polar.

Probl. 4.

To find the Points where the Curve has a given inclination to the Basis.

Then $\frac{m}{n} = t$. Put this equal to the value of t found by Prob. 1. & you have a Curve which describ'd will cut the given Curve in the desired points.



Hence Conics have 1×2 points, those of 3 Dimensions 2×3 points, of 4 Dimens. 3×4 . &c. will satisfy this Probl, & its ever soluble by a line of an inferior Degree.

From a given point (C) to draw a line CB which shall cut any given Curve in a given Angle.

Ex^a. If $dd + \frac{a}{b}xx - yy = 0$, then is $\frac{a}{b}x = v$, and so $mry - mxy + nsy - nyy = \frac{ma}{b}xy - \frac{mas}{b}x + \frac{nar}{b}x - \frac{na}{b}xx$, which is a Conic section passing thro' the Centre of the given Curve. And <76v> it may be made an Hyperbola constantly by substituting dd for $yy - \frac{a}{b}2x$. For the result will be $dd + \frac{nar}{b}x - \frac{mas}{b}x - mry - nsy + nxy - \frac{ma}{b}2y = 0$.

Hence it appears that this Problem may ever be solved by a Curve of the same sort, and sometimes perhaps by one of an inferior.

And also that from a given point to any Curve so many lines may be drawn in a given Angle as the square of its Dimensions abating those that are imaginary, or coincident to the Pole, or should be drawn where some parts of the Curve vanishes ad infinitum.