

School of Engineering and Computer Science

# **Mathematical tools applied to Computer Science**

## **Ch5: Matrix calculus and analysis**

Kamel ATTAR

kamel.attar@epita.fr

Week #10 ♦ 14/NOV/2024 ♦

Week #11 ♦ 21/NOV/2024 ♦



## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Matrices

How to calculate  $A^{-1}$

System of Linear Equations



2

## Solving Systems with Gauss-Jordan Elimination

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Matrices

How to calculate  $A^{-1}$

System of Linear Equations



4

## Solving Systems with Gauss-Jordan Elimination



## 1 Matrices

### Matrix Definitions

### Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

### Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

### Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

## 1 Matrices

### Matrix Definitions

#### Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

#### Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

#### Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

## Definition

A matrix is a rectangular collection of numbers. Generally, matrices are denoted as bold capital letters.

## Example

$$A = \begin{pmatrix} 1 & -5 & 4 \\ 2 & 5 & 3 \end{pmatrix}$$

$A$  is a matrix with two rows and three columns. For that reason, it is called a **2 by 3** matrix. This is called the **dimension** of a matrix.

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix} \text{ is a matrix of size } (m \times n) \text{ over } \mathbb{R}.$$

- $a_{ij} \in \mathbb{R}$  are called the entries of  $A$  (or the  $(i, j)$ -th entry of  $A$ ) that describe the location of the elements of the matrix  $A$
- $a_{ii} \in \mathbb{R}$  are called the diagonal entries of  $A$ , where  $1 \leq i \leq m$  and  $1 \leq j \leq n$ .

## Definition

A matrix  $A$  of size  $(m \times n)$  is said to be

- ① a **row matrix** if  $A$  has just one row (i.e  $m = 1$ ).
- ② a **column matrix** if  $A$  has just one column (i.e  $n = 1$ ).
- ③ a **square matrix** of order  $n$  if  $A$  is a  $(n \times n)$  matrix (i.e number of columns equal to that of rows  $m = n$ ).
- ④ a **zero matrix** if all entries equal to zero (i.e  $a_{ij} = 0$  for any  $1 \leq i \leq m$  and  $1 \leq j \leq n$ ).

## Definition

A matrix  $A$  of size  $(m \times n)$  is said to be

- ① a **row matrix** if  $A$  has just one row (i.e  $m = 1$ ).
- ② a **column matrix** if  $A$  has just one column (i.e  $n = 1$ ).
- ③ a **square matrix** of order  $n$  if  $A$  is a  $(n \times n)$  matrix (i.e number of columns equal to that of rows  $m = n$ ).
- ④ a **zero matrix** if all entries equal to zero (i.e  $a_{ij} = 0$  for any  $1 \leq i \leq m$  and  $1 \leq j \leq n$ ).

## Example

- ①  $A = [1 \ 2 \ 3]$  is a row matrix of size  $(1 \times 3)$ .
- ②  $B = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$  is a column matrix of size  $(3 \times 1)$ .
- ③  $C = \begin{pmatrix} 1 & 3 \\ 0 & 6 \end{pmatrix}$  is a square matrix of size  $(2 \times 2)$ .
- ④  $D = \begin{pmatrix} 1 & 3 & -1 \\ 0 & 6 & 1 \end{pmatrix}$  is a matrix of size  $(2 \times 3)$ .

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

## ★ Transpose a Matrix ★

To take the **transpose** of a matrix, simply switch the rows and column of a matrix. The transpose of  $A$  can be denoted as  $A'$  or  $A^T$ .

### Example

$$\text{If } A = \begin{pmatrix} 1 & 2 & -1 \\ 0 & 3 & 0 \\ 2 & 1 & 2 \end{pmatrix} \text{ the } A^T = \begin{pmatrix} 1 & 0 & 2 \\ 2 & 3 & 1 \\ -1 & 0 & 2 \end{pmatrix}$$

If a matrix is its own transpose, then that matrix is said to be symmetric  $A^T = A$ . Symmetric matrices must be square matrices, with the same number of rows and columns.

### Example

$$A = \begin{pmatrix} 1 & 2 & -1 \\ 2 & 3 & 0 \\ -1 & 0 & 2 \end{pmatrix} \text{ is symmetric because } A^T = \begin{pmatrix} 1 & 2 & -1 \\ 2 & 3 & 0 \\ -1 & 0 & 2 \end{pmatrix} = A$$

## ★ Matrix Addition ★

To perform matrix **addition**, two matrices must have the same dimensions. This means they must have the same number of rows and columns. In that case simply add each individual components, like below.

### Example

$$A + B = \begin{pmatrix} 1 & 0 \\ 0 & 3 \\ 4 & 1 \end{pmatrix} + \begin{pmatrix} 3 & 1 \\ 3 & 4 \\ 5 & 2 \end{pmatrix} = \begin{pmatrix} 1+3 & 0+1 \\ 0+3 & 3+4 \\ 4+5 & 1+2 \end{pmatrix} = \begin{pmatrix} 4 & 1 \\ 3 & 7 \\ 9 & 3 \end{pmatrix}$$

Matrix addition does have many of the same properties as "normal" addition.

- ①  $(A + B) + C = A + (B + C)$ , (Associativity)
- ②  $A + B = B + A$ , (Commutativity)
- ③  $A + O = O + A = A$ .
- ④  $A^T + B^T = (A + B)^T$ .

## ★ Matrix Scalar Multiplication ★

To multiply a matrix by a **scalar**, also known as scalar multiplication, multiply every element in the matrix by the scalar

### Example

If  $A = \begin{pmatrix} 5 & -2 & 3 \\ 2 & -3 & 4 \end{pmatrix}$  and  $\alpha \in \mathbb{R}$  then  $\alpha A = \begin{pmatrix} 5\alpha & -2\alpha & 3\alpha \\ 2\alpha & -3\alpha & 4\alpha \end{pmatrix}$ .

Matrix scalar multiplication has some properties, such as

- ①  $\alpha(A + B) = \alpha A + \alpha B$ ,
- ②  $(\alpha + \beta)A = \alpha A + \beta A$ ,
- ③  $\alpha(\beta A) = (\alpha\beta A)$ .
- ④  $\alpha A^T = (\alpha A)^T$ .

## ★ Matrix Multiplication ★

We start by special case when we have two vectors. To multiply two vectors with the same length together is to take the **dot product**, also called **inner product**. This is done by multiplying every entry in the two vectors together and then adding all the products up.

### Example

$$x \cdot y = xy^T = (1 \ -5 \ 4) \begin{pmatrix} 4 \\ -2 \\ 5 \end{pmatrix} = 1 * 4 + (-5) * (-2) + 4 * 5 = 4 + 10 + 20 = 34$$

- ▶ To perform matrix multiplication, the first matrix must have the same number of columns as the second matrix has rows.


- ▶ To perform matrix multiplication, the first matrix must have the same number of columns as the second matrix has rows.
- ▶ The number of rows of the resulting matrix equals the number of rows of the first matrix, and the number of columns of the resulting matrix equals the number of columns of the second matrix.

- ▶ To perform matrix multiplication, the first matrix must have the same number of columns as the second matrix has rows.
- ▶ The number of rows of the resulting matrix equals the number of rows of the first matrix, and the number of columns of the resulting matrix equals the number of columns of the second matrix.

If  $A$  is an  $m \times r$  matrix and  $B$  is an  $r \times n$  matrix, then  $AB$  will be an  $m \times n$  matrix.

$$\begin{matrix} A & & B & = & AB \\ m \times r & & r \times n & & m \times n \\ \uparrow & \uparrow & \uparrow & & \uparrow \\ \text{insides match} & & & & \\ \uparrow & \uparrow & \uparrow & & \uparrow \\ \text{outdoors give size of } AB & & & & \end{matrix}$$

- ▶ To perform matrix multiplication, the first matrix must have the same number of columns as the second matrix has rows.
- ▶ The number of rows of the resulting matrix equals the number of rows of the first matrix, and the number of columns of the resulting matrix equals the number of columns of the second matrix.
- ▶ To find the entries in the resulting matrix, simply take the dot product of the corresponding row of the first matrix and the corresponding column of the second matrix.



$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad \times \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}_{3 \times 2}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad \times \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}_{3 \times 2}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad \times \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}_{3 \times 2}$$
$$C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ c_{31} & c_{32} \end{bmatrix}_{3 \times 2}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad \times \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}_{3 \times 2}$$
$$C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ c_{31} & c_{32} \end{bmatrix}_{3 \times 2}$$

$$c_{11} = a_{11} \cdot b_{11} + a_{12} \cdot b_{21} + a_{13} \cdot b_{31}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad \times \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}_{3 \times 2}$$

$$C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ c_{31} & c_{32} \end{bmatrix}_{3 \times 2}$$

$$c_{11} = a_{11} \cdot b_{11} + a_{12} \cdot b_{21} + a_{13} \cdot b_{31}$$

$$c_{12} = a_{11} \cdot b_{12} + a_{12} \cdot b_{22} + a_{13} \cdot b_{32}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad \times \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}_{3 \times 2}$$

$$C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ c_{31} & c_{32} \end{bmatrix}_{3 \times 2}$$

$$c_{11} = a_{11} \cdot b_{11} + a_{12} \cdot b_{21} + a_{13} \cdot b_{31}$$

$$c_{12} = a_{11} \cdot b_{12} + a_{12} \cdot b_{22} + a_{13} \cdot b_{32}$$

$$c_{21} = a_{21} \cdot b_{11} + a_{22} \cdot b_{21} + a_{23} \cdot b_{31}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad \times \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}_{3 \times 2}$$

$$C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ c_{31} & c_{32} \end{bmatrix}_{3 \times 2}$$

$$c_{11} = a_{11} \cdot b_{11} + a_{12} \cdot b_{21} + a_{13} \cdot b_{31}$$

$$c_{12} = a_{11} \cdot b_{12} + a_{12} \cdot b_{22} + a_{13} \cdot b_{32}$$

$$c_{21} = a_{21} \cdot b_{11} + a_{22} \cdot b_{21} + a_{23} \cdot b_{31}$$

$$c_{22} = a_{21} \cdot b_{12} + a_{22} \cdot b_{22} + a_{23} \cdot b_{32}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ \textcolor{green}{a_{31}} & \textcolor{green}{a_{32}} & \textcolor{green}{a_{33}} \end{bmatrix}_{3 \times 3} \quad \times \quad B = \begin{bmatrix} \textcolor{red}{b_{11}} & b_{12} \\ \textcolor{red}{b_{21}} & b_{22} \\ \textcolor{red}{b_{31}} & b_{32} \end{bmatrix}_{3 \times 2}$$

$$C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ \textcolor{blue}{c_{31}} & c_{32} \end{bmatrix}_{3 \times 2}$$

$$c_{11} = a_{11} \cdot b_{11} + a_{12} \cdot b_{21} + a_{13} \cdot b_{31}$$

$$c_{12} = a_{11} \cdot b_{12} + a_{12} \cdot b_{22} + a_{13} \cdot b_{32}$$

$$c_{21} = a_{21} \cdot b_{11} + a_{22} \cdot b_{21} + a_{23} \cdot b_{31}$$

$$c_{22} = a_{21} \cdot b_{12} + a_{22} \cdot b_{22} + a_{23} \cdot b_{32}$$

$$c_{31} = a_{31} \cdot b_{11} + a_{32} \cdot b_{21} + a_{33} \cdot b_{31}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad \times \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}_{3 \times 2}$$

$$C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ c_{31} & c_{32} \end{bmatrix}_{3 \times 2}$$

$$c_{11} = a_{11} \cdot b_{11} + a_{12} \cdot b_{21} + a_{13} \cdot b_{31}$$

$$c_{12} = a_{11} \cdot b_{12} + a_{12} \cdot b_{22} + a_{13} \cdot b_{32}$$

$$c_{21} = a_{21} \cdot b_{11} + a_{22} \cdot b_{21} + a_{23} \cdot b_{31}$$

$$c_{22} = a_{21} \cdot b_{12} + a_{22} \cdot b_{22} + a_{23} \cdot b_{32}$$

$$c_{31} = a_{31} \cdot b_{11} + a_{32} \cdot b_{21} + a_{33} \cdot b_{31}$$

$$c_{32} = a_{31} \cdot b_{12} + a_{32} \cdot b_{22} + a_{33} \cdot b_{32}$$

## Example

$$\begin{bmatrix} 2 & 3 & 1 \\ 1 & 2 & 5 \\ 3 & 4 & 2 \end{bmatrix}_{3 \times 3} \begin{bmatrix} 1 & 2 & 4 \\ 5 & 3 & 1 \\ 6 & 2 & 1 \end{bmatrix}_{3 \times 3} = \begin{bmatrix} \square & \square & 12 \\ \square & \square & \square \\ \square & 22 & \square \end{bmatrix}_{3 \times 3}$$

$$(3)(2) + (4)(3) + (2)(2) = 22$$

$$(2)(4) + (3)(1) + (1)(1) = 12$$

## Example

$$\begin{bmatrix} 2 & 3 & 1 \\ 1 & 2 & 5 \\ 3 & 4 & 2 \end{bmatrix}_{3 \times 3} \begin{bmatrix} 1 & 2 & 4 \\ 5 & 3 & 1 \\ 6 & 2 & 1 \end{bmatrix}_{3 \times 3} = \begin{bmatrix} \square & \square & 12 \\ \square & \square & \square \\ \square & 22 & \square \end{bmatrix}_{3 \times 3}$$

The diagram shows the multiplication of two 3x3 matrices. The first matrix has a green arrow pointing from the second column to the third row, and a yellow arrow pointing from the third column to the second row. The second matrix has a yellow arrow pointing from the second row to the first column, and a green arrow pointing from the third row to the second column. The result matrix has a green box around the value 12 at position (3,3) and a yellow box around the value 22 at position (2,2).

$$(3)(2) + (4)(3) + (2)(2) = 22$$

$$(2)(4) + (3)(1) + (1)(1) = 12$$

Matrix multiplication has some of the same properties as "normal" multiplication , such as

- ①  $A(B + C) = AB + AC$  and  $(A + B)C = AC + BC$  (Distributivity)
- ②  $AO = OA = O$  and  $AI = IA = A$ ,
- ③  $(\alpha A)B = A(\alpha B) = \alpha(AB)$ . (Product with a scalar)
- ④  $(AB)^T = B^T A^T$ ,

However matrix multiplication is not commutative  $AB \neq BA$

## Example

We consider the following real matrices

$$A = \begin{pmatrix} 2 & 1 \\ 2 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 2 \\ -2 & -4 \end{pmatrix}$$

Calculate  $AB$ ,  $BA$ . What do you conclude.



## Example

We consider the following real matrices

$$A = \begin{pmatrix} 2 & 1 \\ 2 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 2 \\ -2 & -4 \end{pmatrix}$$

Calculate  $AB$ ,  $BA$ . What do you conclude.

## Solution

$$AB = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \quad BA = \begin{pmatrix} 6 & 3 \\ -12 & -6 \end{pmatrix}.$$

We conclude that  $AB \neq BA$  and  $AB = O_2$  even  $A \neq O_2$  and  $B \neq O_2$ .

⌚ **Wooclap 1.** Let  $A, B, C$  and  $D$  be four matrices such that:

$$A = \begin{pmatrix} 2 & 0 \\ 0 & 1 \\ 4 & -4 \end{pmatrix} \quad B = \begin{pmatrix} 0 & 2 & 5 \\ 0 & 1 & 2 \\ -3 & 0 & 0 \end{pmatrix} \quad C = \begin{pmatrix} 5 & -1 & 3 \\ 2 & 0 & 1 \end{pmatrix}$$

and     $D = \begin{pmatrix} 3 & 2 \\ 1 & -1 \end{pmatrix}.$

Calculate (if possible)  $AB, BA, AD$  and  $DC$ .

⌚ **Wooclap 2.** Find the real numbers  $x, y, z$  and  $t$  in the following cases:

$$\textcircled{1} \quad \begin{bmatrix} 3x + y & x - 3y \\ 4z - 2t & z + t \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 3 & -5 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 2 & 8 \end{bmatrix}$$

$$\textcircled{2} \quad 2 \begin{bmatrix} x & y \\ z & t \end{bmatrix} - 5 \begin{bmatrix} x - 2 & y + 3 \\ z - 2 & t + 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 0 & 5 \end{bmatrix}$$

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

## Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

## ★ Matrix Properties ★

### Identity matrix

An **identity matrix** is a square matrix where every diagonal entry is 1 and all the other entries are 0. The following two matrices are identity matrices and diagonal matrices.

#### Example

$$I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \text{ and } I_4 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

They are called identity matrices because any matrix multiplied with an identity matrix equals itself. The diagonal entries of a matrix are the entries where the column and row number are the same.  $a_{2,2}$  is a diagonal entry but  $a_{3,5}$  is not.

## Definition

Let  $A \in M_n(\mathbb{R})$ . We say that

- ①  $A$  is **lower triangular** matrix if all the elements above the diagonal are zero (i.e.  $a_{ij} = 0, \forall i < j$ ).

$$A \text{ is lower triangular} \iff A = \begin{pmatrix} \color{red}{a_{11}} & 0 & \cdots & 0 \\ a_{21} & \color{red}{a_{22}} & & \vdots \\ \vdots & \vdots & \ddots & 0 \\ a_{n1} & a_{n2} & \cdots & \color{red}{a_{nn}} \end{pmatrix}.$$

## Example

$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 3 & 1 & 0 \end{pmatrix}$  is a lower triangular matrix.

## Definition

Let  $A \in M_n(\mathbb{R})$ . We say that

- ②  $A$  is an **upper triangular** matrix if all the elements below the diagonal are zero (i.e.  $a_{ij} = 0, \forall i > j$ ).

$$A \text{ is upper triangular} \iff A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & a_{nn} \end{pmatrix}.$$

## Example

$$B = \begin{pmatrix} 1 & -1 & -2 \\ 0 & 2 & 3 \\ 0 & 0 & 5 \end{pmatrix} \text{ is an upper triangular matrix.}$$

## Definition

Let  $A \in M_n(\mathbb{R})$ . We say that

- ③  $A$  is a **diagonal** matrix if all the elements above and below the diagonal are zero (i.e.  $a_{ij} = 0, \forall i \neq j$ ).

$$A \text{ is diagonal matrix} \iff A = \begin{pmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & a_{22} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & a_{nn} \end{pmatrix}.$$

## Example

$$C = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -3 \end{pmatrix} \text{ is a diagonal matrix.}$$

## ★ Determinants of a Matrix ★

### Definition

Let  $A$  be a square,  $2 \times 2$  matrix. The determinant of a  $A$  is denoted  $|A|$  and is given by

$$|A| = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{12}a_{21}.$$

The notation  $\det(A)$  is also used for the determinant of  $A$ .

### Example

The determinant of the matrix  $A = \begin{bmatrix} 2 & 4 \\ -3 & 1 \end{bmatrix}$  is

$$|A| = (2 \times 1) - (4 \times (-3)) = 14$$

## Definition (Cofactor)

Let  $A$  be a square matrix. The **cofactor** of the element  $a_{ij}$  is denoted by  $C_{ij}$  and is given by

$$C_{ij} = (-1)^{i+j} M_{ij},$$

where  $M_{ij}$  is the determinant of the matrix  $A$  that remains after deleting row  $i$  and column  $j$ .

## Example

Determine the cofactors of the elements  $a_{11}$  and  $a_{32}$  of  $A = \begin{bmatrix} 1 & 0 & 3 \\ 4 & -1 & 2 \\ 0 & -2 & 1 \end{bmatrix}$ .

## Definition (Cofactor)

Let  $A$  be a square matrix. The **cofactor** of the element  $a_{ij}$  is denoted by  $C_{ij}$  and is given by

$$C_{ij} = (-1)^{i+j} M_{ij},$$

where  $M_{ij}$  is the determinant of the matrix  $A$  that remains after deleting row  $i$  and column  $j$ .

## Example

Determine the cofactors of the elements  $a_{11}$  and  $a_{32}$  of  $A = \begin{bmatrix} 1 & 0 & 3 \\ 4 & -1 & 2 \\ 0 & -2 & 1 \end{bmatrix}$ .

## solution

- The cofactor of  $a_{11}$  :  $C_{11} = (-1)^{1+1} M_{11} = \begin{vmatrix} -1 & 2 \\ -2 & 1 \end{vmatrix} = 3$
- The cofactor of  $a_{32}$  :  $C_{32} = (-1)^{3+2} M_{32} = -\begin{vmatrix} 1 & 3 \\ 4 & 2 \end{vmatrix} = 10.$

## Definition

The determinant of a square matrix is the sum of the products of the elements of any row and their cofactors or any column and their co-adjugates. If  $A$  is  $n \times n$ , then

- $|A| = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in}, \quad \forall 1 \leq i \leq n,$  with respect to the  $i$ th row
- $|A| = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj}, \quad \forall 1 \leq j \leq n,$  with respect to the  $j$ th column

## Example

Evaluate the determinant of matrix  $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 4 & 2 & 1 \end{bmatrix}$  using the first row.

## Definition

The determinant of a square matrix is the sum of the products of the elements of any row and their cofactors or any column and their co-adjugates. If  $A$  is  $n \times n$ , then

- $|A| = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in}, \quad \forall 1 \leq i \leq n,$  with respect to the  $i$ th row
- $|A| = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj}, \quad \forall 1 \leq j \leq n,$  with respect to the  $j$ th column

## Example

Evaluate the determinant of matrix  $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 4 & 2 & 1 \end{bmatrix}$  using the first row.

## solution

Using the elements of the first row and their corresponding cofactors we get

$$\begin{aligned}
 |A| &= a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13} \\
 &= 1(-1)^2 \left| \begin{array}{cc} 0 & 1 \\ 2 & 1 \end{array} \right| + 2(-1)^3 \left| \begin{array}{cc} 3 & 1 \\ 4 & 1 \end{array} \right| + (-1)(-1)^4 \left| \begin{array}{cc} 3 & 0 \\ 4 & 2 \end{array} \right| \\
 &= -6.
 \end{aligned}$$

## Example

Find the determinant of  $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 4 & 2 & 1 \end{bmatrix}$  using the second row. What do you conclude?

## Example

Find the determinant of  $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 4 & 2 & 1 \end{bmatrix}$  using the second row. What do you conclude?

## solution

$$|A| = a_{21}C_{21} + a_{22}C_{22} + a_{23}C_{23} = -3 \left| \begin{array}{cc} 2 & -1 \\ 2 & 1 \end{array} \right| + 0 \left| \begin{array}{cc} 1 & -1 \\ 4 & 1 \end{array} \right| - 1 \left| \begin{array}{cc} 1 & 2 \\ 4 & 2 \end{array} \right| = -6.$$

Note that we have already evaluated this determinant in terms of the first row. As is to be expected, we obtained the same value.

 **Wooclap 4.** Evaluate the determinant of  $\begin{bmatrix} 2 & 1 & 0 & 4 \\ 0 & -1 & 0 & 2 \\ 7 & -2 & 3 & 5 \\ 0 & 1 & 0 & -3 \end{bmatrix}$ .

- If the entries of a row (resp. column) of  $A$  are zero, then  $|A| = 0$ .
- $|AB| = |A||B|$
- $|A^T| = |A|$
- $|A^n| = |A|^n$
- $|cA| = c^n|A|$

## Example

If  $A$  is a  $2 \times 2$  matrix with  $|A| = 4$ , use the properties of determinants to compute  $|A^2|$  and  $|A^T 3A|$ .

- If the entries of a row (resp. column) of  $A$  are zero, then  $|A| = 0$ .
- $|AB| = |A||B|$
- $|A^T| = |A|$
- $|A^n| = |A|^n$
- $|cA| = c^n|A|$

## Example

If  $A$  is a  $2 \times 2$  matrix with  $|A| = 4$ , use the properties of determinants to compute  $|A^2|$  and  $|A^T 3A|$ .

## Solution

- $|A^2| = |A|^2 = 4 \times 4 = 16$ .
- $|A^T 3A| = |A^T||3A| = 3^2|A^T||A| = 144$ .

## Theorem

*The determinant of a triangular matrix is the product of its diagonal elements. Thus*

$$\begin{vmatrix} \color{red}{a_{11}} & a_{12} & \cdots & a_{1n} \\ 0 & \color{red}{a_{22}} & \cdots & a_{2n} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & \color{red}{a_{nn}} \end{vmatrix} = \begin{vmatrix} \color{red}{a_{11}} & 0 & \cdots & 0 \\ a_{21} & \color{red}{a_{22}} & \ddots & \vdots \\ \vdots & \vdots & \ddots & 0 \\ a_{n1} & a_{n2} & \cdots & \color{red}{a_{nn}} \end{vmatrix} = \begin{vmatrix} \color{red}{a_{11}} & 0 & \cdots & 0 \\ 0 & \color{red}{a_{22}} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \color{red}{a_{nn}} \end{vmatrix} \\ = \color{red}{a_{11}} \times \color{red}{a_{22}} \times \cdots \times \color{red}{a_{nn}}.$$

## Example

The determinant of the matrix  $A = \begin{bmatrix} 2 & 0 & -7 & 4 \\ 0 & 3 & 1 & 0 \\ 0 & 0 & -5 & 9 \\ 0 & 0 & 0 & 1 \end{bmatrix}$  is  $|A| = 2 \times 3 \times (-5) \times 1 = -30$

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

## ★ Matrix Inverse ★

### Definition

A squared matrix  $A$  is said to be **invertible** (or **non-singular**) if there exists a matrix  $B$  of same dimension, such that  $AB = BA = I$ .  $B$  is called the inverse of  $A$  and denoted by  $A^{-1}$

- The notion of an inverse matrix only applies to square matrices.
- Not all square matrices are invertible.

### Example

Show that the following pairs of matrices are inverses

a)  $\begin{pmatrix} 3 & 4 \\ 5 & 7 \end{pmatrix}$  and  $\begin{pmatrix} 7 & -4 \\ -5 & 3 \end{pmatrix}$

b)  $\begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & -2 \\ -1 & 3 \end{pmatrix}$

c)  $\begin{pmatrix} 1 & 3 & 3 \\ 1 & 3 & 4 \\ 1 & 4 & 3 \end{pmatrix}$  and  $\begin{pmatrix} 7 & -3 & -3 \\ -1 & 0 & 1 \\ -1 & 1 & 0 \end{pmatrix}$ .

## solution

We can easily calculate the product of two matrices and find  $I$

a)  $\begin{pmatrix} 3 & 4 \\ 5 & 7 \end{pmatrix} \times \begin{pmatrix} -7 & -4 \\ -5 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} -7 & -4 \\ -5 & 3 \end{pmatrix} \times \begin{pmatrix} 3 & 4 \\ 5 & 7 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

b)  $\begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix} \times \begin{pmatrix} 1 & -2 \\ -1 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & -2 \\ -1 & 3 \end{pmatrix} \times \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

c)  $\begin{pmatrix} 1 & 3 & 3 \\ 1 & 3 & 4 \\ 1 & 4 & 3 \end{pmatrix} \times \begin{pmatrix} 7 & -3 & -3 \\ -1 & 0 & 1 \\ -1 & 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$  and  
 $\begin{pmatrix} 7 & -3 & -3 \\ -1 & 0 & 1 \\ -1 & 1 & 0 \end{pmatrix} \times \begin{pmatrix} 1 & 3 & 3 \\ 1 & 3 & 4 \\ 1 & 4 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$

 **Exercise 1.** Let matrix  $A$  be given as:

$$A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$$

And let its inverse,  $A^{-1}$ , be given as:

$$A^{-1} = \begin{bmatrix} a & -\frac{3}{5} \\ b & +\frac{2}{5} \end{bmatrix}$$

Find the values of  $a$  and  $b$ .

Let  $A$  and  $B$  be invertible matrices and  $\alpha$  a nonzero scalar. Then

$$\textcircled{1} \quad (A^{-1})^{-1} = A,$$

$$\textcircled{2} \quad (\alpha A)^{-1} = \frac{1}{\alpha} A^{-1}$$

$$\textcircled{3} \quad AB \text{ is invertible and } (AB)^{-1} = B^{-1}A^{-1}$$

$$\textcircled{4} \quad A^T \text{ is invertible and } (A^T)^{-1} = (A^{-1})^T.$$

$$\textcircled{5} \quad |A^{-1}| = \frac{1}{|A|}.$$

$$\textcircled{6} \quad \left( \begin{array}{cc} a & b \\ c & d \end{array} \right)^{-1} = \frac{1}{ad-bc} \left( \begin{array}{cc} d & -b \\ -c & a \end{array} \right).$$

Let  $A$  and  $B$  be invertible matrices and  $\alpha$  a nonzero scalar. Then

$$\textcircled{1} \quad (A^{-1})^{-1} = A,$$

$$\textcircled{2} \quad (\alpha A)^{-1} = \frac{1}{\alpha} A^{-1}$$

$$\textcircled{3} \quad AB \text{ is invertible and } (AB)^{-1} = B^{-1}A^{-1}$$

$$\textcircled{4} \quad A^T \text{ is invertible and } (A^T)^{-1} = (A^{-1})^T.$$

$$\textcircled{5} \quad |A^{-1}| = \frac{1}{|A|}.$$

$$\textcircled{6} \quad \left( \begin{array}{cc} a & b \\ c & d \end{array} \right)^{-1} = \frac{1}{ad-bc} \left( \begin{array}{cc} d & -b \\ -c & a \end{array} \right).$$

## Example

If  $A = \left( \begin{array}{cc} 1 & 2 \\ 3 & 4 \end{array} \right)$ , then  $A^{-1} = \frac{1}{4-6} \left( \begin{array}{cc} 4 & -2 \\ -3 & 1 \end{array} \right) = -\frac{1}{2} \left( \begin{array}{cc} 4 & -2 \\ -3 & 1 \end{array} \right)$ .

Let  $A$  and  $B$  be invertible matrices and  $\alpha$  a nonzero scalar. Then

$$\textcircled{1} \quad (A^{-1})^{-1} = A,$$

$$\textcircled{2} \quad (\alpha A)^{-1} = \frac{1}{\alpha} A^{-1}$$

$$\textcircled{3} \quad AB \text{ is invertible and } (AB)^{-1} = B^{-1}A^{-1}$$

$$\textcircled{4} \quad A^T \text{ is invertible and } (A^T)^{-1} = (A^{-1})^T.$$

$$\textcircled{5} \quad |A^{-1}| = \frac{1}{|A|}.$$

$$\textcircled{6} \quad \left( \begin{array}{cc} a & b \\ c & d \end{array} \right)^{-1} = \frac{1}{ad-bc} \left( \begin{array}{cc} d & -b \\ -c & a \end{array} \right).$$

## Example

If  $A = \left( \begin{array}{cc} 1 & 2 \\ 3 & 4 \end{array} \right)$ , then  $A^{-1} = \frac{1}{4-6} \left( \begin{array}{cc} 4 & -2 \\ -3 & 1 \end{array} \right) = -\frac{1}{2} \left( \begin{array}{cc} 4 & -2 \\ -3 & 1 \end{array} \right)$ .

For finding the matrix inverse in general, you can use **Gauss-Jordan Algorithm**. However, this is a rather complicated algorithm, so usually one relies upon the computer or calculator to find the matrix inverse.

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

## ★ Calculating $A^{-1}$ by Adjoint Matrix Formula ★

### Definition

Let  $A$  be  $n \times n$  matrix with  $n \geq 2$ . We define the **adjoint matrix** of  $A$  to be the  $n \times n$  matrix over  $\mathbb{R}$

$$\text{adj}(A) = \begin{pmatrix} C_{11} & C_{12} & \cdots & C_{1n} \\ C_{21} & C_{22} & \cdots & C_{2n} \\ \vdots & \vdots & & \vdots \\ C_{n1} & C_{n2} & \cdots & C_{nn} \end{pmatrix}^T$$

where for all  $i, j \in \{1, \dots, n\}$ ,  $C_{ij} = (-1)^{i+j} M_{ij}$  is the  $(i, j)$ -th cofactor of  $A$ , and  $M_{ij}$  is the determinant of the matrix  $A$  that remains.

### Example

Calculate the adjoint of  $A = \begin{bmatrix} 1 & 0 & 3 \\ 2 & 4 & 0 \\ 1 & 0 & 5 \end{bmatrix}$ .

# Solution

We know that

$$\text{adj}(A) = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T = \begin{bmatrix} C_{11} & C_{21} & C_{31} \\ C_{12} & C_{22} & C_{32} \\ C_{13} & C_{23} & C_{33} \end{bmatrix}$$

where  $C_{ij} = (-1)^{i+j} M_{ij}$  and  $M_{ij}$  is the determinant of the matrix  $A$  that remains after deleting row  $i$  and column  $j$  of  $A$ .

$$\begin{aligned} C_{11} &= + \begin{vmatrix} 4 & 0 \\ 0 & 5 \end{vmatrix} = 20 & C_{21} &= - \begin{vmatrix} 0 & 3 \\ 0 & 5 \end{vmatrix} = 0 & C_{31} &= + \begin{vmatrix} 0 & 3 \\ 4 & 0 \end{vmatrix} = -12 \\ C_{12} &= - \begin{vmatrix} 2 & 0 \\ 1 & 5 \end{vmatrix} = -10 & C_{22} &= + \begin{vmatrix} 1 & 3 \\ 1 & 5 \end{vmatrix} = 2 & C_{32} &= - \begin{vmatrix} 1 & 3 \\ 2 & 0 \end{vmatrix} = 6 \\ C_{13} &= + \begin{vmatrix} 2 & 4 \\ 1 & 0 \end{vmatrix} = -4 & C_{23} &= - \begin{vmatrix} 1 & 0 \\ 1 & 0 \end{vmatrix} = 0 & C_{33} &= + \begin{vmatrix} 1 & 0 \\ 2 & 4 \end{vmatrix} = 4 \end{aligned}$$

hence

$$\text{adj}(A) = \begin{bmatrix} 20 & 0 & -12 \\ -10 & 2 & 6 \\ -4 & 0 & 4 \end{bmatrix}$$

 **Exercise 2.** Calculate the adjoint of  $B = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 1 & 4 \\ 0 & 2 & 5 \end{bmatrix}$ .

## Proposition

Let  $A$  be a square matrix of order  $n$ . Then

- ①  $A \cdot \text{adj}(A) = \text{adj}(A) \cdot A = |A| \cdot I_n.$
- ②  $A$  is invertible if and only if  $|A| \neq 0$ .
- ③ If  $n \geq 2$  and  $A$  is invertible, then  $A^{-1} = \frac{1}{|A|} \text{adj}(A)$ .

## Proposition

Let  $A$  be a square matrix of order  $n$ . Then

- ①  $A \cdot \text{adj}(A) = \text{adj}(A) \cdot A = |A| \cdot I_n.$
- ②  $A$  is invertible if and only if  $|A| \neq 0$ .
- ③ If  $n \geq 2$  and  $A$  is invertible, then  $A^{-1} = \frac{1}{|A|} \text{adj}(A).$

## Example

Show that  $B = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 1 & 4 \\ 0 & 2 & 5 \end{bmatrix}$  is invertible then find  $B^{-1}$  by calculating  $\text{adj}(B)$ .

## solution

We have

$$|B| = \begin{vmatrix} 1 & 2 & 3 \\ -1 & 1 & 4 \\ 0 & 2 & 5 \end{vmatrix} = \begin{vmatrix} 1 & 4 \\ 2 & 5 \end{vmatrix} - 2 \begin{vmatrix} -1 & 4 \\ 0 & 5 \end{vmatrix} + 3 \begin{vmatrix} -1 & 1 \\ 0 & 2 \end{vmatrix} = 1 \neq 0$$

Therefore  $B$  is invertible. We have already calculate  $\text{adj}(B) = \begin{bmatrix} -3 & -4 & 5 \\ 5 & 5 & -7 \\ -2 & -2 & 3 \end{bmatrix}$ , then

$$B^{-1} = \frac{1}{|B|} \text{adj}(B) = \begin{bmatrix} -3 & -4 & 5 \\ 5 & 5 & -7 \\ -2 & -2 & 3 \end{bmatrix}$$

 **Exercise 3.** Show that  $C = \begin{bmatrix} 0 & -1 & 3 \\ 2 & 1 & 5 \\ -4 & 3 & -1 \end{bmatrix}$  is invertible then find  $C^{-1}$  by calculating  $\text{adj}(C)$ .

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

## ★ Elementary row operations ★

### ► Purpose of Row Operations:

- Simplify matrices to solve systems of linear equations.
- Find ranks, inverses, and determinants of matrices.

## ★ Elementary row operations ★

► **Purpose of Row Operations:**

- Simplify matrices to solve systems of linear equations.
- Find ranks, inverses, and determinants of matrices.

► **Applications:** Used in Gaussian elimination for solving linear equations.

## ★ Elementary row operations ★

► **Purpose of Row Operations:**

- Simplify matrices to solve systems of linear equations.
- Find ranks, inverses, and determinants of matrices.

► **Applications:** Used in Gaussian elimination for solving linear equations.

► **Overview of Elementary Row Operations**

- **Row Swapping:** Interchanging two rows. Notation:  $R_i \leftrightarrow R_j$
- **Row Multiplication:** Multiplying all entries in a row by a non-zero scalar. Notation:  $kR_i$
- **Row Addition:** Adding or subtracting a multiple of one row to another. Notation:  $R_i + kR_j \rightarrow R_i$

## Example ( Row Swapping)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_3} \rightarrow$$

## Example ( Row Swapping)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \quad R_1 \leftrightarrow R_3 \rightarrow \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

## Example ( Row Swapping)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_1 \leftrightarrow R_3 \rightarrow \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 2 & 4 & 6 \\ 3 & 6 & 9 \\ 1 & 2 & 3 \end{bmatrix} R_2 \rightarrow \frac{1}{3}R_2$$

## Example ( Row Swapping)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_1 \leftrightarrow R_3 \rightarrow \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 2 & 4 & 6 \\ 3 & 6 & 9 \\ 1 & 2 & 3 \end{bmatrix} R_2 \rightarrow \frac{1}{3}R_2 \begin{bmatrix} 2 & 4 & 6 \\ 1 & 2 & 3 \\ 1 & 2 & 3 \end{bmatrix}$$

## Example ( Row Swapping)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_1 \leftrightarrow R_3 \rightarrow \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 2 & 4 & 6 \\ 3 & 6 & 9 \\ 1 & 2 & 3 \end{bmatrix} R_2 \rightarrow \frac{1}{3}R_2 \begin{bmatrix} 2 & 4 & 6 \\ 1 & 2 & 3 \\ 1 & 2 & 3 \end{bmatrix}$$

## Example ( Row Replacement)

$$C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_3 \rightarrow R_3 - 7R_1$$

## Example ( Row Swapping)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_1 \leftrightarrow R_3 \rightarrow \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 2 & 4 & 6 \\ 3 & 6 & 9 \\ 1 & 2 & 3 \end{bmatrix} R_2 \rightarrow \frac{1}{3}R_2 \begin{bmatrix} 2 & 4 & 6 \\ 1 & 2 & 3 \\ 1 & 2 & 3 \end{bmatrix}$$

## Example ( Row Replacement)

$$C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_3 \rightarrow R_3 - 7R_1 \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 0 & -6 & -12 \end{bmatrix}$$

## Example ( Row Swapping)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_1 \leftrightarrow R_3 \rightarrow \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 2 & 4 & 6 \\ 3 & 6 & 9 \\ 1 & 2 & 3 \end{bmatrix} R_2 \rightarrow \frac{1}{3}R_2 \begin{bmatrix} 2 & 4 & 6 \\ 1 & 2 & 3 \\ 1 & 2 & 3 \end{bmatrix}$$

## Example ( Row Replacement)

$$C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_3 \rightarrow R_3 - 7R_1 \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 0 & -6 & -12 \end{bmatrix}$$

## Example (Combination of Operations)

$$D = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 3 & 4 \\ 3 & 5 & 7 \end{bmatrix} R_2 \rightarrow R_2 - 2R_1 \rightarrow R_3 \rightarrow R_3 - 3R_1$$

## Example ( Row Swapping)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_1 \leftrightarrow R_3 \rightarrow \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 2 & 4 & 6 \\ 3 & 6 & 9 \\ 1 & 2 & 3 \end{bmatrix} R_2 \rightarrow \frac{1}{3}R_2 \begin{bmatrix} 2 & 4 & 6 \\ 1 & 2 & 3 \\ 1 & 2 & 3 \end{bmatrix}$$

## Example ( Row Replacement)

$$C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} R_3 \rightarrow R_3 - 7R_1 \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 0 & -6 & -12 \end{bmatrix}$$

## Example (Combination of Operations)

$$D = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 3 & 4 \\ 3 & 5 & 7 \end{bmatrix} R_2 \rightarrow R_2 - 2R_1 \quad R_3 \rightarrow R_3 - 3R_1 \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 2 & 4 \end{bmatrix}$$

## ★ Calculating $A^{-1}$ by Gauss-Jordan Elimination ★

- The elimination steps create the inverse matrix while changing  $A$  to  $I$

Play around with the rows  
(adding, multiplying or swapping)  
until we make Matrix  $A$  into the  
Identity Matrix  $I$


$$\begin{array}{c} \left[ \begin{array}{c|c} A & I \end{array} \right] \\ \text{"Elementary Row Operations"} \\ \downarrow \quad \downarrow \\ \left[ \begin{array}{c|c} I & A^{-1} \end{array} \right] \end{array}$$

And by ALSO doing the changes  
to an Identity Matrix it magically  
turns into the Inverse!

## ★ Calculating $A^{-1}$ by Gauss-Jordan Elimination ★

- The elimination steps create the inverse matrix while changing  $A$  to  $I$

Play around with the rows  
(adding, multiplying or swapping)  
until we make Matrix  $A$  into the  
Identity Matrix  $I$



And by ALSO doing the changes  
to an Identity Matrix it magically  
turns into the Inverse!


- Let  $A = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$ , so to start with Gauss-Jordan on  $A$ , we first write the augmented matrix with  $I_3$ .

$$\left[ \begin{array}{c|ccc} A & : & I_3 \end{array} \right] = \left[ \begin{array}{ccc|ccc} 2 & -1 & 0 & 1 & 0 & 0 \\ -1 & 2 & -1 & 0 & 1 & 0 \\ 0 & -1 & 2 & 0 & 0 & 1 \end{array} \right]$$

## ★ Calculating $A^{-1}$ by Gauss-Jordan Elimination ★

- The elimination steps create the inverse matrix while changing  $A$  to  $I$

Play around with the rows  
(adding, multiplying or swapping)  
until we make Matrix  $A$  into the  
Identity Matrix  $I$



And by ALSO doing the changes  
to an Identity Matrix it magically  
turns into the Inverse!

- Let  $A = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$ , so to start with Gauss-Jordan on  $A$ , we first write the augmented matrix with  $I_3$ .

$$\left[ \begin{array}{c|ccc} A & : & I_3 \end{array} \right] = \left[ \begin{array}{ccc|ccc} 2 & -1 & 0 & 1 & 0 & 0 \\ -1 & 2 & -1 & 0 & 1 & 0 \\ 0 & -1 & 2 & 0 & 0 & 1 \end{array} \right]$$

- Now we do our best to turn  $A$  into an **Identity Matrix** by using the **elementary Row operations**. The goal is to make Matrix  $A$  have **1s** on the diagonal and **0s** elsewhere, and the right hand side comes along for the ride, with every operation being done on it as well.

Our row operations procedure is as follows:

- 1 We keep the first diagonal entry  $a_{11}$  and we make the rest of the **first column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & -1 & 0 & 1 & 0 & 0 \\ -1 & 2 & -1 & 0 & 1 & 0 \\ 0 & -1 & 2 & 0 & 0 & 1 \end{array} \right]$$

Our row operations procedure is as follows:

- 1 We keep the first diagonal entry  $a_{11}$  and we make the rest of the **first column** zero 0.

$$\left[ \begin{array}{ccccc|ccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ -1 & 2 & -1 & \vdots & 0 & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right]$$

Our row operations procedure is as follows:

- 1 We keep the first diagonal entry  $a_{11}$  and we make the rest of the **first column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & -1 & 0 & 1 & 0 & 0 \\ -1 & 2 & -1 & 0 & 1 & 0 \\ 0 & -1 & 2 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + \frac{1}{2}R_1}$$

Our row operations procedure is as follows:

- We keep the first diagonal entry  $a_{11}$  and we make the rest of the **first column** zero 0.

$$\left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ -1 & 2 & -1 & \vdots & 0 & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + \frac{1}{2}R_1} \left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right]$$

Our row operations procedure is as follows:

- We keep the first diagonal entry  $a_{11}$  and we make the rest of the **first column** zero 0.

$$\left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ -1 & 2 & -1 & \vdots & 0 & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + \frac{1}{2}R_1} \left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right]$$

- Then we keep the second diagonal entry  $a_{22}$  and we make all the other entries in the **second column** zero 0.

$$\left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right]$$

Our row operations procedure is as follows:

- We keep the first diagonal entry  $a_{11}$  and we make the rest of the **first column** zero 0.

$$\left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ -1 & 2 & -1 & \vdots & 0 & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + \frac{1}{2}R_1} \left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right]$$

- Then we keep the second diagonal entry  $a_{22}$  and we make all the other entries in the **second column** zero 0.

$$\left[ \begin{array}{ccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right]$$

Our row operations procedure is as follows:

- We keep the first diagonal entry  $a_{11}$  and we make the rest of the **first column** zero 0.

$$\left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ -1 & 2 & -1 & \vdots & 0 & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + \frac{1}{2}R_1} \left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right]$$

- Then we keep the second diagonal entry  $a_{22}$  and we make all the other entries in the **second column** zero 0.

$$\left[ \begin{array}{ccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right] \xrightarrow{\begin{array}{l} R_1 \rightarrow R_1 + \frac{2}{3}R_2 \\ \quad \quad \quad \rightarrow \\ R_3 \rightarrow R_3 + \frac{2}{3}R_2 \end{array}}$$

Our row operations procedure is as follows:

- We keep the first diagonal entry  $a_{11}$  and we make the rest of the **first column** zero 0.

$$\left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ -1 & 2 & -1 & \vdots & 0 & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + \frac{1}{2}R_1} \left[ \begin{array}{cccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right]$$

- Then we keep the second diagonal entry  $a_{22}$  and we make all the other entries in the **second column** zero 0.

$$\left[ \begin{array}{ccccc} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{array} \right] \xrightarrow{\begin{array}{l} R_1 \rightarrow R_1 + \frac{2}{3}R_2 \\ R_3 \rightarrow R_3 + \frac{2}{3}R_2 \end{array}} \left[ \begin{array}{ccccc} 2 & 0 & -\frac{2}{3} & \vdots & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & \vdots & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- ③ Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{ccccccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- ③ Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- ③ Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{cc|c|ccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad \begin{array}{l} R_1 \rightarrow R_1 + \frac{1}{2}R_3 \\ R_2 \rightarrow R_2 + \frac{3}{4}R_3 \end{array}$$

- ③ Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow R_1 + \frac{1}{2}R_3 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

$$R_2 \rightarrow R_2 + \frac{3}{4}R_3$$

- 3 Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow R_1 + \frac{1}{2}R_3 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- 4 The last Gauss-Jordan step is to divide each row by its pivot. The new pivots are 1.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- 3 Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow R_1 + \frac{1}{2}R_3 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- 4 The last Gauss-Jordan step is to divide each row by its pivot. The new pivots are 1.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- 3 Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow R_1 + \frac{1}{2}R_3 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- 4 The last Gauss-Jordan step is to divide each row by its pivot. The new pivots are 1.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow \frac{1}{2}R_1 \quad R_2 \rightarrow \frac{2}{3}R_2 \quad R_3 \rightarrow \frac{3}{4}R_3$$

- 3 Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow R_1 + \frac{1}{2}R_3 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- 4 The last Gauss-Jordan step is to divide each row by its pivot. The new pivots are 1.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow \frac{1}{2}R_1 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & : & \frac{3}{4} & \frac{1}{2} & \frac{1}{4} \\ 0 & 1 & 0 & : & \frac{1}{2} & 1 & \frac{1}{2} \\ 0 & 0 & 1 & : & \frac{1}{4} & \frac{1}{2} & \frac{3}{4} \end{array} \right]$$

- 3 Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow R_1 + \frac{1}{2}R_3 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- 4 The last Gauss-Jordan step is to divide each row by its pivot. The new pivots are 1.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow \frac{1}{2}R_1 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & : & \frac{3}{4} & \frac{1}{2} & \frac{1}{4} \\ 0 & 1 & 0 & : & \frac{1}{2} & 1 & \frac{1}{2} \\ 0 & 0 & 1 & : & \frac{1}{4} & \frac{1}{2} & \frac{3}{4} \end{array} \right] = [I_3 : A^{-1}]$$

- ③ Then we keep the third diagonal entry  $a_{33}$  and we make all the other entries in the **third column** zero 0.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & -\frac{2}{3} & : & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & \frac{3}{2} & -1 & : & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow R_1 + \frac{1}{2}R_3 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right]$$

- ④ The last Gauss-Jordan step is to divide each row by its pivot. The new pivots are 1.

$$\left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & : & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \frac{3}{2} & 0 & : & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \frac{4}{3} & : & \frac{1}{3} & \frac{2}{3} & 1 \end{array} \right] \quad R_1 \rightarrow \frac{1}{2}R_1 \quad \rightarrow \quad \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & : & \frac{3}{4} & \frac{1}{2} & \frac{1}{4} \\ 0 & 1 & 0 & : & \frac{1}{2} & 1 & \frac{1}{2} \\ 0 & 0 & 1 & : & \frac{1}{4} & \frac{1}{2} & \frac{3}{4} \end{array} \right] = [I_3 : A^{-1}]$$

We have reached  $I_3$  in the first half of the matrix, because  $A$  is invertible. The three columns of  $A^{-1}$  are in the second half of  $[I_3 : A^{-1}]$ .

 **Exercise 4.** Show that  $C = \begin{bmatrix} 0 & -1 & 3 \\ 2 & 1 & 5 \\ -4 & 3 & -1 \end{bmatrix}$  is invertible by finding its inverse  $C^{-1}$  using the Gauss-Jordan elimination method.

We will use Gauss-Jordan elimination to compute  $C^{-1}$ .

- **Step 1: Write the augmented matrix** We start with the augmented matrix  $[C \mid I]$ , where  $I$  is the  $3 \times 3$  identity matrix:

$$\left[ \begin{array}{ccc|ccc} 0 & -1 & 3 & 1 & 0 & 0 \\ 2 & 1 & 5 & 0 & 1 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

We will use Gauss-Jordan elimination to compute  $C^{-1}$ .

- **Step 1: Write the augmented matrix** We start with the augmented matrix  $[C \mid I]$ , where  $I$  is the  $3 \times 3$  identity matrix:

$$\left[ \begin{array}{ccc|ccc} 0 & -1 & 3 & 1 & 0 & 0 \\ 2 & 1 & 5 & 0 & 1 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

- **Step 2: Perform row operations to reduce the left-hand side to the identity matrix**

We will use Gauss-Jordan elimination to compute  $C^{-1}$ .

- ▶ **Step 1: Write the augmented matrix** We start with the augmented matrix  $[C \mid I]$ , where  $I$  is the  $3 \times 3$  identity matrix:

$$\left[ \begin{array}{ccc|ccc} 0 & -1 & 3 & 1 & 0 & 0 \\ 2 & 1 & 5 & 0 & 1 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

- ▶ **Step 2: Perform row operations to reduce the left-hand side to the identity matrix**

- Swap  $R_1$  and  $R_2$ : This is because the pivot in  $R_1$  is zero. After the swap:

$$\left[ \begin{array}{ccc|ccc} 2 & 1 & 5 & 0 & 1 & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[ \begin{array}{ccc|ccc} 0 & -1 & 3 & 1 & 0 & 0 \\ 2 & 1 & 5 & 0 & 1 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

We will use Gauss-Jordan elimination to compute  $C^{-1}$ .

- ▶ **Step 1: Write the augmented matrix** We start with the augmented matrix  $[C \mid I]$ , where  $I$  is the  $3 \times 3$  identity matrix:

$$\left[ \begin{array}{ccc|ccc} 0 & -1 & 3 & 1 & 0 & 0 \\ 2 & 1 & 5 & 0 & 1 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

- ▶ **Step 2: Perform row operations to reduce the left-hand side to the identity matrix**

- Swap  $R_1$  and  $R_2$ : This is because the pivot in  $R_1$  is zero. After the swap:

$$\left[ \begin{array}{ccc|ccc} 2 & 1 & 5 & 0 & 1 & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[ \begin{array}{ccc|ccc} 0 & -1 & 3 & 1 & 0 & 0 \\ 2 & 1 & 5 & 0 & 1 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

- Scale  $R_1$  by  $\frac{1}{2}$  to make the pivot 1:

$$\left[ \begin{array}{ccc|ccc} 2 & 1 & 5 & 0 & 1 & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow \frac{1}{2}R_1} \left[ \begin{array}{ccc|ccc} 1 & \frac{1}{2} & \frac{5}{2} & 0 & \frac{1}{2} & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

We will use Gauss-Jordan elimination to compute  $C^{-1}$ .

- ▶ **Step 1: Write the augmented matrix** We start with the augmented matrix  $[C \mid I]$ , where  $I$  is the  $3 \times 3$  identity matrix:

$$\left[ \begin{array}{ccc|ccc} 0 & -1 & 3 & 1 & 0 & 0 \\ 2 & 1 & 5 & 0 & 1 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

- ▶ **Step 2: Perform row operations to reduce the left-hand side to the identity matrix**

- Swap  $R_1$  and  $R_2$ : This is because the pivot in  $R_1$  is zero. After the swap:

$$\left[ \begin{array}{ccc|ccc} 2 & 1 & 5 & 0 & 1 & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[ \begin{array}{ccc|ccc} 2 & 1 & 5 & 0 & 1 & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

- Scale  $R_1$  by  $\frac{1}{2}$  to make the pivot 1:

$$\left[ \begin{array}{ccc|ccc} 1 & \frac{1}{2} & \frac{5}{2} & 0 & 1 & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow \frac{1}{2}R_1} \left[ \begin{array}{ccc|ccc} 1 & \frac{1}{2} & \frac{5}{2} & 0 & 1 & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right].$$

- Eliminate the first column below the pivot: Add  $4R_1$  to  $R_3$ :

$$\left[ \begin{array}{ccc|ccc} 1 & \frac{1}{2} & \frac{5}{2} & 0 & 1 & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ -4 & 3 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_3 \leftrightarrow R_3 + 4R_1} \left[ \begin{array}{ccc|ccc} 1 & \frac{1}{2} & \frac{5}{2} & 0 & 1 & 0 \\ 0 & -1 & 3 & 1 & 0 & 0 \\ 0 & 5 & 9 & 0 & 2 & 1 \end{array} \right].$$

- **Step 3: Continue row reduction** After further row operations, we reduce the left-hand side to the identity matrix. The resulting matrix is:

$$\begin{bmatrix} 1 & 0 & 0 & -\frac{1}{3} & \frac{1}{6} & -\frac{1}{6} \\ 0 & 1 & 0 & -\frac{3}{8} & \frac{1}{4} & \frac{1}{8} \\ 0 & 0 & 1 & \frac{5}{24} & \frac{1}{12} & \frac{1}{24} \end{bmatrix}.$$

- **Step 3: Continue row reduction** After further row operations, we reduce the left-hand side to the identity matrix. The resulting matrix is:

$$\left[ \begin{array}{cccccc} 1 & 0 & 0 & -\frac{1}{3} & \frac{1}{6} & -\frac{1}{6} \\ 0 & 1 & 0 & -\frac{3}{8} & \frac{1}{4} & \frac{1}{8} \\ 0 & 0 & 1 & \frac{5}{24} & \frac{1}{12} & \frac{1}{24} \end{array} \right].$$

- **Step 4: Extract the inverse matrix** The right-hand side of the augmented matrix is  $C^{-1}$ :

$$C^{-1} = \left[ \begin{array}{ccc} -\frac{1}{3} & \frac{1}{6} & -\frac{1}{6} \\ -\frac{3}{8} & \frac{1}{4} & \frac{1}{8} \\ \frac{5}{24} & \frac{1}{12} & \frac{1}{24} \end{array} \right].$$

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

## Definition

A system of  $m$  linear equations in  $n$  variables  $x_1, x_2, x_3, \dots, x_n$  is one that can be written in the form

$$(S) \quad \left\{ \begin{array}{rcl} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n & = & b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n & = & b_2 \\ \vdots & & \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n & = & b_m \end{array} \right.$$

where the coefficients  $a_{ij}$  and  $b_j$  are constants for all  $1 \leq i \leq m$  and  $1 \leq j \leq n$ .

## Definition

A system of  $m$  linear equations in  $n$  variables  $x_1, x_2, x_3, \dots, x_n$  is one that can be written in the form

$$(S) \quad \left\{ \begin{array}{rcl} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n & = & b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n & = & b_2 \\ \vdots & & \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n & = & b_m \end{array} \right.$$

where the coefficients  $a_{ij}$  and  $b_j$  are constants for all  $1 \leq i \leq m$  and  $1 \leq j \leq n$ .

## Example

The following is an example of a system of three linear equations.

$$\left\{ \begin{array}{rcl} x_1 + x_2 + x_3 & = & 2 \\ 2x_1 + 3x_2 + x_3 & = & 3 \\ x_1 - x_2 - 2x_3 & = & -6 \end{array} \right.$$

It can be seen on substitution that  $x_1 = -1, x_2 = 1, x_3 = 2$  is a solution to this system.

The system ( $S$ ), can be written in the form  $AX = B$ , where

$$A = \underbrace{\begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}}_{\text{coefficient matrix}}, B = \underbrace{\begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}}_{\text{variable matrix}} \text{ and } X = \underbrace{\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{pmatrix}}_{\text{constant matrix}}.$$

The system ( $S$ ), can be written in the form  $AX = B$ , where

$$A = \underbrace{\begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}}_{\text{coefficient matrix}}, B = \underbrace{\begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}}_{\text{variable matrix}} \text{ and } X = \underbrace{\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{pmatrix}}_{\text{constant matrix}}.$$

How to write  $A$

- ▶ Write the coefficients of the  $x$ -terms as the numbers down the first column.
- ▶ Write the coefficients of the  $y$ -terms as the numbers down the second column.
- ▶ If there are  $z$ -terms, write the coefficients as the numbers down the third column.

$$\left\{ \begin{array}{lcl} x + 2y - z & = & 3 \\ 2x - y + 2z & = & 6 \\ x - 3y + 3z & = & 4 \end{array} \right. \implies A = \begin{pmatrix} 1 & 2 & -1 \\ 2 & -1 & 2 \\ 1 & -3 & 3 \end{pmatrix}$$

The system ( $S$ ), can be written in the form  $AX = B$ , where

$$A = \underbrace{\begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}}_{\text{coefficient matrix}}, B = \underbrace{\begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}}_{\text{variable matrix}} \text{ and } X = \underbrace{\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{pmatrix}}_{\text{constant matrix}}.$$

How to write  $A$

- ▶ Write the coefficients of the  $x$ -terms as the numbers down the first column.
- ▶ Write the coefficients of the  $y$ -terms as the numbers down the second column.
- ▶ If there are  $z$ -terms, write the coefficients as the numbers down the third column.

$$\left\{ \begin{array}{lcl} x + 2y - z & = & 3 \\ 2x - y + 2z & = & 6 \\ x - 3y + 3z & = & 4 \end{array} \right. \implies A = \begin{pmatrix} 1 & 2 & -1 \\ 2 & -1 & 2 \\ 1 & -3 & 3 \end{pmatrix}$$

How to write  $B$

- ▶ Write the constants raw by raw.

$$\left\{ \begin{array}{lcl} x + 2y - z & = & 3 \\ 2x - y + 2z & = & 6 \\ x - 3y + 3z & = & 4 \end{array} \right. \implies B = \begin{pmatrix} 3 \\ 6 \\ 4 \end{pmatrix}$$

## Definition

We call augmented matrix of the system ( $S$ ), is one that can be written in the form

$$(A|B) = \left( \begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & \vdots & & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{array} \right)$$

## Definition

We call augmented matrix of the system ( $S$ ), is one that can be written in the form

$$(A|B) = \left( \begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & \vdots & & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{array} \right)$$

## Example

The augmented matrix of  $\begin{cases} x_1 + x_2 + x_3 = 2 \\ 2x_1 + 3x_2 + x_3 = 3 \\ x_1 - x_2 - 2x_3 = -6 \end{cases}$  is  $\left( \begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 2 & 3 & 1 & 3 \\ 1 & -1 & -2 & -6 \end{array} \right)$ .

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

- To solve a system of linear equations using an inverse matrix, let  $A$  be the coefficient matrix, let  $X$  be the variable matrix, and let  $B$  be the constant matrix. Thus, we want to solve a system

$$AX = B.$$

For example, look at the following system of equations.

$$\begin{cases} x + y + z &= 1 \\ x + z &= 1 \\ x + 2y + 4z &= 1 \end{cases}$$

- To solve a system of linear equations using an inverse matrix, let  $A$  be the coefficient matrix, let  $X$  be the variable matrix, and let  $B$  be the constant matrix. Thus, we want to solve a system

$$AX = B.$$

For example, look at the following system of equations.

$$\begin{cases} x + y + z &= 1 \\ x + z &= 1 \\ x + 2y + 4z &= 1 \end{cases}$$

- ① From this system, the coefficient matrix is

$$A = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 2 & 4 \end{pmatrix}}_{\text{coefficient matrix}}, \quad X = \underbrace{\begin{pmatrix} x \\ y \\ z \end{pmatrix}}_{\text{variable matrix}} \quad \text{and} \quad B = \underbrace{\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}}_{\text{constant matrix}}.$$

Then  $AX = B$  looks like

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 2 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

- ▶ To solve the system  $AX = B$  for  $X$ , we would simply multiply both sides by the multiplicative inverse (reciprocal) of  $A$ . Thus,

$$\begin{aligned}(A^{-1})AX &= (A^{-1})B \\ ((A^{-1})A)X &= (A^{-1})B \\ IX &= (A^{-1})B \\ X &= (A^{-1})B\end{aligned}$$

- To solve the system  $AX = B$  for  $X$ , we would simply multiply both sides by the multiplicative inverse (reciprocal) of  $A$ . Thus,

$$\begin{aligned}(A^{-1})AX &= (A^{-1})B \\ ((A^{-1})A)X &= (A^{-1})B \\ IX &= (A^{-1})B \\ X &= (A^{-1})B\end{aligned}$$

- ② So first we need to prove that  $A$  is invertible matrix. Using row echelon form we have

$$\begin{aligned}|A| &= \left| \begin{array}{ccc|c} 1 & 1 & 1 & R_2 \rightarrow R_2 - R_1 \\ 1 & 0 & 1 & R_3 \rightarrow R_3 - R_1 \\ 1 & 2 & 4 & \end{array} \right| = \left| \begin{array}{ccc|c} 1 & 1 & 1 & \\ 0 & -1 & 0 & \\ 0 & 1 & 3 & R_3 \rightarrow R_3 + R_2 \end{array} \right| \\ &= \left| \begin{array}{ccc} 1 & 1 & 1 \\ 0 & -1 & 0 \\ 0 & 0 & 3 \end{array} \right| = -3.\end{aligned}$$

Since  $|A| \neq 0$ , then  $A$  is invertible.

③ Now, we need to calculate  $A^{-1}$ . Using any method to calculate the inverse of a 3 by 3 matrix, we have:

$$\text{adj}(A) = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T = \begin{bmatrix} C_{11} & C_{21} & C_{31} \\ C_{12} & C_{22} & C_{32} \\ C_{13} & C_{23} & C_{33} \end{bmatrix}$$

where  $C_{ij} = (-1)^{i+j} M_{ij}$  and  $M_{ij}$  is the determinant of the matrix  $A$  that remains after deleting row  $i$  and column  $j$  of  $A$ .

③ Now, we need to calculate  $A^{-1}$ . Using any method to calculate the inverse of a 3 by 3 matrix, we have:

$$\text{adj}(A) = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T = \begin{bmatrix} C_{11} & C_{21} & C_{31} \\ C_{12} & C_{22} & C_{32} \\ C_{13} & C_{23} & C_{33} \end{bmatrix}$$

where  $C_{ij} = (-1)^{i+j} M_{ij}$  and  $M_{ij}$  is the determinant of the matrix  $A$  that remains after deleting row  $i$  and column  $j$  of  $A$ .

|              |   |              |   |              |   |
|--------------|---|--------------|---|--------------|---|
| $C_{11} = +$ | $\begin{vmatrix} 0 & 1 \\ 2 & 4 \end{vmatrix} = -2$ | $C_{12} = -$ | $\begin{vmatrix} 1 & 1 \\ 1 & 4 \end{vmatrix} = -3$ | $C_{13} = +$ | $\begin{vmatrix} 1 & 0 \\ 1 & 2 \end{vmatrix} = 2$  |
| $C_{21} = -$ | $\begin{vmatrix} 1 & 1 \\ 2 & 4 \end{vmatrix} = -2$ | $C_{22} = +$ | $\begin{vmatrix} 1 & 1 \\ 1 & 4 \end{vmatrix} = 3$  | $C_{23} = -$ | $\begin{vmatrix} 1 & 1 \\ 1 & 2 \end{vmatrix} = -1$ |
| $C_{31} = +$ | $\begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix} = 1$  | $C_{32} = -$ | $\begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} = 0$  | $C_{33} = +$ | $\begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} = -1$ |

③ Now, we need to calculate  $A^{-1}$ . Using any method to calculate the inverse of a 3 by 3 matrix, we have:

$$\text{adj}(A) = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T = \begin{bmatrix} C_{11} & C_{21} & C_{31} \\ C_{12} & C_{22} & C_{32} \\ C_{13} & C_{23} & C_{33} \end{bmatrix}$$

where  $C_{ij} = (-1)^{i+j} M_{ij}$  and  $M_{ij}$  is the determinant of the matrix  $A$  that remains after deleting row  $i$  and column  $j$  of  $A$ .

|              |  |        |              |  |        |              |  |        |
|--------------|--|--------|--------------|--|--------|--------------|--|--------|
| $C_{11} = +$ | $\begin{matrix} 0 & 1 \\ 2 & 4 \end{matrix}$ | $= -2$ | $C_{12} = -$ | $\begin{matrix} 1 & 1 \\ 1 & 4 \end{matrix}$ | $= -3$ | $C_{13} = +$ | $\begin{matrix} 1 & 0 \\ 1 & 2 \end{matrix}$ | $= 2$  |
| $C_{21} = -$ | $\begin{matrix} 1 & 1 \\ 2 & 4 \end{matrix}$ | $= -2$ | $C_{22} = +$ | $\begin{matrix} 1 & 1 \\ 1 & 4 \end{matrix}$ | $= 3$  | $C_{23} = -$ | $\begin{matrix} 1 & 1 \\ 1 & 2 \end{matrix}$ | $= -1$ |
| $C_{31} = +$ | $\begin{matrix} 1 & 1 \\ 0 & 1 \end{matrix}$ | $= 1$  | $C_{32} = -$ | $\begin{matrix} 1 & 1 \\ 1 & 1 \end{matrix}$ | $= 0$  | $C_{33} = +$ | $\begin{matrix} 1 & 1 \\ 1 & 0 \end{matrix}$ | $= -1$ |

Hence  $\text{adj}(A) = \begin{bmatrix} -2 & -2 & 1 \\ -3 & 3 & 0 \\ 2 & -1 & -1 \end{bmatrix}$  and so

$$A^{-1} = \frac{1}{|A|} \text{adj}(A) = \begin{bmatrix} 2/3 & 2/3 & -1/3 \\ 1 & -1 & 0 \\ -2/3 & 1/3 & 1/3 \end{bmatrix}.$$

④ Now we are ready to solve. Multiply both sides of the equation by  $A^{-1}$ . We have

$$AX = B \iff X = A^{-1}B \iff X = \begin{bmatrix} 2/3 & 2/3 & -1/3 \\ 1 & -1 & 0 \\ -2/3 & 1/3 & 1/3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

 **Exercise 5.** Solve the following system using the inverse of a matrix

$$\left\{ \begin{array}{rcl} 5x + 15y + 56z & = & 35 \\ -4x - 11y - 41z & = & -26 \\ -x - 3y - 11z & = & -7 \end{array} \right.$$

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

- ▶ The **Gaussian elimination** method refers to a strategy used to obtain the row-echelon form of a matrix.

- ▶ The **Gaussian elimination** method refers to a strategy used to obtain the row-echelon form of a matrix.
- ▶ The goal is to write matrix  $A$  with the number 1 as the entry down the main diagonal and have all zeros below.

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \xrightarrow{\text{Gaussian elimination}} \begin{bmatrix} 1 & b_{12} & b_{13} \\ 0 & 1 & b_{23} \\ 0 & 0 & 1 \end{bmatrix}$$

- ▶ The **Gaussian elimination** method refers to a strategy used to obtain the row-echelon form of a matrix.
- ▶ The goal is to write matrix  $A$  with the number 1 as the entry down the main diagonal and have all zeros below.

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \xrightarrow{\text{Gaussian elimination}} \begin{bmatrix} 1 & b_{12} & b_{13} \\ 0 & 1 & b_{23} \\ 0 & 0 & 1 \end{bmatrix}$$

- ▶ The first step of the Gaussian strategy includes obtaining a 1 as the first entry, so that row 1 may be used to alter the rows below.

- ▶ The **Gaussian elimination** method refers to a strategy used to obtain the row-echelon form of a matrix.
- ▶ The goal is to write matrix  $A$  with the number 1 as the entry down the main diagonal and have all zeros below.

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \xrightarrow{\text{Gaussian elimination}} \begin{bmatrix} 1 & b_{12} & b_{13} \\ 0 & 1 & b_{23} \\ 0 & 0 & 1 \end{bmatrix}$$

- ▶ The first step of the Gaussian strategy includes obtaining a 1 as the first entry, so that row 1 may be used to alter the rows below.
- 1 The first equation should have a leading coefficient of 1. Interchange rows or multiply by a constant, if necessary.
  - 2 Use row operations to obtain zeros down the first column below the first entry of 1
  - 3 Use row operations to obtain a 1 in row 2, column 2.
  - 4 Use row operations to obtain zeros down column 2, below the entry of 1.
  - 5 Use row operations to obtain a 1 in row 3, column 3.
  - 6 Continue this process for all rows until there is a 1 in every entry down the main diagonal and there are only zeros below.
  - 7 If any rows contain all zeros, place them at the bottom.

 Wooclap 5.

Consider the following linear system .

$$(\mathcal{S}) := \begin{cases} 6x + y + z &= 12 \\ 2x + 4y &= 0 \\ x + 2y + 6z &= 6 \end{cases}$$

- (a) Find the matrices  $A$ ,  $B$  and  $X$  such that  $(\mathcal{S}) \Leftrightarrow AX = B$ .
- (b) Calculate the determinant of  $A$ . Deduce that  $A$  is invertible.
- (c) Calculate the adjoint of  $A$  and find  $A^{-1}$ .
- (d) Deduce the solution of the linear system  $(\mathcal{S})$ .
- (e) Refind the solution of the system  $(\mathcal{S})$  by an another method.

## 1 Matrices

Matrix Definitions

Matrix Arithmetic

Transpose a Matrix

Matrix Addition

Matrix Scalar Multiplication

Matrix Multiplication

Matrix Properties

Determinants of a Matrix

Properties of Determinant

Determinant of Triangular and Diagonal Matrices

Matrix Inverse

## 2 How to calculate $A^{-1}$

Calculating  $A^{-1}$  by Adjoint Matrix Formula

Calculating  $A^{-1}$  by Gauss-Jordan Elimination

Elementary row operations

## 3 System of Linear Equations

Solving Systems with Inverses

Solving Systems with Gaussian Elimination

Solving Systems with Gauss-Jordan Elimination

The Gauss-Jordan elimination method to solve a system of linear equations is described in the following steps.

The Gauss-Jordan elimination method to solve a system of linear equations is described in the following steps.

- 1 Write the augmented matrix of the system.

The Gauss-Jordan elimination method to solve a system of linear equations is described in the following steps.

- 1 Write the augmented matrix of the system.
- 2 Use row operations to transform the augmented matrix in the form described below, which is called the reduced row echelon form (RREF).

The Gauss-Jordan elimination method to solve a system of linear equations is described in the following steps.

- 1 Write the augmented matrix of the system.
- 2 Use row operations to transform the augmented matrix in the form described below, which is called the reduced row echelon form (RREF).
  - (a) The rows (if any) consisting entirely of zeros are grouped together at the bottom of the matrix.

The Gauss-Jordan elimination method to solve a system of linear equations is described in the following steps.

- 1 Write the augmented matrix of the system.
- 2 Use row operations to transform the augmented matrix in the form described below, which is called the reduced row echelon form (RREF).
  - (a) The rows (if any) consisting entirely of zeros are grouped together at the bottom of the matrix.
  - (b) In each row that does not consist entirely of zeros, the leftmost nonzero element is a 1 (called a leading 1 or a pivot).

The Gauss-Jordan elimination method to solve a system of linear equations is described in the following steps.

- 1 Write the augmented matrix of the system.
- 2 Use row operations to transform the augmented matrix in the form described below, which is called the reduced row echelon form (RREF).
  - (a) The rows (if any) consisting entirely of zeros are grouped together at the bottom of the matrix.
  - (b) In each row that does not consist entirely of zeros, the leftmost nonzero element is a 1 (called a leading 1 or a pivot).
  - (c) Each column that contains a leading 1 has zeros in all other entries.

The Gauss-Jordan elimination method to solve a system of linear equations is described in the following steps.

- 1 Write the augmented matrix of the system.
- 2 Use row operations to transform the augmented matrix in the form described below, which is called the reduced row echelon form (RREF).
  - (a) The rows (if any) consisting entirely of zeros are grouped together at the bottom of the matrix.
  - (b) In each row that does not consist entirely of zeros, the leftmost nonzero element is a 1 (called a leading 1 or a pivot).
  - (c) Each column that contains a leading 1 has zeros in all other entries.
  - (d) The leading 1 in any row is to the left of any leading 1's in the rows below it.

The Gauss-Jordan elimination method to solve a system of linear equations is described in the following steps.

- 1** Write the augmented matrix of the system.
- 2** Use row operations to transform the augmented matrix in the form described below, which is called the reduced row echelon form (RREF).
  - (a) The rows (if any) consisting entirely of zeros are grouped together at the bottom of the matrix.
  - (b) In each row that does not consist entirely of zeros, the leftmost nonzero element is a 1 (called a leading 1 or a pivot).
  - (c) Each column that contains a leading 1 has zeros in all other entries.
  - (d) The leading 1 in any row is to the left of any leading 1's in the rows below it.
- 3** Stop process in step **2** if you obtain a row whose elements are all zeros except the last one on the right. In that case, the system is inconsistent and has no solutions. Otherwise, finish step 2 and read the solutions of the system from the final matrix.

The Gauss-Jordan elimination method to solve a system of linear equations is described in the following steps.

- 1 Write the augmented matrix of the system.
- 2 Use row operations to transform the augmented matrix in the form described below, which is called the reduced row echelon form (RREF).
  - (a) The rows (if any) consisting entirely of zeros are grouped together at the bottom of the matrix.
  - (b) In each row that does not consist entirely of zeros, the leftmost nonzero element is a 1 (called a leading 1 or a pivot).
  - (c) Each column that contains a leading 1 has zeros in all other entries.
  - (d) The leading 1 in any row is to the left of any leading 1's in the rows below it.
- 3 Stop process in step 2 if you obtain a row whose elements are all zeros except the last one on the right. In that case, the system is inconsistent and has no solutions. Otherwise, finish step 2 and read the solutions of the system from the final matrix.

Note: When doing step 2, row operations can be performed in any order. Try to choose row operations so that as few fractions as possible are carried through the computation. This makes calculation easier when working by hand.

## Example

Use the Gause-Jordan elimination method to solve the system:

$$\left\{ \begin{array}{rcl} x + 2y - z & = & 4 \\ 2x - y + 5z & = & 5 \\ 3x + 4y + z & = & 9 \end{array} \right. .$$

## Example

Use the Gause-Jordan elimination method to solve the system:

$$\left\{ \begin{array}{rcl} x + 2y - z & = & 4 \\ 2x - y + 5z & = & 5 \\ 3x + 4y + z & = & 9 \end{array} \right. .$$

## solution

$$\left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 2 & -1 & 5 & 5 \\ 3 & 4 & 1 & 9 \end{array} \right)$$

## Example

Use the Gause-Jordan elimination method to solve the system:

$$\left\{ \begin{array}{rcl} x + 2y - z & = & 4 \\ 2x - y + 5z & = & 5 \\ 3x + 4y + z & = & 9 \end{array} \right.$$

## solution

$$\left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 2 & -1 & 5 & 5 \\ 3 & 4 & 1 & 9 \end{array} \right) \quad R_2 \rightarrow R_2 - 2R_1 \quad \rightarrow \\ R_3 \rightarrow R_3 - 3R_1$$

## Example

Use the Gause-Jordan elimination method to solve the system:

$$\left\{ \begin{array}{rcl} x + 2y - z & = & 4 \\ 2x - y + 5z & = & 5 \\ 3x + 4y + z & = & 9 \end{array} \right. .$$

## solution

$$\left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 2 & -1 & 5 & 5 \\ 3 & 4 & 1 & 9 \end{array} \right) \quad R_2 \rightarrow R_2 - 2R_1 \quad \rightarrow \quad \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & -5 & 7 & -3 \\ 0 & -2 & 4 & -3 \end{array} \right)$$

## Example

Use the Gause-Jordan elimination method to solve the system:

$$\left\{ \begin{array}{lcl} x + 2y - z & = & 4 \\ 2x - y + 5z & = & 5 \\ 3x + 4y + z & = & 9 \end{array} \right. .$$

## solution

$$\left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 2 & -1 & 5 & 5 \\ 3 & 4 & 1 & 9 \end{array} \right) \quad R_2 \rightarrow R_2 - 2R_1 \quad \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & -5 & 7 & -3 \\ 0 & -2 & 4 & -3 \end{array} \right) \quad R_3 \rightarrow R_3 - \frac{2}{5}R_2$$

$$\rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & -5 & 7 & -3 \\ 0 & 0 & \frac{6}{5} & -\frac{9}{5} \end{array} \right)$$

## Example

Use the Gause-Jordan elimination method to solve the system:

$$\left\{ \begin{array}{lcl} x + 2y - z & = & 4 \\ 2x - y + 5z & = & 5 \\ 3x + 4y + z & = & 9 \end{array} \right. .$$

## solution

$$\begin{array}{ccc}
 \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 2 & -1 & 5 & 5 \\ 3 & 4 & 1 & 9 \end{array} \right) & R_2 \rightarrow R_2 - 2R_1 & \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & -5 & 7 & -3 \\ 0 & -2 & 4 & -3 \end{array} \right) \\
 & R_3 \rightarrow R_3 - 3R_1 & & R_3 \rightarrow R_3 - \frac{2}{5}R_2 \\
 \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & -5 & 7 & -3 \\ 0 & 0 & \frac{6}{5} & -\frac{9}{5} \end{array} \right) & R_2 \rightarrow -\frac{1}{5}R_2 & \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & 1 & -\frac{7}{5} & \frac{3}{5} \\ 0 & 0 & 1 & -\frac{3}{2} \end{array} \right)
 \end{array}$$

## Example

Use the Gause-Jordan elimination method to solve the system:

$$\left\{ \begin{array}{l} x + 2y - z = 4 \\ 2x - y + 5z = 5 \\ 3x + 4y + z = 9 \end{array} \right.$$

## solution

$$\begin{array}{ccc|c}
 1 & 2 & -1 & 4 \\
 2 & -1 & 5 & 5 \\
 3 & 4 & 1 & 9
 \end{array}
 \begin{array}{l}
 R_2 \rightarrow R_2 - 2R_1 \\
 R_3 \rightarrow R_3 - 3R_1
 \end{array}
 \rightarrow
 \begin{array}{ccc|c}
 1 & 2 & -1 & 4 \\
 0 & -5 & 7 & -3 \\
 0 & -2 & 4 & -3
 \end{array}
 \begin{array}{l}
 R_3 \rightarrow R_3 - \frac{2}{5}R_2
 \end{array}
 \\
 \rightarrow
 \begin{array}{ccc|c}
 1 & 2 & -1 & 4 \\
 0 & -5 & 7 & -3 \\
 0 & 0 & \frac{6}{5} & -\frac{9}{5}
 \end{array}
 \begin{array}{l}
 R_2 \rightarrow -\frac{1}{5}R_2 \\
 R_3 \rightarrow \frac{5}{6}R_3
 \end{array}
 \rightarrow
 \begin{array}{ccc|c}
 1 & 2 & -1 & 4 \\
 0 & 1 & -\frac{7}{5} & \frac{3}{5} \\
 0 & 0 & 1 & -\frac{3}{2}
 \end{array}
 \begin{array}{l}
 R_1 \rightarrow R_1 + R_3 \\
 R_2 \rightarrow R_2 + \frac{7}{5}R_3
 \end{array}
 \\
 \rightarrow
 \begin{array}{ccc|c}
 1 & 2 & 0 & \frac{5}{2} \\
 0 & 1 & 0 & -\frac{3}{2} \\
 0 & 0 & 1 & -\frac{3}{2}
 \end{array}
 \end{array}$$

## Example

Use the Gause-Jordan elimination method to solve the system:

$$\left\{ \begin{array}{l} x + 2y - z = 4 \\ 2x - y + 5z = 5 \\ 3x + 4y + z = 9 \end{array} \right.$$

## solution

$$\begin{array}{ccc}
 \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 2 & -1 & 5 & 5 \\ 3 & 4 & 1 & 9 \end{array} \right) & R_2 \rightarrow R_2 - 2R_1 & \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & -5 & 7 & -3 \\ 0 & -2 & 4 & -3 \end{array} \right) \\
 & R_3 \rightarrow R_3 - 3R_1 & & R_3 \rightarrow R_3 - \frac{2}{5}R_2 \\
 \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & -5 & 7 & -3 \\ 0 & 0 & \frac{6}{5} & -\frac{9}{5} \end{array} \right) & R_2 \rightarrow -\frac{1}{5}R_2 & R_1 \rightarrow R_1 + R_3 \\
 & R_3 \rightarrow \frac{5}{6}R_3 & \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & 1 & -\frac{7}{5} & \frac{3}{5} \\ 0 & 0 & 1 & -\frac{3}{2} \end{array} \right) & R_2 \rightarrow R_2 + \frac{7}{5}R_3 \\
 \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & 0 & \frac{5}{2} \\ 0 & 1 & 0 & -\frac{3}{2} \\ 0 & 0 & 1 & -\frac{3}{2} \end{array} \right) & R_1 \rightarrow R_1 - 2R_2 & \rightarrow \left( \begin{array}{ccc|c} 1 & 0 & 0 & \frac{11}{2} \\ 0 & 1 & 0 & -\frac{3}{2} \\ 0 & 0 & 1 & -\frac{3}{2} \end{array} \right)
 \end{array}$$

## Example

Use the Gause-Jordan elimination method to solve the system:

$$\left\{ \begin{array}{l} x + 2y - z = 4 \\ 2x - y + 5z = 5 \\ 3x + 4y + z = 9 \end{array} \right.$$

## solution

$$\begin{array}{ccc}
 \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 2 & -1 & 5 & 5 \\ 3 & 4 & 1 & 9 \end{array} \right) & R_2 \rightarrow R_2 - 2R_1 & \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & -5 & 7 & -3 \\ 0 & -2 & 4 & -3 \end{array} \right) \\
 & R_3 \rightarrow R_3 - 3R_1 & & R_3 \rightarrow R_3 - \frac{2}{5}R_2 \\
 \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & -5 & 7 & -3 \\ 0 & 0 & \frac{6}{5} & -\frac{9}{5} \end{array} \right) & R_2 \rightarrow -\frac{1}{5}R_2 & R_1 \rightarrow R_1 + R_3 \\
 & R_3 \rightarrow \frac{5}{6}R_3 & \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & 1 & -\frac{7}{5} & \frac{3}{5} \\ 0 & 0 & 1 & -\frac{3}{2} \end{array} \right) & R_2 \rightarrow R_2 + \frac{7}{5}R_3 \\
 \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & 0 & \frac{5}{2} \\ 0 & 1 & 0 & -\frac{3}{2} \\ 0 & 0 & 1 & -\frac{3}{2} \end{array} \right) & R_1 \rightarrow R_1 - 2R_2 & \rightarrow \left( \begin{array}{ccc|c} 1 & 0 & 0 & \frac{11}{2} \\ 0 & 1 & 0 & -\frac{3}{2} \\ 0 & 0 & 1 & -\frac{3}{2} \end{array} \right)
 \end{array}$$

Thus  $x = 11/2$ ,  $y = -3/2$ ,  $z = -3/2$ .

 **Exercise 6.** Use the Gause-Jordan elimination method to solve the system:

$$\begin{cases} x + y - 2z = 9 \\ 2x + 4y - 3z = 1 \\ 3x + 6y - 5z = 0 \end{cases} .$$