

①

MAT 2384 - Solutions to Suggested Problems
Exact ODEs - Integrating factors

$$1) (e^y - ye^x) dx + (xe^y - e^x) dy = 0$$

$$M(x,y) = e^y - ye^x, \quad N(x,y) = xe^y - e^x$$

$$\frac{\partial M}{\partial y} = e^y - e^x, \quad \frac{\partial N}{\partial x} = e^y - e^x. \quad \text{The equation is exact since}$$

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}. \quad \text{we look for a function } F(x,y) \text{ such that}$$

$$\frac{\partial F}{\partial x} = M(x,y) = e^y - ye^x \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x,y) = xe^y - e^x$$

$$\frac{\partial F}{\partial x} = e^y - ye^x \Rightarrow F(x,y) = \int (e^y - ye^x) dx + g(y) \quad \text{where}$$

$g(y)$ is a function of y only.

$$F(x,y) = xe^y - ye^x + g(y) \quad \text{which gives} \quad \frac{\partial F}{\partial y} = xe^y - e^x + g'(y)$$

$$\text{Now } \frac{\partial F}{\partial y} = xe^y - e^x \Rightarrow g'(y) = 0 \Rightarrow g(y) = A = \text{Constant}.$$

$$F(x,y) = xe^y - ye^x + A. \quad \text{The solution is } F(x,y) = B \Rightarrow$$

$$xe^y - ye^x = C.$$

Let us check via implicit differentiation:

$$e^y - xe^y y' - y'e^x - ye^x = 0 \Rightarrow (e^y - ye^x) + (xe^y - e^x) y' = 0$$

$$\Rightarrow (e^y - ye^x) dx + (xe^y - e^x) dy = 0 \quad \text{by rewriting } y' \text{ as } \frac{dy}{dx}.$$

$$2) -\pi \sin(\pi x) \sinh y \, dx + \cos(\pi x) \cosh(y) \, dy = 0 \quad (2)$$

$$\frac{\partial M}{\partial y} = -\pi \sin(\pi x) \cosh y \quad \text{and} \quad \frac{\partial N}{\partial x} = -\pi \sin(\pi x) \cosh(y)$$

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} \quad , \quad \text{the equation is then exact and the solution is of}$$

$$\text{the form } F(x, y) = C \quad \text{where} \quad \frac{\partial F}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x, y)$$

$$\frac{\partial F}{\partial x} = -\pi \sin(\pi x) \cosh y \Rightarrow F = \int -\pi \sin(\pi x) \cosh(y) \, dx + g(y)$$

$$\Rightarrow F(x, y) = \cos(\pi x) \cosh(y) + g(y)$$

$$\Rightarrow \frac{\partial F}{\partial y} = \cos(\pi x) \sinh(y) + g'(y)$$

$$\text{Comparing with } \frac{\partial F}{\partial y} = N = \cos(\pi x) \cosh(y), \text{ we get } g'(y) = 0$$

$$\Rightarrow g(y) = A = \text{Constant.}$$

$$\text{So } F(x, y) = \cos(\pi x) \cosh(y) + A \quad \text{and the solution is}$$

$$\cos(\pi x) \cosh(y) = C. \quad \text{Let us check by implicit Differentiation}$$

$$-\pi \sin(\pi x) \cosh(y) + \cos(\pi x) \sinh(y) \frac{dy}{dx} = 0 \Leftrightarrow$$

$$-\pi \sin(\pi x) \cosh(y) \, dx + \cos(\pi x) \sinh(y) \, dy = 0.$$

$$3) \, q_x \, dx + 4y \, dy = 0 : \text{Check for exactness:}$$

$$\frac{\partial M}{\partial y} = 0, \quad \frac{\partial N}{\partial x} = 0 : \text{Exact ODE. Look for } F(x, y) \text{ such}$$

$$\text{that } \frac{\partial F}{\partial x} = q_x \quad \text{and} \quad \frac{\partial F}{\partial y} = 4y$$

$$F = \int q_x \, dx + g(y) = \frac{q}{2} x^2 + g(y) \Rightarrow \frac{\partial F}{\partial y} = g'(y)$$

(3)

$\Rightarrow g'(y) = 4y \Rightarrow g(y) = 2y^2 + A$. The solution is then

$$F(x, y) = C \Rightarrow \frac{1}{2}x^2 + 2y^2 = C \text{ or } 9x^2 + 4y^2 = K;$$

Let us check : $18x + 8y \frac{dy}{dx} = 0 \Rightarrow 18x dx + 8y dy = 0 \Rightarrow$

$$9x dx + 4y dy = 0$$

$$4) e^{-2\theta} dr - 2r e^{-2\theta} d\theta = 0 \Leftrightarrow$$

$$\frac{\partial M}{\partial \theta} = -2e^{-2\theta}, \quad \frac{\partial N}{\partial r} = -2e^{-2\theta}; \text{ Exact equation.}$$

We look for $F(r, \theta)$ such that $\frac{\partial F}{\partial r} = e^{-2\theta}$ and $\frac{\partial F}{\partial \theta} = -2r e^{-2\theta}$

$$\frac{\partial F}{\partial r} = e^{-2\theta} \Rightarrow F(r, \theta) = \int e^{-2\theta} dr + g(\theta) = r e^{-2\theta} + g(\theta)$$

$$\Rightarrow \frac{\partial F}{\partial \theta} = -2r e^{-2\theta} + g'(\theta). \text{ Comparing with } \frac{\partial F}{\partial \theta} = -2r e^{-2\theta},$$

We get that $g'(\theta) = 0 \Rightarrow g(\theta) = A = \text{Constant.}$

The solution is then $r e^{-2\theta} = C$ or $r = C e^{2\theta}$

$$\text{check } e^{-2\theta} - 2r e^{-2\theta} \frac{d\theta}{dr} = 0 \Leftrightarrow e^{-2\theta} dr - 2r e^{-2\theta} d\theta = 0$$

$$5) \left[-\frac{y}{x^2} + 2 \cos 2x \right] dx + \left[\frac{1}{x} - 2 \sin(2y) \right] dy = 0$$

$$\frac{\partial M}{\partial y} = -\frac{1}{x^2}, \quad \frac{\partial N}{\partial x} = -\frac{1}{x^2}; \text{ Exact Equation.}$$

Look for $F(x, y)$ such that $\frac{\partial F}{\partial x} = -\frac{y}{x^2} + 2 \cos(2x)$

$$\frac{\partial F}{\partial y} = \frac{1}{x} - 2 \sin(2y)$$

using the second equation, we get

$$F = \int \left[\frac{1}{x} - 2 \sin(2y) \right] dy + g(x) = y/x + \cos(2y) + g(x) \quad (4)$$

$$\Rightarrow \frac{\partial F}{\partial x} = -\frac{y}{x^2} + g'(x) \text{ . Comparing with } \frac{\partial F}{\partial x} = -\frac{y}{x^2} + 2 \cos 2x$$

$$\text{we get that } g'(x) = 2 \cos(2x) \Rightarrow g(x) = \int 2 \cos(2x) dx = \sin(2x) + K$$

$$\text{so } F(x, y) = y/x + \cos(2y) + \sin(2x) + K \text{ and the solution}$$

$$\text{is of the form } \frac{y}{x} + \cos(2y) + \sin(2x) = C$$

Check by implicit differentiation:

$$\frac{y'x - y}{x^2} - 2y' \sin(2y) + 2 \cos(2x) = 0 \Rightarrow$$

$$\frac{y'}{x} - \frac{y}{x^2} - 2y' \sin(2y) + 2 \cos(2x) = 0 \Rightarrow$$

$$[-y/x^2 + 2 \cos(2x)] + \left[\frac{1}{x} - 2 \sin(2y) \right] y' = 0 \Leftrightarrow$$

$$[-y/x^2 + 2 \cos(2x)] dx + \left[\frac{1}{x} - 2 \sin(2y) \right] dy = 0 \text{ by rewriting}$$

$$y' \text{ as } \frac{dy}{dx}.$$

$$b) -2xy \sin(x^2) dx + \cos(x^2) dy = 0$$

$$M_y = -2x \sin(x^2) \text{ , } N_x = -2x \sin(x^2) \text{ : Exact Equation.}$$

$$\text{Look for } F(x, y) \text{ such that } \frac{\partial F}{\partial x} = -2xy \sin(x^2)$$

$$\frac{\partial F}{\partial y} = \cos(x^2)$$

$$\text{using } \frac{\partial F}{\partial y} = \cos(x^2) \text{ , we get } F = \int \cos(x^2) dy + g(x) \Rightarrow$$

(5)

$$F(x,y) = y \cos(x^2) + g(x) \Rightarrow \frac{\partial F}{\partial x} = -2xy \sin(x^2) + g'(x).$$

Comparing with $\frac{\partial F}{\partial x} = -2xy \sin(x^2)$, we get that $g'(x) = 0 \Rightarrow$

$g(x) = A$ and $F(x,y) = y \cos(x^2) + A$. The solution looks like:

$$y \cos(x^2) = C \quad \text{or} \quad y = \frac{C}{\cos(x^2)}.$$

Check $y' \cos(x^2) - 2xy \sin(x^2) = 0 \Leftrightarrow -2xy \sin(x^2) dx + \cos(x^2) dy = 0$

7) $-y dx + x dy = 0 \Rightarrow M_y = -1, N_x = 1$; Not exact.

$$\frac{M_y - N_x}{N} = -\frac{2}{x} = f(x), \quad \text{no } \mu(x) = e^{\int f(x) dx} = e^{\int -\frac{2}{x} dx}$$

$$= e^{-2 \ln|x|} = e^{\ln(\frac{1}{x^2})} = \frac{1}{x^2} \text{ is an integrating factor.}$$

Multiplying our DE with $\frac{1}{x^2}$, we get

$$\underbrace{-\frac{y}{x^2} dx}_{M^*} + \underbrace{\frac{1}{x} dx}_{N^*} = 0 \quad ! \quad M_y^* = -\frac{1}{x^2}, \quad N_x^* = -\frac{1}{x^2} \text{ which}$$

is now exact. Look for $F(x,y)$ such that

$$\frac{\partial F}{\partial x} = -\frac{y}{x^2}, \quad \frac{\partial F}{\partial y} = \frac{1}{x}. \quad \text{The second equation gives}$$

$$F(x,y) = \int \frac{1}{x} dy + g(x) = \frac{y}{x} + g(x) \Rightarrow \frac{\partial F}{\partial x} = -\frac{y}{x^2} + g'(x)$$

Comparing with $\frac{\partial F}{\partial x} = -\frac{y}{x^2}$, we get that $g'(x) = 0 \Rightarrow g(x) = A$.

So the solution looks like $\frac{y}{x} = C$ or

$$y = kx$$

8) $(x^4 + y^2) dx - xy dy = 0$, $y(2) = 1$

(6)

$M_y = 2y$, $N_x = -y$; $M_y \neq N_x$, the equation is not exact.

Since $\frac{M_y - N_x}{N} = \frac{3y}{-xy} = -\frac{3}{x} = f(x)$, an integrating factor

is $e^{\int f(x) dx} = e^{\int -\frac{3}{x} dx} = e^{-3 \ln|x|} = e^{\ln|\frac{1}{x^3}|} = \frac{1}{x^3}$.

Multiplying our DE with $\frac{1}{x^3}$, we get:

$\underbrace{(x + \frac{y^2}{x^3})}_{M^*} dx - \underbrace{\frac{y}{x^2}}_{N^*} dy = 0$:

$M^*_y = \frac{2y}{x^3}$, $N^*_x = \frac{2y}{x^3}$ The new equation is exact.

Look for $F(x, y)$ such that $\frac{\partial F}{\partial x} = x + \frac{y^2}{x^3}$, $\frac{\partial F}{\partial y} = -\frac{y}{x^2}$

$\frac{\partial F}{\partial y} = -\frac{y}{x^2} \Rightarrow F(x, y) = -\int \frac{y}{x^2} dy + g(x) = -\frac{y^2}{2x^2} + g(x)$

$\Rightarrow \frac{\partial F}{\partial x} = \frac{y^2}{x^3} + g'(x)$. Comparing with $\frac{\partial F}{\partial x} = x + \frac{y^2}{x^3}$ gives

$g'(x) = x \Rightarrow g(x) = \frac{1}{2} x^2$. The solution looks like

$-\frac{y^2}{2x^2} + \frac{1}{2} x^2 = C$. Multiplying with 2 gives

$-\frac{y^2}{x^2} + x^2 = K$. We look for the value of K

using the initial Condition $y(2) = 1$:

$-\frac{1}{4} + 4 = K \Rightarrow K = \frac{15}{4} = 3.75$. The particular

solution is $-\frac{y^2}{x^2} + x^2 = 3.75$.

9) $-y \sin(xy) dx - x \sin(xy) dy = 0, \quad y(1) = \pi$ ⑦

$$M_y = -\sin(xy) - xy \cos(xy), \quad N_x = -\sin(xy) - xy \cos(xy):$$

Exact Equation. Look for $F(x, y)$ satisfying:

$$\frac{\partial F}{\partial x} = -y \sin(xy) \quad (1) \quad \text{and} \quad \frac{\partial F}{\partial y} = -x \sin(xy) \quad (2):$$

$$F(x, y) = \int -y \sin(xy) dx = \cos(xy) + g(y)$$

$$\frac{\partial F}{\partial y} = -x \sin(xy) + g'(y). \text{ Comparing with } \frac{\partial F}{\partial y} = -x \sin(xy)$$

gives $g'(y) = 0 \Rightarrow g(y) = A = \text{constant}.$

The general solution looks like $\cos(xy) = C$

Check $-\sin(xy) [y + xy'] = 0 \Leftrightarrow -y \sin(xy) dx - x \sin(xy) dy = 0.$
using the condition $y(1) = \pi$;

$$\cos(\pi) = C \Rightarrow C = -1$$

The particular solution is $\cos(xy) = -1$

10) $[\cos(2x) + 2 \sin(2x)] dx + e^x dy = 0, \quad y(0) = 1$

$$M_y = 0, \quad N_x = e^x : \text{the equation is not exact.}$$

Since $\frac{M_y - N_x}{N} = \frac{0 - e^x}{e^x} = -1 = f(x)$, on integrating

factor is $e^{\int (-1) dx} = e^{-x}$. Multiplying the ODE by e^{-x} gives

$$\underbrace{e^{-x} [\cos(2x) + 2 \sin(2x)] dx}_{M^*} + \underbrace{1 dy}_{N^*} = 0$$

(8)

$M_y^* = 0 = N_x^*$: Now exact. Look for $F(x, y)$ such that

$$\frac{\partial F}{\partial x} = e^{-x} [\cos(2x) + 2 \sin(2x)] \text{ and } \frac{\partial F}{\partial y} = 1. \text{ The second equation}$$

$$\text{gives } F = \int 1 dy + f(x) = y + f(x)$$

$$\frac{\partial F}{\partial x} = f'(x) = e^{-x} \cos(2x) + 2e^{-x} \sin(2x) \Rightarrow f(x) = \int e^{-x} \cos(2x) dx + 2 \int e^{-x} \sin(2x) dx$$

For $\int e^{-x} \cos(2x) dx$, use integration by parts:

$$u = e^{-x}, \quad v' = \cos(2x) \\ u' = -e^{-x}, \quad v = \frac{1}{2} \sin(2x)$$

$$\begin{aligned} \int e^{-x} \cos(2x) dx &= uv - \int u'v dx = \frac{1}{2} e^{-x} \sin(2x) - \int (-e^{-x}) \frac{1}{2} \sin(2x) dx \\ &= \frac{1}{2} e^{-x} \sin(2x) + \frac{1}{2} \int e^{-x} \sin(2x) dx. (*) \end{aligned}$$

For $\int e^{-x} \sin(2x) dx$, use integration by parts again:

$$u = e^{-x}, \quad v' = \sin(2x) \\ u' = -e^{-x}, \quad v = -\frac{1}{2} \cos(2x)$$

$$\begin{aligned} \int e^{-x} \sin(2x) dx &= -\frac{1}{2} e^{-x} \cos(2x) - \int (-e^{-x}) \left(-\frac{1}{2} \cos(2x)\right) dx \\ &= -\frac{1}{2} e^{-x} \cos(2x) - \frac{1}{2} \int e^{-x} \cos(2x) dx \end{aligned}$$

Going back to line (*) we get

$$\int e^{-x} \cos(2x) dx = \frac{1}{2} e^{-x} \sin(2x) + \frac{1}{4} e^{-x} \cos(2x) - \frac{1}{4} \int e^{-x} \cos(2x) dx$$

(9)

$$\Rightarrow \frac{5}{4} \int e^{-x} \cos(2x) dx = \frac{1}{2} e^{-x} \sin(2x) - \frac{1}{4} e^{-x} \cos(2x) \text{ giving}$$

$$\int e^{-x} \cos(2x) = \frac{1}{5} e^{-x} [2 \sin(2x) - \cos(2x)] .$$

$$\text{In a similar way, we get } \int e^{-x} \sin(2x) dx = -\frac{e^{-x}}{5} [\sin(2x) - 2 \cos(2x)]$$

$$\text{Therefore } f(x) = \int e^{-x} \cos(2x) dx + 2 \int e^{-x} \sin(2x) dx = e^{-x} \cos(2x)$$

$$\text{and } F(x, y) = y + f(x) = y + e^{-x} \cos(2x)$$

$$\text{The general solution is } y + e^{-x} \cos(2x) = C$$

$$\text{Check } y' - e^{-x} \cos(2x) - 2e^{-x} \sin(2x) = 0 \Rightarrow$$

$$-e^{-x} (\cos(2x) + 2 \sin(2x)) dx + dy = 0 \text{ or}$$

$$[\cos(2x) + 2 \sin(2x)] dx + e^x dy = 0$$

$$y(0) = 1 \Rightarrow 1 + 1 = C \Rightarrow C = 2$$

$$\text{The particular solution is } y + e^{-x} \cos(2x) = 2$$

$$\text{or } y = 2 - e^{-x} \cos(2x)$$

11) $y \cos(x+y) dx + [3 \sin(x+y) + y \cos(x+y)] dy = 0, \quad y(0) = \pi/2$ (10)

$$M_y = \cos(x+y) - y \sin(x+y); \quad N_x = 3 \cos(x+y) - y \sin(x+y)$$

$M_y \neq N_x$, the equation is NOT exact

Now $\frac{M_y - N_x}{M} = \frac{-2 \cos(x+y)}{y \cos(x+y)} = -\frac{2}{y} = g(y)$, an integrating factor

is given by
$$e^{\int g(y) dy} = e^{-\int \frac{2}{y} dy} = e^{-2 \ln|y|} = y^{-2} = \frac{1}{y^2}$$

Multiplying with y^2 gives:

$$\underbrace{y^3 \cos(x+y)}_{M^*} dx + \underbrace{[3y^2 \sin(x+y) + y^3 \cos(x+y)]}_{N^*} dy = 0$$

$$\left. \begin{aligned} M_y^* &= 3y^2 \cos(x+y) - y^3 \sin(x+y) \\ N_x^* &= 3y^2 \cos(x+y) - y^3 \sin(x+y) \end{aligned} \right\} \begin{aligned} M_y^* &= N_x^*, \text{ The new} \\ D.E. &\text{ is Exact.} \end{aligned}$$

Look for $F(x,y)$ such that $\frac{\partial F}{\partial x} = y^3 \cos(x+y)$ (1)

$$\frac{\partial F}{\partial y} = 3y^2 \sin(x+y) + y^3 \cos(x+y) \quad (2)$$

$$(1) \Rightarrow F(x,y) = \int y^3 \cos(x+y) dx + h(y) = y^3 \sin(x+y) + h(y)$$

$$\Rightarrow \frac{\partial F}{\partial y} = 3y^2 \sin(x+y) + y^3 \cos(x+y) + h'(y). \text{ Comparing with (2)}$$

gives $h'(y) = 0 \Rightarrow h(y) = k$. The general solution is

$$y^3 \sin(x+y) = C$$

$$y(0) = \pi/2 \Rightarrow (\pi/2)^3 \sin(0 + \pi/2) = C \Rightarrow C = \frac{\pi^3}{8}$$

The particular solution is $y^3 \sin(x+y) = \frac{\pi^3}{8}$.

(11)

$$12) [\sin(y) \cos(y) + x \cos^2 y] dx + x dy = 0$$

$$M_y = \cos^2 y - \sin^2 y - 2x \cos y \sin y, N_x = 1 \quad \therefore \text{Not exact.}$$

$$M_y - N_x = \cos^2 y - \sin^2 y - 2x \cos y \sin y - 1 = -2\sin^2 y - 2x \cos y \sin y$$

$$\text{since } \cos^2 y - 1 = -\sin^2 y.$$

$$\frac{M_y - N_x}{N} = \frac{-2 \sin y (\sin y + x \cos y)}{\cos y [\sin y + x \cos y]} = -2 \tan y = g(y).$$

$$\begin{aligned} \text{An integrating factor is } e^{-\int g(y) dy} &= e^{-\int -2 \tan y dy} = e^{2 \int \tan y dy} \\ &= e^{-2 \ln |\cos y|} = e^{\ln \left(\frac{1}{\cos^2 y} \right)} = \frac{1}{\cos^2 y} \quad \text{since } \int \tan y dy = -\ln |\cos x|. \end{aligned}$$

Multiplying by $\frac{1}{\cos^2 y}$ gives:

$$\underbrace{(\tan y + x)}_{M^*} dx + \underbrace{\frac{x}{\cos^2 y}}_{N^*} dy = 0$$

$$M_y^* = \frac{1}{\cos^2 y}, N_x^* = \frac{1}{\cos^2 y}. \quad \text{The new equation is Exact.}$$

$$\text{Find } F(x, y) \text{ such that } \frac{\partial F}{\partial x} = \tan y + x \text{ and } \frac{\partial F}{\partial y} = \frac{x}{\cos^2 y}$$

$$F = \int (\tan y + x) dx + g(y) \Rightarrow F(x, y) = x \tan y + \frac{1}{2} x^2 + g(y)$$

$$\Rightarrow \frac{\partial F}{\partial y} = \frac{x}{\cos^2 y} + g'(y) \text{ which gives } g'(y) = 0 \text{ and } g(y) = A$$

The general solution is

(12)

$$x \tan y + \frac{1}{2} x^2 = C$$

$$13) (Ax + By) dx + (Cx + Dy) dy = 0$$

$M_y = B$, $N_x = C$. The equation is Exact if $M_y = N_x \Leftrightarrow B = C$.

In this case, we look for $F(x, y)$; $\frac{\partial F}{\partial x} = Ax + By$, $\frac{\partial F}{\partial y} = Bx + Dy$

$$F = \frac{A}{2} x^2 + Bxy + g(y) \Rightarrow \frac{\partial F}{\partial y} = Bx + g'(y) = Bx + Dy \Rightarrow$$

$g'(y) = Dy \Rightarrow g(y) = \frac{1}{2} Dy^2$. The general solution is

$$\boxed{\frac{1}{2} Ax^2 + Bxy + \frac{1}{2} Dy^2 = C}$$