

Midterm2

Langchen Liu

2024-04-02

Load Data and pre-processing

```
load('data/labeled_points.Rdata')

labeled = labeled %>%
  select(ID, landcover)

d = labeled_train %>%
  left_join(labeled, by = 'ID')

d = d %>%
  mutate(veg = ifelse(landcover %in% c('natforest', 'orchard', 'cropland'),
    1, 0),
    NDVI100 = NDVI * 100)
head(d,2)

## # A tibble: 2 x 22
##   B1    B2    B3    B4    B5 B6_VCID_1 B6_VCID_2    B7    B8    lat    lon
##   <dbl> <dbl> <dbl> <dbl> <dbl>   <dbl>     <dbl> <dbl> <dbl> <dbl> <dbl>
## 1    66    60    73    68    93     146     176    76    69  11.2  3.41
## 2    64    59    69    72    89     147     178    72    69  11.2  3.41
## # i 11 more variables: year <int>, month <int>, day <int>, date <date>,
## #   ID <int>, NDVI <dbl>, NDBI <dbl>, EVI <dbl>, landcover <chr>, veg <dbl>,
## #   NDVI100 <dbl>

dm = d %>%
  group_by(ID) %>%
  summarise(
    mean.NDVI100 = mean(NDVI100, na.rm = T),
    landcover = unique(landcover),
    veg = unique(veg)) %>%
  as.data.frame()
head(dm,2)

##   ID mean.NDVI100 landcover veg
## 1 2043     15.06720  orchard   1
## 2 2069     13.42962  orchard   1
```

Fit a logistic model

```
m1 = glm(veg ~ mean.NDVI100,
  data = dm,
```

```

    family = binomial(link = logit))
summary(m1)

##
## Call:
## glm(formula = veg ~ mean.NDVI100, family = binomial(link = logit),
##      data = dm)
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  -0.1318      0.2599  -0.507   0.612
## mean.NDVI100  0.7953      0.1029   7.729 1.09e-14 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for binomial family taken to be 1)
##
##      Null deviance: 449.87  on 399  degrees of freedom
## Residual deviance: 105.95  on 398  degrees of freedom
## AIC: 109.95
##
## Number of Fisher Scoring iterations: 8

```

a. Data exploration

Find another predictor involving NDVI (other than mean(NDVI100)) that might be useful in predicting vegetation.

I might begin with variance(NDVI100)

```


dm = d %>%
  group_by(ID) %>%
  summarise(
    mean.NDVI100 = mean(NDVI100, na.rm = T),
    var.NDVI100 = var(NDVI100, na.rm = T),
    landcover = unique(landcover),
    veg = unique(veg)) %>%
  as.data.frame()

### plot the variance and make a smooth curve

ggplot(dm, aes(x = var.NDVI100, y = veg)) +
  geom_point() +
  geom_smooth(method = 'glm',
              method.args = list(family = 'binomial')) +
  labs(title = 'Variance of NDVI100 vs. Vegetation')

## `geom_smooth()` using formula = 'y ~ x'

```



From the plot, we can see that the variance of NDVI100 is a good predictor of vegetation. The reason is $\text{veg} = 0$ is most likely to have a smaller variance in NDVI100, while the $\text{veg} = 1$ points have a significantly greater variance in NDVI100.

b. Fit a logistic model

```
m2 = glm(veg ~ var.NDVI100 + mean.NDVI100,
  data = dm,
  family = binomial(link = logit))

summary(m2)
```

```
##
## Call:
## glm(formula = veg ~ var.NDVI100 + mean.NDVI100, family = binomial(link = logit),
##      data = dm)
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept) -4.453135   1.140102  -3.906 9.39e-05 ***
## var.NDVI100  0.033005   0.008608   3.834 0.000126 ***
## mean.NDVI100 0.404484   0.132986   3.042 0.002354 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for binomial family taken to be 1)
##
##      Null deviance: 449.868  on 399  degrees of freedom
```

```
## Residual deviance: 84.101 on 397 degrees of freedom
## AIC: 90.101
##
## Number of Fisher Scoring iterations: 8
```

We can see that the new predictor (var.NDVI100) significantly improves the previous model.

First, the intercept and the coefficients of the new model has a good significant level, while the intercept of the previous model is not as significant.

Second, both model share the same null deviance but the new model has a much smaller residual deviance, this means that the new model has a greater difference in residual and null deviance, which means the new model explains the statistical relationship of the predictors and the response variables better.

Third, the AIC of the new model is smaller than the previous model, which means the new model has a better fit.

As a result, the new model is better than the previous model. I will recommend adding this new predictor (var.NDVI100) to the model.