

Theorem 1 (Residue Theorem). Let f be analytic in the region G except for the isolated singularities a_1, a_2, \dots, a_m . If γ is a closed rectifiable curve in G which does not pass through any of the points a_k and if $\gamma \approx 0$ in G then

$$\frac{1}{2\pi i} \int_{\gamma} f = \sum_{k=1}^m n(\gamma; a_k) \operatorname{Res}(f; a_k).$$

Theorem 2 (Maximum Modulus). Let G be a bounded open set in \mathbb{C} and suppose that f is a continuous function on G^- which is analytic in G . Then

$$\max\{|f(z)| : z \in G^-\} = \max\{|f(z)| : z \in \partial G\}.$$

A
Λ
Δ
∇
B
C
D
Σ
E
F
Γ
G
H
I
J
K
L
M
N
O
Θ
Ω
ϐ
P
Φ
Π
Ξ
Q
R
S
T
U
V
W
X
Y
Υ
Ψ
Z
1
2
3
4
5
6
7
8
9
0

a
α
b
β
c
δ
d
δ
e
ε
f
ζ
ξ
g
γ
h
ħ
h
i
i
j
j
k
κ
κ
l
ℓ
λ
m
n
η
θ
θ
o
σ
σ
φ
φ
Ϙ
p
r
r
r
q
r
s
t
τ
π
ι
μ
ν
υ
w
ω
ω
x
χ
y
ψ
z
∞
∝
∅
∅
d
d̃
ə