Notation

This section provides a concise reference describing the notation used throughout this book. If you are unfamiliar with any of the corresponding mathematical concepts, we describe most of these ideas in chapters 2–4.

Numbers and Arrays

- a A scalar (integer or real)
- a A vector
- A A matrix
- A A tensor
- I_n Identity matrix with n rows and n columns
- I Identity matrix with dimensionality implied by context
- $e^{(i)}$ Standard basis vector [0..., 0, 1, 0, ..., 0] with a 1 at position i
- diag(a) A square, diagonal matrix with diagonal entries given by a
 - a A scalar random variable
 - a A vector-valued random variable
 - A A matrix-valued random variable

χi

CONTENTS

R	The set of real numbers
{0,	1) The set containing 0 and 1
{0, 1,	,n) The set of all integers between 0 and n
[a,	b] The real interval including a and b
(a,	b] The real interval excluding a but including b
A\E	Set subtraction, i.e., the set containing the elements of A that are not in B
G	A graph
Ра _G	(x_i) The parents of x_i in G
	Indexing
aį	Element i of vector a, with indexing starting at 1
a_{-i}	All elements of vector a except for element i
$A_{i,j}$	Element i, j of matrix A
$A_{i,:}$	Row i of matrix A
A :,i	Column i of matrix A
$\boldsymbol{A}_{i,j,k}$	Element (i, j, k) of a 3-D tensor A
$A_{:,:,i}$	2-D slice of a 3-D tensor
aį	Element i of the random vector a
	Linear Algebra Operations
Α□	Transpose of matrix A
A +	Moore-Penrose pseudoinverse of A
А□В	Element-wise (Hadamard) product of A and B
det(A)	Determinant of A
	хіі
	• • • • • • • • • • • • • • • • • • •
ITENITO	
NTENTS	

CON

d), (Calculus
$\frac{dy}{dx}$	Derivative of y with respect to x
$\frac{\partial y}{\partial x}$	Partial derivative of y with respect to x
∇×У	Gradient of y with respect to x
Δхλ	Matrix derivatives of y with respect to X
₩VV	Tensor containing derivatives of v with respect to

Χ ∂f Jacobian matrix $J \in R^{m \times n}$ of $f : R^n \to R^m$ дх $\nabla_x^2 f(x)$ or H(f)(x)The Hessian matrix of f at input point x Definite integral over the entire domain of x \Box f(x)dx \Box f(x)dx Definite integral with respect to x over the set S Probability and Information Theory a⊥b The random variables a and b are independent a⊥blc They are conditionally independent given c P(a) A probability distribution over a discrete variable p(a) A probability distribution over a continuous variable, or over a variable whose type has not been specified a ~ P Random variable a has distribution P $\mathsf{E}_{\mathsf{x},\mathsf{P}}[\mathsf{f}(\mathsf{x})]$ or $\mathsf{E}\mathsf{f}(\mathsf{x})$ Expectation of f(x) with respect to P(x)Var(f(x))Variance of f(x) under P(x)Cov(f(x), g(x))Covariance of f(x) and g(x) under P(x)H(x)Shannon entropy of the random variable x D_{KL}(PQ) Kullback-Leibler divergence of P and Q Gaussian distribution over x with mean μ and $N(x; \mu, \Sigma)$ covariance S

xiii

CONTENTS

Functions

$f:A\to B$	The function f with domain A and range B
$f \circ g$ $f(x; \theta)$	Composition of the functions f and g A function of x parametrized by θ . (Sometimes we write f (x) and omit the argument θ to lighten notation)
log x	Natural logarithm of x
σ(x)	Logistic sigmoid, $\frac{1}{1 + \exp(-x)}$
ζ(x)	Softplus, $log(1 + exp(x))$
llxll p	L ^p norm of x
llxll	1 2 norm of x

 x^+ Positive part of x, i.e., max(0, x)

1 condition is 1 if the condition is true, 0 otherwise

Sometimes we use a function whose argument is a scalar but apply it to a vector, matrix, or tensor: f(x), f(X), or f(X). This denotes the application of f to the array element-wise. For example, if $C = \sigma(X)$, then $C_{i,j,k} = \sigma(X_{i,j,k})$ for all valid values of i, j and k.

Datasets and Distributions

	Datasets and Distributions
p _{data}	The data generating distribution
ρ̂ _{data}	The empirical distribution defined by the training set
Χ	A set of training examples
x ⁽ⁱ⁾	The i-th example (input) from a dataset
y ⁽ⁱ⁾ or y ⁽ⁱ⁾	The target associated with $\mathbf{x}^{(i)}$ for supervised learning
X	The $m \times n$ matrix with input example $x^{(i)}$ in row $X_{i,:}$