

Section 1. Notes on koala spacing

Fencing, predator control and revegetation remain the main koala recovery actions. The koala (*Phascolarctos cinereus*) shelters in eucalypt canopy across coastal and inland woodland. Field officers recorded 330 koala observations during the reporting period. The koala (*Phascolarctos cinereus*) shelters in eucalypt canopy across coastal and inland woodland. The department will review koala monitoring protocols by 30 June 2027. Field officers recorded 69 koala observations during the reporting period.

Field officers recorded 18 koala observations during the reporting period. Survey effort for the koala increased by 27 per cent on the previous year. Survey effort for the koala increased by 27 per cent on the previous year.

Run-in label in bold

Body text of the same size directly beneath it

A larger line closing the page

Section 2. Notes on quokka spacing

The quokka (*Setonix brachyurus*) shelters in dense scrub across Rottnest Island and the south-west. Survey effort for the quokka increased by 11 per cent on the previous year. The quokka (*Setonix brachyurus*) shelters in dense scrub across Rottnest Island and the south-west.

Field officers recorded 119 quokka observations during the reporting period. The department will review quokka monitoring protocols by 30 June 2027. Fencing, predator control and revegetation remain the main quokka recovery actions.

The quokka (*Setonix brachyurus*) shelters in dense scrub across Rottnest Island and the south-west. Funding of \$27.2 million supports quokka habitat restoration. Community groups reported that quokka numbers were stable in WA.

Run-in label in bold

Body text of the same size directly beneath it

A larger line closing the page

Section 3. Notes on platypus spacing

Survey effort for the platypus increased by 22 per cent on the previous year. The department will review platypus monitoring protocols by 30 June 2027. The department will review platypus monitoring protocols by 30 June 2027.

Survey effort for the platypus increased by 25 per cent on the previous year. Funding of \$77.5 million supports platypus habitat restoration. Community groups reported that platypus numbers were stable in VIC.

The platypus (*Ornithorhynchus anatinus*) shelters in creek banks across freshwater streams of the east. Funding of \$41.2 million supports platypus habitat restoration. The platypus (*Ornithorhynchus anatinus*) shelters in creek banks across freshwater streams of the east.

Run-in label in bold

Body text of the same size directly beneath it

A larger line closing the page

Section 4. Notes on echidna spacing

Funding of \$46.6 million supports echidna habitat restoration. Community groups reported that echidna numbers were stable in NSW. The department will review echidna monitoring protocols by 30 June 2028.

The department will review echidna monitoring protocols by 30 June 2028. Field officers recorded 442 echidna observations during the reporting period. The echidna (*Tachyglossus aculeatus*) shelters in log hollows across almost every Australian habitat.

The department will review echidna monitoring protocols by 30 June 2028. Funding of \$32.7 million supports echidna habitat restoration. Survey effort for the echidna increased by 25 per cent on the previous year.

Run-in label in bold

Body text of the same size directly beneath it

A larger line closing the page

Section 5. Notes on bilby spacing

Community groups reported that bilby numbers were stable in TAS. The bilby (*Macrotis lagotis*) shelters in spiral burrows across arid and semi-arid grassland. Field officers recorded 248 bilby observations during the reporting period.

Funding of \$66.9 million supports bilby habitat restoration. Fencing, predator control and revegetation remain the main bilby recovery actions. The department will review bilby monitoring protocols by 30 June 2026.

Fencing, predator control and revegetation remain the main bilby recovery actions. Field officers recorded 59 bilby observations during the reporting period. Fencing, predator control and revegetation remain the main bilby recovery actions.

Run-in label in bold

Body text of the same size directly beneath it

A larger line closing the page

Section 6. Notes on numbat spacing

The numbat (*Myrmecobius fasciatus*) shelters in fallen logs across wandoo woodland. The department will review numbat monitoring protocols by 30 June 2028. The department will review numbat monitoring protocols by 30 June 2028.

Fencing, predator control and revegetation remain the main numbat recovery actions.

Community groups reported that numbat numbers were stable in TAS. Field officers recorded 219 numbat observations during the reporting period.

Field officers recorded 40 numbat observations during the reporting period. Fencing, predator control and revegetation remain the main numbat recovery actions. The department will review numbat monitoring protocols by 30 June 2027.

Run-in label in bold

Body text of the same size directly beneath it

A larger line closing the page