

GE23131-Programming Using C-2024

Quiz navigation

[Show one page at a time](#)[Finish review](#)

Status	Finished
Started	Monday, 23 December 2024, 5:33 PM
Completed	Tuesday, 26 November 2024, 9:43 AM
Duration	27 days 7 hours

Question 1

Correct

Marked out of 3.00

☐ Flag question

A set of N numbers (separated by one space) is passed as input to the program. The program must identify the count of numbers where the number is odd number.

Input Format:

The first line will contain the N numbers separated by one space.

Boundary Conditions:

$$3 \leq N \leq 50$$

The value of the numbers can be from -99999999 to 99999999

Output Format:

The count of numbers where the numbers are odd numbers.

Example Input / Output 1:

Input:

5 10 15 20 25 30 35 40 45 50

Output:

5

Explanation:

Answer: (penalty regime: 0 %)

	Input	Expected	Got	
	5 10 15 20 25 30 35 40 45 50	5	5	

Passed all tests!

Question **2**

Correct

Marked out of 5.00

☐ Flag question

Given a number N, return true if and only if it is a *confusing number*, which satisfies the following condition:

We can rotate digits by 180 degrees to form new digits. When 0, 1, 6, 8, 9 are rotated 180 degrees, they become 0, 1, 9, 8, 6 respectively. When 2, 3, 4, 5 and 7 are rotated 180 degrees, they become invalid. A *confusing number* is a number that when rotated 180 degrees becomes a **different** number with each digit valid.

Example 1:

6 -> 9

Input: 6

Output: true

Explanation:

We get 9 after rotating 6, 9 is a valid number and $9 \neq 6$.

Example 2:

89 -> 68

Input: 89

We get 68 after rotating 89, 86 is a valid number and $86 \neq 89$.

Example 3:

11 -> 11

Input: 11

Output: false

Explanation:

We get 11 after rotating 11, 11 is a valid number but the value remains the same, thus 11 is not a confusing number.

Note:

1. $0 \leq N \leq 10^9$
2. After the rotation we can ignore leading zeros, for example if after rotation we have 0008 then this number is considered as just 8.

Answer: (penalty regime: 0 %)

	Input	Expected	Got	
	6	true	true	
	89	true	true	
	25	false	false	

Passed all tests!

Question **3**
Correct

Flag
question

has. For example, food item with value 1 has 1 macronutrient, food item with value 2 has 2 macronutrients, and incrementing in this fashion.

The nutritionist has to recommend the best combination to patients, i.e. maximum total of macronutrients. However, the nutritionist must avoid prescribing a particular sum of macronutrients (an 'unhealthy' number), and this sum is known. The nutritionist chooses food items in the increasing order of their value. Compute the highest total of macronutrients that can be prescribed to a patient, without the sum matching the given 'unhealthy' number.

Here's an illustration:

Given 4 food items (hence value: 1,2,3 and 4), and the unhealthy sum being 6 macronutrients, on choosing items 1, 2, 3 -> the sum is 6, which matches the 'unhealthy' sum. Hence, one of the three needs to be skipped. Thus, the best combination is from among:

- $2 + 3 + 4 = 9$
- $1 + 3 + 4 = 8$
- $1 + 2 + 4 = 7$

Since $2 + 3 + 4 = 9$, allows for maximum number of macronutrients, 9 is the right answer.

Complete the code in the editor below. It must return an integer that represents the maximum total of macronutrients, modulo 1000000007 ($10^9 + 7$).

It has the following:

n : an integer that denotes the number of food items

k : an integer that denotes the unhealthy number

Constraints

- $1 \leq n \leq 2 \times 10^9$
- $1 \leq k \leq 4 \times 10^{15}$

Input Format For Custom Testing

The first line contains an integer, n , that denotes the number of food items.

The second line contains an integer, k , that denotes the unhealthy number.

Sample Input 0

2
2

Sample Output 0

Explanation 0

The following sequence of $n = 2$ food items:

1. Item 1 has 1 macronutrients.
2. $1 + 2 = 3$; observe that this is the max total, and having avoided having exactly $k = 2$ macronutrients.

Sample Input 1

2
1

Sample Output 1

2

Explanation 1

1. Cannot use item 1 because $k = 1$ and $sum \equiv k$ has to be avoided at any time.
2. Hence, max total is achieved by $sum = 0 + 2 = 2$.

Sample Case 2

Sample Input For Custom Testing**Sample Input 2**

3
3

Sample Output 2

5

Explanation 2

$2 + 3 = 5$, is the best case for maximum nutrients.

REC-CIS

Answer. (penalty regime: 0 %)

	Input	Expected	Got	
	2 2	3	3	
	2 1	2	2	
	3 3	5	5	

Passed all tests!

[Save the state of the flags](#)[Finish review](#)