

Client-ready insight report

WEEK 4 ----- ANA 4

PREPARED BY: DEEPAK BARANWAL (Data Analyst Intern)

PROJECT NAME: AI MENTAL HEALTH UK

DATE: 28 – 30 MAY, 2026

Executive Summary:

Background:

Mental health remains one of the most significant public health challenges in the United Kingdom. Increasing demand for mental health services, crisis intervention support, and community-based care has placed substantial pressure on healthcare systems. Understanding service utilization patterns, crisis trends, and regional variations is essential for effective resource allocation and evidence-based decision making.

This report presents key insights derived from the analysis of publicly available NHS Mental Health Statistics, Samaritans suicide statistics, Office for National Statistics (ONS) datasets, and Republic of Ireland (ROI) suicide data. The purpose of the analysis is to provide an overview of current mental health trends and highlight areas requiring continued monitoring and intervention.

Objective:

The primary objective of this report is to evaluate crisis-related mental health indicators and identify significant trends that may support healthcare planning and policy development. The analysis focuses on:

- NHS Mental Health Key Performance Indicators (KPIs)
- Mental health service utilization patterns
- Crisis-related support service activity
- Long-term suicide trends in the UK
- Regional variation in suicide rates
- Comparative analysis between the UK and the Republic of Ireland

Scope of Analysis:

The analysis was conducted using NHS Mental Health Bulletin datasets, ONS suicide statistics, Samaritans public statistics, and Republic of Ireland suicide data. Data processing and visualization were performed using Python and GoogleColab.

The study examined four major areas:

1. NHS Mental Health KPIs, including beds, contacts, crisis calls, admissions, and referral-to-treatment indicators.
2. Executive dashboard metrics designed to support performance monitoring and decision-making.
3. Long-term crisis and suicide trends across England and Wales.
4. Regional and comparative analyses to identify geographical differences and broader mental health patterns.

Overall Summary:

The findings indicate continued demand for mental health and crisis intervention services across the UK. Community-based crisis services, particularly Crisis Resolution Teams and 24/7 Crisis Response Lines, recorded substantial levels of activity. Long-term suicide trend analysis highlights persistent mental health challenges, while regional analysis reveals significant geographical variation in outcomes.

The evidence suggests that targeted regional interventions, continued investment in crisis response services, and enhanced monitoring of mental health indicators remain important priorities for healthcare providers and policymakers. The insights presented in this report can support strategic planning and contribute to the ongoing improvement of mental health service delivery across the NHS.

Key Findings:

1. NHS Mental Health KPI Findings:

Five key performance indicators (KPIs) were extracted from NHS Mental Health Services Data Set (MHSDS) resources, including beds, contacts, crisis calls, admissions, and referral-to-treatment (RTT) measures. These indicators provide a high-level overview of service demand and healthcare system activity.

The analysis identified consistently high levels of mental health service utilization across multiple service categories. Contact-related metrics represented one of the largest areas of activity, highlighting the increasing demand for community-based mental health support. Crisis-related service indicators also demonstrated substantial utilization, emphasizing the importance of maintaining accessible crisis intervention pathways.


Admissions and bed utilization indicators reflected the ongoing requirement for inpatient mental health services, while RTT indicators highlighted the importance of monitoring timely access to care. Together, these KPIs provide a comprehensive overview of NHS mental health service performance and operational demand.

2. Dashboard Insights and Service Utilization Trends:

The executive dashboard developed during the analysis consolidated key mental health indicators into a single monitoring framework. The dashboard was designed to support performance monitoring, service planning, and strategic decision-making.

The dashboard findings indicated that mental health service activity remains consistently high across multiple service areas. Contact-based services accounted for a significant proportion of recorded activity, demonstrating continued demand for support and intervention. Crisis-related indicators also showed substantial utilization levels, reflecting the importance of rapid access mental health services.


The dashboard enables healthcare stakeholders to monitor service performance, identify emerging trends, and evaluate changes in demand over time. Such monitoring tools are valuable for supporting evidence-based decision-making and resource allocation within mental health services.



3. UK Crisis Trend Analysis:


Analysis of suicide occurrence data for England and Wales revealed long-term changes in crisis-related outcomes between 2001 and 2024. Suicide occurrences remained relatively stable during the early years of the study period but increased noticeably after 2017. The highest occurrence levels were recorded between 2018 and 2022, indicating growing pressure on mental health systems and crisis intervention services.

Although a decline was observed in the most recent years, the latest figures should be interpreted cautiously because recent data may be provisional and subject to future revision. Overall, the trend highlights the continued importance of suicide prevention initiatives and accessible mental health support services.



4. Regional Variation Analysis:

Regional analysis identified substantial variation in suicide rates across England. The North East recorded the highest suicide rate at 15.1 deaths per 100,000 population, while London recorded the lowest rate at 8.3 deaths per 100,000 population. Northern regions generally exhibited higher suicide rates than southern regions. These findings suggest that geographical, socioeconomic, demographic, and healthcare accessibility factors may contribute to differences in mental health outcomes across regions. The results highlight the importance of region-specific interventions and targeted public health strategies.




5. Republic of Ireland Comparative Analysis

A comparative assessment of Republic of Ireland (ROI) suicide statistics provided additional context for understanding mental health trends beyond the UK. The ROI dataset showed fluctuations in suicide deaths over time, with elevated levels observed during the late 2000s and early 2010s. While the overall trend shares similarities with the UK pattern, the ROI data demonstrated greater year-to-year variability. The comparison reinforces the broader challenges associated with suicide prevention and highlights the importance of sustained investment in mental health services across neighbouring regions.

6. Crisis Service Contact Volume Analysis:

Analysis of NHS crisis service contact data revealed significant differences in utilization across service types. Crisis Resolution Team/Home Treatment Services recorded the highest activity level, with approximately 4.59 million contacts. The 24/7 Crisis Response Line recorded more than 427,000 contacts, indicating strong demand for immediate crisis support.



Crisis House Services and Walk-in Crisis Assessment Units reported lower utilization levels compared with larger community-based crisis services. These findings demonstrate that community-focused crisis intervention services remain a critical component of mental health support infrastructure and play a significant role in reducing pressure on inpatient services.

Data Sources and Limitations:

Data Sources:

This analysis was conducted using multiple publicly available datasets obtained from trusted healthcare and statistical organizations. The use of multiple sources enabled a broader understanding of mental health service utilization, crisis trends, and suicide-related outcomes.

NHS Mental Health Bulletin 2024–25

The NHS Mental Health Bulletin served as the primary source for mental health service activity indicators. Data extracted from the bulletin included care contacts, crisis service utilization, admissions, bed occupancy measures, and other operational performance indicators. These datasets provided valuable insights into the demand for NHS mental health services and crisis intervention pathways.

NHS Mental Health Services Data Set (MHSDS)

MHSDS resources were used to identify and evaluate key performance indicators including beds, contacts, crisis calls, admissions, and referral-to-treatment measures. These indicators formed the basis of KPI extraction and dashboard development activities.

Office for National Statistics (ONS)

ONS suicide statistics were used to analyze long-term suicide occurrence trends and regional variations across England and Wales. The datasets provided reliable and nationally recognized indicators for understanding changes in suicide-related outcomes over time.

Central Statistics Office (CSO) Ireland

Republic of Ireland suicide statistics obtained from CSO Ireland were used to perform comparative analysis between the UK and ROI. This additional perspective strengthened the overall assessment by identifying similarities and differences in long-term suicide trends across neighbouring regions.

Limitations:

While the analysis provides valuable insights into mental health service utilization and crisis trends, several limitations should be considered when interpreting the findings.

1. Annual Reporting Structure:

Many of the publicly available datasets were reported at an annual level rather than monthly or quarterly intervals. As a result, detailed seasonal pattern analysis could not be performed. The absence of higher-frequency observations limits the ability to identify short-term fluctuations in service demand.

2. Limited Patient-Level Information:

The datasets used in this study were aggregated public datasets and did not contain patient-level information. Consequently, detailed demographic, clinical, and individual outcome analyses could not be conducted.

3. Provisional and Recently Reported Data

Some of the most recent statistics may be provisional and subject to future revisions. Therefore, recent trends should be interpreted with caution until final validated figures become available.

4. Differences Across Data Sources

The analysis combined information from multiple organizations including NHS, ONS, Samaritans, and CSO Ireland. Differences in reporting methodologies, definitions, and publication schedules may influence direct comparisons between datasets.

5. Public Data Availability Constraints

The analysis was limited to publicly accessible datasets. Additional insights may be obtainable through more detailed internal NHS operational data that are not publicly available.

6. Data Quality Considerations

Despite these limitations, all datasets were obtained from recognized public institutions and underwent basic validation and preprocessing procedures before analysis. The combined use of multiple authoritative sources improves the reliability of the findings and provides a robust foundation for strategic decision-making and service planning.

Recommendations and Conclusion:

Recommendations:

Based on the findings obtained from KPI analysis, dashboard monitoring, crisis trend assessment, and service utilization analysis, several recommendations are proposed to support future mental health service planning and policy development.

Recommendation 1: Strengthen Crisis Resolution Team Capacity

The analysis identified Crisis Resolution Team/Home Treatment Services as the most heavily utilized crisis service, recording approximately 4.59 million contacts. This demonstrates the critical role these services play in supporting individuals experiencing acute mental health difficulties.

Increasing workforce capacity, expanding service coverage, and improving resource allocation within Crisis Resolution Teams may help maintain service quality while addressing growing demand.

Recommendation 2: Expand 24/7 Crisis Response Services

The 24/7 Crisis Response Line recorded more than 427,000 contacts, highlighting substantial demand for immediate crisis support. Ensuring continuous accessibility to crisis response services remains essential for individuals requiring urgent mental health assistance.

Additional investment in crisis response infrastructure, staffing, and digital support channels may improve service accessibility and response effectiveness.

Recommendation 3: Prioritize High-Risk Regions

Regional analysis identified significant geographical variation in suicide rates across England. The North East reported the highest suicide rate, while London reported the lowest.

Healthcare providers and policymakers should consider implementing targeted interventions in high-risk regions. Region-specific prevention strategies may improve outcomes by addressing local social, economic, and healthcare-related challenges.

Recommendation 4: Improve Mental Health Data Monitoring

The analysis demonstrated the value of dashboards and KPI monitoring in identifying service demand and performance trends. Expanding real-time monitoring capabilities may support faster decision-making and improve operational planning.

The continued development of integrated reporting systems and interactive dashboards can enhance visibility of key mental health indicators across healthcare organizations.

Recommendation 5: Enhance Early Intervention and Prevention Programs

Long-term suicide trend analysis indicates that mental health challenges remain a significant public health concern. Early intervention programs, community outreach initiatives, and preventative mental health services can play an important role in reducing crisis events and improving long-term outcomes.

Investment in preventative strategies may help reduce demand on crisis services while improving overall population wellbeing.

Conclusion:

This report examined mental health service utilization, crisis-related activity, suicide trends, and regional variations using publicly available NHS, ONS, Samaritans, and CSO Ireland datasets.

The findings indicate that mental health services continue to experience substantial demand, particularly within crisis intervention pathways. Community-based services such as Crisis Resolution Teams and 24/7 Crisis Response Lines play a vital role in supporting individuals experiencing mental health crises.

Long-term trend analysis highlighted the ongoing importance of suicide prevention efforts, while regional analysis revealed notable geographical disparities that may require targeted intervention. Comparative analysis between the United Kingdom and the Republic of Ireland further demonstrated that mental health challenges remain a shared concern across neighbouring regions.

Overall, the evidence supports continued investment in crisis response services, regional mental health initiatives, data-driven decision making, and preventative care strategies. Strengthening these areas will contribute to improved service delivery, enhanced patient outcomes, and more effective mental health support systems in the future.

Report Prepared For

NHS Mental Health Analytics – Client Insight Report

Prepared By

Deepak Baranwal

Data Analytics Internship – Week 4 ANA-4